



UNDERSTANDING TUNGSTEN ACCUMULATION DURING ICRH OPERATION ON WEST

Patrick Maget, Pierre Manas, Rémi Dumont, Clemente Angioni, Jean François Artaud, Clarisse Bourdelle, Laurent Colas, Pascal Devynck, Fajardo D., Nicolas Fedorczak, et al.

► To cite this version:

Patrick Maget, Pierre Manas, Rémi Dumont, Clemente Angioni, Jean François Artaud, et al.. UNDERSTANDING TUNGSTEN ACCUMULATION DURING ICRH OPERATION ON WEST. 48th EPS conference on plasma physics, Jun 2022, On line, France. cea-03736601

HAL Id: cea-03736601

<https://cea.hal.science/cea-03736601>

Submitted on 22 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNDERSTANDING TUNGSTEN ACCUMULATION DURING ICRH OPERATION ON WEST



P. Maget, P. Manas, R. Dumont, C. Angioni¹, J-F Artaud,
C. Bourdelle, L. Colas, P. Devynck, D. Fajardo¹, N. Fedorczak,
M. Goniche, J. Hillairet, Ph. Huynh, J. Morales, V. Ostuni, D.
Vézinet and the WEST team[#]

CEA IRFM F-13108 Saint Paul-lez-Durance, France.

¹ Max-Planck-Institut für Plasmaphysik, D-85748 Garching, Germany.

see <http://west.cea.fr/WESTteam>

48th EPS 2022 - Online
June 30th 2022
04.114



Impurity peaking is a concern in tokamaks with metallic walls → core radiation

- ▶ Expected in presence of strong poloidal asymmetries [Angioni'14], mainly driven by :
 - Toroidal rotation (centrifugal force)
 - ICRH heating (electrostatic force)
- ▶ Operation on WEST : no external torque, no core particle source → ITER-relevant conditions
 - Several observations of radiative collapses
 - During LHCD [Ostuni, O4.113], LHCD+ICRH [Goniche'21], and *pure ICRH operation [this talk]*

ICRH impact tungsten profiles through 3 main channels

- ▶ Direct : electrostatic force
- ▶ Non-linear : effect of tungsten peaking on the electrostatic force (i.e. on the direct channel)
- ▶ Indirect : ICRH induced rotation

Modeling tungsten behavior in ICRH operation mobilizes a wide range of physics issues

- ▶ Finite Orbit Width (FOW) effects
- ▶ Fast Hydrogen Temperature Screening (HTS) effect [Casson'15,Casson'20]
- ▶ Ripple losses
- ▶ ... with consequences on the electrostatic force and electron heating

Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels

Tungsten peaking with ICRH and plasma rotation

Meeting the collapse condition

Summary and conclusion

Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels

Tungsten peaking with ICRH and plasma rotation

Meeting the collapse condition


Summary and conclusion

Observation of tungsten accumulation & radiative collapse (L-mode)

- ▶ Core electron temperature falls above some ICRH power
- ▶ Bolometry inversion shows fast increase of core radiation
- ▶ Radiation peaking does not precede the collapse
 - Reversed temperature screening generates the large peaking

Modelling tools

- ▶ Equilibrium from **METIS** interpretative run [Artaud'18]
- ▶ ICRH deposition from **EVE/AQL** computations [Dumont'09 & '12]
- ▶ Turbulent transport computed with **QuaLiKiz** [Bourdelle'07]
 - Indicates nearly zero turbulent level inside $r/a=0.3$
- ▶ Collisional transport of tungsten → steady-state profile
 - **FACIT** [Maget'20&'22,Fajardo'22]: Fast tool, analytical
 - Includes neoclassical & classical transport
 - Does not cover Hydrogen Temperature Screening (HTS)
 - **NEO** [Belli'08]: kinetic tool with accurate collision operator
 - Includes neoclassical transport only
 - Covers Hydrogen Temperature Screening (HTS)



Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels

Tungsten peaking with ICRH and plasma rotation

Meeting the collapse condition

Summary and conclusion

Direct channel : (horizontal) asymmetry of the electrostatic potential


$$\delta_\phi = \frac{\epsilon}{1 + Z_{eff} T_e / T_i} \left[f_H \left(\frac{T_\perp}{T_\parallel} - 1 \right) + m_i \frac{(R_0 \Omega)^2}{T_i} \right]$$

[Reinke'12]

ICRH drive

(f_H : minority fraction)

- ▶ ICRH drives High Field Side tungsten asymmetry
- ▶ This direct drive is very sensitive to
 - Finite Orbit Width (FOW) effect
 - Pitch angle scattering of hydrogen species on impurities




[Odstrcil'18]

Finite Orbit Width (FOW) effects reduce electron power source and temperature anisotropy

► Surrogate FOW model implemented in EVE

- Large orbits broaden the power deposition & temperature anisotropy profiles [Casson'20]
 - when compared to Zero Orbit Width (ZOW)



Finite Orbit Width (FOW) effect :

Less core electron heating

Less temperature anisotropy (reduction of the direct channel)

Finite Orbit Width (FOW) effects reduce electron power source and temperature anisotropy


- ▶ Surrogate FOW model implemented in EVE: broad power & temperature anisotropy profiles

FOW effects reduce Hydrogen Temperature Screening (HTS)

- ▶ A negative radial gradient of T_{\perp} drives outward pinch [Casson'14]
- ▶ With FOW effect, reduced gradient → reduced impact of HTS
- ▶ NEO computations confirms this
 - Large core tungsten removal in Zero Orbit Width (ZOW) limit
 - Not consistent with bolometry measurement
 - Small HTS effect in FOW limit
 - FACIT computation can be used

*Strong HTS effect:
Not consistent
with experiment*

Small HTS effect



Nonlinear channel: the parallel hydrogen temperature increases with Z_{eff}

- ▶ Pitch angle scattering of hydrogen species on impurities
- ▶ Parallel temperature increases with Z_{eff} [Stix'75]

$$T_{\parallel} \approx \frac{E_{\gamma}}{4}$$


$$E_{\gamma} \propto \left[\frac{2A_i^{1/2}}{n_e} \sum_{\beta} n_{\beta} Z_{\beta}^2 \right]^{2/3} T_e$$

- ▶ EVE/AQL computations in good agreement with the formula :

$$T_{\parallel} = T_{\parallel}^0 [Z_{\text{eff}}]^{2/3}$$


- ▶ The electron heat source (p_e) is not modified
- ▶ *Tungsten accumulation reduces the direct ICRH drive*
- ▶ *Self-regulating mechanism*
- ▶ *This nonlinear channel has not been considered before*

$$\frac{T_{\perp}}{T_{\parallel}}$$



Indirect channel : ICRH induced rotation

- ▶ WEST is a pure RF heated tokamak : no extrinsic momentum source
- ▶ ICRH induced rotation reported on tokamaks
 - Various physics mechanisms involved, difficult to disentangle [Rice'07]
 - **ICRH rotation spin up depends on the minority fraction** [Chouli'14]
 - Large counter rotation spin up at low f_H
 - Moderate co-rotation spin up at large f_H
- ▶ No core rotation measurement available for our experimental case
 - ICRH acceleration evidenced by MHD frequency measurement
 - located on $q=4$
 - Rotation measurements will be produced in future campaigns
 - *Here, toroidal rotation in the core is a free variable*
 - *Can be constrained indirectly by bolometry measurements*
 - through P_{rad} and n_W at $r/a=0.3$



Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels


Tungsten peaking with ICRH and plasma rotation

Meeting the collapse condition

Summary and conclusion

Map of tungsten peaking as a function of ICRH power and toroidal rotation (before collapse)

- ▶ Tungsten profile reconstructed in the collisional limit (no turbulence) with FACIT




Map of tungsten peaking as a function of ICRH power and toroidal rotation (before collapse)

► Two regions of large tungsten peaking


- At **low rotation** and **high ICRH power**
- At **high rotation** and **moderate ICRH power**

► A valley of flat tungsten profile separates the two



Map of tungsten peaking as a function of ICRH power and toroidal rotation before collapse

- ▶ Two regions of large tungsten peaking
 - At **low rotation** and **high ICRH power**
 - At **high rotation** and **moderate ICRH power**
- ▶ A valley of flat tungsten profile separates the two



Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels

Tungsten peaking with ICRH and plasma rotation



Meeting the collapse condition

Summary and conclusion

The collapse can be initiated if the electron heat source is balanced by radiation



- ▶ Tungsten profile → radiative power density, $P_{rad}(r/a=0.3)$, p_{rad}^{max}
- ▶ Electron heat source $p_e = p_e^\Omega + p_e^{ICRH}$
- ▶ Bolometry constraint on $P_{rad}(0.3)$: two branches (*)
 - **Low rotation**: tungsten peaking driven by ICRH
 - **Larger rotation**: tungsten peaking driven by centrifugal force
- ▶ Collapse condition is met on the **higher rotation branch only**
 - But **effective ICRH power must be lower than computed**
 - A reduction of p_e by more than 50% is required
 - The EVE/AQL results are then only indicative

} Collapse criterion : p_{rad}^{max}/p_e



Collapse condition consistent with experiment if large reduction of effective ICRH power

- ▶ At nominal power the injected power on electrons is largely above the radiated power density
- ▶ With an artificial decrease by 60%, the collapse condition is met at $t = 5.75\text{s}$ as observed
 - Only the “high” velocity branch remains
- ▶ Large losses of fast minority ions in the ripple could explain this situation (on-going investigations)



Context

A case of tungsten accumulation during ICRH operation on WEST

Physics of tungsten peaking with ICRH : the 3 channels

Tungsten peaking with ICRH and plasma rotation

Meeting the collapse condition

Summary and conclusion

Radiative collapse are key events for quantifying heat sources & sinks

- ▶ Of particular importance for developing robust heating schemes in tungsten environment

ICRH is a potential drive for tungsten accumulation

- ▶ Due to the induced poloidal asymmetry of the electrostatic potential

ICRH drive is modified by several effects

- ▶ Finite Orbit Widths : Reduced Hydrogen Temperature Screening & broader deposition
- ▶ Non-linear mitigation by the increase of the parallel hydrogen temperature (*new mechanism*)
- ▶ Induced toroidal rotation

ICRH impact on W transport in torque-free discharges, no core particle source

- ▶ Overall tungsten peaking is moderate (except in collapse phases)
 - on WEST, as well as on Alcator C-Mod [Loarte'15]
- ▶ The low rotation branch (ICRH-driven tungsten peaking) remains possibly unexplored
 - non-linear mitigation sufficient to prevent operational limitations ?

Thank you for your attention



Contact: patrick.maget@cea.fr

<https://westusers.partenaires.cea.fr>




partners





Histogram of tungsten peaking in WEST database

► Plateaus database (see Ostuni O4.113, this conference)



Mitigation of collapses at higher density & higher current


- ▶ Series of experiments at 0.5 and 0.7 MA
- ▶ Higher density allows operating at higher ICRH power without radiative collapse
 - EVE/AQL indicates larger core electron heating & lower ICRH drive
 - Qualitatively favorable for lowering ripple losses & tungsten sources
- ▶ Larger plasma current could also be favorable
 - Larger good confinement region with respect to ripple losses




Collapse condition consistent with experiment if large reduction of effective ICRH power

- At nominal power the injected power on electrons is largely above the radiated power density
- With an artificial decrease by 60%, the collapse condition is met at $t = 5.75\text{s}$ as observed

Nominal ICRH power



ICRH power reduced by 60%



Mechanisms leading to an accelerated Tungsten accumulation

- ▶ Response of the Tungsten profile to a decrease of the ion temperature gradient
- ▶ Accelerated radiative collapse :
 - when a hollow temperature leads to a larger Tungsten peaking
 - Otherwise, the collapse is mitigated
- ▶ The collapse process is favored at low rotation

$$T_e(x) = T_e^{\text{ref}}(x) \times (1 - \gamma \exp(-30x^2))$$

$$T_i(x) = T_i^{\text{ref}}(x) \times (1 - (\gamma/2) \exp(-30x^2))$$

