



**HAL**  
open science

## Production of Long-Lived Ultracold Li<sub>2</sub> Molecules from aFermi gas

J. Cubizolles, Thomas Bourdel, S. J.J.M.F. Kokkelmans, G. V. Shlyapnikov,  
C. Salomon

► **To cite this version:**

J. Cubizolles, Thomas Bourdel, S. J.J.M.F. Kokkelmans, G. V. Shlyapnikov, C. Salomon. Production of Long-Lived Ultracold Li<sub>2</sub> Molecules from aFermi gas. 2003. hal-00000540

**HAL Id: hal-00000540**

**<https://hal.science/hal-00000540v1>**

Preprint submitted on 1 Aug 2003

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Production of Long-Lived Ultracold Li<sub>2</sub> Molecules from a Fermi gas

J. Cubizolles<sup>1</sup>, T. Bourdel<sup>1</sup>, S. J. J. M. F. Kokkelmans<sup>1</sup>, G. V. Shlyapnikov<sup>1,2,3</sup> and C. Salomon<sup>1</sup>

<sup>1</sup>Laboratoire Kastler Brossel, Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris 05, France

<sup>2</sup>FOM Institute AMOLF, Kruislaan 407, 1098 SJ Amsterdam, The Netherlands

<sup>3</sup>Russian Research Center Kurchatov Institute, Kurchatov Square, 123182 Moscow, Russia

(Dated: August 1, 2003)

We create weakly-bound Li<sub>2</sub> molecules from a degenerate two component Fermi gas by sweeping a magnetic field across a Feshbach resonance. The atom-molecule transfer efficiency can reach 85% and is studied as a function of magnetic field and initial temperature. The bosonic molecules remain trapped for 0.5s and their temperature is within a factor of 2 from the Bose-Einstein condensation temperature. A thermodynamical model reproduces qualitatively the experimental findings.

PACS numbers: 03.75.Ss, 05.30.Fk, 32.80.Pj, 34.50.-s

Feshbach resonances constitute a unique tool to tune the microscopic interactions in ultracold bosonic and fermionic gases [1, 2, 3]. These resonances arise when the total energy of a pair of colliding atoms matches the energy of a bound state of another hyperfine manifold, leading to the resonant occupation of this state during the collision. Thus, by means of an external magnetic field, one is able to change the magnitude and sign of the scattering length. In bosonic samples, the collapse of Bose-Einstein condensates (BEC) for negative scattering length [4], soliton formation [5, 6], and coherent oscillations between an atomic condensate and molecules [7] have been observed. For fermions with attractive interaction, the superfluid transition temperature is predicted to be maximum near a Feshbach resonance [8, 9, 10, 11, 12].

In this letter we give another striking example of the control of interactions in a Fermi gas. We perform time-dependent experiments near a Feshbach resonance to produce in a reversible manner ultracold and trapped molecules from a quantum degenerate fermionic <sup>6</sup>Li gas. The production efficiency exceeds 80% and the observed molecule lifetime reaches half a second. The phase-space density of these bosonic molecules is on the order of one, the highest value reported so far. Using a similar method with fermionic <sup>40</sup>K atoms, the JILA group recently reported molecule production with a lifetime of 1 ms, atom to molecule conversion efficiency of 50% and direct measurement of the molecular binding energy [13]. Molecules have also recently been produced from <sup>87</sup>Rb and <sup>133</sup>Cs condensates [14, 15] and also from a cold <sup>133</sup>Cs cloud [16]. Molecule formation has also been achieved through one-photon or two-photon photoassociation but with a considerably lower phase-space density than reported here [17]. Our work paves the way to Bose-Einstein condensation of molecules and to the study of the crossover between the regime of molecular BEC and the regime of superfluid BCS pairing in Fermi systems [8, 9, 11, 12].

In <sup>6</sup>Li a broad ( $\simeq 100$  Gauss) Feshbach resonance exists between the two Zeeman sublevels of the hyperfine ground state :  $|1/2, -1/2\rangle$  and  $|1/2, 1/2\rangle$  at a magnetic field of 810 Gauss, see Fig.1 [18, 19]. In a recent study

of this Feshbach resonance, we have reported an anomalous negative value for the gas interaction energy between 700 Gauss and 810 gauss, i.e. below resonance, where the scattering length  $a$  is positive [19]. We suggested that this could be due to the presence of weakly bound molecules confined simultaneously with the cloud of ultracold fermions, and a recent theoretical paper explains our results [20].




FIG. 1: Calculated scattering length  $a$  versus magnetic field for the <sup>6</sup>Li  $|F, m_F\rangle = |1/2, 1/2\rangle, |1/2, -1/2\rangle$  mixture near the 810 Gauss Feshbach resonance. Scanning over the resonance from 1 to 2 in 50 ms produces cold and trapped molecules with an efficiency up to 85%, resulting in an almost complete disappearance of the atomic signal at 2. Reversing then the scan to the initial position re-establishes the initial atomic signal. In all cases, atoms are detected and imaged at position 4 after abrupt switch-off of the  $B$  field.

The method used here to produce molecules is illustrated in Fig.1 and was suggested for Bose gases in [21, 22, 23]. It consists in scanning over a Feshbach resonance from the region of attractive interaction ( $a < 0$ ) to region 2 in Fig.1 where  $a$  is large and positive, and where a weakly bound molecular state exists with energy  $E_b = -\hbar^2/ma^2$  (where  $m$  is the atomic mass). It is thus energetically favorable to populate this bound state. Having twice the polarisability of individual atoms, the

molecules are expected to be confined in an optical dipole trap with the same oscillations frequencies as atoms and twice the trap depth.

Experimentally, using the setup described in [19], we prepare  $N_1 = N_{\uparrow} + N_{\downarrow} = 1.5 \cdot 10^5$  atoms in a 50(10)% mixture of  $|1/2, -1/2\rangle$  and  $|1/2, 1/2\rangle$  in a crossed Nd:YAG dipole trap at  $B = 1060 G$  where the scattering length between  $|1/2, -1/2\rangle$  and  $|1/2, 1/2\rangle$  is large and negative  $a = -150 \text{ nm}$  (position 1 in Fig. 1). For a power of 1.53 W (resp. 4.3 W) in the horizontal (resp. vertical) trapping beam propagating along Ox (resp. Oy), oscillation frequencies are  $\omega_x/2\pi = 2.9(3) \text{ kHz}$ ,  $\omega_y/2\pi = 5.9(5) \text{ kHz}$  and  $\omega_z/2\pi = 6.5(6) \text{ kHz}$ . By evaporation in the optical trap, the temperature  $T$  of the gas mixture can be tuned between 0.2 and  $0.5 T_F$  where  $T_F$  is the Fermi temperature defined by  $T_F = \hbar\bar{\omega} (3N_1)^{1/3}/k_B$  and  $\bar{\omega} = (\omega_x\omega_y\omega_z)^{1/3}$ .

The Fermi gas quantum degeneracy is measured through analysis of absorption images after abrupt ( $20 \mu\text{s}$ ) switch-off of the magnetic field and time of flight expansion of the cloud. Free  $^6\text{Li}$  atoms are detected at zero magnetic field, for which  $a \simeq 0$ , using laser light tuned to the D2 line, position 4 in Fig. 1. Having prepared the degenerate Fermi gas at position 1, we sweep the magnetic field in 50 ms to position 2 where  $a$  is large and positive, the region where a weakly-bound molecular state exists. Absorption images indeed reveal that the number of atoms  $N_2$  as counted after  $B$  switch-off to position 4 as explained above, has dramatically decreased to  $\simeq 20\%$  of the initial number  $N_1$ . After letting the system at position 2 for a variable wait time  $t_w$  between 0 and 5 s, we sweep the magnetic field back to position 1 in 50 ms and count the number of atoms,  $N_3$ , after this round-trip through resonance.

Surprisingly, when position 2 is at 689 Gauss and  $t_w = 0$ , we find  $N_3 \simeq N_1$ , indicating that no loss has occurred in 100 ms. This proves that the decrease in detected atom number at position 2 is not due to losses. We are thus led to conclude that: (i) at position 2, 80% of the atoms were in a molecular bound state which is not detected using light resonant with the atoms at position 4, (ii) the atom-to-molecule formation process is reversible.  $(N_3 - N_2)/2$  represents thus the number of molecules at position 2.

In comparison to other recent experiments on bosons and fermions [7, 13, 14, 15], our 50 ms scan over the Feshbach resonance is very slow and, in particular, much longer than the collision time in the cloud. This allows the system to relax towards thermal equilibrium between atoms and molecules during the magnetic field ramp.

The relative fraction of atoms bound in a molecule,  $(N_3 - N_2)/N_3$ , and the temperature of the atoms at position 2 and 3 are plotted in Fig. 2 as a function of the magnetic field at position 2. The initial parameters at 1060 G for this experiment are  $T = 4.7 \mu\text{K}$ ,  $T_F = 11 \mu\text{K}$ , and  $N_1 = 8 \cdot 10^4$ . The frequencies of the trap are  $\omega_x/2\pi = 2.2(2) \text{ kHz}$ ,  $\omega_y/2\pi = 4.6(4) \text{ kHz}$ ,  $\omega_z/2\pi = 5.1(5) \text{ kHz}$ .




FIG. 2: a: fraction of atoms in molecular bound state,  $(N_3 - N_2)/N_3$ , versus magnetic field at position 2 in Fig. 1. b: Corresponding atom temperature at position 2 (squares) and 3 (circles). Solid (dotted) line corresponds to the non-interacting (interacting) thermodynamical model described in text.

One observes that very few molecules are detected above 0.77 kG. When decreasing the magnetic field, the fraction of molecules increases up to 60%. Below 650 G, losses become important and the molecular fraction decreases, an effect that we will study below.

A key parameter in our detection scheme is the switch off time ( $20 \mu\text{s}$ ) of the Feshbach magnetic field from position 2. Indeed, the molecules may or may not dissociate in this process. Molecules will dissociate if the relative rate of change of the binding energy  $dE_b/E_b dt$  is greater than the oscillation pulsation  $E_b/\hbar$ . So, the dissociation of molecules is governed by the parameter  $\alpha = \hbar dE_b/E_b^2 dt$ . If  $\alpha \gg 1$ , they dissociate and appear as free atoms in the absorption image. Our detection scheme is unable to prove molecule formation. On the contrary, if the binding energy is sufficiently large at position 2 ( $\alpha \ll 1$ ), the molecular state follows adiabatically the change of  $B$  to a deeper bound state which is off-resonant with the detection laser. In this case,  $N_3 > N_2$  is the signature of molecule formation.

For  $B = 700 \text{ G}$ , one has  $\alpha \simeq 1$ . Reducing the rate

$dB/dt$  by a factor of  $\simeq 500$ , we observe a shift of the threshold for molecule detection to 800 G, as expected from the expression of  $\alpha$  and the position of the Feshbach resonance at 810 G [19]. As a second consequence of the role of the switch-off time, we attribute the decrease of the molecular fraction above 720 G in Fig. 2a to the crossover between the adiabatic and non adiabatic regimes.

The temperature of the atoms at position 2 and 3 is plotted in Fig. 2b. A decrease of the magnetic field increases the binding energy of the molecules. Therefore, the molecular fraction increases and so does the temperature. However, sweeping back the magnetic field to the initial position, the temperature returns close to its initial value, demonstrating the reversibility of the molecular formation process. According to [24], in our experiment the rate of 3-body recombination of the weakly bound state is greater than  $10^4$  Hz (for  $B > 665$  G). Moreover, the atom-atom, atom-molecule and molecule-molecule [25] collision rates are also large compared to the rate at which the resonance is scanned. Therefore the system remains close to thermal equilibrium. During the molecule formation process, the released binding energy must be converted into kinetic energy, therefore the increase of temperature in Fig. 2b, as  $a$  decreases, is expected.

Interestingly, the efficiency of molecule formation strongly depends on the trap depth and reaches values up to 80 % in Fig. 3, the highest efficiency reported so far. The trap depth is slowly (800 ms) reduced at  $B = 1060$  G, before the magnetic field sweep through resonance. The gas temperature and the Fermi temperature decrease by evaporation and adiabatic cooling due to the reduction of the trap oscillation frequencies. At 689 G,  $a = 78$  nm,  $E_b/k_B = 12 \mu\text{K}$ . When reducing the trap depth,  $T$  and  $T_F$  become small compared to  $E_b$  and the molecular fraction increases towards 1. Other experiments show that both evaporation (reduction of the ratio  $T/T_F$ ), and adiabatic cooling (reduction of  $T$  and  $T_F$  by the same factor) effectively lead to an increase of the molecular fraction. For instance, at constant  $T_F = 11 \mu\text{K}$ , as the temperature is decreased from  $0.5 T_F$  to  $0.3 T_F$ , we find that the molecular fraction at 689 G increases from 40 % to 65 %. Similarly, at constant  $T/T_F = 0.3$ , as  $T$  varies from  $4.5 \mu\text{K}$  to  $1.8 \mu\text{K}$ , it increases from 50 % to 85 %.

Our results can be understood within a thermodynamical model assuming that atoms and molecules are in equilibrium in a trap during the magnetic field sweep [20]. We assume that the sweep is adiabatic, so the entropy is conserved and it is calculated for the initial situation where no molecules are present at  $B = 1060$  G. The atom-atom, atom-molecule, molecule-molecule interactions were either neglected (solid line in Fig. 2), or included via a mean field contribution proportional to  $a$  [24, 25], in the regime where it is allowed (dotted line). The molecular fraction is then determined by a set of two




FIG. 3: Fraction of atoms bound in a molecule as a function of the depth of the dipole trap for  $B = 689$  G,  $a = 78$  nm,  $E_b = 12 \mu\text{K}$ .

parameters  $T/T_F$  and  $E_b/T_F$ . As the binding energy gets more negative, the fraction of molecules increases since the molecular state becomes energetically more favorable. Moreover, the temperature increases as each molecule formation event transfers the binding energy into kinetic energy. Both models agree fairly well with our data in the region where both the detection of molecules is efficient and losses are unimportant ( $660 \lesssim B \lesssim 720$  G). In the interacting case, a perfect agreement is achieved if we allow for a 10% mismatch in the quality of the spin mixture. In the strongly interacting region where  $na^3 \geq 1$ , the non-interacting model is shown with a dashed line because we do not expect our model of separated gases of fermionic atoms and bosonic molecules to be valid. Those models also account qualitatively for the increase in the molecular fraction with the evaporation in Fig. 3. Indeed, as the trap depth is lowered,  $T$  decreases, the ratio  $E_b/T$  increases and the bound state will then have a higher occupancy.




FIG. 4: Decay of  $\text{Li}_2$  molecules in the optical trap for two values of the magnetic field: (a)  $B = 689$  G,  $a = 78$  nm, (b)  $B = 636$  G,  $a = 35$  nm. Note the difference in time scales. Solid lines are fits with 2-body and 1-body processes giving initial time constants of 500 ms in (a) and 20 ms in (b).

In order to investigate prospects for the formation of a

molecular BEC, two questions immediately arise: what are the lifetime and degeneracy parameter of the produced molecular cloud? The decay of the molecules in the optical trap is found by measuring the quantity  $(N_3 - N_2)/2$  as a function of the wait time  $t_w$  at position 2 before returning to 3 for the detection process. Fig. 4 presents the number of remaining molecules as a function of the time elapsed at point 2 in a trap with frequencies  $\omega_x/2\pi = 0.95(10)$  kHz,  $\omega_y/2\pi = 2.0(2)$  kHz,  $\omega_z/2\pi = 2.2(2)$  kHz. In Fig. 4 a and b, the initial temperatures at position 1 are the same,  $1.1 \mu\text{K} = 0.3 T_F$ . Fig 4a corresponds to  $B = 689 \text{ G}$ ,  $a = 78 \text{ nm}$ ,  $E_b/k_B = 12 \mu\text{K}$ , whereas Fig. 4b corresponds to  $B = 636 \text{ G}$ ,  $a = 35 \text{ nm}$ , and  $E_b/k_B = 60 \mu\text{K}$ . The initial fraction of unpaired atoms (not engaged in a molecule) is 17% in (a) and 25% in (b). For a change in  $a$  by the modest factor of 2.2, the two samples exhibit strikingly different lifetimes. Decays are well fitted by 2-body + 1-body loss processes with initial lifetimes of  $\simeq 500 \text{ ms}$  in (a) (689 G) and 20 ms in (b) whereas the 1-body decay exceeds 4 seconds.

A strong decrease in the lifetime of molecules with decreasing  $a$  is expected assuming that they undergo collisional relaxation to deep bound states [25]. However, the observed factor of 25 in lifetimes is surprisingly large. At 689 G, we can evaluate the 2-body loss rate to be  $G = 2.4_{-1.6}^{+3.2} 10^{-13} \text{ cm}^3 \cdot \text{s}^{-1}$ . At 636 G, we have initially  $N_3 \simeq 0.3N_1$ . These fast losses indicate that very likely some evaporation is involved in the limited lifetime and that the system has not reached thermal equilibrium. Therefore, the 2-body loss rate cannot be safely estimated.

Let us now evaluate the phase space density of the trapped molecules. In Fig. 2, at 689 G, there are  $1.8 \cdot 10^4$  molecules, confined with  $3.3 \cdot 10^4$  atoms. The measured atom temperature is  $T_{at} = 6.7 \mu\text{K}$ . Assuming thermal equilibrium between atoms and molecules,  $T_{at} = T_m$  for magnetic fields above  $\simeq 650 \text{ G}$ , we obtain the peak density of molecules  $n_m \simeq 4 \cdot 10^{13} \text{ cm}^{-3}$ . Then the critical temperature for molecule condensation is reduced due to the interactions and is  $3.5 \mu\text{K}$  giving  $T/T_C \simeq 2$ . Since  $n_m a_m^3 \simeq 5 \cdot 10^{-3} \ll 1$  with  $a_m = 0.6 a = 47 \text{ nm}$  ([25]), the molecular gas is in the dilute regime and the mean distance between molecules is larger than the typical size of a molecule  $a$ . In fact, for all data between 675 G and 750 G, the phase space density of the molecules is not far from the condensation point. Since, the lifetime of the molecules, 500 ms, is long in this region when compared to the molecule-molecule elastic collision time  $1/(n_m 8\pi a_m^2 v_m) \simeq 3 \mu\text{s}$ , (for  $a_m = 47 \text{ nm}$ ), it should be possible to evaporate the molecules further to reach the Bose-Einstein condensation threshold.

In summary, we have produced long-lived and trapped  $\text{Li}_2$  dimers. The atom-molecule conversion efficiency can approach 1 when the Fermi quantum degeneracy is strong. The lifetime of the trapped molecules strongly depends on the scattering length. Current research concen-

trates on evaporation towards molecular BEC. Prospects for producing superfluid Fermi mixtures and for investigating the transition between molecular condensates and superfluid Fermi gases are promising [11, 12, 25, 26].

We are grateful to D. S. Petrov, S. Stringari, Y. Castin, R. Combescot and J. Dalibard for useful discussions. This work was supported by CNRS, Collège de France, and Région Ile de France. S. Kokkelmans acknowledges a Marie Curie grant from the E.U. under contract number MCFI-2002-00968. Laboratoire Kastler Brossel is *Unité de recherche de l'Ecole Normale Supérieure et de l'Université Pierre et Marie Curie, associée au CNRS*.

- 
- [1] H. Feshbach, Ann. Phys. (N.Y.) **5**, 357 (1958), **19**, 287 (1962).
  - [2] See for instance, *ultracold matter*, nature insight, **416** (2002).
  - [3] S. Inouye, *et al.*, Nature **392**, 151 (1998).
  - [4] E. A. Donley, *et al.*, Nature **412**, 295 (2001).
  - [5] L. Khaykovich, *et al.*, Science **296**, 1290 (2002).
  - [6] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, Nature **417**, 150 (2002).
  - [7] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Wieman, Nature **417**, 529 (2002).
  - [8] P. Nozières and S. Schmitt-Rink, J. Low temp. Phys. **59**, 195 (1985).
  - [9] C. A. R. Sá de Melo, M. Randeria, and J. R. Engelbrecht, Phys. Rev. Lett. **71**, 3202 (1993).
  - [10] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiofalo, and R. Walser, Phys. Rev. Lett. **87** 120406 (2001).
  - [11] Y. Ohashi, and A. Griffin, Phys. Rev. Lett. **89**, 130402 (2002).
  - [12] J. N. Milstein, S. J. J. M. F. Kokkelmans, and M. J. Holland, Phys. Rev. A **66**, 043604 (2002).
  - [13] C. A. Regal, C. Ticknor, J. L. Bohn, and D. S. Jin, Nature **424**, 47 (2003).
  - [14] J. Herbig, *et al.*, submitted to Science.
  - [15] S. Dürr, T. Volz, A. Marte, and G. Rempe, Cond-mat/0307440.
  - [16] C. Chin, A. J. Kerman, V. Vuletic, and S. Chu, Phys. Rev. Lett. **90**, 033201 (2003).
  - [17] R. Wynar, R. Freeland, D. Han, C. Ryu, and D. Heinzen, Science **287**, 1016 (2000).
  - [18] S. Gupta, *et al.*, Science **300**, 1723 (2003).
  - [19] T. Bourdel, *et al.*, Phys. Rev. Lett. **91**, 020402 (2003).
  - [20] S. J. J. M. F. Kokkelmans, G. V. Shlyapnikov, and C. Salomon, to be published.
  - [21] E. Timmermans, P. Tommasini, R. Cote, M. Hussein and A. J. Kerman, cond-mat/9805323.
  - [22] F. A. van Abeelen and B. J. Verhaar, Phys. Rev. Lett. **83**, 1550 (1999).
  - [23] F. H. Mies, E. Tiesinga, and P. S. Julienne, Phys. Rev. A **61**, 022721 (2000).
  - [24] D. S. Petrov, Phys. Rev. A **67**, 010703(R) (2003).
  - [25] D. S. Petrov *et al.*, to be published.
  - [26] L. D. Carr, Y. Castin, and G. V. Shlyapnikov, to be published.