



HAL
open science

Deformation of grain boundaries in polar ice

Gaël Durand, François Graner, Jérôme Weiss

► **To cite this version:**

Gaël Durand, François Graner, Jérôme Weiss. Deformation of grain boundaries in polar ice. 2004. hal-00000573v2

HAL Id: hal-00000573

<https://hal.science/hal-00000573v2>

Preprint submitted on 29 Apr 2004 (v2), last revised 22 Jul 2004 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deformation of grain boundaries in polar ice

G. DURAND¹, F. GRANER²[*], AND J. WEISS¹

¹ *Laboratoire de Glaciologie et de Géophysique de l'Environnement (FRE 2192 CNRS - Université J. Fourier Grenoble 1), BP 96, F-38402 Saint Martin d'Hères cedex, France*

² *Laboratoire de Spectrométrie Physique (UMR 5588 CNRS - Université J. Fourier Grenoble 1), BP 87, F-38402 Saint Martin d'Hères cedex, France*

PACS. 83.80.Nb – Geological materials: Earth, magma, ice, rocks, etc... .

PACS. 81.70.-q – Nondestructive materials testing and analysis: optical methods.

PACS. 62.20.Fe – Deformation and plasticity (including yield, ductility, and superplasticity).

Abstract. –

The ice microstructure (grain boundaries) is a key feature used to study ice evolution and to investigate past climatic changes. We studied a deep ice core, in Dome Concordia, Antarctica, which records past mechanical deformations. We measured a “texture tensor” which characterizes the pattern geometry and reveals local heterogeneities of deformation along the core. These results question key assumptions of the current models used for dating.

Motivations. – Polar ice cores are the focus of many investigations because they record the history of climatic changes. Owing to snow accumulation, snow to ice transformation and slow ice sheet flow ($\sim 10^{-12} \text{s}^{-1}$), a journey down to the deep layers of the ice sheet is a journey back to several hundred of thousands of years into the past [1].

A crucial step of paleoclimatic studies from ice cores is dating. In Antarctica, counting annual layers is impossible [2]: absolute dating is only possible for the very top of the ice cores where ice layers containing volcanic impurities can be related to historical volcanic eruptions. Below, dating relies on ice sheet flow models of the evolution of ice layer thinning with depth [2]. Such models are loosely constrained by the identification of large climatic transitions. For the sake of simplicity, these models assume a smooth and monotonous increase of the thinning with depth, hence ignore any possible localization of the deformation [2, 3].

In this letter, we question this essential assumption. We present a method to extract geometrical information (such as thinning, shear, localization of the deformation) from pictures of a cellular pattern using local spatial averages of the “texture tensor” [4]. We apply this analysis to the grain boundaries (the so-called “microstructure”) of ice samples from a deep ice core.

Samples. – Dome Concordia, Antarctica (75° 06' 04" S, 123° 20' 52" E, elevation 3233 m a.s.l) is at the summit of an Antarctic ice dome. It has been chosen because it is usually assumed that the ice flow is axisymmetric around the vertical (z) axis, and isotropic within the




Fig. 1 – Two thin sections of ice imaged between crossed polarizers in white light. Each grain has an almost uniform crystallographic orientation, visualized by its color. (a) At 362 m depth, the microstructure looks isotropic; (b) at 2629 m depth the microstructure is visibly anisotropic (the white scratches do not affect the image analysis, which ignores them). Superimposed on (a): notations used in the measurement of $\overline{\overline{M}}$, here *e.g.* at the site labeled by a red dot. We note $\vec{\ell}$ a vector linking it to one of its neighbours. There are 3 such vectors in the first shell, $p = 1$ (purple); 6 in shell $p = 2$ (green); 12 in shell $p = 3$ (orange).

horizontal plane. In February 2003, the European ice core drilling program (EPICA) reached the depth of 3201.65 m below the ice surface, close to the ground (depth 3309 ± 22 m).

Samples have been extracted and transported according to standard procedures [5]. We analyzed ~ 11 cm high, ~ 0.1 mm thin sections (Fig. 1), with a sampling interval varying from 2 to 25 m. The vertical axis on a thin section always corresponds to the true *in situ* vertical axis z ; its horizontal axis, labelled x in what follows, has an unknown and variable orientation within the true *in situ* horizontal plane.

Data analysis. – By processing pictures of ice under crossed polarizers, we determine the grain boundaries [5], then the sites (“vertices”) where three boundaries meet, see Fig. (1a). For each pair of neighbouring sites, we draw the vector which links them and denote it by $\vec{\ell}$. We then construct the tensor $\vec{\ell} \otimes \vec{\ell}$: its coordinates are $(\ell_i \ell_j)$, where i, j are here x or z ; this tensor is not sensitive to the sign of the vector (it is invariant under $\vec{\ell} \rightarrow -\vec{\ell}$), but it characterizes its length ℓ and its direction.




Fig. 2 – Analysis of the images presented in Fig. 1. The texture tensor $\overline{\overline{M}}$ measured at each site is represented as an ellipse, with its axes along the eigenvectors, and its half-axes proportional to eigenvalues λ_1, λ_2 respectively. A site around which the pattern is isotropic is represented by a circle; conversely, a strong anisotropy is represented as an elongated ellipse. The size of the ellipse (same scale for each ellipse) represents the local length of the grain boundaries, *i.e.* quantifies the local grain size. Due to the definition of $\overline{\overline{M}}$ (eq. 1), we exclude sites closer than p grains from the image boundary; here $p = 3$.

To statistically characterize the pattern, ref. [4] proposed to define the *texture tensor* $\overline{\overline{M}}$, as the average of $\vec{\ell} \otimes \vec{\ell}$ over a box of fixed size. Such box should be smaller than the image (in order to visualize local details), but still large enough to include a number $N \gg 1$ of wall vectors (to have relevant statistics) [6]. Here, grains have variable size (grains grow with time, hence old grains in deep ice are much larger than young grains near the dome surface): it would be difficult to select such a fixed box size.

We thus chose a local averaging and define $\overline{\overline{M}}$ at each given site as:

$$\overline{\overline{M}} = \left\langle \left(\begin{array}{cc} \ell_x^2 & \ell_x \ell_z \\ \ell_z \ell_x & \ell_z^2 \end{array} \right) \right\rangle_p = \frac{1}{N} \sum_{k=1}^N \vec{\ell}(k) \otimes \vec{\ell}(k) = \mathcal{R} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \mathcal{R}^{-1}. \quad (1)$$

Here, \mathcal{R} is the rotation which diagonalizes $\overline{\overline{M}}$, and (λ_1, λ_2) are the corresponding eigenvalues; $\langle \cdot \rangle_p$ denotes the average over N vectors, up to the p -th neighbours (Fig. 1a). Hence, at each site we include approximately the same number of vectors, $N \approx 3 + 6 + \dots + 3 \times 2^{p-1}$. A




Fig. 3 – Histogram of U_{xz} (left) and of U_{zz} (right) measured with $p = 3$ on Fig. (2a): 362 m deep sample (dots), and the corresponding gaussian distribution (same mean and standard deviation) (thin solid line); and on Fig. (2b): 2629 m deep sample (thick solid line).

smaller p explores local details, a larger p (hence a larger scale) improves the statistics.

This tensor, independent from crystallographic information, does not require any knowledge or hypothesis regarding the past history of the material. The diagonal components M_{xx} and M_{zz} of this tensor are both of order of the average square distance $\langle \ell^2 \rangle$ between sites. Conversely, the off-diagonal component $M_{xz} = M_{zx}$ is much smaller, and even vanishes when the pattern is isotropic. Hence $\lambda_1, \lambda_2 > 0$: the largest eigenvalue corresponds to the direction in which grains are most elongated (Fig. 2).

While many other quantitative descriptors exist [7], few are adapted to the determination of the anisotropy of such microstructure pattern. Previous studies on polar ice calculate the aspect ratio of the grains from the linear intercept method [5, 8]. Other use the eigenvalues of the grain inertia tensor [9, 10]; this is a true tensor, with mathematical advantages: for instance, its value does not depend on the particular choice of axes, hence undergoes less artifacts when examining the 2D cut of a 3D pattern.

The tensor $\overline{\overline{M}}$ has the same advantages; but also an additional one, thanks to its definition (eq. 1) being quadratic in $\vec{\ell}$: it has a physical signification in terms of mechanical deformations [4]. More precisely, its variations (with respect to a reference $\overline{\overline{M}}_0$ measured in an isotropic, relaxed state) define a statistical strain tensor $\overline{\overline{U}}$:

$$\overline{\overline{U}} = \frac{\log(\overline{\overline{M}}) - \log(\overline{\overline{M}}_0)}{2}, \quad (2)$$

which exists even during large deformations and, at least during elastic deformations, coincides with the classical definition of strain [4].

For the 2D pattern studied here, $\overline{\overline{M}}_0$ is unknown, and processes other than deformation (such as grain growth) modify the microstructure. We first assume that the (viscoplastic) deformation is isochore (incompressibility of ice), so that $Tr \overline{\overline{U}} = 0$ and $\lambda_1^0 \lambda_2^0 = \lambda_1 \lambda_2$. We also assume that the reference state is isotropic; hence $\overline{\overline{M}}_0$ is isotropic too, so that $\lambda_1^0 = \lambda_2^0 =$




Fig. 4 – Localization of deformation inside the ice core. Each point corresponds to one of 329 samples; same scale for both figures. (a) $|U_{xz}|$ versus depth. Closed circles highlight samples where $|U_{xz}| > 3\sigma$. (b) U_{zz} versus depth. Two large closed circles linked by a thick solid line indicate two successive measures which 3σ confidence intervals do not overlap. A large open circle indicates a sample with a large significant shear ($U_{xz} > 3\sigma$, from (a)). Vertical lines indicate zones, associated to climatic transitions, where the grain growth rate changes significantly (see Discussion).

$\sqrt{\lambda_1 \lambda_2}$. We then can measure $\overline{\overline{U}}$ through the following equation:

$$\overline{\overline{U}} = \frac{1}{2} \mathcal{R} \begin{pmatrix} \log\left(\frac{\lambda_1}{\sqrt{\lambda_1 \lambda_2}}\right) & 0 \\ 0 & \log\left(\frac{\lambda_2}{\sqrt{\lambda_1 \lambda_2}}\right) \end{pmatrix} \mathcal{R}^{-1} = \frac{1}{4} \mathcal{R} \begin{pmatrix} \log\left(\frac{\lambda_1}{\lambda_2}\right) & 0 \\ 0 & \log\left(\frac{\lambda_2}{\lambda_1}\right) \end{pmatrix} \mathcal{R}^{-1}. \quad (3)$$

Hence $\overline{\overline{U}}$ has two independent components U_{zz} and U_{xz} , both largely insensitive to the grain size fluctuations.

To estimate the statistical variability of the measure, we generate 150 isotropic microstructures from a 2D Potts model of normal grain growth [11]. We check that (i) each component of $\overline{\overline{U}}$ fluctuates around zero; (ii) its distribution is gaussian (see an example with $p = 3$ on Fig. 3); (iii) its standard deviation σ only depends on the number N of vectors, *i.e.* on the scale of observation p ; and (iv) $\sigma(N) \propto N^{-1/2}$: more precisely, $\sigma \approx 0.34 N^{-1/2}$.

Results. – We first measure the deformation and its heterogeneity within one sample. Fig. (2a) shows that the upper sample (depth 362 m) is almost homogeneous. Its anisotropy is small and, as expected from a sample taken at a dome, it mainly reflects a uniaxial compression along the vertical axis. Conversely (Fig. 2b), the anisotropy of the lower sample, at 2629 m: is stronger; is localized in space; and breaks the axisymmetry around z (see Discussion).

Quantitatively, the distributions of U_{xz} (vertical compression) and U_{zz} (horizontal shear) of both samples differ significantly (Fig. 3). The upper sample is not sheared ($\langle U_{xz} \rangle = 0.00$), and is slightly flattened onto the horizontal plane ($\langle U_{zz} \rangle = -0.05$). The lower sample is significantly sheared ($\langle U_{xz} \rangle = 0.13$) and strongly flattened ($\langle U_{zz} \rangle = -0.25$). Note that for $p = 3$ ($N = 21$) the intrinsic value of σ (see above) is 0.074; here, the statistical deviation (0.1) is comparable, but slightly larger, reflecting the heterogeneity of deformation at small scales ($10^{-2} - 10^{-1}$ m).

We now turn to heterogeneities at large scale ($1 - 10^2$ m), by measuring the global $\overline{\overline{U}}$ (integrated over all the grain boundaries of a given sample), versus the sample depth (Fig. 4). Whereas U_{xz} remains around zero, as expected in a dome situation, U_{zz} increases (in absolute value) with depth.

The essential point is the localization: namely, the observation of a deformed (sheared or compressed) layer immediately close to less deformed ones. As mentioned, the grain size increases with depth, therefore the number of vectors per sample decreases, hence σ increases. The localization is nevertheless significant, and is increasingly frequent at increasing depth (closed circles on Fig. 4).

Discussion. – Since the pictures are 2D cuts of actual 3D grains, the apparent average grain size could be misestimated; however, since $\overline{\overline{U}}$ is dimensionless, it should remain unaffected. When, and only when, the pattern is axisymmetric, we can refine the above analysis (eq. 3). In that case, the eigenvectors are (x, y, z) , and the corresponding eigenvalues are $(\lambda_1, \lambda_1, \lambda_2)$. The eigenvalues of $\overline{\overline{U}}$ are now $\log(\lambda_1/\lambda_2)/6$ and $\log(\lambda_2/\lambda_1)/3$. We have checked that the results corresponding to the 2D analysis (Figs. 3, 4) and the 3D one (data not shown) are completely similar, with even slightly more heterogeneities using the latter analysis. Of course, care is required for the interpretation of an individual sheared sample (for instance regarding U_{zz} at 2629 m, Fig. 3b). But on average, the breaking of the axisymmetry hypothesis (large open circles on Fig. 4b)) does not introduce significant artifacts on the compression heterogeneities we recorded (closed circles on Fig. 4b).

Processes such as normal grain growth counteract the anisotropy of grains induced by deformation. Hence the strain recorded by the microstructure $\overline{\overline{U}}$ underestimates the strain $\overline{\overline{\epsilon}}$ actually experienced by the material. The growth rate itself could undergo some fluctuations. At few depths, near the bottom of the core, a correlation has been observed between small grain sizes (related to a large amount of dust in the ice [12]) and large strains. In these few cases, highlighted by the thick vertical lines in figure 4b, the large differences of U_{zz} observed between two adjacent layers could be partly explained by the difference in the grain growth rate. However, this effect is unable to explain most heterogeneities (closed circles on Fig. 4b). In addition, it does not explain the large number of shear layers (closed circles on Fig. 4a). We thus argue that the heterogeneity of anisotropy, observed at both small and large scales, has a mechanical origin.

Conclusion and perspectives. – A dating chart is the relation $t(z)$ between depth z and age t of ice: it requires a model, and hypotheses. The ice core is drilled exactly at the summit of a dome, in order to assume that the flow is axisymmetric; so that the ice thinning results from vertical compression only, without horizontal shear. Current models further assume a smooth and monotonous increase of the thinning of ice defined as the ratio $e(z)/e(0)$, where e is the thickness of the annual ice layer at depth z .

The second assumption has already been shown to be wrong in the Greenland ice cores GRIP and GISP2 ([13, 14, 15]). Flow disturbances have been reported within at least the 30% deepest part of the GISP2 core, based on the observations of wavy ash layers, crystal stripings visible by eyes, and anomalous fabrics [13]. The dating has been particularly questioned by the observation of a folding, *i.e.* local inversion of ice layers [16].

Here, our method is more accurate and applies to ice itself, without requiring markers nor extreme events. We show that flow disturbances are detectable from almost the top of the ice sheet, and increase in number and intensity with depth. In fact, both dating model assumptions are contradicted by our results: the strain gradient is variable and not even always positive (Fig. 4b); and although the flow at a dome is axisymmetric on average, in detail there is a symmetry breaking due to shear, especially in deep layers (Fig. 4a). This suggests to reconsider current standard dating charts.

As a first step, we might replace in the dating model the assumed strain gradient by the value of $\partial U_{zz}/\partial z$ observed in the microstructure. Unfortunately, as mentioned above, this

still underestimates the real strain, and the proportionality coefficient could vary with the grain growth rate.

In the future, we expect to correlate the grain boundary pattern with the *c*-axis orientations, to improve our understanding of their coupling.

* * *

We thank M. Aubouy for his support and useful discussions, M. Glock for reading the manuscript, and M. Krinner and M. Manouvrier for their technical support. This work was partially supported by CNRS ATIP 0693. This work is a contribution to the “European project for Ice Coring in Antarctica” (EPICA), a joint ESF (European Science Foundation)/EC scientific program, funded by the European Commission under the Environment and Climate Programme (1994-1998) contract ANV4-CT95-0074 and by national contributions from Belgium, Denmark, France, Germany, Italy, the Netherlands, Norway, Sweden, Switzerland and the United Kingdom.

REFERENCES

- [*] Author for correspondence at graner@ujf-grenoble.fr. Fax: (+33) 4 76 63 54 95.
- [1] PETIT J.R., JOUZEL J., RAYNAUD D., BARKOV N.I., BARNOLA J.M., BASILE I., BENDER M., CHAPPELLAZ J., DAVIS M., DELAYGUE G., DELMOTTE M., KOTLYAKOV V.M., LEGRAND M., LIPENKOV V.Y., LORIUS C., PEPIN L., RITZ C., SALTZMAN E. and STIEVENARD M., *Nature*, **399** (1999) 429.
- [2] SCHWANDER J., JOUZEL J., HAMMER C.U., PETIT J.R., UDISTI R. AND WOLFF E. , *Geophysical research letters*, **28** (2001) 4243.
- [3] HAMMER C.U., CLAUSEN H.B., DANSGAARD W., GUNDESTRUP N., JOHNSEN S.J. and REEH N., *J. Glacio.*, **20** (1978) 3.
- [4] AUBOUY M., JIANG Y., GLAZIER J. A. and GRANER F., *Granular Matt.*, **5** (2003) 67
- [5] GAY M., and WEISS J., *J. Glaciol.*, **45** (1999) 547.
- [6] ASIPAUSKAS M., AUBOUY M., GLAZIER J. A., GRANER F. and JIANG Y., *Granular Matt.*, **5** (2003) 71.
- [7] MECKE K. and STOYAN D. (Editors), *Morphology of Condensed Matter - Physics and Geometry of Spatially Complex Systems, Lecture Notes in Physics*, Vol. **600** (Springer, Heidelberg) 2002
- [8] ARNAUD L., WEISS J., GAY M. and DUVAL P., *Annals Glacio.*, **30** (2000) 8.
- [9] N. AZUMA, Y. WANG, K. MORI, H. NARITA, T. HONDOH, H. SHOJI and O. WATANABE, *Ann. Glaciol.*, **29** (1999) 163
- [10] WILEN L.A., DIPRINZIO C.L., ALLEY R.B. and AZUMA N., *Micro. Res. and Tech.*, **62** (2003) 2.
- [11] ANDERSON M.P., GRETT G.S. and SROLOVITZ D.J., *Philos. Mag. B*, **59** (1989) 293.
- [12] WEISS J., VIDOT J., GAY M., ARNAUD L., DUVAL P. and PETIT J.R., *Annals Glacio.*, **35** (2002) 552.
- [13] ALLEY R.B., GOW A.J., MEESE D.A., FITZPATRICK J.J., WASHINGTON E.D. and BOLZAM J.F., *J. Geophys. Res.*, **102** (1997) 26819.
- [14] GOW A.J., MEESE D.A., ALLEY R.B., FITZPATRICK J.J., ANANDAKRISHNAN S., WOODS G.A. and ELDER B.C., *J. Geophys. Res.*, **102** (1997) 26559.
- [15] DAHL-JENSEN D., THORSTEINSSON T., ALLEY R. and SHOJI H., *J. Geophys. Res.*, **102** (1997) 26831.
- [16] GROOTES P.M., STUIVER M., WHITE J.W.C., JOHNSEN S. and JOUZEL J., *Nature*, **366** (1993) 552.