



**HAL**  
open science

## About "On certain incomplete statistics" by Lima et al

Michel Pezeril, Alain Le Méhauté, Qiuping A. Wang

► **To cite this version:**

Michel Pezeril, Alain Le Méhauté, Qiuping A. Wang. About "On certain incomplete statistics" by Lima et al. 2003. hal-00000825v1

**HAL Id: hal-00000825**

**<https://hal.science/hal-00000825v1>**

Preprint submitted on 12 Nov 2003 (v1), last revised 7 May 2004 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# About “On certain incomplete statistics” by Lima et al

M. Pezeril,

*Laboratoire de Physique de l'état Condensé,  
Université du Maine, 72000 Le Mans, France*

A. Le Méhauté, and Q.A. Wang

*Institut Supérieur des Matériaux et Mécaniques Avancés,  
44, Avenue F.A. Bartholdi, 72000 Le Mans, France*

## Abstract

Lima et al recently claim that (*Chaos, Solitons & Fractals*, 2004;19:1005) the entropy for the incomplete statistics based on the peculiar normalization  $\sum_i p_i^q = 1$  should be  $S = -\sum_i p_i^{2q-1} \ln_q p_i$  instead of  $S = -\sum_i p_i^q \ln_q p_i$  initially proposed by Wang. We indicate here that this conclusion is a result of erroneous use of temperature definition for the incomplete statistics.

PACS : 05.20.-y, 05.70.-a, 02.50.-r

## 1 Introduction

In a recent work[1] addressing the incomplete statistics (IS) based upon the normalization  $\sum_i p_i^q = 1$  proposed by Wang[2, 3, 4], Lima et al calculated the entropy for IS with a method based on the usual thermodynamic relations preserved within the nonextensive statistical mechanics (NSM)[5, 6, 7] derived from Tsallis entropy  $S = -\sum_i p_i^q \ln_q p_i$  (let Boltzmann constant  $k$  be equal to unity) where  $\ln_q p_i = \frac{p_i^{1-q} - 1}{1-q}$  is the so called generalized logarithm and where  $q$  is the nonadditive index. They ended with an entropy  $S_{IS} = -\sum_i p_i^{2q-1} \ln_q p_i$  and concluded that  $S$  was intrinsically connected to the complete distribution normalization  $\sum_i p_i = 1$ .

Their calculation is based on the following relationships. 1) The first law :  $dU = TdS - PdV$  or  $dF = -SdT - PdV$ ,  $F = U - TS = -T \ln_q Z$ , and  $S = -\left(\frac{\partial F}{\partial T}\right)_V$  where  $T = 1/\beta = \left(\frac{\partial U}{\partial S}\right)_V$  is the temperature and  $Z$  is the partition function of IS. The calculations of [1] are correct as long as the above relationships hold, as indicated by Lima et al. By this comment, we would like to indicate that these conventional thermodynamic relationships are formally preserved within IS but with a deformed entropy and a physical or generalized temperature  $T_p$  different from the  $T$  given above. So the conclusion of [1] should be modified.

Due to the fact that there are several versions for NSM proposed from different statistics or information considerations, thermodynamic functions do not in general have the same nonadditive nature in different versions of NSM. This has led to different definitions of physical temperature  $\beta_p$  which is sometimes equal to  $\beta$ [6, 8], sometimes equal to  $\beta$  multiplied by a function of the partition function  $Z^{q-1}$  or  $Z^{1-q}$ [3]. Within IS using the energy expectation  $U = \sum_i p_i^q E_i$  in respecting its nonadditivity[9], the maximum of  $S$  leads to the distribution function  $p_i = \frac{1}{Z} [1 - (1 - q)\beta_p E_i]^{\frac{1}{1-q}}$  with the partition function  $Z^q = \sum_i [1 - (1 - q)\beta_p E_i]^{\frac{1}{1-q}}$  where  $\beta_p$  is given by[3] :

$$\beta_p = Z^{1-q} \frac{\partial S}{\partial U} = Z^{1-q} \beta. \quad (1)$$

On the other hand, the introduction of the distribution  $p_i$  into Tsallis entropy gives

$$S = \frac{Z^{q-1} - 1}{q - 1} + \beta_p Z^{q-1} U \quad (2)$$

or  $S_p = Z^{1-q} S = \ln_q Z + \beta_p U$  where  $S_p$  is a ‘‘deformed entropy’’ introduced in [3] to write heat as  $dQ = T_p dS_p$  and first law as  $dU = T_p dS_p + dW$  or as  $dF = -S_p dT_p + dW$  where

$$F = U - T_p S_p = -T_p \ln_q Z \quad (3)$$

is the Helmholtz free energy and  $dW$  is the work done to the system. We have here

$$S_p = -\left(\frac{\partial F}{\partial T_p}\right). \quad (4)$$

Thanks to the work of [1], this useful ‘‘entropy’’ whose probability dependence

was unknown has been calculated and given by :

$$S_p = - \sum_i p_i^{2q-1} \ln_q p_i = - \sum_i p_i^q \frac{p_i^{q-1} - 1}{q-1}. \quad (5)$$

which is not the original entropy  $S$  of IS[2].  $S_p$  is concave only for  $q > 1/2$  so that not to be maximized to get distribution functions for NSM although its maximum formally leads to  $p'_i \propto [1 - (q-1)\beta_p E_i]^{\frac{1}{q-1}}$ . Notice that this latter is not the original distribution function  $p_i$  of IS. It should also be noticed that the method for calculating  $S_p$  cannot be used for calculating  $S$  by using  $\beta$  or  $T$  within this version of IS because  $S \neq -\left(\frac{\partial F}{\partial T}\right)$  although we can formally write  $F = U - T_p S_p = U - TS$ . In addition,  $Z$  is not derivable with respect to  $\beta$  since it is a self-referential function when written as a function of  $\beta$ .

$S$  can be calculated in this way only when  $\beta_p = \beta$ , which is possible within NSM only when “unnormalized expectation” is used[8]. For IS, if one use the unnormalized expectation  $U = \sum_i p_i E_i$  (or normalized by  $U = \frac{\sum_i p_i E_i}{\sum_i p_i}$ ), the maximum of  $S$  will lead to the distribution function  $p_i = \frac{1}{Z} [1 - (q-1)\beta_p E_i]^{\frac{1}{q-1}}$  with the partition function  $Z^q = \sum_i [1 - (q-1)\beta_p E_i]^{\frac{1}{q-1}}$ . It is easy to prove  $\sum_i p_i = Z^{q-1} + (q-1)\beta_p U$  leading to

$$S = \frac{Z^{q-1} - 1}{q-1} + \beta_p U \quad (6)$$

which implies  $\beta_p = \frac{\partial S}{\partial U} = \beta$ . Now we can naturally write  $dQ = TdS$  and  $dU = TdS + dW$  or  $dF = -SdT + dW$  with  $F = U - TS = -T \ln'_q Z$  where  $\ln'_q x = \frac{x^{q-1}-1}{q-1}$  is another generalized logarithm. It is easy to prove that  $S = -\left(\frac{\partial F}{\partial T}\right)$  with the normalization  $\sum_i p_i^q = 1$  leads to Tsallis entropy.

Let us give here a summary of the definitions of temperature for different versions of NSM found in the literature. We have  $\beta_p = Z^{q-1}\beta = Z^{q-1}\frac{\partial S}{\partial U}$  for the normalized expectations  $U = \sum_i p_i E_i$  or  $U = \sum_i p_i^q E_i / \sum_i p_i^q$  with  $\sum_i p_i = 1$ ;  $\beta_p = Z^{1-q}\beta = Z^{1-q}\frac{\partial S}{\partial U}$  for the normalized expectations  $U = \sum_i p_i^q E_i$  with  $\sum_i p_i^q = 1$ ; and  $\beta_p = \beta = \frac{\partial S}{\partial U}$  if and only if unnormalized expectation  $U = \sum_i p_i^q E_i$  with  $\sum_i p_i = 1$  or  $U = \sum_i p_i E_i$  with  $\sum_i p_i^q = 1$  is used. The conclusion of this comment is that  $S = -\left(\frac{\partial F}{\partial T}\right)$  is valid only for  $\beta_p = \beta$  and that Tsallis entropy can be derived with this formula for both  $\sum_i p_i = 1$  and  $\sum_i p_i^q = 1$ .

## Acknowledgments

We would like to thank J.A.S. Lima for useful discussions and for sending us important references.

## References

- [1] J.A.S. Lima, J.R. Bezerra and R. Silva, *Chaos, Solitons & Fractals*, 19;2004:1005
- [2] Q.A. Wang, *Chaos, Solitons & Fractals*, 12;2001:1431
- [3] Q.A. Wang, *Euro. Phys. J. B*, 26;2002:357
- [4] Q.A. Wang, *Chaos, Solitons & Fractals*, 19;2004:220
- [5] C. Tsallis, *J. Stat. Phys.*, 52;1988:479
- [6] E.M.F. Curado, and C. Tsallis, *J. Phys. A: Math. Gen.*, 24;1991:L69
- [7] F. Pennini, A.R. Plastino and A. Plastino, *Physica A*, 258;1998:446  
C. Tsallis, R.S. Mendes and A.R. Plastino, *Physica A*, 261;1999:534
- [8] Q.A. Wang and A. Le Méhauté, *Chaos, Solitons & Fractals*, 15;2003:537
- [9] Q.A. Wang, L. Nivanen, A. Le Méhauté and M. Pezeril, *J. Phys. A*, 35;2002:7003