



**HAL**  
open science

## Quantum calculations of Coulomb reorientation for sub-barrier fusion

Cédric Simenel, Philippe Chomaz, Gilles de France

► **To cite this version:**

Cédric Simenel, Philippe Chomaz, Gilles de France. Quantum calculations of Coulomb reorientation for sub-barrier fusion. 2004. hal-00001050v2

**HAL Id: hal-00001050**

**<https://hal.science/hal-00001050v2>**

Preprint submitted on 5 May 2004 (v2), last revised 27 Jul 2004 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Quantum calculations of Coulomb reorientation for sub-barrier fusion

C. Simenel <sup>a,b)</sup>, Ph. Chomaz <sup>a)</sup> and G. de France <sup>a)</sup>

<sup>a)</sup> GANIL, BP 55027, F-14076 Caen Cedex 5, France

<sup>b)</sup> DSM/DAPNIA CEA SACLAY, F-91191 Gif-sur-Yvette, France

(May 5, 2004)

Classical mechanics and Time Dependent Hartree-Fock (TDHF) calculations of heavy ions collisions are performed to study the rotation of a deformed nucleus in the Coulomb field of its partner. This reorientation is shown to be *independent* on charges and relative energy of the partners. It only depends upon the deformations and inertias. TDHF calculations predict an increase by 30% of the induced rotation due to quantum effects while the nuclear contribution seems negligible. This reorientation modifies strongly the fusion cross-section around the barrier for light deformed nuclei on heavy collision partners. For such nuclei a hindrance of the sub-barrier fusion is predicted.

Tunneling, the slow "quantum leak" through a classical barrier, is an intriguing phenomenon in nature. In 1928, Gamow discovered this effect looking for an explanation of the alpha radioactivity [1]. However, the tunneling of complex systems remains to be understood. As in the Gamow times nuclear physics is providing one of the most challenging field to understand the tunneling phenomenon. In particular, fusion reaction cross sections involving massive nuclei at, or below, the Coulomb barrier are, in some cases, orders of magnitude over expectations from one dimensional quantum tunneling predictions. Couplings between the internal degrees of freedom and the relative motion deeply modifies the tunneling phenomenon [2]. Neutron transfer, excitation of low-lying vibrational and rotational states, neck formation, zero-point motion and polarization of collective surface vibration as well as static deformation have been identified as key inputs in the understanding of this sub-barrier fusion enhancement [3].

As pointed out in [4,5], for nuclei with a significant static quadrupole deformation, the main effects are i) on the barrier height (geometrical effect) since the barrier is lower in the elongated direction and ii) on the reorientation of the deformed nucleus (rotational effect) under the torque produced by the long-range Coulomb force.

In [6–12], fusion excitation functions were measured for  $^{16}\text{O}$  (spherical) +  $^{144-154}\text{Sm}$  reactions around the Coulomb barrier.  $^{144}\text{Sm}$  is spherical whereas  $^{154}\text{Sm}$  is prolate ( $\beta_2 \approx 0.3$ ). The data were interpreted as arising from the different orientations of the deformed collision partner. An enhancement of the fusion probability is observed when the deformation axis of the prolate nucleus is parallel to the collision axis ("parallel collision") and a hindrance when the two axis are nearly perpendicular ("perpendicular collision"). In these studies, however, the assumption of an *isotropic orientation distribution* of the deformed nucleus at contact was made. In an other hand, authors of ref. [13] recently argued that the Coulomb force systematically rotates the de-

formed nucleus toward the stable perpendicular configuration. This contradicts the classical calculations reported in refs. [14,15] which show partial reorientation.

From the quantum mechanics point of view, the reorientation is a consequence of the excitation of rotational states. Computational techniques have been developed in the past to solve coupled channel equations for multiple Coulomb excitation [16–20] but a good understanding of the Coulomb reorientation dynamics during the approach phase is still required.

In this work we obtain a deeper insight in the Coulomb reorientation revisiting the classical result, solving analytically and numerically the equations of motion of a rigid body, and performing novel quantum approaches describing the deformed projectile within the time dependent Hartree-Fock (TDHF) approach. Then we use these reorientation results in a calculation of the fusion cross-sections to discuss the induced effects.

Assuming first a classical treatment of nuclear orientation (the various orientations do not interfere) and ignoring any reorientation effects, the fusion cross section is given by the average orientation formula (a.o.f.) [21,22]

$$\sigma_{fus.} \approx \int_{\varphi=0}^{\frac{\pi}{2}} \sigma(\varphi) \sin \varphi d\varphi \quad (1)$$

where  $\varphi$  is the angle between the deformation and the collision axis and  $\sigma(\varphi)$  is the associated cross section. However, the Coulomb force induces a torque which, integrated over the whole history, up to the distance of closest approach  $D_0$ , rotates the initial angle  $\varphi_\infty$  into  $\varphi_0$ . Because of reorientation  $\Delta\varphi = \varphi_0 - \varphi_\infty \neq 0$ , the distribution of  $\varphi_0$  loses its isotropy and the  $\sin \varphi$  term in Eq. 1 has to be modified.

To estimate  $\Delta\varphi$ , we first consider the classical motion of a deformed rigid projectile in the Coulomb field of the target. We assume that the projectile of mass  $A_p$  presents a sharp surface at a radius  $R(\theta) = R_0 \sqrt{\alpha^{-4} \cos^2 \theta + \alpha^2 \sin^2 \theta}$  where  $R_0 = r_0 A_p^{\frac{1}{3}}$ ,  $r_0 = 1.2$  fm,  $\alpha = 1 - \varepsilon$  and  $\varepsilon = \sqrt{5/16\pi\beta_2}$ , the deformation parameter.




FIG. 1. Classical calculations (started at  $D_\infty = 241$  fm) of the reorientation  $\Delta\varphi$  of the projectile as function of its initial orientation  $\varphi_\infty$  for the reactions  $^{24}\text{Mg}+^{208}\text{Pb}$  (solid line) and  $^{154}\text{Sm}+^{16}\text{O}$  (dashed line) at the barrier.

Fig. 1 shows the evolution of the reorientation as function of the initial orientation for central collisions at the barrier  $B = Z_p Z_t e^2 / r_0 (A_t^{1/3} + A_p^{1/3})$  where  $Z_p$  ( $A_p$ ) and  $Z_t$  ( $A_t$ ) are the projectile and target number of protons (nucleons). The figure presents two typical asymmetric reactions of a prolate projectile on a spherical target:  $^{24}\text{Mg}(\beta_2 \approx 0.4)+^{208}\text{Pb}$  and  $^{154}\text{Sm}(\beta_2 \approx 0.3)+^{16}\text{O}$ . For symmetry reasons  $\Delta\varphi = 0^\circ$  for  $\varphi_\infty = 0^\circ$  and  $90^\circ$ . The maximal reorientation  $\Delta\varphi_{max}$  occurs around  $45^\circ$ . For the heavy deformed projectile,  $^{154}\text{Sm}$ ,  $\Delta\varphi_{max}$  is less than  $2^\circ$  whereas for  $^{24}\text{Mg}$  it is large ( $\sim 23^\circ$ ).

To understand this difference and get a deeper insight into the phenomenon, an analytical expression for the maximum reorientation can be derived following the approximations made in ref. [23] i.e. assuming that  $\varepsilon$  and  $\Delta\varphi$  are small. Computing the torque leads to following the equation of motion for  $\varphi(t)$

$$\ddot{\varphi}(t) \approx \frac{9Z_p Z_t e^2 \varepsilon}{2mA_p D(t)^3} \sin(2\varphi(t)) \quad (2)$$

where  $m$  is the nucleon mass and  $D(t)$  the distance between the two nuclei. Replacing the time variable by  $\xi(t) = D(t)/D_0$  and neglecting deformation and rotation on the dynamics of  $D(t)$ , Eq. 2 becomes

$$\frac{\partial\varphi(\xi)}{\partial\xi} + 2\xi(\xi - 1) \frac{\partial^2\varphi(\xi)}{\partial\xi^2} = \frac{9\varepsilon}{2\xi} \frac{A_t}{A_p + A_t} \sin(2\varphi_\infty) \quad (3)$$

where  $\sin(2\varphi(t))$  have been replaced by  $\sin(2\varphi_\infty)$  treating the reorientation perturbatively (see [23]). Only the factor  $A_t/(A_p + A_t)$  remains since the initial center of mass energy  $E$  and the charges have been taken into account in  $D_0 = e^2 Z_p Z_t / E$ . The solution of Eq. 3 is

$$\varphi(\xi) = \varphi_\infty + \frac{3\varepsilon A_t}{A_p + A_t} \sin(2\varphi_\infty) (\xi(2 - \zeta) - \ln\zeta + \delta) \quad (4)$$

where  $\zeta = 1 + \sqrt{1 - \xi^{-1}}$  and  $\delta = \ln 2 - 1/2$ . Solved up to the distance of closest approach ( $\xi = 1$ ), it leads to

$$\Delta\varphi = 3\varepsilon \frac{A_t}{A_p + A_t} \sin(2\varphi_\infty) \left( \frac{1}{2} + \ln 2 \right). \quad (5)$$

It can be shown by performing the time integral introduced in ref. [24] that Eq. 5 is equivalent to the one reported in ref. [24] except that with this formulation it is clear that the reorientation depends neither on projectile and target charges nor on the initial energy and excitation energy of the first rotational state but only on the deformation and the mass ratio. This counter-intuitive result, which has been exhaustively checked numerically, can be understood: the increase of the Coulomb interaction with charges (like  $Z_p Z_t$ ) is compensated by an increase of  $D_0$ . A similar balance occurs with incident energy. An increase in  $E$  reduces  $D_0$  and the time to interact leading to a zero net effect on the integrated reorientation. The strong difference between the two systems shown in Fig. 1 is thus only due to the difference in  $A_t/(A_p + A_t)$ , and not to the difference in the Coulomb forces.

Fig. 2 shows the evolution of the orientation as a function of  $D$  for the central reaction  $^{24}\text{Mg}+^{208}\text{Pb}$  at the barrier. Results from the numerical solution of the classical dynamic of a deformed rigid body (dotted line) and approximated analytical expression Eq. 4 (dashed line) are very close. The small difference observed at the turning point can be attributed to the higher orders terms in  $\varepsilon$  not taken into account in Eq. 4. The difference at large distance is due to the fact that the numerical simulation only starts at a finite value of the initial distance  $D_\infty$  (here  $D_\infty = 241$  fm) while the analytical result integrates the effects from  $D_\infty \rightarrow \infty$ .

To take into account the quantal nature of the nuclei and to avoid the rigid body approximation, we have performed TDHF calculations [25–29] of this nuclear reaction. TDHF theory is optimized for the prediction of the average values of one body observables and so the deformation and the orientation should be well estimated by TDHF. In the TDHF approach, the evolution of the one-body density matrix  $\rho = \sum_{n=1}^N |\varphi_n\rangle \langle \varphi_n|$  is determined by a Liouville equation,  $i\hbar \partial_t \rho = [h(\rho), \rho]$  where  $h(\rho)$  is the mean-field Hamiltonian. We have used the code built by P. Bonche and coworkers [30] with an effective Skyrme mean-field [31]. Two different parametrizations were used,  $\text{SkM}^*$  [32] and  $\text{SLy4}$  [33], in order to control that the conclusions are almost independent on the force. Because of the long range nature of the Coulomb interaction, the calculation must be started much before the turning point typically for an initial distance around 200 fm. However, for reactions below the barrier we can separate the dynamics of the target from the one of the projectile. Therefore, we have modify the TDHF code in order to compute the evolution of the nuclei separately

in their center of mass frame. We assume that the centers of mass follow Rutherford trajectories and we add the Coulomb field of the partner.




FIG. 2. For the central collision of  $^{24}\text{Mg}$  on  $^{208}\text{Pb}$  at the barrier the orientation  $\varphi$  of the  $^{24}\text{Mg}$  as function of the relative distance  $D$  predicted by TDHF with the SkM\* force (solid line) or by classical simulation (dotted line) and its analytic approximation (dashed line).

As an example we consider the central collision of a  $^{24}\text{Mg}$  projectile on a  $^{208}\text{Pb}$  target at the Coulomb barrier with the initial conditions ( $\varphi_\infty = 45^\circ$ ;  $D_\infty = 241$  fm). Fig. 2 shows that the evolutions of  $\varphi$  as function of  $D$  for classical and TDHF calculations have the same behavior. The maximum reorientation predicted by this TDHF calculation is  $\Delta\varphi = 28.2^\circ$  (resp.  $33.6^\circ$ ), with the SkM\* (resp. SLy4) force. Both Skyrme parametrizations give the same order of magnitude ( $\sim 30^\circ$ ), however the classical expectation was  $\Delta\varphi \sim 23^\circ$ . This 30%–difference in the reorientation, which is independent of the initial orientation as we have numerically checked, indicates a smaller moment of inertia in TDHF as compared to the rigid-body classical approximation. This reduced inertia can be attributed to a spherical core in the N-body wavefunction of the  $^{24}\text{Mg}$  which does not participate to the rotation.

Experimentally, the question is: what is the effect of the reorientation on fusion cross-sections? We have used the code CCDEF [34] to estimate the fusion cross-section  $\sigma_{fus}(E)$  for the reaction  $^{24}\text{Mg}+^{208}\text{Pb}$ . CCDEF takes into account the shape of the nucleus on the basis of the a.o.f.. We then go beyond this assumption by including in CCDEF the reorientation obtained with TDHF for central collisions. A commonly used way to present this excitation function is to compute the so-called bar-

rier distribution  $B(E) = \partial_{E^2}^2 (\sigma_{fus}(E) \cdot E)$  [35]. Fig. 3 shows barrier distributions extracted from CCDEF without shape effect (i.e. the 1D barrier, solid line). The width of the peak results from quantum tunneling. A prolate deformation  $\beta_2 = 0.4$  of  $^{24}\text{Mg}$  with an isotropic distribution of orientation (dashed line) flattens considerably the barrier distribution with a prominent part on the high energy tail. A low energy shoulder extending down to  $\sim 5$  MeV below the 1D barrier maximum is responsible for sub-barrier fusion enhancement (as compared to the single barrier case). Classically, the low energy part of the barrier distribution can be interpreted as coming from "parallel collisions" whereas the high energy part comes from "perpendicular collisions". The high energy component dominates because a prolate nucleus has one elongated direction (low barrier) for two short axis (high barriers).




FIG. 3. Barrier distribution for the reaction  $^{24}\text{Mg}+^{208}\text{Pb}$  i) assuming spherical nuclei (solid line); ii) considering a prolate deformed  $^{24}\text{Mg}$  ( $\beta_2 = 0.4$ ) and the a.o.f. (dashed line); iii) and including reorientation with (dotted-dashed line) and without the rotational energy (dotted line).

The barrier distributions including the reorientation predicted by TDHF are plotted in Fig. 3 both neglecting (dotted line) and taking into account the rotational energy (dotted-dashed line) which is a second order correction in  $\epsilon$  and then increases only slightly the barrier. Compared to the a.o.f. prediction the TDHF results exhibit reduced low energy shoulder and increased high energy peak. This arises from that the Coulomb reorientation increases  $\varphi$  and thus increases the barrier height. Consequently, the sub-barrier fusion enhancement observed for reactions involving a light nucleus on a deformed heavy ion like  $^{154}\text{Sm}$  is expected to be reduced when the deformed nucleus is light and its collision partner heavy. Experimentally, the effect of the reorientation should then be studied by comparing the excitation functions of reactions with deformed projectiles such as  $^{24}\text{Mg}$  on different targets. To simplify the understanding of the reaction doubly-magic spherical targets such as  $^{16}\text{O}$ ,

$^{40}\text{Ca}$  and  $^{208}\text{Pb}$  might be first tested. Eq. 5 shows that the reorientation should increase with the mass of the target reducing the sub-barrier fusion cross section.

Finally one may worry about the effect of the nuclear interaction at short distance. To evaluate its importance on the reorientation we have simulated the fusion of two nuclei in TDHF. Since TDHF does not allow directly tunneling, we have studied the modulation of the threshold energy  $B$  of fusion reaction as a function of the initial orientation  $\varphi$ :  $\Delta B(\varphi) = B(\varphi) - B(0)$ . As a deformed nucleus we have considered again  $^{24}\text{Mg}$ . To focus on the nuclear contribution we have chosen a light spherical target, the  $^{16}\text{O}$  and we have started the reaction at short distance. The observed variation of the fusion threshold energy appears to follow, within the numerical error due to the considered energy step, the expected  $\sin^2 \varphi$  modulation expected for a quadrupole deformed projectile in absence of reorientation. Indeed, we get  $\Delta B(45^\circ)/\Delta B(90^\circ) = 0.45(5)$  i.e. an almost negligible deviation from the  $\sin^2$  law which would predict  $\Delta B(45^\circ)/\Delta B(90^\circ) = 1/2$ . Inverting the problem in order to extract the reorientation due to the observe small reduction of the barrier leads to  $\Delta\varphi(45^\circ) \sim -3^\circ$ . Considering the short range nature of the nuclear force this effect is expected to not vary much with the target. Compared with the expected magnitude of the Coulomb effect this nuclear reorientation appears to be negligible. This can be related to the range of the forces. Indeed, the Coulomb interaction is a long range force so the induced torque has time to rotate the nucleus and to produce a large reorientation which is proportional to an average angular velocity times the average time it is rotating. Conversely, the nuclear field acts over a very short time so that even if it contributes to the excitation of rotational states, the nucleus has hardly enough time to actually rotate.

To summarize, we have studied the reorientation effect of a deformed projectile on a spherical target. Our analytical results for the classical dynamics of a rigid body, confirmed by exact classical simulations, show that, in contrast to a naive expectation, the Coulomb reorientation depends neither on charges nor on relative energy. The relevant observables are the deformation parameter and the mass ratio. The reorientation is then expected to be maximum when the deformed nucleus is light and its collision partner heavy. Those conclusions have been extended to quantum dynamics using TDHF. These calculations show that the nuclear contribution are negligible and they exhibit a sizeable increase of the reorientation as compared to classical calculations, interpreted in terms of a smaller moment of inertia for the quantum system as compared to the rigid body approximation. Calculations of barrier distributions show that sub-barrier fusion is partially hindered by the reorientation process. We also suggest experiments to measure the effect of the reorientation on excitation functions.

J.P. Wieleczko, N. Rowley, V. Yu Denisov and E. Pol-

lacco are thanked for fruitful discussions, M. Faure for his programming work and P. Bonche for providing his TDHF code.

- 
- [1] G. Gamow, *Z. Phys.* **51**, 204 (1928); **52**, 510 (1928).
  - [2] C.H. Dasso, S. Landowne and A. Winther, *Nucl. Phys.* **A405**, 381 (1983).
  - [3] M. Dasgupta *et al.*, *Annu. Rev. Nucl. Part. Sci.* **48**, 401 (1998).
  - [4] P.W. Riesenfeldt and T.D. Thomas, *Phys. Rev.* **C2**, 711 (1970).
  - [5] A.S. Jensen and C.Y. Wong, *Phys. Rev.* **C1**, 1321 (1970).
  - [6] R.G. Stokstad *et al.*, *Phys. Rev. Lett.* **41**, 465 (1978).
  - [7] R.G. Stokstad *et al.*, *Phys. Rev.* **C21**, 2427 (1980).
  - [8] D.E. DiGregorio *et al.*, *Phys. Lett.* **B176**, 322 (1986).
  - [9] D.E. DiGregorio *et al.*, *Phys. Rev.* **C39**, 516 (1989).
  - [10] J.X. Wei *et al.*, *Phys. Rev. Lett.* **67**, 3368 (1991).
  - [11] J.R. Leigh *et al.*, *Phys. Rev.* **C47**, R437 (1993).
  - [12] C.R. Morton *et al.*, *Phys. Rev. Lett.* **72**, 4074 (1994).
  - [13] V. Yu Denisov and W. Nörenberg, *Eur. Phys. J.* **A15**, 375 (2002).
  - [14] H. Holm *et al.*, *Phys. Lett.* **B29**, 473 (1969).
  - [15] L. Wilets *et al.*, *Phys. Rev.* **156**, 1349 (1967).
  - [16] M. Rhoades-Brown, M.H. Macfarlane and S.C. Pieper, *Phys. Rev.* **C21**, 2417 (1980).
  - [17] L.D. Tolsma *Phys. Rev.* **C20**, 592 (1979).
  - [18] F. Rösler *et al.*, *Comp. Phys. Com.* **8**, 35 (1974).
  - [19] G.H. Rawitscher and C.H. Rasmussen, *Comp. Phys. Com.* **11**, 183 (1976).
  - [20] L.D. Tolsma, *Phys. Rev.* **C35**, 177 (1987).
  - [21] T. Rumin, K. Hagino and N. Takigawa, *Phys. Rev.* **C63**, 044603 (2001).
  - [22] K. Hagino *et al.*, *Phys. Rev.* **C51**, 3190 (1995).
  - [23] K. Alder and A. Winther, *electromagnetic excitation* (north-holland/american elsevier, 1978), Eq. VIII.6.4.
  - [24] R.A. Broglia and A. Winther, *Heavy-Ion Reactions* (Addison-Wesley, New York, 1991), Eq. II.7.7.
  - [25] D.R. Hartree, *Proc. Camb. Phil. Soc.* **24**, 89 (1928).
  - [26] V.A. Fock, *Z. Phys.* **61**, 126 (1930).
  - [27] D. Vautherin and D.M. Brink, *Phys. Rev.* **C5**, 626 (1972).
  - [28] P. Bonche, S. Koonin and J.W. Negele, *Phys. Rev.* **C13**, 1226 (1976).
  - [29] J.W. Negele, *Rev. Mod. Phys.* **54**, 913 (1982).
  - [30] K.-H. Kim *et al.*, *J. Phys.* **G23**, 1267 (1997).
  - [31] T. Skyrme, *Phil. Mag.* **1**, 1043 (1956).
  - [32] J. Bartel *et al.*, *Nucl. Phys.* **A386**, 79 (1982).
  - [33] E. Chabanat *et al.*, *Nucl. Phys.* **A635**, 231 (1998).
  - [34] J. Fernandez-Niello, C.H. Dasso and S. Landowne, *Comp. Phys. Com.* **54**, 409 (1989).
  - [35] N. Rowley *et al.*, *Phys. Lett.* **B254**, 25 (1991).