



HAL
open science

Nuclear liquid-gas phase transition and supernovae evolution

Jerome Margueron, Jesus Navarro, Patrick Blottiau

► **To cite this version:**

Jerome Margueron, Jesus Navarro, Patrick Blottiau. Nuclear liquid-gas phase transition and supernovae evolution. 2004. hal-00001067v1

HAL Id: hal-00001067

<https://hal.science/hal-00001067v1>

Preprint submitted on 26 Jan 2004 (v1), last revised 11 Mar 2004 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nuclear liquid-gas phase transition and supernovae evolution

Jérôme Margueron ^{a,*},¹ Jesús Navarro ^b Patrick Blottiau ^c

^aGANIL CEA/DSM - CNRS/IN2P3 BP 5027 F-14076 Caen Cedex 5, France

^bIFIC (CSIC - Universidad de Valencia) Apdo. 22085, E-46.071-Valencia, Spain

^cCEA/DIF DPTA BP12 91680 Bruyères-le-Châtel Cedex, France

Abstract

It is shown that the large density fluctuations appearing at the onset of the nuclear liquid-gas phase transition can play an important role in the supernovae evolution. Due to these fluctuations, the neutrino gas may be trapped inside a thin layer of matter near the proto-neutron star surface. The resulting increase of pressure may induce strong particle ejection a few hundred milliseconds after the bounce of the collapse, contributing to the revival of the shock wave. The Hartree-Fock+RPA scheme, with a finite-range nucleon-nucleon effective interaction, is employed to estimate the effects of the neutrino trapping due to the strong density fluctuations, and to discuss qualitatively the consequences of the suggested new scenario.

Key words: neutrinos, supernovae, phase transition, nuclear matter

PACS: 97.60.Bw, 26.50.+x, 25.30.Pt, 21.60.Jz

The first simulations of Colgate and White [1] and Arnett [2] have settled the general scenario of explosive supernovae, which schematically goes as follows. Stars with more than about ten times the Sun mass develop central iron cores which eventually become unstable and collapse to neutron stars, due to electron capture and photodisintegration onto iron-group nuclei. The interior of neutron stars is denser than nuclear matter and initially extremely hot. Above nuclear density, the equation of state becomes stiff enough to produce a bounce of the core, and a shock wave is formed, moving to the infalling outer core and envelope. Particle reactions at such conditions create a huge number of

* Corresponding author.

Email address: margueron@tphys.physik.uni-tuebingen.de (Jérôme Margueron).

¹ Present address: Institut für Theoretische Physik, Universität Tübingen, D-72076 Tübingen, Germany

neutrinos, which eventually escape from the dense neutron star, transferring energy to the matter in the still infalling outer layers of the star. This neutrino heating is believed to cause the violent disruption of the star in a supernova explosion. However, from the very beginning it was found that the resulting shock wave is not energetic enough. Recently, an enormous effort has been done to describe the neutrino production and interactions in great detail [3], including the effects of stellar rotation and of violent anisotropic plasma motions. Convective processes in the supernova core had been recognized to accelerate the energy transport inside the neutron star and to enhance the deposition of energy by neutrinos in the outer stellar layers, thus supporting the explosion of the star (the "delayed" mechanism). Along some decades, numerical simulations were performed, starting from the previous 1D codes [4,5] and the flux-limited neutrino transport scheme [6] to reach the multi-dimensional calculations, showing the importance of convection and Rayleigh-Taylor instabilities, and the most refined neutrino transport methods [7,8]. Nevertheless, the outcome of these worldwide most elaborate supernova simulations is disappointing, as no explosions could be obtained. This negative result shatters the widely accepted view of how the explosion starts.

In this letter we point out that large density fluctuations associated to the nuclear liquid-gas phase transition, can play an important role in the neutrino trapping. Indeed, the scattering of neutrinos is hugely increased at the onset of strong density fluctuations, so that neutrinos should be trapped inside a thin layer of the proto-neutron star (PNS) for densities in the range between about $0.1\rho_0$ and $0.6\rho_0$, where ρ_0 is the nuclear matter density at saturation. The resulting increase in the neutrino pressure may induce strong particle ejection a few hundred milliseconds after the bounce of the collapse, contributing to the revival of the shock wave.

The neutrino-nucleon cross section is obtained from a phase space integration of the nuclear response function. In the non-relativistic limit, the neutrino mean-free path λ is given by the well-known expression [9]

$$1/\lambda(k_\nu, T) = \frac{G_F^2}{16\pi^2} \int d\mathbf{k}_3 \left(c_V^2 (1 + \cos\theta) \mathcal{S}^{(0)}(q, T) + c_A^2 (3 - \cos\theta) \mathcal{S}^{(1)}(q, T) \right), \quad (1)$$

where k_ν is the neutrino energy-momentum, T is the temperature, G_F is the Fermi constant, c_V (c_A) the vector (axial) coupling constant, \mathbf{k}_3 the final neutrino momenta, $q = k_\nu - k_3$ the transferred energy-momentum, and $\cos\theta = \hat{\mathbf{k}}_\nu \cdot \hat{\mathbf{k}}_3$. The dynamical structure factors $\mathcal{S}^{(S)}(q, T)$ describe the response of nuclear matter to excitations induced by neutrinos, and they contain the relevant information on the medium. The vector (axial) part of the neutral




Fig. 1. The adimensional Landau parameters F_0 are plotted as a function of the density at several values of temperature. The left pannel corresponds to symmetric nuclear matter, and the right pannel to an asymmetric parameter $x_p = 0.3$. In the asymmetric case, the definition $F_0 = (F_0^{nn} + F_0^{pp} + 2F_0^{np})/2$ has been taken and the coefficients $F_0^{\tau\tau'}$ are the second derivative of the free energy density. In both cases, the onset for the spinodal instability is given by $F_0 = -1$.

current gives rise to density (spin-density) fluctuations, corresponding to the $S = 0$ ($S = 1$) spin channel. Density and spin-density responses contribute to the scattering of neutrinos, while the isospin-flip response contribute to neutrino absorption process, which is known to dominate the total cross section [10]. In the following, we shall show that large isoscalar fluctuations can give the dominant contribution at subnuclear densities.

Clearly, an enhancement of the response function in either channel will result in a quenching of the neutrino mean-free path. It has been shown [11] that at densities higher than ρ_0 a ferromagnetic instability can appear, leading consequently to a zero neutrino mean-free path [12]. However, the density at which such an instability could appear is not yet a settled question. For instance, whereas all Skyrme effective interactions predict a ferromagnetic transition for densities less than $\sim 3.5\rho_0$ [13], recent microscopic calculations [14,15] push it to higher densities, in case it exists.

At present, a large nuclear community is currently studying the liquid-gas phase transition which occurs during multi-fragmentation experiments in heavy-ion collisions [16]. Such a transition appears below the saturation density ρ_0 of nuclear matter, at around $\sim 0.6\rho_0$, i.e. $\sim 0.1 \text{ fm}^{-3}$. Heavy nuclei are located on the liquefaction curve for $T=0 \text{ MeV}$ and the critical temperature is located between 15 and 20 MeV, depending on the specific nuclear model used. The onset of liquid-gas coexistence phase is associated to the spinodal instability, related to a negative curvature in the nucleonic free energy density such that the speed of sound vanishes. In the density regime close to the spinodal, the density response function will be enhanced, thus resulting in a reduced




Fig. 2. Pressures (left panel) and particle fractions (right panel) plotted as a function of the density for asymmetric nuclear matter at temperature $T=10$ MeV and leptonic fraction $Y_l=0.285$, calculated with the Gogny D1P effective interaction. The total pressure (solid line) is the sum of the nuclear (dotted line), the electronic (dashed line) and the neutrino (dot-dashed line) pressures. On the right panel, solid, dashed and dot-dashed lines refer to the particle fraction of neutrons, charged particles (i.e. protons and electrons), and neutrinos, respectively.

neutrino mean-free path.

In the following we shall describe the density fluctuations of nuclear matter within the Hartree-Fock (HF) plus Random Phase Approximation (RPA) scheme, using the Gogny D1P finite-range nucleon-nucleon effective interaction [17]. The calculation of the RPA response functions will be performed in the Landau approximation, ignoring the Landau coefficients beyond the dipole one in both density and spin channels [18]. It is worth recalling that some values of Landau parameters are related to different instabilities. In particular, the density at which the dimensionless Landau parameter F_0 equals -1 indicates the onset of the spinodal instability [19]. In Fig. 1, we show F_0 versus the total density for symmetric nuclear matter and asymmetric nuclear matter with $x_p \equiv Z/(N+Z) = 0.3$. The two panels look very similar and we clearly see that out of the spinodal region, F_0 is attractive for a wide range of densities and temperatures.

In Fig. 2 we plot the pressures and particle fractions as a function of the density for asymmetric nuclear matter at a temperature of $T=10$ MeV and leptonic fraction $Y_l=0.285$, defined as $(\rho_e + \rho_\nu)/\rho$. In the low density region $\rho < \rho_0$, the total pressure is mainly supported by electrons, and the contribution of neutrinos is about 20% or less [20]. Hence, an increase of the neutrino pressure by a factor of 5 will increase by a factor of 2 the total pressure.

Let us assume the extreme hypothesis that, after a few hundred milliseconds, all the neutrinos contained inside the star are trapped inside the thin layer




Fig. 3. The density-density response function $\mathcal{S}^{(0)}(q, T)$ is plotted as a function of the transferred energy for two values of the density: (a) ρ_0 and (b) $0.75\rho_0$. The calculations have been done for momentum $q = 10$ MeV, temperature $T = 10$ MeV and leptonic fraction $Y_l=0.285$. The solid and dashed lines stand for the HF and the RPA results respectively.

near the surface, and that the initial distribution of neutrino density is quasi-uniform. Therefore, the increase of the neutrino pressure inside the thin layer is given by the ratio of the PNS radius R over the width ϵ of the thin layer and writes

$$\frac{P_{\text{layer}}}{P_{\text{inside}}} \sim \left(\frac{R}{3\epsilon}\right)^{4/3} \quad (2)$$

where the neutrino pressure is polytropic versus the density with the power $4/3$. Hence, for a typical radius of 10 km and a width of 100 m, the neutrino pressure increases by a factor greater than 200. Of course, the Pauli principle should strongly moderate this factor, but this schematic calculation points out the possible efficiency of the neutrino trapping.

This qualitative calculation assumes that all species are independent from each other. In fact, neutrinos are in β -equilibrium, so that a modification of their number implies a corresponding change of the number of other particles, in particular the electrons. If the neutrino density is increased at constant temperature and constant baryonic density, then the β equilibrium shifts in the direction to raise the number of electrons and protons. This is the consequence of Le Chatelier's principle of minimum constraint extended to quantum systems. The same principle predicts that when the temperature is increased, the system tends to move towards the endothermic direction, thus increasing the number of electrons and protons. Consequently, raising both neutrino density and temperature produces electrons and protons. As electrons are mainly responsible for the pressure at low density, the overall effect of neutrino trapping is the increase of the total pressure.




Fig. 4. The neutrino mean-free path is plotted as a function of the density for asymmetric nuclear matter at $T=10$ MeV and $Y_l=0.285$. Left panel: HF result, right panel: ratio between RPA and HF mean-free paths.

Now we concentrate on the mechanism responsible for neutrino trapping. In the following, we will approach the liquid-gas phase transition from the homogeneous nuclear matter. This can be done by decreasing either the density at constant temperature or the temperature at constant density. In both cases, we should keep the density beyond the spinodal one or the temperature beyond the critical one. The density-density response function $\mathcal{S}^{(0)}(q, T)$ is represented in Fig. 3 for density values approaching the spinodal point, namely ρ_0 (panel a) and $0.75\rho_0$ (panel b). The solid line stands for the HF calculation while the dashed line includes the RPA correlations, with a particle-hole interaction described in terms of the monopolar and dipolar Landau parameters. In the left panel, we see two symmetric collective states inside the continuum strength. These states indicate that near the Fermi surface, the residual interaction is repulsive. It can be seen that when the system approaches the liquid-gas phase transition, a zero energy mode becomes increasingly enhanced. In fact, the peak diverges at the spinodal density. This is a signature of a strong attractive residual interaction, which indeed induces the phase transition.

The neutrino mean-free path as a function of the density is plotted in Fig. 4 for a temperature $T=10$ MeV and a leptonic fraction $Y_l = 0.285$. Left and right panels correspond to HF and RPA results, respectively. One can see that the RPA correlations induce a strong reduction of the neutrino mean-free path inside the liquid-gas phase transition. Hence, neutrinos could be massively trapped inside the thin external layer with densities near the densities of the liquid-gas phase transition, and a radius close to the radius of the proto-neutron star. The spinodal density is close to $0.6\rho_0$ in symmetric nuclear matter and is nearly constant in asymmetric nuclear matter until $\rho_x/\rho \sim 0.1$ ($x = n$ or p) [19].

Of course, the RPA scheme can be questionable in such a large fluctuations

regime. The result of effects as short-range correlations, possible contributions of two or more loop processes at these densities have to be elucidated, as well as the response to neutrino probes of inhomogeneous nuclear matter inside the phase coexistence. Nevertheless, we believe that our approach reflects the basic physics near the spinodal point.

Let us now analyze the physical consequences for the supernovae-II mechanism due to the interplay between neutrino propagation and the liquid-gas phase transition. During the iron core collapse, neutrinos are trapped inside the core. The energy gained by the collapse is essentially carried away by the neutrinos after the bounce. Neutrinos are massively trapped for densities below the saturation one, and hence those neutrinos which are diffusing from the inner part of the PNS cannot reach the outer part of the star. An important fraction of neutrinos accumulates inside a thin layer close to the surface of the PNS. The neutrino Fermi energy increases and shifts the beta equilibrium to the production of protons and electrons. The neutrino and electronic pressures increase as well as the temperature inside the thin layer. We believe that this effect may lead to a partial conversion of the energy carried away by the neutrino gas to the matter through the beta equilibrium. According to the Euler transport equation, a local increase of pressure will produce an ejection of matter. This radiated matter will encounter the shock wave and may contribute to its revival. Indeed, the energy carried away by neutrinos represents about 98% of the collapse energy while the shock wave transports only 2% of it. If the ejection of particles carries away only 2% of neutrino energy, the total kinetic energy of matter will increase by a factor of 2. This is the reason why we believe that this new scenario may contain an efficient mechanism for the explosion.

This scenario may modify the neutrino emission signal from supernovae. Indeed, at the beginning of the collapse (for $\rho \sim 10^{10} \text{g/cm}^3$), an electronic neutrino burst should come from the electronic capture. Then neutrinos are trapped and do not radiate from the PNS. After the bounce, if the proposed mechanism is strong enough to produce a second shock wave, the neutrinos "englued" inside the thin layer would be massively liberated. In this case, a second neutrino burst, mixing the leptonic flavors, would be observed. The observation of two neutrino bursts could be a signature of the second explosion. However, this scenario does not necessarily imply a second shock wave but could produce only a huge ejection of matter. Then, we stress that the non-observation of two neutrino bursts is not in contradiction with this scenario.

In this letter we have explored the consequences that large density fluctuations inside the liquid-gas phase transition could have for neutrino propagation and supernovae-II explosion. As far as we know, this effect has never been studied. The calculations presented here are very schematic and apply only to the homogeneous phase close to the coexistence region, either for densities higher

than the spinodal one, or temperatures higher than the critical one. Two directions should now be explored: inside the homogeneous matter, we should now include more correlations to improve the calculation of the response function approaching the phase transition, and we should also get an estimation of the response of inhomogeneous nuclear matter to neutrino probes. For instance, close to the critical temperature, we expect to observe the coherent scattering of neutrinos in analogy with the well-known critical opalescence phenomenon (with respect to photons) observed in condensed matter. In conclusion, this new scenario should now be included in an hydrodynamical simulation to get the precise time scale as well as to estimate the balance effects of neutrino Pauli blocking and the shift of the beta equilibrium and obviously possible "feedback" effects. Work in this direction is now in progress

Acknowledgments

Interesting discussions with M. Baldo, Ph. Chomaz, H. Mütter, A. Pérez-Cañellas and A. Sedrakian are gratefully acknowledged. J.M. wants also to thank N. Van Giai and late D. Vautherin for introducing him into the subject of neutrino propagation in nuclear matter. J.N. is supported by MCyT/FEDER (Spain), grant number BMF2001-0262, and Generalitat Valenciana, grant number GV01-216.

References

- [1] S.A. Colgate and R.H. White, *Astrophys. J.* **143** (1966) 626.
- [2] W.D. Arnett, *Can. J. Phys.* **44** (1966) 2553.
- [3] R. Buras, M. Rampp, H.-Th. Janka, and K. Kifonidis, *Phys. Rev. Lett.* **90** (2003) 241101.
- [4] H.A. Bethe and J.R. Wilson, *Astrophys. J.* **295** (1985) 14.
- [5] P. Blottiau, Ph. D. thesis, Paris VII University (1989).
- [6] S. Bruenn, *Astrophys. J. Suppl.* **58** (1985) 771.
- [7] P. Mellor, J.P. Chièze and J.L. Basdevant, *Astron. Astrophys.* **197** (1988) 123.
- [8] M. Rampp and H.T. Janka, *Astron. Astrophys.* **396** (2002) 361.
- [9] N. Iwamoto and C.J. Pethick, *Phys. Rev.* **D 25** (1982) 313.
- [10] S. Reddy, M. Prakash and J.M. Lattimer, *Phys. Rev D* **58** (1998) 013009.
- [11] A. Vidaurre, J. Navarro and J. Bernabeu, *Astron. Astrophys.* **135** (1984) 361.
- [12] J. Navarro, E.S. Hernández and D. Vautherin, *Phys. Rev. C* **60** (1999) 045801.

- [13] J. Margueron, J. Navarro and Nguyen Van Giai, Phys. Rev. C **66** (2002) 014303.
- [14] S. Fantoni, A. Sarsa and K.E. Schmidt, Phys. Rev. Lett. **87** (2001) 181101.
- [15] I. Vidaña and I. Bombaci, Phys. Rev. C **66** (2002) 045801.
- [16] Ph. Chomaz et al., Phys. Rep. **389** (2004) 263.
- [17] M. Farine et al., J. Phys .G: Nucl.Part.Phys. **25** (1999) 863.
- [18] J. Margueron, Ph.D. Thesis, Paris XI University.
- [19] J. Margueron and Ph. Chomaz, Phys. Rev. C **67** (2003) 041602(R).
- [20] E. Suraud, Ph.D. thesis, Paris VI University.