



HAL
open science

On Cremona transformations of prime order

Arnaud Beauville

► **To cite this version:**

| Arnaud Beauville. On Cremona transformations of prime order. 2004. hal-00001113v2

HAL Id: hal-00001113

<https://hal.science/hal-00001113v2>

Preprint submitted on 13 Feb 2004 (v2), last revised 21 Sep 2004 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On Cremona transformations of prime order

Arnaud BEAUVILLE

1. The result

The *Cremona group* $\text{Bir}(\mathbb{P}^2)$ is the group of birational transformations of $\mathbb{P}_{\mathbb{C}}^2$, or equivalently the group of \mathbb{C} -automorphisms of the field $\mathbb{C}(x, y)$. There is an extensive classical literature about this group, in particular about its finite subgroups; we refer to the introduction of [dF] for a list of references.

The classification of conjugacy classes of transformations of prime order p has been given a modern treatment in [B-B] for $p = 2$ and in [dF] for $p \geq 3$. Let us recall the main results. The linear transformations of given order are contained in a unique conjugacy class. Apart from these there are 3 continuous families of conjugacy classes of involutions (the famous de Jonquières, Bertini and Geiser involutions), and 2 continuous families of transformations of order 3. Every transformation of prime order ≥ 7 is conjugate to a linear automorphism.

We prove in this note the following result, which completes the classification:

Theorem. — *There is a unique conjugacy class of order 5 in the Cremona group apart from the class containing linear automorphisms.*

Most of the work has been done already in [dF], where it is proved that there can be only one more conjugacy class of order 5, namely that of the transformation $\gamma : (x, y) \mapsto (1 - y, 1 - \frac{y}{x})$. What remains to be proved is that γ is conjugate to a linear automorphism.

Fix a primitive 5-th root of unity ζ . The problem is readily translated into the existence of a rational function $F(x, y)$ satisfying the functional equation

$$F(\zeta x, y)F(\zeta^{-1}x, y) + F(x, y) = 1 \quad (\star)$$

and such that the rational map $(x, y) \mapsto (F(x, y), 1 - F(\zeta x, y))$ is birational. At this point Don Zagier showed to me a very simple way to construct solutions to (\star) ; it turns out that essentially the most economical choice satisfies indeed the second condition and thus leads to a proof of the theorem.

2. The proof

Let us first recall why linear automorphisms of the same (finite) order are conjugate:

Proposition. — *Let n be a positive integer. Two linear automorphisms of order n are conjugate in $\text{Bir}(\mathbb{P}^2)$.*

Proof: Let $T = \mathbb{C}^* \times \mathbb{C}^*$ be the standard maximal torus of PGL_3 . We can view T as a Zariski open subset of \mathbb{P}^2 , so its automorphism group $\mathrm{GL}(2, \mathbb{Z})$ embeds naturally in $\mathrm{Bir}(\mathbb{P}^2)$: an element $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of $\mathrm{GL}(2, \mathbb{Z})$ corresponds to the Cremona transformation $(x, y) \mapsto (x^a y^b, x^c y^d)$. The image of $\mathrm{GL}(2, \mathbb{Z})$ in $\mathrm{Bir}(\mathbb{P}^2)$ normalizes T , and its action on T by conjugation is just the original action of $\mathrm{GL}(2, \mathbb{Z})$ on T .

We want to prove that two elements of order n of T are conjugate under this action. The kernel of the multiplication by n in T is $(\mathbb{Z}/n)^2$; any element of order n in the \mathbb{Z}/n -module $(\mathbb{Z}/n)^2$ is part of a basis, which implies that it is conjugate to $(1, 0)$ under the action of $\mathrm{SL}(2, \mathbb{Z}/n)$. Since the natural map $\mathrm{SL}(2, \mathbb{Z}) \rightarrow \mathrm{SL}(2, \mathbb{Z}/n)$ is surjective, this proves our result. ■

Proof of the theorem

We want to prove that $\gamma = h \circ \ell \circ h^{-1}$, with $h \in \mathrm{Bir}(\mathbb{P}^2)$ and $\ell \in \mathrm{PGL}_3$. By the Proposition we can take for ℓ the map $(x, y) \mapsto (\zeta x, y)$, where ζ is a primitive 5-th root of unity. Write $h(x, y) = (F(x, y), G(x, y))$. The equality $h = \gamma \circ h \circ \ell^{-1}$ becomes

$$F(x, y) = 1 - G(\zeta^{-1}x, y) \quad G(x, y) = 1 - \frac{G(\zeta^{-1}x, y)}{F(\zeta^{-1}x, y)}.$$

The first equality gives $G(x, y) = 1 - F(\zeta x, y)$, and the second becomes

$$F(\zeta x, y)F(\zeta^{-1}x, y) + F(x, y) = 1. \quad (\star)$$

Conversely, if F satisfies (\star) and the map $h : (x, y) \mapsto (F(x, y), 1 - F(\zeta x, y))$ is birational, we have $\gamma = h \circ \ell \circ h^{-1}$.

The way to solve equation (\star) was told to me by Don Zagier; it is related to the famous 5-term relation for the dilogarithm, see [Z]. We will use for the cross-ratio of distinct elements x_1, \dots, x_4 in a field K the normalization

$$[x_1 : \dots : x_4] = \frac{(x_1 - x_4)(x_2 - x_3)}{(x_1 - x_3)(x_2 - x_4)}.$$

Lemma.— *Let K be a field, and σ an automorphism of order 5 of K . Let t be an element of K with $\sigma(t) \neq t$. The cross-ratio $r = [\sigma(t) : \dots : \sigma^4(t)]$ satisfies*

$$\sigma(r)\sigma^{-1}(r) + r = 1.$$

Proof: Let t_0, \dots, t_4 be distinct elements of $\mathbb{P}^1(K)$. The cross-ratios

$$r_i = [t_{i+1} : t_{i+2} : t_{i+3} : t_{i+4}],$$

where the index i runs in $\mathbb{Z}/5$, satisfy

$$r_{i-1}r_{i+1} + r_i = 1 \quad \text{for each } i ;$$

this is easily seen either by direct computation, or by using the PGL_2 -invariance of the cross-ratio to reduce to the case $(t_0, \dots, t_4) = (\lambda, 0, \infty, 1, \mu)$.

Coming back to the statement of the lemma we put $t_i = \sigma^i(t)$; then $r_i = \sigma^i(r)$, hence the result. ■

We apply this lemma to the field $\mathbb{C}(x, y)$, with the automorphism σ which fixes y and maps x to ζx . Starting with any rational function $t(x, y)$ with $t(x, y) \neq t(\zeta x, y)$, the cross-ratio $F(x, y) = [t(\zeta x, y) : \dots : t(\zeta^4 x, y)]$ is a solution to our equation (\star) ; we want moreover the map $h : (x, y) \mapsto (F(x, y), 1 - F(\zeta x, y))$ to be birational. Observe that this excludes functions t of the form $x^p a(y) + x^q b(y)$ (and more generally transforms of such functions under PGL_2), because for that choice h factors through the map $(x, y) \mapsto x^{q-p} b(y) a(y)^{-1}$. This leads us to try $t(x, y) = x + x^2 + x^3 y$; putting $\omega = \zeta + \zeta^{-1}$ we find:

$$F(x, y) = \omega \frac{1 - x - x^2(y - 1) + 2x^3 y - x^4 y^2}{1 - x - \omega^2 x^2(y - 1) + \omega x^3 y - \omega^2 x^4 y^2} .$$

We want to invert the system $F(x, y) = u$, $1 - F(\zeta x, y) = v$. Together with the functional equation (\star) for F it implies

$$F(\zeta^i x, y) = u_i \quad (0 \leq i \leq 3), \quad \text{with } u_0 = u, \quad u_1 = 1 - v, \quad u_2 = \frac{v}{u}, \quad u_3 = \frac{u - v}{u(1 - v)} ;$$

that is, explicitly:

$$x \zeta^i (\omega^{-1} u_i - 1) + x^2 (y - 1) \zeta^{2i} (\omega u_i - 1) + x^3 y \zeta^{3i} (2 - u_i) + x^4 y^2 \zeta^{4i} (\omega u_i - 1) = \omega^{-1} u_i - 1 .$$

This gives a system of 4 linear equations in $x, x^2(y - 1), x^3 y, x^4 y^2$. If the corresponding determinant is nonzero we can express x and $x^3 y$, and thus also y , as rational functions of (u, v) , so that h is birational.

Thus it remains to prove that the determinant of the system

$$\det \begin{pmatrix} \omega^{-1} u_0 - 1 & \omega u_0 - 1 & 2 - u_0 & \omega u_0 - 1 \\ \zeta(\omega^{-1} u_1 - 1) & \zeta^2(\omega u_1 - 1) & \zeta^3(2 - u_1) & \zeta^4(\omega u_1 - 1) \\ \zeta^2(\omega^{-1} u_2 - 1) & \zeta^4(\omega u_2 - 1) & \zeta(2 - u_2) & \zeta^3(\omega u_2 - 1) \\ \zeta^3(\omega^{-1} u_3 - 1) & \zeta(\omega u_3 - 1) & \zeta^4(2 - u_3) & \zeta^2(\omega u_3 - 1) \end{pmatrix}$$

is nonzero. It is enough to prove that it does not vanish for particular values of u, v such that u_0, \dots, u_3 are finite. We choose $u = \omega^{-1}$, $v = -\omega$. This gives $u_0 = u_1 = \omega^{-1}$, $u_2 = \omega - 1$, $u_3 = 3\omega - 1$; our determinant becomes:

$$\begin{aligned}
& \det \begin{pmatrix} \omega^{-1} & 0 & \omega^2 & 0 \\ \zeta\omega^{-1} & 0 & \zeta^3\omega^2 & 0 \\ -\zeta^2\omega^{-1} & -2\zeta^4\omega & \zeta(3-\omega) & -2\zeta^3\omega \\ \zeta^3\omega^2 & \zeta(2-4\omega) & 3\zeta^4\omega^2 & \zeta^2(2-4\omega) \end{pmatrix} \\
&= -\det \begin{pmatrix} \omega^{-1} & \omega^2 \\ \zeta\omega^{-1} & \zeta^3\omega^2 \end{pmatrix} \det \begin{pmatrix} -2\zeta^4\omega & -2\zeta^3\omega \\ \zeta(2-4\omega) & \zeta^2(2-4\omega) \end{pmatrix} \\
&= 2\zeta^2\omega^2(4\omega-2)(\omega^2-4) \neq 0,
\end{aligned}$$

which proves the theorem. ■

REFERENCES

- [B-B] L. BAYLE, A. BEAUVILLE: *Birational involutions of \mathbb{P}^2* . Kodaira's issue, Asian J. Math. **4** (2000), 11–17.
- [dF] T. DE FERNEX: *On planar Cremona maps of prime order*. Nagoya Math. J. **174** (2004). Preprint math.AG/0302175.
- [Z] D. ZAGIER: *The remarkable dilogarithm*. J. Math. Phys. Sci. **22** (1988), 131–145. Reprinted in “Number theory and related topics (Bombay, 1988)”, 231–249, Tata Inst. Fund. Res. Stud. Math. **12** (1989).

Arnaud BEAUVILLE
 Institut Universitaire de France
 &
 Laboratoire J.-A. Dieudonné
 UMR 6621 du CNRS
 UNIVERSITÉ DE NICE
 Parc Valrose
 F-06108 NICE Cedex 2