



A short note about Morozov's formula

Bertrand Eynard

► To cite this version:

| Bertrand Eynard. A short note about Morozov's formula. 2004. hal-00002150

HAL Id: hal-00002150

<https://hal.science/hal-00002150>

Preprint submitted on 25 Jun 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A short note about Morozov's formula

*B. Eynard*¹

Service de Physique Théorique de Saclay,
F-91191 Gif-sur-Yvette Cedex, France.

Abstract

The purpose of this short note, is to rewrite Morozov's formula for correlation functions over the unitary group, in a much simpler form, involving the computation of a single determinant.

1 Introduction

The main result of this paper is given in 2.13.

In [1], A. Morozov proposed a formula for correlation functions of unitary matrices with Itzykson-Zuber's type measure:

$$\langle |U_{ji}|^2 \rangle_{U(N)} := \frac{1}{I(X, Y)} \int_{U(N)} dU |U_{ji}|^2 e^{\text{tr}(XU^\dagger YU)} \quad (1.1)$$

where dU is the Haar measure over the unitary group $U(N)$ (appropriately normalized, so that 1.4 below holds with prefactor 1), and X and Y are two given diagonal complex matrices:

$$X = \text{diag}(x_1, \dots, x_N) \quad , \quad Y = \text{diag}(y_1, \dots, y_N) \quad (1.2)$$

and the normalization factor $I(X, Y)$ is the so-called Harish-Chandra-Itzykson-Zuber integral:

$$I(X, Y) := \int_{U(N)} dU e^{\text{tr}(XU^\dagger YU)} \quad (1.3)$$

¹E-mail: eynard@sph.t.saclay.cea.fr

It is well-known [4, 5] that:

$$I(X, Y) = \frac{\det E}{\Delta(X)\Delta(Y)} \quad (1.4)$$

where E is the matrix with entries:

$$E_{ij} := e^{x_i y_j} \quad (1.5)$$

and where $\Delta(X)$ and $\Delta(Y)$ are the Vandermonde determinants:

$$\Delta(X) := \prod_{i < j} (x_i - x_j) \quad , \quad \Delta(Y) := \prod_{i < j} (y_i - y_j) . \quad (1.6)$$

Here we shall assume that $\det E \neq 0$ and $\Delta(X) \neq 0$ and $\Delta(Y) \neq 0$.

Morozov's formula was proven in [2], from Shatashvili's formula [3]. Morozov's formula was originally written as follows, for any arbitrary sequences of complex numbers a_i, b_j one has:

$$\begin{aligned} & \sum_{i,j=1}^N a_i b_j \int_{U(N)} dU |U_{ji}|^2 e^{\text{tr}(X U^\dagger Y U)} = \frac{1}{\Delta(X)\Delta(Y)} \sum_{\rho \in S_N} \epsilon(\rho) e^{\sum x_\ell y_{\rho(\ell)}} \\ & \det \begin{pmatrix} 1 & \cdots & 1 \\ x_{i_1} & \cdots & x_{i_{n+1}} \\ \vdots & & \vdots \\ x_{i_1}^{n-1} & \cdots & x_{i_{n+1}}^{n-1} \\ a_{i_1} & \cdots & a_{i_{n+1}} \end{pmatrix} \det \begin{pmatrix} 1 & \cdots & 1 \\ y_{\rho(i_1)} & \cdots & y_{\rho(i_{n+1})} \\ \vdots & & \vdots \\ y_{\rho(i_1)}^{n-1} & \cdots & y_{\rho(i_{n+1})}^{n-1} \\ b_{\rho(i_1)} & \cdots & b_{\rho(i_{n+1})} \end{pmatrix}, \\ & \sum_{n=0}^{N-1} (-1)^n \sum_{i_1 < i_2 < \dots < i_{n+1}} \frac{\det \begin{pmatrix} 1 & \cdots & 1 \\ x_{i_1} & \cdots & x_{i_{n+1}} \\ \vdots & & \vdots \\ x_{i_1}^n & \cdots & x_{i_{n+1}}^n \end{pmatrix} \det \begin{pmatrix} 1 & \cdots & 1 \\ y_{\rho(i_1)} & \cdots & y_{\rho(i_{n+1})} \\ \vdots & & \vdots \\ y_{\rho(i_1)}^n & \cdots & y_{\rho(i_{n+1})}^n \end{pmatrix}}{\det \begin{pmatrix} 1 & \cdots & 1 \\ x_{i_1} & \cdots & x_{i_{n+1}} \\ \vdots & & \vdots \\ x_{i_1}^n & \cdots & x_{i_{n+1}}^n \end{pmatrix} \det \begin{pmatrix} 1 & \cdots & 1 \\ y_{\rho(i_1)} & \cdots & y_{\rho(i_{n+1})} \\ \vdots & & \vdots \\ y_{\rho(i_1)}^n & \cdots & y_{\rho(i_{n+1})}^n \end{pmatrix}}, \end{aligned} \quad (1.7)$$

where S_N is the symmetric group of rank N .

2 Rewriting Morozov's formula

The purpose of this short note is to rewrite this formula in a much simpler form, in a way very similar to what was done in [2].

For any two complex numbers x and y (such that $|x| > \max |x_i|$ and $|y| > \max |y_i|$), consider the choice:

$$a_i = \frac{1}{x - x_i} = \sum_{r=0}^{\infty} \frac{x_i^r}{x^{r+1}} \quad , \quad b_i = \frac{1}{y - y_i} = \sum_{s=0}^{\infty} \frac{y_i^s}{y^{s+1}} \quad (2.8)$$

Introduce the Schur polynomials (corresponding to hook diagramms):

$$\begin{aligned}
S_r(x_{i_1}, \dots, x_{i_{n+1}}) &:= \frac{\det \begin{pmatrix} 1 & \cdots & 1 \\ x_{i_1} & \cdots & x_{i_{n+1}} \\ \vdots & & \vdots \\ x_{i_1}^{n-1} & \cdots & x_{i_{n+1}}^{n-1} \\ x_{i_1}^r & \cdots & x_{i_{n+1}}^r \end{pmatrix}}{\det \begin{pmatrix} 1 & \cdots & 1 \\ x_{i_1} & \cdots & x_{i_{n+1}} \\ \vdots & & \vdots \\ x_{i_1}^n & \cdots & x_{i_{n+1}}^n \end{pmatrix}} \\
&= \sum_{a_1 \leq a_2 \leq \dots \leq a_{r-n}} \prod_{k=1}^{r-n} x_{i_{a_k}} = \sum_{j_1 + \dots + j_{n+1} = r-n} x_{i_1}^{j_1} \cdots x_{i_{n+1}}^{j_{n+1}}. \quad (2.9)
\end{aligned}$$

The formal generating function of these Schur polynomials is:

$$\sum_{r=0}^{\infty} \frac{1}{x^{r+1}} S_r(x_{i_1}, \dots, x_{i_{n+1}}) = \prod_{k=1}^{n+1} \frac{1}{x - x_{i_k}} \quad (2.10)$$

Inserting that into 1.7, one gets:

$$\begin{aligned}
&I(X, Y) \sum_{i,j} \frac{1}{x - x_i} \frac{1}{y - y_j} \langle |U_{ji}|^2 \rangle_{U(N)} \\
&= \int_{U(N)} dU \operatorname{tr} \left(\frac{1}{x - X} U \frac{1}{y - Y} U^\dagger \right) e^{\operatorname{tr}(XU^\dagger YU)} \\
&= \frac{1}{\Delta(X)\Delta(Y)} \sum_{r,s=0}^{\infty} \sum_{\rho \in S_N} \epsilon(\rho) e^{\sum x_\ell y_{\rho(\ell)}} \\
&\quad \sum_{n=0}^{N-1} (-1)^n \sum_{i_1 < i_2 < \dots < i_{n+1}} \frac{S_r(x_{i_1}, \dots, x_{i_{n+1}})}{x^{r+1}} \frac{S_s(y_{\rho(i_1)}, \dots, y_{\rho(i_{n+1})})}{y^{s+1}} \\
&= \frac{1}{\Delta(X)\Delta(Y)} \sum_{\rho \in S_N} \epsilon(\rho) e^{\sum x_\ell y_{\rho(\ell)}} \sum_{n=0}^{N-1} (-1)^n \sum_{i_1 < i_2 < \dots < i_{n+1}} \prod_{k=1}^{n+1} \frac{1}{x - x_{i_k}} \prod_{l=1}^{n+1} \frac{1}{y - y_{\rho(i_l)}} \\
&= \frac{1}{\Delta(X)\Delta(Y)} \sum_{\rho \in S_N} \epsilon(\rho) e^{\sum x_\ell y_{\rho(\ell)}} \left[1 - \prod_{i=1}^N \left(1 - \frac{1}{x - x_i} \frac{1}{y - y_{\rho(i)}} \right) \right] \\
&= \frac{1}{\Delta(X)\Delta(Y)} \sum_{\rho \in S_N} \epsilon(\rho) \left[\prod_{i=1}^N e^{x_i y_{\rho(i)}} - \prod_{i=1}^N \left(e^{x_i y_{\rho(i)}} - \frac{1}{x - x_i} e^{x_i y_{\rho(i)}} \frac{1}{y - y_{\rho(i)}} \right) \right] \\
&= \frac{1}{\Delta(X)\Delta(Y)} \left[\det(e^{x_i y_j}) - \det \left(e^{x_i y_j} - \frac{1}{x - x_i} e^{x_i y_j} \frac{1}{y - y_j} \right) \right]
\end{aligned} \quad (2.11)$$

Thus, Morozov's formula can be rewritten:

$$\int_{U(N)} dU \operatorname{tr} \left(\frac{1}{x - X} U \frac{1}{y - Y} U^\dagger \right) e^{\operatorname{tr}(XU^\dagger YU)} = \frac{\det E - \det \left(E - \frac{1}{x-X} E \frac{1}{y-Y} \right)}{\Delta(X)\Delta(Y)} \quad (2.12)$$

or:

$$\begin{aligned}
 \left\langle \text{tr} \left(\frac{1}{x-X} U \frac{1}{y-Y} U^\dagger \right) \right\rangle &= 1 - \frac{\det \left(E - \frac{1}{x-X} E \frac{1}{y-Y} \right)}{\det E} \\
 &= 1 - \det \left(1 - \frac{1}{x-X} E \frac{1}{y-Y} E^{-1} \right)
 \end{aligned} \tag{2.13}$$

3 Concluding remarks

From that expression of Morozov's formula, it is rather easy to recover any individual correlator by taking residues:

$$\left\langle U_{ij} U_{ji}^\dagger \right\rangle = \underset{x \rightarrow x_i}{\text{Res}} \underset{y \rightarrow x_j}{\text{Res}} \left\langle \text{tr} \left(\frac{1}{x-X} U \frac{1}{y-Y} U^\dagger \right) \right\rangle \tag{3.14}$$

Notice also that 2.13 is very similar to what was found in [2] after integration over X and Y .

Notice that by expanding the determinant in 2.13 along its last column, one can find a recursion relation relating $U(N)$ to $U(N-1)$ integrals, which is equivalent to Shatashvili's approach [3].

Acknowledgements: the author wants to thank M. Bertola and J. Harnad for discussions about that topic. The author thinks that this short note should be seen as an addendum to the article [2].

References

- [1] A. Morozov, "Pair correlator in the Itzykson–Zuber Integral", *Modern Phys. Lett. A* **7**, no. 37 3503–3507 (1992).
- [2] M. Bertola, B. Eynard, "Mixed Correlation Functions of the Two-Matrix Model", SPHT T03/028, CRM-2961 (2003). *J. Phys. A* **36** (2003) 7733–7750, xxx, hep-th/0303161.
- [3] S. L. Shatashvili, "Correlation Functions in the Itzykson–Zuber Model", *Comm. Math. Phys.* **154** 421–432 (1993).
- [4] C. Itzykson and J.B. Zuber, "The planar approximation (II)", *J. Math. Phys.* **21**, 411 (1980).
- [5] M.L. Mehta, *Random Matrices*, 2nd edition, (Academic Press, New York, 1991).