



HAL
open science

Extremality for the Vafa-Witten bound on the sphere

Marc Herzlich

► **To cite this version:**

Marc Herzlich. Extremality for the Vafa-Witten bound on the sphere. *Geometric And Functional Analysis*, 2005, 15, pp.1153-1161. 10.1007/s00039-005-0536-5 . hal-00002414v2

HAL Id: hal-00002414

<https://hal.science/hal-00002414v2>

Submitted on 29 Nov 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXTREMALITY FOR THE VAFA–WITTEN BOUND ON THE SPHERE

MARC HERZLICH

ABSTRACT. We prove that the round metric on the sphere has the largest first eigenvalue of the Dirac operator among all metrics that are larger than it. As a corollary, this gives an alternative proof of an extremality result for scalar curvature due to M. Llarull.

1. INTRODUCTION

C. Vafa and E. Witten have shown in [18] that there exists a common upper bound for the smallest eigenvalue of all *twisted* Dirac operators on a given Riemannian manifold. The common upper bound they found depends strongly on the choice of the metric on the base. Using an analogous method, H. Baum [3] exhibited later an explicit upper bound for the first eigenvalue of the (untwisted) Dirac operator on an even-dimensional Riemannian manifold that can be sent on a sphere by a map of (high) non-zero degree. This bound depends on the Lipschitz norm of the map from the manifold to the round sphere.

The goal of this short note is to elaborate further on H. Baum's results [3]. Indeed, we prove below that some *optimal* upper bound can be obtained for the bottom of the spectrum of the (untwisted) Dirac operator of a large class of metrics on the sphere. This extends [3] in two different ways, first by providing an optimal bound, and secondly by extending it to odd dimensions.

Theorem 1. *Let g be any Riemannian metric and b be the round metric of constant curvature 1 on the sphere \mathbb{S}^n . If $g \geq b$ pointwise, then there is an eigenvalue of the Dirac operator of g in $[-\frac{n}{2}, \frac{n}{2}]$. Moreover, if no eigenvalue lies in the open interval $]-\frac{n}{2}, \frac{n}{2}[$, then g is isometric to the round metric b .*

As an interesting corollary, one gets an alternative proof of M. Llarull's extremality result for scalar curvature on the sphere [14]. Indeed, the classical first eigenvalue estimate for the Dirac operator due to Th. Friedrich [7] states that the smallest eigenvalue $\lambda_1(g)$ (in absolute value) of the Dirac operator satisfies:

$$|\lambda_1(g)|^2 \geq \frac{n}{4(n-1)} \inf \text{Scal}_g .$$

1991 *Mathematics Subject Classification.* 53C27, 58J50, 58J60.

Key words and phrases. Dirac operator, eigenvalue estimate, scalar curvature.

The author is a member of the EDGE Research Training Network HPRN-CT-2000-00101 of the European Union and is supported in part by an ACI program of the French Ministry of Research.

Hence one concludes:

Theorem 2 (Llarull [14]). *Let g be any metric and b be the round metric of constant curvature 1 on the sphere \mathbb{S}^n . If $g \geq b$ pointwise, then*

$$\inf \text{Scal}_g \leq n(n-1) ,$$

the inequality being strict if g is distinct from b .

The proof of Theorem 1 is close to the original approach of Vafa and Witten, see also M. Atiyah [1] and H. Baum [3]. Since we look for optimal bounds, we have however to pay a special attention to all the estimates involved. As a result, this implies that two different proofs are required in the even- and odd-dimensional cases (see [6] for another occurrence of this problem in a related context). The first one is obtained from an index argument, whereas the second one needs a slightly more delicate proof using spectral flow considerations.

Acknowledgements. It was Maung Min-Oo who pointed out during a lecture in July 2001 [16] that Vafa–Witten was optimal on a round sphere and that a strong link might exist with Llarull extremality. This idea appears (in a somewhat more hidden way) on pages 84–85 in M. Gromov’s long and rich paper [11]. The author is then happy to acknowledge the origin of the ideas behind this note. He thanks Maung Min-Oo again, Andrei Moroianu and Uwe Semmelmann for useful comments, H el ene Davaux for her careful reading of a first draft of this paper, and the referee for his remarks.

2. BACKGROUND MATERIAL

For any metric g on \mathbb{S}^n , we let Σ_g be its spin bundle. Using the idea introduced by J. P. Bourguignon and P. Gauduchon [4], one may identify the spin bundle Σ_g with the spin bundle Σ_b through a lift of the principal oriented orthonormal frame bundles isomorphism

$$H : P_{SO}\mathbb{S}^n(b) \longrightarrow P_{SO}\mathbb{S}^n(g)$$

induced by the unique symmetric positive definite map H such that

$$g(H\cdot, H\cdot) = b(\cdot, \cdot).$$

If $g \geq b$, this implies that H has operator norm relative to b bounded from above by 1. This allows to transfer the Clifford action relative to g on the bundle Σ_b . Denoting by ℓ_b (resp. ℓ_g) : $T\mathbb{S}^n \rightarrow \text{End}(\Sigma_b)$ the Clifford actions for b and g , they are related as:

$$\ell_b(\cdot) = \ell_g(H\cdot)$$

The Levi-Civita connection for any metric g can be transferred in the same way on Σ_b as a metric connection (but with torsion). In what follows, we will always assume that all spinor bundles on the sphere have been identified to those relative to b , as above.

The following facts are extracted from classical texts on spinor geometry, see also [2, 3, 8] which are useful references.

In even dimensions $n = 2m$, the spin bundle Σ_g splits into two *half-spin* bundles Σ_g^+ and Σ_g^- ; on the sphere \mathbb{S}^{2m} , the whole spin bundle, or equivalently the sum of both half-spin bundles, is a trivial bundle obtained by restricting the spinor bundle of \mathbb{R}^{2m+1} to the sphere; namely:

$$\mathbb{T} := \Sigma_b = \Sigma_b^+ \oplus \Sigma_b^- = \mathbb{S}^{2m} \times \mathbb{C}^{2^m}.$$

Moreover, projections on each factor Σ_b^\pm can be explicitly described at a point x on the sphere as:

$$\Pi_\pm(x) = \frac{1}{2}(1 \pm i\ell_0(x))$$

where ℓ_0 is the algebraic Clifford action of Cl_{2m+1} on \mathbb{C}^{2^m} (as a result, none of the half-spin bundles is trivial). Using this expression, it is easy to relate the trivial connection ∇^0 (of flat space) acting on \mathbb{T} with the Levi-Civita connection of b acting on Σ_b^\pm . If ψ is any spinor field in Σ_b^\pm , then:

$$\nabla_X^\pm \psi = \Pi_\pm \nabla_X^0 \Pi_\pm \psi = \nabla_X^0 \psi - \frac{1}{2} \ell_0(x) \ell_0(X) \psi$$

and the full Levi-Civita connection for b on $\Sigma_b = \Sigma_b^+ \oplus \Sigma_b^-$ is $\nabla^b = \nabla^+ \oplus \nabla^-$ (note that actions ℓ_0 on vectors tangent to \mathbb{S}^n and ℓ_b may be identified).

Moreover, the bundle $\mathbb{T} \otimes \mathbb{T} = \Sigma_b \otimes \Sigma_b$ is the (complex) differential form bundle $\Lambda^\bullet \mathbb{S}^n \otimes \mathbb{C}$ of the sphere. On this bundle, two different Clifford actions exist. The first is the usual one for twisted spinor bundles, with the Clifford algebra acting on the left factor of the tensor product; we will continue to denote this one below by ℓ , as it is defined on a decomposed element $\sigma \otimes \tau$ of $\Sigma_b \otimes \Sigma_b$ by:

$$\ell_b(X)(\sigma \otimes \tau) = (\ell_b(X)\sigma) \otimes \tau.$$

But there is another, right-handed, one that we shall denote by r and which is defined by:

$$r_b(X)(\sigma \otimes \tau) = \sigma \otimes (\ell_b(X)\tau).$$

Both Clifford actions can be explicitly described when one identifies $\Sigma_b \otimes \Sigma_b$ with $\Lambda^\bullet \mathbb{S}^{2m}$ as follows: for any p -form ω , and any 1-form α

$$\ell_b(\alpha^\sharp)\omega = \alpha \wedge \omega - i_{\alpha^\sharp}\omega, \quad r_b(\alpha^\sharp)\omega = (-1)^p (\alpha \wedge \omega + i_{\alpha^\sharp}\omega)$$

(musical isomorphism \sharp referring, as always, to b). For more information on these points, the reader is referred to the book [13] (see also the paper [17] where a concise account of these facts is given). The reader should also be careful about tensor product connections, as they might not coincide: for instance, $\nabla^b \otimes 1 + 1 \otimes \nabla^b$ is the Levi-Civita connection on differential forms on the sphere whereas $\nabla^0 \otimes 1 + 1 \otimes \nabla^0$ is the trivial connection induced from \mathbb{R}^{2m+1} .

In odd dimensions $n = 2m - 1$, the Clifford algebra is isomorphic to the sum of two copies of the matrix algebra $\text{End}(\mathbb{C}^{2^{m-1}})$. Hence it has two inequivalent representations which lead to two different Clifford bundles Σ_b and Σ'_b on the sphere

\mathbb{S}^{2m-1} . They are equivalent as bundles with structure group $\text{Spin}(2m-1)$ but the Clifford actions differ. Their sum is again a trivial bundle:

$$\mathbb{T} := \Sigma_b \oplus \Sigma'_b = \mathbb{S}^{2m-1} \times \mathbb{C}^{2^m} ;$$

as above, it is the restriction to the sphere of the full spinor bundle of flat space \mathbb{R}^{2m} . This bundle itself splits in two trivial subbundles of rank 2^{m-1}

$$\mathbb{T} = \mathbb{T}^+ \oplus \mathbb{T}^-$$

obtained by taking the $(\pm i)$ -eigenspaces of the volume form of \mathbb{R}^{2m} . Here we have to take care that these bundles are different from the bundles Σ_b and Σ'_b on the sphere already alluded to a few lines above: for instance, \mathbb{T}^+ and \mathbb{T}^- are exchanged when Clifford multiplied by vectors, whereas Σ_b and Σ'_b are preserved by the Clifford action. Notice that, contrarily to what happens if $n = 2m$, the bundles \mathbb{T}^\pm are preserved by the flat connection ∇^0 . As final remark, we recall that $\mathbb{T} \otimes \mathbb{T}$ is again isomorphic to the full differential form bundle of \mathbb{R}^{2m} .

From now on and to avoid confusion, we will use notations \mathbb{T} , Σ_b and Σ_g in the following sense : they will always denote the corresponding bundle relative to the metric b , but endowed either with the trivial connection ∇^0 , or the Levi-Civita connections ∇^b or ∇^\pm of b on spinors, or the Levi-Civita connection ∇^g of g on spinors.

3. THE PROOF: EVEN-DIMENSIONAL CASE

We consider the bundle $S = \Sigma_g \otimes \Sigma_b^+$; as spin bundles for different metrics always are identified, this means that we take the tensor product of the whole spin bundle Σ_b with the half-spin Σ_b^+ , but endowed with the *non-trivial* tensor product connection $\nabla^g \otimes 1 + 1 \otimes \nabla^+$, where ∇^g is the Levi-Civita connection of the metric g and ∇^+ has been defined in the previous section.

We now apply Atiyah-Singer index theorem to the twisted Dirac operator

$$(3.1) \quad \mathcal{D}^{g+} : \Sigma_g^+ \otimes \Sigma_b^+ \longrightarrow \Sigma_g^- \otimes \Sigma_b^+.$$

One of the most important consequences of index theory is the topological invariance of the index. The index of the following model ('round') Dirac operator:

$$\mathcal{D}^{b+} : \Sigma_b^+ \otimes \Sigma_b^+ \longmapsto \Sigma_b^- \otimes \Sigma_b^+$$

equals 1 (it is the operator whose index equals one half of the Euler number in dimension $4k+2$ and one half of the sum of the Euler number and the signature in dimension $4k$; hence this value on the even-dimensional sphere, see for instance [8, p. 95–96] for an explicit derivation), so that the index of the Dirac operator \mathcal{D}^{g+} built with the connection $\nabla^g \otimes 1 + 1 \otimes \nabla^+$ on the twisted spin bundle $S = \Sigma_g \otimes \Sigma_b^+$ is also 1. Its kernel is then non-zero.

Let us now consider the tensor product bundle $\Sigma_g \otimes \mathbb{T}$, endowed this time with the connection $\nabla^g \otimes 1 + 1 \otimes \nabla^0$. As the pair (\mathbb{T}, ∇^0) is a *trivial* flat bundle on the sphere, the spectrum of the twisted Dirac operator attached to this connection is

the same as the spectrum of the Dirac operator on Σ_g , but with each eigenvalue repeated 2^m times its multiplicity. Applying standard perturbation theory, one gets the first part of the statement of Theorem 1 if one is able to bound the difference

$$\bar{L} := \mathcal{D}^{g^0} - (\mathcal{D}^{g^+} \oplus \mathcal{D}^{g^-}) : \Sigma_g \otimes \Sigma_b \longrightarrow \Sigma_g \otimes \Sigma_b.$$

This is easily computed on a decomposed section $\sigma \otimes \tau$ of $\Sigma_g \otimes \Sigma_b$: if $\{e_i\}$ is any g -orthonormal basis on the sphere,

$$\begin{aligned} \bar{L}(\sigma \otimes \tau) &= \sum_{i=1}^n \ell_g(e_i) (\nabla_{e_i}^g \sigma \otimes \tau + \sigma \otimes \nabla_{e_i}^0 \tau - \nabla_{e_i}^g \sigma \otimes \tau - \sigma \otimes \nabla_{e_i}^+ \tau) \\ &= \frac{1}{2} \sum_i \ell_g(e_i) \sigma \otimes \ell_0(x) \ell_0(e_i) \tau \end{aligned}$$

This shows that

$$\bar{L} = \frac{1}{2} r_0(x) L,$$

where L is the (pointwise) linear map of vector bundles given on a spinor field ψ in $\Sigma_g \otimes \Sigma_b$ by

$$L\psi = \sum_i \ell_g(e_i) r_b(e_i) \psi.$$

We now choose a b -orthonormal basis $\{\varepsilon_i\}$ that diagonalizes H . Eigenvalues μ_i live in $]0, 1]$ and $\{e_i = \mu_i \varepsilon_i\}$ is a g -orthonormal basis, and moreover,

$$L\psi = \sum_i \mu_i^{-1} \ell_b(e_i) r_b(e_i) \psi = \sum_i \mu_i \ell_b(\varepsilon_i) r_b(\varepsilon_i) \psi.$$

As Clifford multiplication by a b -unit vector has b -operator norm equal to 1, one immediately obtains that

$$|L\psi|_b \leq \left(\sum \mu_i \right) |\psi|_b \leq n |\psi|_b,$$

with equality if and only if $\mu_i = 1$ for all i . This concludes the proof of Theorem 1 in the even-dimensional case. \square

Remark. Using the explicit expressions of ℓ_b and r_b on differential forms, one gets

$$\ell_b(\varepsilon_k) r_b(\varepsilon_k) (\varepsilon_{i_1} \wedge \cdots \wedge \varepsilon_{i_p}) = \begin{cases} (-1)^p \varepsilon_{i_1} \wedge \cdots \wedge \varepsilon_{i_p} & \text{if } k \in \{i_1, \dots, i_p\}, \\ (-1)^{p+1} \varepsilon_{i_1} \wedge \cdots \wedge \varepsilon_{i_p} & \text{if not,} \end{cases}$$

and this of course leads to an explicit form of the previous estimate.

4. THE PROOF: ODD-DIMENSIONAL CASE

Our treatment of the odd-dimensional case is much inspired by [1, 6]. Starting from the Dirac operator on the spin bundle Σ_g , our goal is to find a trivial twisting bundle endowed with a non-flat connection whose associated twisted Dirac operator has a non-zero kernel. As index considerations are useless in odd dimensions, we have to rely on a spectral flow argument.

Let the dimension be $n = 2m - 1$. Our choice of bundle on \mathbb{S}^n is

$$\Sigma_g \otimes \mathbb{T}^+ = \Sigma \otimes \mathbb{C}^{2^{m-1}}$$

(notations as in section 2), so that, as above, the Dirac operator \mathcal{D}^{g0} associated to $\nabla^g \otimes 1 + 1 \otimes \nabla^0$ on $\Sigma_g \otimes \mathbb{T}^+$ has the same spectrum as \mathcal{D}^g on Σ_g (up to multiplicity). Note that this is the only possible choice of connection at this stage since \mathbb{T}^\pm is not preserved by the Levi-Civita connection of (\mathbb{S}^n, b) .

We now fix e unit in \mathbb{R}^{2m} and we define for each x in \mathbb{S}^{2m-1} an endomorphism $u(x)$ of the fiber of \mathbb{T}^+ at x as follows: we start with $(e \cdot x \cdot)$ which is an element \mathcal{Cl}_{2m} that preserves both factors \mathbb{T}^\pm . Hence we let:

$$u(x) = \text{the projection of } (e \cdot x \cdot) \text{ in } \text{End}(\mathbb{T}^+) \text{ (= End}(\mathbb{C}^{2^{m-1}})\text{)}.$$

As each x is skew-hermitian and of square -1 , $u(x)$ is unitary for each x and one can define the unitarily (gauge) equivalent connection

$$\nabla^u = u^{-1} \circ \nabla^0 \circ u$$

on \mathbb{T}^+ . The path of connections

$$t \in [0, 1] \longmapsto \nabla^g \otimes 1 + 1 \otimes \nabla^t \text{ with } \nabla^t = (1 - t)\nabla^0 + t\nabla^u .$$

gives rise to a path of Dirac operators \mathcal{D}^t on $\Sigma_g \otimes \mathbb{T}^+$ with unitarily equivalent operators at $\{t = 0\}$ and $\{t = 1\}$. The spectral flow of this family can be computed using the index theorem on $\mathbb{S}^{2m-1} \times \mathbb{S}^1$, applied to the Dirac operator acting on the spinor bundle Σ_b twisted by the bundle obtained by identifying \mathbb{T}^+ at $\{t = 0\}$ and $\{t = 1\}$ through u .

This spectral flow is non-zero: as a matter of fact, its value is

$$sf = \int_{\mathbb{S}^{2m-1} \times \mathbb{S}^1} \hat{A}(\mathbb{S}^{2m-1} \times \mathbb{S}^1) \text{ch}(\tilde{\mathbb{T}}^+) \quad ,$$

where $\tilde{\mathbb{T}}^+$ is the bundle obtained on $\mathbb{S}^{2m-1} \times \mathbb{S}^1$ from $\mathbb{T}^+ \times [0, 1]$ by identifying through u at $\{t = 0\}$ and $\{t = 1\}$. Moreover,

$$\hat{A}(\mathbb{S}^{2m-1} \times \mathbb{S}^1) = 1,$$

and $\text{ch}(\tilde{\mathbb{T}}^+)$ can be computed with the curvature 2-form

$$\Omega = dt \wedge u^{-1} du - (t - t^2) u^{-1} du \wedge u^{-1} du$$

obtained by writing the connection $\nabla^t = \nabla^0 + \omega^t$, where ω^t is the 1-form with values in $\text{End}(\mathbb{T}^+)$ equal to $t u^{-1}(du \cdot)$; hence

$$\begin{aligned} \text{ch}(\Omega) &= \left(\frac{i}{2\pi} \right)^m \text{Tr}(\Omega^m) \\ &= C (t - t^2)^{m-1} dt \wedge \text{Tr}(u^{-1} du \wedge \cdots \wedge u^{-1} du) \end{aligned}$$

with C a non-zero constant. This computation shows that

$$sf = \int_{\mathbb{S}^{2m-1} \times \mathbb{S}^1} \text{ch}(\Omega) = C \int_{[0,1]} (t - t^2)^{m-1} dt \int_{\mathbb{S}^{2m-1}} \text{Tr}(x \cdot e_1 \cdots e_{2m-1}) d\text{vol}_{\mathbb{S}^{2m-1}}$$

and this last result is non-zero by the very definition of \mathbb{T}^\pm as the $(\pm i)$ -eigenspaces of the volume form of \mathbb{R}^{2m} (see [5, 6] for instance for more detailed computations in an analogous case).

As a result, there exists a value of t in $[0, 1]$ for which the kernel of \mathcal{D}^t is non-zero. Suppose $t \leq 1/2$, then, as in the previous section, what is needed to conclude is an estimate of the difference

$$\bar{L}_t = \mathcal{D}^t - \mathcal{D}^{g^0}$$

(in case $t > 1/2$, one should use $u^{-1} \circ \mathcal{D}^{g^0} \circ u$ rather than \mathcal{D}^{g^0} but this is harmless since both operators have the same spectrum). An easy computation shows that this difference of operators acts as the following linear map on $\Sigma_g \otimes \mathbb{T}$ seen as a sub-bundle of the whole differential form bundle $\mathbb{T} \otimes \mathbb{T}$ of \mathbb{R}^{2m} :

$$\bar{L}_t = r_b(tx) \circ \left(\sum_{i=1}^{2m-1} \mu_i \ell_b(\varepsilon_i) r_b(\varepsilon_i) \right)$$

(same notations as in the previous section). The end of the proof is then entirely analogous to that in even dimensions. \square

5. FINAL COMMENTS

It is known that scalar curvature extremality extends to a large family of Riemannian manifolds; among these, one can find complex projective spaces, Kähler manifolds of positive Ricci curvature, non-negatively curved locally symmetric spaces (see the foundational [11, §6] and [9, 10, 12, 14, 15]). In view of Theorem 2, it seems then natural to ask the following obvious questions:

Question 1. Can one find other examples of eigenvalue-extremal metrics ?

Question 2. In particular, if g is a metric on $\mathbb{C}\mathbb{P}^m$ (with m odd, so that it is spin) that is larger than the Fubini-Study metric f , does one have $|\lambda_1(g)| \leq |\lambda_1(f)|$ with equality if and only if g is isometric to f ?

REFERENCES

1. M. Atiyah, *Eigenvalues of the Dirac operator*, Arbeitstagung Bonn 1984, Lect. Notes in Math., vol. 1111, Springer, 1985, pp. 251–260.
2. H. Baum, *Complete Riemannian manifolds with imaginary Killing spinors*, Ann. Glob. Anal. Geom. **7** (1989), 205–226.
3. ———, *An upper bound for the first eigenvalue of the Dirac operator on compact spin manifolds*, Math. Z. **206** (1991), 409–422.
4. J. P. Bourguignon and P. Gauduchon, *Spineurs, opérateurs de Dirac et variations de métriques*, Commun. Math. Phys. **144** (1992), 581–599.
5. H. Davaux, *K-aire et courbure scalaire des variétés riemanniennes*, Ph.D. thesis, Univ. Montpellier II, 2002.
6. ———, *An optimal inequality between scalar curvature and spectrum of the laplacian*, Math. Ann. **327** (2003), 271–292.
7. T. Friedrich, *Der erste Eigenwert des Dirac-Operators einer kompakten Riemannschen Mannigfaltigkeit nichtnegativer Krümmung*, Math. Nachr. **97** (1980), 117–146.
8. P. B. Gilkey, *Invariance theory, the heat equation and the Atiyah-Singer index theorem*, Publish and Perish, 1984, electronic edition 1996, available through EMIS Elec. Library <http://www.emis.de/monographs/gilkey>.
9. S. Goette and U. Semmelmann, *Spin^c structures and scalar curvature estimates*, Ann. Glob. Anal. Geom. **20** (2001), 301–324.
10. ———, *Scalar curvature estimates for compact symmetric spaces*, Diff. Geom. Appl. **16** (2002), 65–78.
11. M. Gromov, *Positive curvature, macroscopic dimension, spectral gaps and higher signatures*, Functional analysis on the eve of the 21st century, Vol. II (New Brunswick, NJ, 1993) (S. Gindikin, J. Lepowski, and R. Wilson, eds.), Progress in Math., vol. 132, Birkhäuser, Boston, 1996, pp. 1–213.
12. W. Kramer, *The scalar curvature on totally geodesic fiberings*, Ann. Global Anal. Geom. **18** (2000), 589–600.
13. H. B. Lawson and M. L. Michelsohn, *Spin geometry*, Princeton Math. Series, vol. 38, Princeton Univ. Press, Princeton, 1989.
14. M. Llarull, *Sharp estimates and the Dirac operator*, Math. Ann. **310** (1998), 55–71.
15. M. Min-Oo, *Scalar curvature rigidity of certain symmetric spaces*, Geometry, topology and dynamics, Proc. Lect. Notes, vol. 15, CRM Montréal, 1998, pp. 127–136.
16. ———, *The Dirac operator in geometry and physics*, Global Riemannian geometry: curvature and topology (V. Palmer, ed.), Adv. Courses Math. CRM Barcelona, Birkhäuser, 2003, pp. 55–87.
17. I. Prokhorenkov and K. Richardson, *Perturbation of Dirac operators*, preprint (2003), available as math.DG/0307251.
18. C. Vafa and E. Witten, *Eigenvalue inequalities for fermions in gauge theories*, Commun. Math. Phys. **95** (1984), 257–276.

INSTITUT DE MATHÉMATIQUES ET MODÉLISATION DE MONTPELLIER, UMR 5149 DU CNRS,
UNIVERSITÉ MONTPELLIER II, FRANCE

E-mail address: herzlich@math.univ-montp2.fr