



The roots of any polynomial equation

Geert-Jan Uytendewilligen

► To cite this version:

| Geert-Jan Uytendewilligen. The roots of any polynomial equation. 2004. <hal-00002529v2>

HAL Id: hal-00002529

<https://hal.science/hal-00002529v2>

Preprint submitted on 20 Aug 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



The roots of any polynomial equation

G.A.Uytendewilligen,

Bergen op Zoomstraat 76, 5652 KE Eindhoven. g.a.uytendewilligen@zonnet.nl

Abstract

We provide a method for solving the roots of the general polynomial equation

$$a_n \cdot x^n + a_{n-1} \cdot x^{n-1} + \dots + a_1 \cdot x + s = 0 \quad (1)$$

To do so, we express x as a powerseries of s , and calculate the first $n-1$ coefficients. We turn the polynomial equation into a differential equation that has the roots as solutions. Then we express the powerseries' coefficients in the first $n-1$ coefficients. Then the variable s is set to a_0 . A free parameter is added to make the series convergent. © 2004 G.A.Uytendewilligen. All rights reserved.

Keywords: Algebraic equation

The method

The method is based on [1]. Let's take the first $n-1$ derivatives of (1) to s . Equate these derivatives to zero.

Then find $\frac{d^i}{ds^i} x(s)$ in terms of $x(s)$ for i from 1 to $n-1$. Now make a new differential equation

$$m_1 \cdot \frac{d^{n-1}}{ds^{n-1}} x(s) + m_2 \cdot \frac{d^{n-2}}{ds^{n-2}} x(s) + \dots + m_n \cdot x(s) + m_{n+1} = 0 \quad (2)$$

and fill in our $\frac{d^i}{ds^i} x(s)$ in (2). Multiply by the denominator of the expression. Now we have a polynomial in $x(s)$ of degree higher than n . Using (1) as property, we simplify this polynomial to the degree of n . Set it equal to (1) and solve $m_1 \dots m_{n+1}$ in terms of s and $a_1 \dots a_n$. Substituting these in (2) gives a differential equation that has the zeros of (1) among its solutions. We then insert

$$x(s) = y(s) - \frac{a_{n-1}}{n \cdot a_n} \quad (3)$$

in (2). Multiplying by the denominator we get a differential equation of the linear form:

$$p_1 \cdot \frac{d^{n-1}}{ds^{n-1}} y(s) + p_2 \cdot \frac{d^{n-2}}{ds^{n-2}} y(s) + \dots + p_n \cdot y(s) = 0 \quad (4)$$

With $p_1(s) \dots p_n(s)$ polynomials in s . If we substitute our powerseries, all the coefficients are determined by the first $n-1$ coefficients. The first coefficients are calculated as follows: A powerseries is filled in in (1).

$$x(s) = \sum_{i=0}^{n-2} b_i \cdot s^i - \frac{a_n}{n \cdot a_{n-1}} \quad (5)$$

and it should be zero for all s . From this, we calculate b_i for i from 0 to $n-2$. b_0 is a root of an $n-1$ degree polynomial and the other b_i are expressed in b_0 . Now a powerseries is inserted in (4):

$$y(s) = \sum_{i=0}^{\infty} b_i \cdot s^i \quad (6)$$

and we get an equation of the form:

$$q_1(i) \cdot c_1 \cdot b_i + q_2(i) \cdot c_2 \cdot b_{i+1} + \dots + q_n(i) \cdot c_n \cdot b_{i+n-1} = 0 \quad (7)$$

where $q_m(i)$ are polynomials in i of degree $n-1$. c_m Are constants.

We define b_{n-1} as the determinant of a matrix A

$$A = \begin{bmatrix} \frac{c_{n-1} \cdot q_{n-1}(0)}{c_n \cdot q_n(0)} & \dots & \frac{c_2 \cdot q_2(0)}{c_n \cdot q_n(0)} & \frac{c_1 \cdot q_1(0)}{c_n \cdot q_n(0)} & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & 0 & 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 & 1 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 0 & 0 & 0 & 0 & \dots & 1 & 1 \\ 0 & \dots & 0 & 1 & b_0 & 0 & 0 & \dots & 0 & 0 \\ 0 & \dots & 1 & 0 & 0 & -b_1 & 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 & 0 & b_2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & \dots & 0 & -1^{n-2} \cdot b_{n-2} \end{bmatrix} \quad (8)$$

and for the rest of the coefficients

$$b_{i+n-1} = \begin{bmatrix} \frac{-c_{n-1} \cdot q_{n-1}(i)}{c_n \cdot q_n(i)} & \frac{-c_{n-2} \cdot q_{n-2}(i)}{c_n \cdot q_n(i)} & \frac{-c_{n-3} \cdot q_{n-3}(i)}{c_n \cdot q_n(i)} & \dots & \frac{-c_1 \cdot q_1(i)}{c_n \cdot q_n(i)} & 0 & 0 & 0 & 0 & \dots \\ -1 & \frac{-c_{n-1} \cdot q_{n-1}(i-1)}{c_n \cdot q_n(i-1)} & \frac{-c_{n-2} \cdot q_{n-2}(i-1)}{c_n \cdot q_n(i-1)} & \dots & \frac{-c_2 \cdot q_2(i-1)}{c_n \cdot q_n(i-1)} & \frac{c_1 \cdot q_1(i-1)}{c_n \cdot q_n(i-1)} & 0 & 0 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & -1 & \frac{-c_{n-1} \cdot q_{n-1}(1)}{c_n \cdot q_n(1)} & \frac{c_{n-2} \cdot q_{n-2}(1)}{c_n \cdot q_n(1)} & \dots & \frac{c_1 \cdot q_1(1)}{c_n \cdot q_n(1)} & 0 & \dots \\ 0 & 0 & 0 & 0 & -1 & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix} \quad (9)$$

The series (6) can be proven to be convergent [2] if for a constant E

$$s \leq \frac{1}{E} \cdot |c_n| \quad (10)$$

and if all the absolute values of the coefficients of (1) are chosen smaller than 1. This is done by dividing a polynomial by (more than) the maximum of the absolute values of the coefficients.

To make the series convergent, we transform s to $e \cdot s$. That is, if we insert the powerseries it is not in s but in $e \cdot s$. Writing out the terms of the sum, we find that each term d_i has a factor $s^i \cdot e^{i-n+1}$. Setting

$$e = \frac{|c_n|}{E} \quad (12)$$

We still need $s < 1$, which is why we set s to a_0 and $a_0 < 1$.

References

- [1] R. Harley, On the theory of the Transcendental Solution of Algebraic Equations, Quart. Journal of Pure and Applied Math, Vol. 5 p.337. 1862
- [2] E. Kreyszig, Advanced Engineering Mathematics, John Wiley & Sons, Inc. p.789. 1993.