



Analytic cliffordian functions

Guy Laville, Eric Lehman

► To cite this version:

| Guy Laville, Eric Lehman. Analytic cliffordian functions. 2005. hal-00003213

HAL Id: hal-00003213

<https://hal.science/hal-00003213>

Preprint submitted on 4 Feb 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ANALYTIC CLIFFORDIAN FUNCTIONS

by

Guy LAVILLE and Eric LEHMAN
 Université de Caen - CNRS UMR 6139
 Laboratoire Nicolas ORESME
 14032 Caen France

e-mail : guy.laville@math.unicaen.fr ; lehman@math.unicaen.fr

Abstract

In classical function theory, a function is holomorphic if and only if it is complex analytic. For higher dimensional spaces it is natural to work in the context of Clifford algebras. The structures of these algebras depend on the parity of the dimension n of the underlying vector space. The theory of holomorphic Cliffordian functions reflects this dependence. In the case of odd n the space of functions is defined by an operator (the Cauchy-Riemann equation) but not in the case of even n . For all dimensions the powers of identity (z^n, x^n) are the foundation of function theory.

Key words : Non commutative analysis, Clifford analysis, analytic functions, holomorphic Cliffordian functions, iterated Laplacians.

AMS classification : 30 G 35, 15 A 66.

I. Introduction

A complex analytic function $f(z)$ may be defined as being locally the sum of a convergent power series $f(z) = \sum_{N=1}^{\infty} a_N z^{N-1}$ or as being holomorphic,

that is such that $\left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y}\right)f(x + iy) = 0$. A real analytic function $u(x)$ may be defined as being locally the sum of a convergent power series or by $u(x) = \overline{f(x + i0)}$ where f is a complex holomorphic function such that $f(\bar{z}) = \overline{f(z)}$. The main difference between the complex and the real case is the existence or non existence of a differential relation characterizing holomorphy.

We extend the definitions of analyticity and holomorphy to functions defined on Clifford algebras $\mathbb{R}_{0,n}$ distinguishing between the case of odd n , $n = 2m + 1$, and the case of even n , $n = 2m$. We show that the equivalence between analyticity and holomorphy still holds. The cases of odd n and even

n interrelate in a way that reflects the difference between the structures of the algebras $\mathbb{R}_{0,2m}$ and $\mathbb{R}_{0,2m+1}$. In particular the center of $\mathbb{R}_{0,2m}$ is \mathbb{R} although the center of $\mathbb{R}_{0,2m+1}$ is $\mathbb{R} \oplus \mathbb{R}e_{12\dots 2m+1}$, where $e_{12\dots 2m+1}$ is a pseudoscalar.

II. Notations

Let V_n be an anti-Euclidean vector space of dimension n . For any orthonormal basis e_1, \dots, e_n of V_n we have for all distinct i and j in $\{1, \dots, n\}$

$$e_i^2 = -1 \quad \text{and} \quad e_i e_j = -e_j e_i.$$

If $I \subset \{1, \dots, n\}$ and $I = \{i_1, \dots, i_k\}$ with $i_1 < \dots < i_k$ we set $e_I = e_{i_1} e_{i_2} \dots e_{i_k}$. For $I = \emptyset$, we set $e_\emptyset = e_0 = 1$. Then $(e_I)_{I \subset \{1, \dots, n\}}$ is a basis of the Clifford algebra $\mathbb{R}_{0,n}$ seen

as a real vector space. If $A = \sum_{I \subset \{1, \dots, n\}} A_I e_I$, with $A_I \in \mathbb{R}$, is an element of

$\mathbb{R}_{0,n}$ we call $A_0 = A_\emptyset$ the scalar part of A and denote it by $A_0 = S(A)$. Following the $\mathbb{R} - \mathbb{C}$ case and also Leutwiler and Eriksson-Bique [EL2], we introduce the decomposition

$$\mathbb{R}_{0,n} = \mathbb{R}_{0,n-1} \oplus e_n \mathbb{R}_{0,n-1}$$

(For convenience in our computations we have chosen $e_n \mathbb{R}_{0,n-1}$ instead of $\mathbb{R}_{0,n-1} e_n$). This decomposition means that given a vector e_n there are two maps from $\mathbb{R}_{0,n}$ to $\mathbb{R}_{0,n-1}$, denoted \mathcal{R} and \mathcal{J} , such that for any A in $\mathbb{R}_{0,n}$ we have

$$A = \mathcal{R}A + e_n \mathcal{J}A.$$

We have chosen notation \mathcal{R} and \mathcal{J} and the following notation for conjugation to stress the fact that for $n = 1$, we have $\mathbb{R}_{0,1} = \mathbb{C}$, $\mathbb{R}_{0,0} = \mathbb{R}$ which yield the usual relations between \mathbb{C} and \mathbb{R} . If $A \in \mathbb{R}_{0,n}$, we call the conjugate of A and denote by \overline{A} the element of $\mathbb{R}_{0,n}$ defined by

$$\overline{A} = \mathcal{R}A - e_n \mathcal{J}A$$

If z is a paravector, that is an element of $\mathbb{R} \oplus V_n$, we have

$$z = z_0 + z_1 e_1 + \dots + z_n e_n \quad \text{with} \quad z_0 \in \mathbb{R}, z_1 \in \mathbb{R}, \dots, z_n \in \mathbb{R}.$$

We denote by $|z|$ the positive real number such that $|z|^2 = z_0^2 + z_1^2 + \dots + z_n^2$ and by $x = \mathcal{R}z = z_0 + z_1 e_1 + \dots + z_{n-1} e_{n-1}$ the paravector in $\mathbb{R} \oplus V_{n-1}$ such that

$$z = x + z_n e_n \quad \text{and} \quad \overline{z} = x - z_n e_n.$$

We define z_* by : $z_* = z_0 - z_1 e_1 - \dots - z_n e_n$. Then $|z|^2 = zz_* = z_* z$.

We introduce the differential operators

$$D = \frac{\partial}{\partial z_0} + e_1 \frac{\partial}{\partial z_1} + \dots + e_n \frac{\partial}{\partial z_n}, \quad D_* = \frac{\partial}{\partial z_0} - e_1 \frac{\partial}{\partial z_1} - \dots - e_n \frac{\partial}{\partial z_n}$$

and

$$\Delta = DD_* = D_* D.$$

Note that Δ is the usual Laplacian.

If $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n) \in (\mathbb{N} \cup \{0\})^{n+1}$ is a multi-index we denote its length by $|\alpha| = \alpha_0 + \alpha_1 + \dots + \alpha_n$. The elementary multi-index ε_k is defined by $\varepsilon_k = (\delta_{k0}, \delta_{k1}, \dots, \delta_{kn})$ where δ_{ij} is the Kronecker symbol equal to 1 if $i = j$ and to 0 if $i \neq j$.

The order on the set of multi-indexes is the lexicographical order.

III. Analytic Cliffordian polynomials

Laville and Ramadanoff [LR1] have defined holomorphic Cliffordian polynomials for odd n . The same definitions can also be used for even n . We will see that these homogeneous polynomials are the building blocks of analytic Cliffordian functions in both cases. Therefore we call them analytic Cliffordian polynomials. Polynomials with the same structure were introduced by Heinz Leutwiler in [Le1] and also in [EL1]. Monogenic polynomials are particular cases of analytic Cliffordian polynomials [BDS], [DSS].

III.1. Three classes of analytic Cliffordian polynomials

Definition 1.a.- Let a be a paravector and $N \in \mathbb{N}$. We call elementary analytic monomial and denote by $M_N^a(z)$ the homogeneous monomial function of degree $N - 1$ defined by

$$M_N^a(z) = (az)^{N-1}a = a(za)^{N-1}.$$

Remark 1.- We may also define $M_N^a(z)$ for all integers by $M_0^a(z) = z^{-1}$ and for $N < 0$, $M_N^a(z) = M_{1-N}^{z^{-1}}(a^{-1})$. It is often convenient to write $\not{a}z \not{a}^{N-1}$ instead of $M_N^a(z)$. We have then for $N \in \mathbb{Z}$ and \sqrt{a} a paravector such that $(\sqrt{a})^2 = a$

$$\not{a}z \not{a}^N = \sqrt{a}(\sqrt{a}z\sqrt{a})^N \sqrt{a}.$$

These polynomials are close to similar ones in [Le2].

Remark 2.- $M_N^a(z) = |a|^N M_N^{a/|a|}(z) = |z|^{N-1} M_N^a(z/|z|)$.

Proposition 1.- $M_N^a(z) \in \mathbb{R} \oplus V_n$ and $|M_N^a(z)| = |a|^N |z|^{N-1}$.

Proof.- Computing aza explicitly, we get $aza \in \mathbb{R} \oplus V_n$ and $|aza| = |a|^2 |z|$. Note that

$$(*) \quad M_N^a(z) = aM_{N-1}^z(a)a.$$

Proposition 2.- $M_N^a(z) = D_* \frac{1}{N} S((az)^N)$.

Proof.- Let us define θ by $|a||z| \cos \theta = S(az)$, $0 \leq \theta \leq \pi$. Then $2S(az) = az + z_* a_*$ and $aza = (az + z_* a_*)a - z_* a_* a = 2S(az)a - |a|^2 z_* = 2|a||z| \cos \theta a - |a|^2 |z|^2 z^{-1}$. A simple recursion using (*) yields

$$M_N^a(z) = |a|^{N-1} |z|^{N-1} \frac{\sin N\theta}{\sin \theta} a - |a|^N |z|^N \frac{\sin(N-1)\theta}{\sin \theta} z^{-1}.$$

Since $2S((az)^N) = M_N^a(z)z + z_* M_N^a(z)_*$, we get

$$S((az)^N) = |a|^N |z|^N \cos(N\theta).$$

From the definitions of $|z|$ and θ , we get $D_*|z| = |z|z^{-1}$ and $D_*\theta = \frac{\cos \theta}{\sin \theta} z^{-1} - |a|^{-1} |z|^{-1} \frac{1}{\sin \theta} a$. Finally

$$D_* S(az)^N = N|a|^N |z|^N z^{-1} \cos(N\theta) - N|a|^N |z|^N \sin N\theta D_*\theta = NM_N^a(z).$$

Corollary.- $z^N = D_* \frac{1}{N+1} S(z^{N+1})$.

Proof.- Chose $a = e_0$ and replace N by $N+1$ in proposition 2.

Remark.- Let $T_N(x)$ and $U_N(x)$, $x \in \mathbb{R}$, be the classical Tchebycheff polynomials of the first and second kind. Recall that

$$T_N(x) = \cos(N \operatorname{Arc} \cos x)$$

$$U_N(x) = \frac{\sin((N+1) \operatorname{Arc} \cos x)}{\sin(\operatorname{Arc} \cos x)}.$$

Thus, when $|a|=1$, $|z|=1$, we get

$$S((az)^N) = T_N(S(az)) \text{ and } M_N^a(z) = U_{N-1}(S(az))a - U_{N-2}(S(az))z^{-1}.$$

Definition 1.b.- Let a_1, \dots, a_k be paravectors and let N_1, \dots, N_k be integers belonging to $\mathbb{N} \cup \{0\}$. We set $N = N_1 + \dots + N_k$ and denote

by $\mathcal{P}_{N_1, \dots, N_k}$ the set of partitions I of $\{1, \dots, N\}$ into a union of disjoint subsets $I = (I_1, \dots, I_k)$ such that $\text{Card } I_1 = N_1, \dots, \text{Card } I_k = N_k$. For $I \in \mathcal{P}_{N_1, \dots, N_k}$ and $\nu \in \{1, \dots, N\}$, we define b_ν^I by $b_\nu^I = a_j$ where j is the element of $\{1, \dots, k\}$ such that $\nu \in I_j$. We define the a_1, \dots, a_k symmetrical analytic homogeneous polynomial of degree $N - 1$ in z and N_j in a_j by

$$S_{N_1, \dots, N_k}^{a_1, \dots, a_k}(z) = \sum_{I \in \mathcal{P}_{N_1, \dots, N_k}} \left(\prod_{\nu=1}^{N-1} (b_\nu^I z) \right) b_N^I.$$

Proposition 3.- $S_{N_1, \dots, N_k}^{a_1, \dots, a_k}$ is a real linear combination of elementary analytic Cliffordian monomials of degree $N_1 + \dots + N_k - 1$.

Proof.- For any real λ , we have

$$M_N^{a+\lambda b}(z) = \sum_{p+q=N} \lambda^q S_{p,q}^{a,b}(z).$$

Choose $N + 1$ different values for λ ; one gets a Van der Monde matrix which is invertible. This shows the result for $k = 2$. We can iterate the same argument noting that for any real λ

$$S_{N_1, \dots, N_{k-1}, N_k}^{a_1, \dots, a_{k-1}, a_k + \lambda a_{k+1}}(z) = \sum_{p+q=N_k} \lambda^q S_{N_1, \dots, N_{k-1}, p, q}^{a_1, \dots, a_{k-1}, a_k, a_{k+1}}(z).$$

Definition 2.- For each multi-index $\alpha \in (\mathbb{N} \cup \{0\})^{n+1}$ we define Q_α as the homogeneous polynomial in z_0, \dots, z_n of degree $|\alpha| - 1$, by

$$Q_\alpha(z) = \partial_\alpha z^{2|\alpha|-1}$$

where ∂_α is the differential operator of order $|\alpha|$: $\partial_\alpha = \frac{\partial^{|\alpha|}}{\partial z_0^{\alpha_0} \partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}$.

Definition 3.- For each multi-index $\alpha \neq (0, 0, \dots, 0)$ we define the analytic Cliffordian polynomial P_α by

$$P_\alpha(z) = \sum_{\sigma \in S_\alpha} \left(\prod_{\nu=1}^{|\alpha|-1} (e_{\sigma(\nu)} z) \right) e_{\sigma(|\alpha|)}$$

where S_α is the set of maps σ from $\{1, \dots, |\alpha|\}$ to $\{0, 1, \dots, n\}$ such that $\text{Card}(\sigma^{-1}(\{k\})) = \alpha_k$ for all k in $\{0, 1, \dots, n\}$.

III.2. Relations among the $M_N^a(z)$, the $Q_\alpha(z)$ and the $P_\alpha(z)$

For any N in \mathbb{N} , the real linear space generated by the elementary analytic monomials M_N^a , the real linear space generated by the Q_α with $|\alpha| = N$ and the real linear space generated by the P_α with $|\alpha| = N$ are identical.

Proposition 4.- $Q_\alpha(z) = \partial_\alpha M_{2|\alpha|}^1(z)$.

Corollary.- For any multi-index α , $Q_\alpha(z) \in \mathbb{R} \oplus V_n$ and there exists a scalar polynomial $q_\alpha(z)$ homogeneous of degree $|\alpha|$ such that $Q_\alpha(z) = D_* q_\alpha(z)$.

Proof.- ∂_α and D_* commute and $M_{2|\alpha|}^1|z| = z^{2|\alpha|-1}$. Use proposition 1 and corollary of proposition 2.

Proposition 5.-

$$Q_\alpha(z) = k_\alpha P_\alpha(z) + \sum_{\substack{\alpha' > \alpha \\ |\alpha'| = |\alpha|}} \lambda_{\alpha\alpha'} P_{\alpha'}(z)$$

where

$$k_\alpha = \binom{2|\alpha| - 1}{\alpha_0} \alpha_0! \alpha_1! \dots \alpha_n!$$

and

$$\lambda_{\alpha\alpha'} \in \mathbb{Z}.$$

Proof.- Let $\beta = (\alpha_1, \dots, \alpha_n) \in (\mathbb{N} - \{0\})^n$ be the multi-index such that $\alpha = (\alpha_0, \beta)$. From the definition of Q_α follows :

$$Q_\alpha(z) = \binom{2|\alpha| - 1}{\alpha_0} \alpha_0! \frac{\partial^{|\beta|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}} z^{2|\beta| - 1 + \alpha_0}.$$

Consider $z^{2|\beta| - 1 + \alpha_0}$ as an explicit product of $2|\beta| - 1 + \alpha_0$ factors each equal to z , that is : $z^{2|\beta| - 1 + \alpha_0} = z \cdot z \cdot z \cdot \dots \cdot z$. To apply $\frac{\partial}{\partial x_k}$ is equivalent to replace each z once by e_k and to add all the products obtained. If we never derivate two successive z then we get $\alpha_1! \dots \alpha_n! P_\alpha(z)$. If we derivate two successive z , we get factors $e_k e_h$ which annihilate if $k \neq h$ because $e_k e_h = -e_h e_k$ and else -1 if $k = h$. Since $1 = e_0$ we get the terms of $P_{\alpha'}$ with $\alpha' > \alpha$ in the lexicographical order.

Corollary 1.- $P_\alpha(z) = k_\alpha^{-1} Q_\alpha(z) + \sum_{\substack{\alpha' > \alpha \\ |\alpha'| = |\alpha|}} \mu_{\alpha\alpha'} Q_{\alpha'}(z)$ where $\mu_{\alpha\alpha'} \in \mathbb{Q}$.

Corollary 2.- For any multi-index α , $P_\alpha(z) \in \mathbb{R} \oplus V_n$ and there exists a scalar polynomial $p_\alpha(z)$ homogeneous of degree $|\alpha|$ such that $P_\alpha(z) = D_* p_\alpha(z)$.

Proposition 6.- $M_N^a(z) = \sum_{|\alpha|=N} a^\alpha P_\alpha(z)$, where $a^\alpha := a_0^{\alpha_0} a_1^{\alpha_1} \dots a_n^{\alpha_n}$.

Proposition 7.- For $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n)$, we have $P_\alpha(z) = S_{\alpha_0, \alpha_1, \dots, \alpha_n}^{e_0, e_1, \dots, e_n}(z)$.

Proof.- The definition of P_α is the definition of S in which the k paravectors $a_1 \dots a_k$ are the $(n+1)$ elements of a basis of $S \oplus V : e_0, e_1, \dots, e_n$.

Corollary.- The real linear space generated by the P_α with $|\alpha| = N$ is independent of the basis e_1, \dots, e_n of V_n .

Remarks.- In the case of odd n , this is already known.

Note that generally the polynomials $P_\alpha(z)$ with $|\alpha| = N$ are not \mathbb{R} -linearly independent. For example, if $n = 3$ and $N = 4$ we have

$$3P_{4000} + 3P_{0400} + 3P_{0040} + 3P_{0004} + P_{2200} + P_{2020} + P_{2002} + P_{0220} + P_{0202} + P_{0022} = 0.$$

III.3. Some properties of the polynomials P_α

Proposition 8.- For $\alpha = (\alpha_0, \dots, \alpha_n)$, the polynomial $p_\alpha(z)$ is even in z_k if α_k is even and odd in z_k if α_k is odd.

Proof.- We know that $P_\alpha(z) \in \mathbb{R} \oplus V_n$. Then we can write

$$P_\alpha(z) = A_0(z) + A_1(z)e_1 + \dots + A_n(z)e_n.$$

Suppose α_k even. $P_\alpha(z)$ is a sum of terms like

$$e_{i_1} z e_{i_2} z \dots z e_{i_{|\alpha|}}$$

in which e_k occurs α_k times, that is an even number of times. Write $z = z_0 + z_1 e_1 + \dots + z_n e_n$ and develop all the terms. The terms which contribute to $A_k(z)e_k$ must contain an odd number of times the vector e_k and then $z_k e_k$ has to appear an odd number of times. $A_k(z)$ is then a sum of terms which are all odd in z_k and $A_k(z)$ is odd in z_k . Since $P_\alpha(z) = D_* p_\alpha(z)$, we have $A_k(z) = -\frac{\partial}{\partial z_k} p_\alpha(z)$. Since $p_\alpha(z)$ is homogeneous, we can conclude that $p_\alpha(z)$ is even in z_k .

If α_k is odd the proof is the same.

Examples : $p_{(2,0,1,0)}(z) = (3z_0^2 - z_1^2 - z_2^2 - z_3^2)z_2$ is even in z_0, z_1 and z_3 and odd in z_2 . $p_{(1,1,1,0)}(z) = 8z_0 z_1 z_2$ is odd in z_0, z_1 and z_2 and even in z_3 .

Corollary.- If α_n is even, then $P_\alpha(\bar{z}) = \overline{P_\alpha(z)}$; if α_n is odd then $P_\alpha(\bar{z}) = -\overline{P_\alpha(z)}$.

Proof.- Let us write

$$P_\alpha(z) = A_0(z) + A_1(z)e_1 + \dots + A_{n-1}(z)e_{n-1} + A_n(z)e_n.$$

If α_n is even, then $p_\alpha(z)$ is even in z_n and the polynomials $A_k = \pm \frac{\partial}{\partial z_k} p_\alpha(z)$ for $k \neq n$, are all even in z_n , but A_n is odd in z_n . Thus

$$P_\alpha(\bar{z}) = A_0(z) + A_1(z)e_1 + \dots + A_{n-1}(z)e_{n-1} - A_n(z)e_n = \overline{P_\alpha(z)}.$$

If α_n is odd, $A_0, A_1 \dots A_{n-1}$ are odd in z_n and A_n is even in z_n , so that

$$P_\alpha(\bar{z}) = -A_0(z) - A_1(z)e_1 - \dots - A_{n-1}(z)e_{n-1} + A_n(z)e_n = -\overline{P_\alpha(z)}.$$

Remark.- $\frac{\partial}{\partial z_i} P_\alpha(z) = e_i^2 \left\{ 2|\alpha| P_{\alpha-\varepsilon_i}(z) - (\alpha_i + 1) \sum_{k=0}^n P_{\alpha+\varepsilon_i-2\varepsilon_k}(z) \right\}.$

IV. Analytic Cliffordian functions and holomorphic Cliffordian functions

Definition 1.- Let Ω be a domain of $\mathbb{R} \oplus V_n$ and $f : \Omega \rightarrow \mathbb{R}_{0,n}$. We say that f is a left analytic Cliffordian function if any ω in $\mathbb{R} \oplus V_n$ has a neighbourhood Ω_ω in Ω such that for any z in Ω_ω , $f(z)$ is the sum of a convergent series

$$f(z) = \sum_{N=1}^{\infty} \sum_{a \in A_N} M_N^a(z - \omega) C_a$$

where for each N in \mathbb{N} , A_N is a finite subset of $\mathbb{R} \oplus V_n$, for each a in A_N , $C_a \in \mathbb{R}_{0,n}$ and $\sum_{N=1}^{\infty} \sum_{a \in A_N} |a|^N |z - \omega|^{N-1} |C_a|$ is convergent in Ω_ω .

Remark 1.- The relation $M_N^a(z - \omega) = \sum_{p+q=N} (-1)^q S_{p,q}^{a, a\omega a}(z)$ and the proposition 3 prove the consistency of the definition with respect to translations. Consequently we will restrict ourselves to the case $\omega = 0$.

Remark 2.- The above definition is obviously intrinsic, but we get an equivalent definition if we replace the monomials M_N^a by the polynomials P_α ; the function $f : \mathbb{R} \oplus V_n \rightarrow \mathbb{R}_{0,n}$ is left analytic Cliffordian in a neighbourhood Ω of 0 if for every z in Ω , $f(z)$ is the sum of a convergent series

$$f(z) = \sum_{N=1}^{\infty} \sum_{|\alpha|=N} P_\alpha(z) c_\alpha$$

where α are multi-indexes belonging to $(\{0\} \cup \mathbb{N})^{1+n}$, and for each α we have $c_\alpha \in \mathbb{R}_{0,n}$ and $\sum_{N=1}^{\infty} \sum_{|\alpha|=N} |P_\alpha(z)| |c_\alpha|$ is convergent.

Definition 2.- Let Ω be a domain in $\mathbb{R} \oplus V_n$. A fonction $u : \mathbb{R} \oplus V_n \rightarrow \mathbb{R}_{0,n}$ is called a left holomorphic Cliffordian function

- (i) for odd n , if $D\Delta^m u = 0$ where $m = (n-1)/2$
- (ii) for even n , if for any $\omega \in \Omega$ a neighbourhood Λ_ω in $\mathbb{R} \oplus V_{n+1}$ and a left holomorphic Cliffordian function f defined on Λ_ω exist such that
 - for all z in Λ_ω : \bar{z} is in Λ_ω and $f(\bar{z}) = \overline{f(z)}$
 - for all x in $\Lambda_\omega \cap (\mathbb{R} \oplus V_n)$, $u(x) = f(x)$.

$$\mathbb{R}e_n$$

$$\mathbb{R} \oplus V_n$$

$$\mathbb{R} \oplus V_{n+1}$$

Theorem.- Let Ω be a domain of $\mathbb{R} \oplus V_n$. A function $f : \Omega \rightarrow \mathbb{R}_{0,n}$ is left analytic Cliffordian if and only if it is left holomorphic Cliffordian.

For odd n , the theorem has already been proven in [LR1]. Let n be even, $n = 2m$. Let u be a left analytic Cliffordian function on a neighbourhood Ω of 0 and let $S_n(r)$ be a sphere of center 0 and radius $r > 0$ included in Ω . For $|x| < r$ we have

$$u(x) = \sum_{N=1}^{\infty} \sum_{a \in A_N} M_N^a(x) C_a$$

where $\sum_{N=1}^{\infty} \sum_{a \in A_N} |a|^N |x|^{N-1} |C_a|$ is convergent. Chose $\Lambda_0 = S_{n+1}(r)$ the interior of the sphere of center 0 and radius r in $\mathbb{R} \oplus V_{n+1}$ and let $f : S_{n+1}(r) \rightarrow \mathbb{R}_{0,n+1}$ be defined by

$$f(z) = \sum_{N=1}^{\infty} \sum_{a \in A_N} M_N^a(z) C_a.$$

Then f is left holomorphic Cliffordian on $S_{n+1}(r)$ and $f(x) = u(x)$. Since $a \in \mathbb{R} \oplus V_n$, one gets $M_N^a(\bar{z}) = \overline{M_N^a(z)}$. And since $C_a \in \mathbb{R} \oplus V_n$, we have $f(\bar{z}) = \overline{f(z)}$.

Conversely, let f be a left holomorphic Cliffordian function defined on a neighbourhood Λ_0 of 0 in $\mathbb{R} \oplus V_{n+1}$ such that $f(\bar{z}) = \overline{f(z)}$. We want to show that $u : \Lambda_0 \cap \mathbb{R} \oplus V_n \rightarrow \mathbb{R}_{0,n}$, $x \mapsto u(x) = f(x)$ is left analytic Cliffordian.

Since $n+1$ is odd, f is analytic Cliffordian and we can write

$$f(z) = \sum_{N=1}^{\infty} \sum_{|\beta|=N} P_{\beta}(z) c_{\beta} = D_* \sum_{N=1}^{\infty} \sum_{|\beta|=N} p_{\beta}(z) c_{\beta}.$$

Let H_N be the real linear space of scalar homogeneous polynomials in z_0, \dots, z_n, z_{n+1} of total degree N and $(m+1)$ -harmonic generated by $(p_{\beta})_{|\beta|=N}$ (the dimension of H_N is $C_{N+n}^n - C_N^n$). We can extract a subset B_N of $\{\beta \in (\{0\} \cup \mathbb{N})^{n+2} / |\beta| = N\}$ such that $(p_{\beta})_{\beta \in B_N}$ is a basis of H_N . D_* is a linear map from H_N to $\mathbb{R} \otimes V_{n+1}$ and $\text{Ker } D_*$ is a subspace of H_N . Let $(\psi_j)_{j \in \mathcal{J}}$ be a basis of $\text{Ker } D_*$; since $(p_{\beta})_{\beta \in B_N}$ is a basis of H_N there is a subset B_N° of B_N such that $((\psi_j)_{j \in \mathcal{J}}, (p_{\beta})_{\beta \in B_N^{\circ}})$ is a basis of H_N . Then $((\psi_j \otimes e_I)_{j \in \mathcal{J}, I \subset \{1, \dots, n+1\}}, (p_{\beta} \otimes e_I)_{\beta \in B_N^{\circ}, I \subset \{1, \dots, n+1\}})$ is a basis of the real linear space $H_N \otimes \mathbb{R}_{0,n+1}$, and there are unique real numbers $\theta_{j,I}$ and $d_{\beta,I}$ such that

$$\sum_{|\beta|=N} p_{\beta}(z) c_{\beta} = \sum_{j \in \mathcal{J}} \sum_{I \subset \{1, \dots, n+1\}} \theta_{j,I} \psi_j(z) e_I + \sum_{\beta \in B_N^{\circ}} \sum_{I \subset \{1, \dots, n+1\}} d_{\beta,I} p_{\beta}(z) e_I.$$

Let us write $d_{\beta} = \sum_I d_{\beta,I} e_I$, using homogeneity we get for any analytic cliffordian function f the existence and unicity of the coefficients d_{β} in $\mathbb{R}_{0,n+1}$ such that

$$f(z) = \sum_{N=1}^{\infty} \sum_{\beta \in B_N^{\circ}} P_{\beta}(z) d_{\beta}.$$

Let $B_N^{\circ+}$ be the set of multi-indexes $\beta = (\beta_0, \dots, \beta_n, \beta_{n+1})$ where $\beta \in B_N^{\circ}$ and β_{n+1} is even and $B_N^{\circ-} = B_N^{\circ} - B_N^{\circ+}$. If $\beta \in B_N^{\circ+}$, the corollary of proposition 8 implies $P_{\beta}(\bar{z}) = \overline{P_{\beta}(z)}$ and if $\beta \in B_N^{\circ-}$ we have $P_{\beta}(\bar{z}) = -\overline{P_{\beta}(z)}$. The relation $f(\bar{z}) = \overline{f(z)}$ becomes then

$$\sum_{N=1}^{\infty} \left\{ \sum_{\beta \in B_N^{\circ+}} \overline{P_{\beta}(z)} d_{\beta} - \sum_{\beta \in B_N^{\circ-}} \overline{P_{\beta}(z)} d_{\beta} \right\} = \sum_{N=1}^{\infty} \left\{ \sum_{\beta \in B_N^{\circ+}} \overline{P_{\beta}(z)} \bar{d}_{\beta} + \sum_{\beta \in B_N^{\circ-}} \overline{P_{\beta}(z)} \bar{d}_{\beta} \right\}$$

By homogeneity we get

$$\sum_{\beta \in B_N^{\circ+}} \overline{P_{\beta}(z)} (\bar{d}_{\beta} - d_{\beta}) + \sum_{\beta \in B_N^{\circ-}} \overline{P_{\beta}(z)} (\bar{d}_{\beta} + d_{\beta}) = 0$$

and by conjugation

$$\sum_{\beta \in B_N^{\circ+}} P_{\beta}(z) (d_{\beta} - \bar{d}_{\beta}) + \sum_{\beta \in B_N^{\circ-}} P_{\beta}(z) (d_{\beta} + \bar{d}_{\beta}) = 0.$$

By unicity of the coefficients of the P_β for $\beta \in B_N^\circ$, we get $d_\beta = \bar{d}_\beta$ if $\beta \in B_N^{\circ+}$ and $\bar{d}_\beta = -d_\beta$ if $\beta \in B_N^{\circ-}$. Let us write $a_\beta = d_\beta = \bar{d}_\beta$ if $\beta \in B_N^{\circ+}$ and $e_{n+1}b_\beta = d_\beta = -\bar{d}_\beta$ if $\beta \in B_N^{\circ-}$, we get

$$f(z) = \sum_{N=1}^{\infty} \left\{ \sum_{\beta \in B_N^{\circ+}} P_\beta(z) a_\beta + \sum_{\beta \in B_N^{\circ-}} P_\beta(z) e_{n+1} b_\beta \right\},$$

where $a_\beta \in \mathbb{R}_{0,n}$ and $b_\beta \in \mathbb{R}_{0,n}$. The two following lemmas will then prove the theorem.

Lemma 1.- Let $n = 2m$ and $\beta = (\beta_0, \dots, \beta_n, \beta_{n+1})$. If β_{n+1} is even then the restriction of P_β to $\mathbb{R} \oplus V_n$ is analytic Cliffordian from $\mathbb{R} \oplus V_n$ to $\mathbb{R}_{0,n}$.

Proof.- First we show that if $x \in \mathbb{R} \oplus V_n$ then $P_\beta(x) \in \mathbb{R}_{0,n}$ or better $P_\beta(x) \in \mathbb{R} \oplus V_n$. We know that $P_\beta(z) = D_* p_\beta(z)$ in $\mathbb{R}_{0,n+1}$ and $p_\beta(z)$ is even in z_{n+1} since β_{n+1} is even. So $\left[\frac{\partial}{\partial z_{n+1}} p_\beta(z) \right]_{z_{n+1}=0} = 0$ and $P_\beta(x) \in \mathbb{R} \oplus V_n$ for $x = \mathcal{R}z$.

Secondly we show that $P_\beta(x)$ is analytic Cliffordian. We know that

$$P_\beta(x) = \sum_{\sigma \in S_\beta} \left(\prod_{\nu=1}^n (e_{\sigma(\nu)} x) \right) e_{\sigma(n+1)},$$

which means that $P_\beta(x)$ is the sum of all different polynomials deduced from

$$(*) \quad \underbrace{e_0 x e_0 \dots e_0}_{\beta_0 \text{ times } e_0} x \underbrace{e_1 x e_1 \dots e_1}_{\beta_1 \text{ times } e_1} x e_2 x e_2 \dots e_n x \underbrace{e_{n+1} x e_{n+1} x \dots x e_{n+1}}_{\beta_{n+1} \text{ times } e_{n+1}}$$

by permutations of the e'_i s. Note that

$$e_{n+1} = (-1)^m e_{12\dots n e_{12\dots n+1}}$$

and that the pseudo scalar $e_{12\dots 2m 2m+1}$ belongs to the center of $\mathbb{R}_{0,2m+1}$. Since β_{n+1} is even and since $(e_{12\dots n+1})^2 \in \{1, -1\}$, we deduce that up to a sign we can replace (*) by :

$$(**) \quad \underbrace{e_0 x e_0 \dots e_0}_{\beta_0 \text{ times } e_0} x \underbrace{e_1 x e_1 \dots e_1}_{\beta_1 \text{ times } e_1} x e_2 x e_2 \dots e_n x \underbrace{e_{12\dots n} x e_{12\dots n} \dots x e_{12\dots n}}_{\beta_{n+1} \text{ times } e_{12\dots n}}$$

Let us chose a basis in V_n such that $x = x_0 e_0 + x_1 e_1$. Then e_1 commutes with x and for $i \geq 2$, $e_i e_{i+1}$ commutes with e_1 , x and of course e_0 . Suppose $\beta_2 > 0$, for each term of the form

$$A e_2 B e_{12\dots n} C$$

we have another term equal to $Ae_{12\dots n}Be_nC$. But then the commutation rules give since $n = 2m$

$$Ae_2Be_{12\dots n}C + Ae_{1e\dots n}Be_2C = 0.$$

Thus $\beta_2 = 0$. And similarly $\beta_3 = \dots = \beta_n = 0$. We get :

$$P_\beta(x) = \sum_{\sigma \in S_\beta} \left(\prod_{\nu=1}^{|\beta|-1} E_{\sigma(\nu)} x \right) E_{\sigma(|\beta|)}$$

where $E_0 = e_0$, $E_1 = e_1$ and $E_{n+1} = e_{12\dots n}$. Commuting systematically e_1 and $e_3e_4, e_5e_6, \dots, e_{n-1}e_n$ from E_{n+1} to the right

$$P_\beta(x) = \left(\sum_{\sigma \in S_\alpha} \left(\prod_{\nu=1}^{|\alpha|-1} e_{\sigma(\nu)} x \right) e_{\sigma(|\alpha|)} \right) (e_1)^{\beta_{n+1}} (e_{34\dots n})^{\beta_{n+1}}$$

where $\alpha = (\beta_0, \beta_1, \beta_{n+1}, 0, \dots, 0) \in (\{0\} \cup \mathbb{N})^{n+1}$. Thus $P_\beta(x) = \pm P_\alpha(x)$ with P_α analytic Cliffordian.

Lemma 2.- Let $n = 2m$ and $\beta = (\beta_0, \dots, \beta_n, \beta_{n+1})$. If β_{n+1} is odd then the restriction of $P_\beta e_{n+1}$ to $\mathbb{R} \oplus V_n$ is analytic Cliffordian from $\mathbb{R} \oplus V_n$ to $\mathbb{R}_{0,n}$.

Proof.- Now $p_\beta(z)$ is odd in z_{n+1} so $\left[\frac{\partial}{\partial z_i} p_\beta(z) \right]_{z_{n+1}=0} = 0$ for $i = 0, \dots, n$ and $P_\beta(x) = - \left[\frac{\partial}{\partial z_{n+1}} p_\beta(z) \right]_{z_{n+1}=0} e_{n+1}$ and $P_\beta(x)e_{n+1}$ is a scalar. For the second part of the proof we reduce the sum

$$P_\beta(x)e_{n+1} = \sum_{\sigma \in S_\beta} \left(\prod_{\nu=1}^{|\beta|-1} e_{\sigma(\nu)} x \right) e_{\sigma(|\beta|)} e_{n+1}$$

as in Lemma 1.

Corollary 1.- The space of left analytic Cliffordian functions is an \mathbb{R} -vector space and an $\mathbb{R}_{0,n}$ -right module, closed relatively to scalar derivations.

To deduce corollary 2 and 3 from the above theorem, the following lemma is convenient.

Lemma 3.- If $v \in \mathbb{R} \oplus V_{2m}$, then $\sum_{i=0}^{2m} e_i v e_i = (1 - 2m)v_*$.

Corollary 2.- If f is left analytic Cliffordian, then Df is also left analytic Cliffordian.

Proof.- If n is odd let $n = 2m + 1$. Since f is left analytic Cliffordian it is left holomorphic Cliffordian and $D\Delta^m f = 0$. But since D commutes with D and Δ we have

$$D\Delta^m(Df) = D(D\Delta^m f) = 0.$$

Thus Df is left holomorphic Cliffordian or left analytic Cliffordian.

If n is even let $n = 2m$. Let us write $x = z_0 + z_1 e_1 + \dots + z_{2m} e_{2m}$ and $D' = \sum_{i=0}^{2m} e_i \frac{\partial}{\partial z_i}$. Then we have $D = D' + e_{2m+1} \frac{\partial}{\partial z_{2m+1}}$.

We want to show that if u is left analytic Cliffordian then $D'u$ is also left analytic Cliffordian. In fact, we need only to show this for $u(x) = M_N^a(x)$ for any $a \in \mathbb{R} \oplus V_{2m}$ and any $N \in \mathbb{N}$. A straightforward computation gives

$$D' M_N^a(x) = \sum_{k=1}^{N-1} \sum_{i=0}^{2m} e_i M_k^a(x) e_i M_{N-k}^a(x)$$

and since $M_k^a(x) \in \mathbb{R} \oplus V_{2m}$, lemma 3 gives us

$$D' M_N^a(x) = -(2m-1) \sum_{k=1}^{N-1} [M_k^a(x)]_* M_{N-k}^a(x).$$

If N is odd, let $N = 2M + 1$; we have

$$D' M_{2M+1}^a(x) = -2(2m-1) \sum_{k=1}^M |a|^{2k} |x|^{2k-2} S((xa)^{2M-2k+1}).$$

If N is even, let $N = 2M$; we have

$$D' M_{2M}^a(x) = -(2m-1) \left\{ |a|^{2M} |x|^{2M-2} + 2 \sum_{k=1}^M |a|^{2k} |x|^{2k-2} S((xa)^{2M-2k}) \right\}.$$

The same computations in $\mathbb{R}_{0,2m+1}$ give

$$D M_{2M+1}^a(z) = -2(2m) \sum_{k=1}^M |a|^{2k} |z|^{2k-2} S((za)^{2M-2k+1})$$

and

$$D M_{2M}^a(z) = -(2m) \left\{ |a|^{2M} |z|^{2M-2} + 2 \sum_{k=1}^M |a|^{2k} |z|^{2k-2} S((za)^{2M-2k}) \right\}.$$

Then we have for any N

$$D' M_N^a(x) = \frac{2m-1}{2m} [D M_N^a(z)]_{z=x}.$$

Since $a \in \mathbb{R}_{0,2m}$, we have $S((za)^{N-2k}) = S((xa)^{N-2k})$, and since $|\bar{z}| = |z|$, we have $D M_N^a(\bar{z}) = D M_N^a(z) = \overline{D M_N^a(z)}$. The theorem then proves that $D' M_N^a(x)$ is left analytic Cliffordian.

Corollary 3.- If f is left analytic Cliffordian, then $D_* f_*$ is also left analytic Cliffordian.

Proof.- Let us write f as

$$f(z) = \sum_{N=1}^{\infty} \sum_{|\alpha|=N} P_{\alpha}(z) c_{\alpha} = \sum_{N=1}^{\infty} \sum_{|\alpha|=N} D_* p_{\alpha}(z) c_{\alpha}$$

and define the left analytic Cliffordian function \tilde{f} by

$$\tilde{f}(z) = \sum_{N=1}^{\infty} \sum_{|\alpha|=N} P_{\alpha}(z) c_{\alpha*}.$$

Then we have

$$D_* f_*(z) = (Df(z))_* = \sum_{N=1}^{\infty} \sum_{|\alpha|=N} \Delta p_{\alpha}(z) c_{\alpha*} = D \tilde{f}(z).$$

By corollary 1, $D \tilde{f}$ is left analytic Cliffordian, thus $D_* f_*$ is also left analytic Cliffordian.

V. Cauchy's problem and boundary data

V.1. Aim

We intend to generalize to Clifford algebras of any dimensions the method of extending real analytic functions into complex holomorphic functions. Let Ω be a domain of $\mathbb{R} \oplus V_{2m}$ and $u : \Omega \rightarrow \mathbb{R}_{0,2m}$. The Cauchy problem

$$\begin{cases} D \Delta^m f = 0 \\ f|_{\Omega} = u \end{cases}$$

where the unknown function f has to be defined on an open set Λ containing Ω , seems not well defined since the partial differential equations are of order

$2m + 1$. The Cauchy-Kowalewski theorem tells us that we need the normal derivatives of f to $\mathbb{R} \oplus V_{2m}$ up to the order $2m$. We will see that the algebraic and analytic properties of f and u enable us to compute these derivatives uniquely given the function u .

We will denote by \mathcal{A}_{2m} the linear space of left analytic Cliffordian functions defined on Ω and taking their values in $\mathbb{R}_{0,2m}$. Similarly, \mathcal{A}_{2m+1} is the linear space of left analytic Cliffordian functions defined on Λ and with values in $\mathbb{R}_{0,2m+1}$.

V.2. The operator $(A \mid \nabla_n)$

Definition.- Let $A \in \mathbb{R}_{0,n}$. We define $(A \mid \nabla_n) : \mathcal{A}_n \longrightarrow \mathcal{A}_n$ by:

- (i) $\forall a \in \mathbb{R} \oplus V_n \quad \forall N \in \mathbb{N} \quad (A \mid \nabla_n)M_N^a(z) = \sum_{k=1}^{N-1} M_k^a(z) A M_{N-k}^a(z)$
- (ii) $\forall f \in \mathcal{A}_n \quad \forall K \in \mathbb{R}_{0,n} \quad (A \mid \nabla_n)(f(z)K) = (A \mid \nabla_n)f(z)K$
- (iii) $(A \mid \nabla_n)$ is \mathbb{R} -linear and continuous.

Consequence.- If $f(z) = \sum_{N=1}^{\infty} \sum_{a \in A_n} M_N^a(z) C_a$, then

$$(A \mid \nabla_n)f(z) = \sum_{N=1}^{\infty} \sum_{a \in A_N} ((A \mid \nabla_n)M_N^a(z)) C_a.$$

Proposition 9.- If $f \in \mathcal{A}_n$, then $\frac{\partial}{\partial z_k} f(z) = (e_k \mid \nabla_n)f(z)$.

Proof.- We need only to verify the proposition for $f(z) = M_N^a(z)$. We have

$$\frac{\partial}{\partial z_k} a z a z \dots z a = a e_k a z \dots z a + a z a e_k \dots z a + \dots + a z a z \dots e_k a = (e_k \mid \nabla_n) a z a z \dots z a.$$

Proposition 10.- If $f \in \mathcal{A}_n$ and if λ belongs to the center of $\mathbb{R}_{0,n}$ then

$$(\lambda A \mid \nabla_n)f = \lambda(A \mid \nabla_n)f.$$

Lemma.- If $v \in \mathbb{R} \oplus V_{2m}$, then

$$\sum_{i=0}^{2m} e_i v e_i = (-1)^m (1 - 2m) e_{1 \dots 2m} v e_{1 \dots 2m}.$$

Proof.- Both sides of the equality are equal to $(1 - 2m)v_*$.

Proposition 11.- If $u \in \mathcal{A}_{2m}$, then

$$e_{12\dots 2m}(e_{12\dots 2m} \mid \nabla_{2m})u = \frac{(-1)^{m+1}}{2m-1} Du.$$

Proof.- We need only verify the relation for $u(x) = M_N^a(x)$. Using the definition and the lemma, we get

$$\begin{aligned} (-1)^m(1-2m)e_{12\dots 2m}(e_{12\dots 2m} \mid \nabla_{2m})u &= \sum_{k=1}^{N-1} \sum_{i=0}^{2m} e_i M_k^a(x) e_i M_{N-k}^a(x) \\ &= \sum_{i=0}^{2m} e_i \sum_{k=1}^{N-1} M_k^a(x) e_i M_{N-k}^a(x) \\ &= \sum_{i=0}^{2m} e_i \frac{\partial}{\partial x_i} M_N^a(x) = D M_N^a(x). \end{aligned}$$

Proposition 12.- For $\mu \in \{0\} \cup \mathbb{N}$, we have

$$\begin{aligned} (e_{12\dots 2m} \mid \nabla_{2m})^{2\mu} u &= \frac{(-1)^{m\mu}}{(2m-1)^{2\mu}} \Delta^\mu u \\ (e_{12\dots 2m} \mid \nabla_{2m})^{2\mu+1} u &= \frac{(-1)^{1+m\mu}}{(2m-1)^{2\mu+1}} e_{12\dots 2m} D \Delta^\mu u. \end{aligned}$$

Proof.- From proposition 3, we get

$$\begin{aligned} (e_{12\dots 2m} \mid \nabla_{2m})u &= \frac{-1}{2m-1} e_{12\dots 2m} Du \\ &= \frac{-1}{2m-1} D_* u_* e_{12\dots 2m} \end{aligned}$$

The corollary 2 of the theorem of last paragraph shows that $D_* u_* e_{12\dots 2m}$ is left analytic Cliffordian. Thus we may apply the operator as many times as we want. We get

$$\begin{aligned} ((e_{12\dots 2m} \mid \nabla_{2m})^2 u &= \frac{-1}{2m-1} e_{12\dots 2m} D \left\{ \frac{-1}{2m-1} D_* u_* e_{12\dots 2m} \right\} \\ &= \frac{1}{(2m-1)^2} e_{12\dots 2m} \Delta u_* e_{12\dots 2m} \\ &= \frac{1}{(2m-1)^2} (e_{12\dots 2m})^2 \Delta u \\ &= \frac{(-1)^m}{(2m-1)^2} \Delta u. \end{aligned}$$

A simple recursion gives then the general formulae.

V.3. Analytic extention

Theorem.- Let Ω be a domain of $\mathbb{R} \oplus V_{2m}$ and let $u : \Omega \rightarrow \mathbb{R}_{0,2m}$ be left analytic Cliffordian. There exist a domain Λ in $\mathbb{R} \oplus V_{2m+1}$ with $\Omega \subset \Lambda$ and a unique left holomorphic Cliffordian function f defined on $\Lambda \subset \mathbb{R} \oplus V_{2m+1}$, such that $f|_{\Omega} = u$. That function f is such that $f(\bar{z}) = \overline{f(z)}$ and if we denote by $\frac{\partial}{\partial n}$ the normal derivative to Ω , we have for any μ in $\{0\} \cup \mathbb{N}$

$$\begin{aligned} \left(\frac{\partial}{\partial n}\right)^{2\mu} f|_{\Omega} &= \frac{(-1)^{\mu}}{(2m-1)^{2\mu}} \Delta^{\mu} u \\ \left(\frac{\partial}{\partial n}\right)^{2\mu+1} f|_{\Omega} &= \frac{(-1)^{\mu+1}}{(2m-1)^{2\mu+1}} e_{2m+1} D\Delta^{\mu} u. \end{aligned}$$

Proof.- 1°) Suppose f is a solution. We use the usual notations

$$z = z_0 + z_1 e_1 + \dots + z_{2m} e_{2m} + z_{2m+1} e_{2m+1} = x + z_{2m+1} e_{2m+1}.$$

Thus we have

$$\left(\frac{\partial}{\partial n}\right)^j f = \left(\frac{\partial}{\partial z_{2m+1}}\right)^j f.$$

Since f is left analytic Cliffordian, we have $\frac{\partial}{\partial z_{2m+1}} f = (e_{2m+1} | \nabla_{2m+1}) f$ and thus

$$\left(\frac{\partial}{\partial n}\right)^j f = (e_{2m+1} | \nabla_{2m+1})^j f.$$

Note that $e_{2m+1} = (-1)^m e_1 2 \dots 2m 2m+1 e_1 2 \dots 2m$ and that $(-1)^m e_1 2 \dots 2m 2m+1$ belongs to the center of $\mathbb{R}_{0,2m+1}$. Using proposition 10 we get

$$\left(\frac{\partial}{\partial n}\right)^j f = ((-1)^m e_1 2 \dots 2m 2m+1)^j (e_1 2 \dots 2m | \nabla_{2m+1})^j f.$$

Taking the restriction to $z = x \in \Omega$, we get

$$\left(\frac{\partial}{\partial n}\right)^j f|_{\Omega}(x) = (-1)^{mj} (e_1 2 \dots 2m 2m+1)^j (e_1 2 \dots 2m | \nabla_{2m})^j u(x).$$

Proposition 12 gives us then for $j = 2\mu$ and $j = 2\mu + 1$

$$\left(\frac{\partial}{\partial n}\right)^{2\mu} f|_{\Omega}(x) = \frac{(-1)^{\mu}}{(2m-1)^{2\mu}} \Delta^{\mu} u(x)$$

and

$$\left(\frac{\partial}{\partial n}\right)^{2\mu+1} f|_{\Omega}(x) = \frac{(-1)^{\mu+1}}{(2m-1)^{2\mu+1}} e_{2m+1} D\Delta^{\mu} u(x).$$

2°) Using the above conditions for $j \leq 2m$, the theorem of Cauchy-Kowalewski proves the existence of $\Lambda' \subset \mathbb{R} \oplus V_{2m+1}$ with $\Lambda' \cap (\mathbb{R} \oplus V_{2m}) = \Omega$ and the existence and unicity of f in Λ' .

3°) Knowing the existence of f , the usual Taylor formula gives us

$$f(z) = \sum_{\mu=0}^{\infty} \frac{(-1)^{\mu}}{(2m-1)^{2\mu}(2\mu)!} (z_{2m+1})^{2\mu} \Delta^{\mu} u(x) \\ + e_{2m+1} \sum_{\mu=0}^{\infty} \frac{(-1)^{\mu+1}}{(2m-1)^{2\mu+1}(2\mu+1)!} (z_{2m+1})^{2\mu+1} D\Delta^{\mu} u(x).$$

This formula shows, since $Du(x) \in \mathbb{R}_{0,2m}$, that $f(\bar{z}) = \overline{f(z)}$ in a subdomain Λ of $\mathbb{R} \oplus V_{2m+1}$ such that $\Omega \subset \Lambda \subset \Lambda'$.

VI. Fueter's method

Fueter's method is well known and widely used to construct functions connected to monogenic functions [Fu], [De], [Qi]. It is known to be effective to construct holomorphic Cliffordian functions in the case of odd n . We show that it is still valid for our definition in the case of even n .

Theorem.- Let φ be a complex holomorphic function defined on D_{φ} an open subset of the upper half-plane and let p and q be the real functions of two variables defined by

$$\forall \zeta = \xi + i\eta \in D_{\varphi} \quad \varphi(\zeta) = p(\xi, \eta) + iq(\xi, \eta).$$

Let $\vec{z} = z_1 e_1 + \dots + z_n e_n$, for $z_0 + i |\vec{z}| \in D_{\varphi}$ we define $u(z_0 + \vec{z})$ by :

$$u(z_0 + \vec{z}) = p(z_0, |\vec{z}|) + \frac{\vec{z}}{|\vec{z}|} q(z_0, |\vec{z}|).$$

Then u is a (left and right) holomorphic Cliffordian function.

Proof.- For odd n , this result is already known [LR1].

If n is even, let $n = 2m$. Let $x = z_0 + \vec{z}$ and $z = x + z_{2m+1} e_{2m+1}$. Define f by $f(z) = p(z_0, |\vec{z} + z_{2m+1} e_{2m+1}|) + \frac{\vec{z} + z_{2m+1} e_{2m+1}}{|\vec{z} + z_{2m+1} e_{2m+1}|} q(z_0, |\vec{z} + z_{2m+1} e_{2m+1}|)$.

From the case of odd n , we know that f is a left and right holomorphic Cliffordian function. The theorem of paragraphe IV shows then that u is a left and right holomorphic Cliffordian function, since we have $f(\bar{z}) = \overline{f(z)}$ and $u(x) = f(x)$.

Bibliographie

- [BDS] F. BRACKS, R. DELANGHE, F. SOMMEN - Clifford analysis ; *Pitman* (1982) .
- [De] C. DEAVORS - The quaternion calculus ; *Am. Math. monthly* (1973), 995-1008 .
- [DSS] R. DELANGHE, F. SOMMEN, V. SOUCEK - Clifford algebra and spinor-valued functions ; *Kluwer* (1992).
- [EL1] S.L. ERIKSSON-BIQUE, H. LEUTWILER - On modified quaternionic analysis in \mathbb{R}^3 ; *Arch. Math.* 70 (1998), 228-234.
- [EL2] S.L. ERIKSSON-BIQUE, H. LEUTWILER - Hyperholomorphic functions ; *to appear*.
- [Fu] R. FUETER - Die Functionentheorie der Differengleichungen $\Delta u = 0$ and $\Delta\Delta u = 0$ mit vier reellen Variablen ; *Comm. Math. Helv.* 7 (1935), 307-330.
- [Le1] H. LEUTWILER - Modified quaternionic analysis in \mathbb{R}^3 ; *Complex variables Vol.20* (1992), 19-51.
- [Le2] H. LEUTWILER - Rudiments of a function theory in \mathbb{R}^3 ; *Expo. Math.* 14 (1996), 97-123.
- [LR1] G. LAVILLE, I. RAMADANOFF - Holomorphic Cliffordian functions ; *Advances in Applied Clifford algebras* 8, n° 2 (1998), 321-340.
- [LR2] G. LAVILLE, I. RAMADANOFF - Elliptic Cliffordian functions ; *Complex variables Vol. 45, n° 4* (2001), 297-318.
- [Qi] T. QIAN - Generalization of Fueter's result to \mathbb{R}^{n+1} ; *Rend. Math. Acc. Lincei* 8 (1997), 111-117.