



HAL
open science

The L-R Correlator and its Chiral Condensates in the MHA and MHA+V' Approximations to Large-Nc QCD

Samuel Friot

► **To cite this version:**

Samuel Friot. The L-R Correlator and its Chiral Condensates in the MHA and MHA+V' Approximations to Large-Nc QCD. Nuclear Physics B Proceedings Supplements, 2006, 152. hal-00003302

HAL Id: hal-00003302

<https://hal.science/hal-00003302>

Submitted on 16 Nov 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The L-R Correlator and its Chiral Condensates in the MHA and MHA+V' Approximations to Large- N_c QCD

Samuel Friot ^{a b*}

^aCentre de Physique Théorique CNRS-Luminy, Case 907 F-13288 Marseille Cedex 9, France
CNRS UMR 6207

^bLaboratoire de Mathématiques Université Paris-Sud - Bât. 425, 91405 Orsay Cedex, France
CNRS UMR 8628

In this talk, we describe part of a recent work [1] on the correlation function of a $V - A$ current with a $V + A$ current in the framework of QCD in the limit of a large number of colours N_c . The discussion takes place within two successive approximations of this theory, called MHA and MHA+V'. Results concerning the evaluation of chiral condensates of dimension six and eight, as well as matrix elements of the Q_7 and Q_8 electroweak penguin operators, are given.

1. Introduction

The correlation function of a left-handed current with a right-handed current

$$L^\mu(x) = \bar{u}(x)\gamma^\mu \frac{1 - \gamma_5}{2} d(x)$$

and

$$R^\nu(0) = \bar{d}(0)\gamma^\nu \frac{1 + \gamma_5}{2} u(0),$$

in QCD and in the chiral limit, depends only on one invariant amplitude $\Pi_{LR}(Q^2)$ of the euclidean momentum squared $Q^2 = -q^2$, with q the momentum flowing through the two-point function:

$$\begin{aligned} 2i \int d^4x e^{iq \cdot x} \langle 0 | T \{ L^\mu(x) R^\nu(0) \} | 0 \rangle \\ = (q^\mu q^\nu - q^2 g^{\mu\nu}) \Pi_{LR}(Q^2). \end{aligned} \quad (1)$$

Here, we shall be particularly concerned with the study of $\Pi_{LR}(Q^2)$ in the limit of a large number of colors N_c in QCD. We want to compute the lowest dimension operator product expansion (OPE) condensates of this function in

*Talk given at the 11th High-Energy Physics International Conference on Quantum Chromodynamics, 5-10 July (2004), Montpellier (France).

two successive approximations to the large- N_c limit: the so-called *minimal hadronic approximation* (MHA) [2] consisting of a spectrum of the pion state, a vector state and an axial vector state; and the improved approximation where an extra higher vector state is added.

Part of our motivation comes from the fact that in the literature the values of these condensates are rather controversial. Phenomenological analyses in refs. [2,3,4,5] find for example opposite sign for the first two condensates, in contradistinction to the results in refs. [6,7,8] which find the same sign.

In this talk, we give results only for the two lowest dimension condensates and the matrix elements of Q_7 and Q_8 (see however [1] for more results and details).

In practice, we shall be working with the dimensionless complex function $W_{LR}[z]$ defined as

$$W_{LR}[z] = -z \Pi_{LR}(zM_\rho^2), \quad \text{with} \quad \text{Re } z = \frac{Q^2}{M_\rho^2},$$

and use the mass of the lowest massive state, the $\rho(770 \text{ MeV})$, to normalize quantities with dimensions. In large- N_c QCD the function $W_{LR}[z]$ is a meromorphic function and, therefore, in full generality, it can be approximated by successive par-

tial fractions of the type

$$W_{LR}[z] = A_N \prod_{i=1}^P \frac{1}{(z + \rho_i)} \prod_{j=1}^N (z + \sigma_j), \quad (2)$$

with $\rho_1 = 1$ and $\rho_i \neq \rho_k$ for $i \neq k$; where P (and N) get larger and larger, but finite. A_N is the overall normalization and in what follows, we shall often use the notation $\rho_2 \equiv \rho_A$ and $\rho_3 \equiv \rho_{V'}$.

On the other hand, in QCD, the OPE of the two currents in Eq. (1) fixes the large- Q^2 fall off in $1/Q^2$ -powers of the invariant function $\Pi_{LR}(Q^2)$ [9] to

$$\begin{aligned} \Pi_{LR}(Q^2) &\underset{Q^2 \rightarrow \infty}{=} \sum_{n=1}^{\infty} c_{2n+4}(Q^2, \mu^2) \\ &\times \langle \mathcal{O}_{2n+4}(\mu^2) \rangle (Q^2)^{-(n+2)} = \frac{1}{2} \sum_{n=1}^{\infty} \frac{\langle \mathcal{O}_{2n+4} \rangle}{(Q^2)^{n+2}}. \end{aligned} \quad (3)$$

Matching the leading asymptotic behaviour for large- z in Eq. (2) to the one of the OPE in Eq. (3), restricts the number of zeros N and the number of poles P in Eq. (2) to obey the constraint

$$N - P = -2. \quad (4)$$

The case where $N = 0$ corresponds to the MHA while the MHA+ V' case is obtained for $N = 1$.

2. Π_{LR} in the MHA and $\pi-V-A-V'$ Spectra

It is claimed by some of the authors of refs. [6, 7,8] that the reason why their phenomenological analysis of the chiral condensates give the same sign for $\langle \mathcal{O}_6 \rangle$ and $\langle \mathcal{O}_8 \rangle$ is due to the fact that the hadronic τ -decay spectrum is sensitive to the presence of the ρ' , while the MHA ignores all higher states beyond the first axial state. Partly motivated by this claim, we want to analyze here the case, beyond the MHA, where an extra vector state V' , and therefore one zero σ , are also included. Let us collect some relevant equations describing this case (the corresponding expressions in the MHA case can be obtained by putting $\rho_{V'} = \sigma$ in the MHA+ V' equations).

2.1. The Correlation Function

With a spectrum of the pion pole, and V , A , and V' states, the relevant correlation function is

$$W_{LR}[z] = A_1 \frac{z + \sigma}{(z + 1)(z + \rho_A)(z + \rho_{V'})},$$

where $A_1 \frac{\sigma}{\rho_A \rho_{V'}} = \frac{F_0^2}{M_V^2} \equiv \rho_F$ and therefore

$$\langle \mathcal{O}_6 \rangle = \frac{-2}{\sigma} F_0^2 M_A^2 M_{V'}^2 = -M_V^6 \frac{2}{\sigma} \rho_F \rho_A \rho_{V'}. \quad (5)$$

Because of the positivity of $W_{LR}[z]$ for $\text{Re } z \geq 0$, the position of the zero has to be in the Minkowski axis and, then, $\sigma > 0$.

2.2. The Linear Constraint

This is a very interesting constraint, key equation of our analysis, which already provides a semi-quantitative argument in favor of the opposite sign option for the condensates $\langle \mathcal{O}_6 \rangle$ and $\langle \mathcal{O}_8 \rangle$. It simply follows by expanding Eq. (2) to first non-trivial order in inverse powers of z

$$\sum_{j=1}^N \sigma_j - \sum_{i=1}^P \rho_i = \frac{1}{M_V^2} \frac{\langle \mathcal{O}_8 \rangle}{\langle \mathcal{O}_6 \rangle}. \quad (6)$$

In the case corresponding to the MHA, where by definition there are no zeros, this constraint simply becomes:

$$1 + \rho_A = -\frac{1}{M_V^2} \frac{\langle \mathcal{O}_8 \rangle}{\langle \mathcal{O}_6 \rangle}, \quad (7)$$

implying that, in the MHA $\langle \mathcal{O}_8 \rangle$ and $\langle \mathcal{O}_6 \rangle$ must have *opposite signs*.

Now, in the MHA+ V' case, Eq. (6) reduces to

$$\sigma - (1 + \rho_A + \rho_{V'}) = \frac{1}{M_V^2} \frac{\langle \mathcal{O}_8 \rangle}{\langle \mathcal{O}_6 \rangle}. \quad (8)$$

This equation states that for $\langle \mathcal{O}_8 \rangle$ to have the same sign as $\langle \mathcal{O}_6 \rangle$, the position of the zero has to be *far beyond the largest V' -pole*:

$$\sigma > 1 + \rho_A + \rho_{V'}. \quad (9)$$

Fixing the position of the poles at the values of the observed spectrum (and ignoring errors for

the purpose of the discussion), one has $M_V = 0.776$ GeV, $M_A = 1.230$ GeV ($\rho_A = 2.5$) and $M_{V'} = 1.465$ GeV ($\rho_{V'} = 3.6$); which means that for the equal sign requirement option to be satisfied, one must have $\sigma > 7.2$. In GeV units this corresponds to a mass of 2.1 GeV. Now, in writing a large- N_c ansatz for the $W_{LR}[z]$ function, one is implicitly assuming an effective cancellation between the extra poles and zeros in the complex z -plane which lie beyond a disc of radius s_0 covering all the poles and zeros retained in that approximation. The result $\sigma > 7.2$ implies that the radius in question has to be $\sqrt{s_0} > 2.1$ GeV. A priori that *seems a good thing* because the OPE-matching is now applied at $Q^2 \geq s_0$; i.e. in a more asymptotic region than in the case of the MHA ansatz; however, it also implies that there are no further poles in the region between $M_{V'} \simeq 1.5$ and the effective mass $M_\sigma \simeq 2.1$ GeV corresponding to the zero at $\sigma \simeq 7.2$. This, however, is in contradiction with the observed a_1 -like state at $M_{A'} \simeq 1.64$ GeV and ρ -like states at $M_{V''} \simeq 1.72$ GeV and $M_{V'''} \simeq 1.9$ GeV below $M_\sigma \simeq 2.1$ GeV. Alternatively, if one excludes those three states A' , V'' and V''' as all the phenomenological analyses using τ -data do in fact, then the position of the zero σ should be $\sigma \lesssim \frac{M_{A'}^2}{M_V^2} \approx 4.5$, implying according to Eq. (9), that $\langle \mathcal{O}_8 \rangle$ and $\langle \mathcal{O}_6 \rangle$ must have opposite signs, in contradiction with the claims of refs. [6,7,8].

3. Numerical Analyses and Conclusions

Confronting the MHA and MHA+V' approximations to the experimental values of some observables that we introduce in the following will allow us to test their consistency and adjust their free parameters (ρ_F and ρ_A in the MHA case, ρ_F , ρ_A , $\rho'_{V'}$ and σ in the MHA+V' case) in order to make predictions for the condensates and matrix elements. We use as input the following set of experimental data

$$\delta m_\pi = 4.5936 \pm 0.0005 \text{ MeV}, \quad \text{ref. [11]}$$

$$L_{10} = (-5.13 \pm 0.19) \times 10^{-3}, \quad \text{ref. [3]}$$

$$\Gamma_{\rho \rightarrow e^+ e^-} = (6.77 \pm 0.32) \times 10^{-3} \text{ MeV}, \quad \text{ref. [11]}$$

$$\Gamma_{a \rightarrow \pi \gamma} = (640 \pm 246) \times 10^{-3} \text{ MeV}, \quad \text{ref. [11]}$$

$$L_9 = (6.9 \pm 0.7) \times 10^{-3}, \quad \text{ref. [12]}$$

$$M_\rho = (775.9 \pm 0.5) \text{ MeV}; \quad \text{ref. [11]}$$

In fact, some of these observables, when expressed in terms of the MHA or MHA+V' parameters, depend not only on these ones but also on $(m_{\pi^+} + m_{\pi^0})$ and/or M_V . Therefore, it is more appropriate for our purposes to use in our fit procedure the dimensionless quantities:

$$\frac{m_{\pi^+} + m_{\pi^0}}{M_\rho^2} \delta m_\pi = \frac{3}{4} \frac{\alpha}{\pi} \frac{\rho_A \log(\rho_A)}{\rho_A - 1}, \quad (10)$$

$$L_{10} = -\frac{1}{4} \rho_F \left(1 + \frac{1}{\rho_A} \right), \quad (11)$$

$$\frac{1}{M_\rho} \Gamma_{\rho \rightarrow e^+ e^-} = \frac{4\pi\alpha^2}{3} \frac{\rho_A}{\rho_A - 1}, \quad (12)$$

$$\frac{1}{M_\rho} \Gamma_{a \rightarrow \pi \gamma} = \frac{\alpha}{24} \frac{\sqrt{\rho_A}}{\rho_A - 1}, \quad (13)$$

$$L_9 = \frac{1}{2} \rho_F. \quad (14)$$

For lack of place we gave the formulae only in the case of the MHA, the corresponding MHA+V' equations can be found in [1].

A standard χ^2 statistical regression method permits to fit the parameters.

3.1. The case of the MHA spectrum

For this spectrum, we find $\rho_F = (12.36 \pm 0.35) \times 10^{-3}$ and $\rho_A = 1.464 \pm 0.004$, with a $\chi^2_{\min} = 1.21$ for 3 degrees of freedom (dof). The covariance matrix is given by

$$\text{cov}(\rho_F, \rho_A) = \begin{pmatrix} 1.21 & 1.36 \\ 1.36 & 162 \end{pmatrix} \times 10^{-7}. \quad (15)$$

We deduce from these results two conclusions: first that the MHA framework is statistically relevant and second that the fitted free parameters have small statistical errors.

3.2. The case of a $\pi - V - A - V'$ spectrum

Eq. (14) becomes now a function of g_V as explained in [1]. The way we treat this is by con-

sidering the observable²:

$$\frac{1}{M_\rho} \Gamma_{\rho \rightarrow \pi\pi} = \frac{1}{48\pi} \times \frac{1}{\rho_F^2} \frac{\sigma (\rho_F - 2L_9 \rho_{V'})^2 (\rho_A - 1)(\rho_{V'} - 1)}{(1 - \rho_{V'})^2 \rho_F \rho_A \rho_{V'} (\sigma - 1)},$$

as a function of L_9 which has an error itself, and is added as an extra parameter in our fit. The number of dof does not change since L_9 is also taken as an observable. For this case, we impose a criterion of rejection through the ordering:

$$\rho_A < \rho_{V'} < \sigma < \rho_0 \doteq \frac{s_0}{M_V^2}.$$

The first and second inequalities reflect the knowledge that the new state has a higher mass than the axial, that it is a V -like pole and, therefore, its residue contributes positively to the $W_{LR}(z)$ -function; the third inequality follows from the requirement that the perturbative threshold s_0 already lies beyond the radius where the analytic structure of the poles and zeros retained satisfies the leading OPE constraint.

The χ^2 regression leads to

$$\rho_F = (12.36 \pm 0.03) \times 10^{-3}, \quad (16)$$

$$\rho_A = 1.466 \pm 0.003, \quad (17)$$

$$\rho_{V'} = 2.63 \pm 0.01, \quad (18)$$

$$\sigma = 2.64 \pm 0.01. \quad (19)$$

The results in Eqs. (16) to (19) correspond to a value: $L_9 = (6.44 \pm 0.02) \times 10^{-3}$, with a $\chi_{\min}^2 = 0.60$ for 1 dof.

We conclude that the parameters ρ_F and ρ_A are statistically stable when compared to those found in the MHA case. We also find that $\rho_{V'} \approx \sigma$, which is consistent with the fact that the MHA approximation already seems to have the bulk of the full large- N_c information. In other words, adding an extra V' -pole appears to be compensated, at a very good approximation, by the position of the nearby zero.

3.3. predictions

Using Eqs. (5), (8) (and corresponding equations in the MHA case) as well as expressions

²The decay width of $\rho \rightarrow \pi\pi$ is $\Gamma_{\rho \rightarrow \pi\pi} = (150.4 \pm 1.3) \text{MeV}$ [11].

of the matrix elements of four-quark operators ($[1],[10]$), we find

	$\langle \mathcal{O}_6 \rangle$ $\times 10^3 \text{ GeV}^6$	$\langle \mathcal{O}_8 \rangle$ $\times 10^3 \text{ GeV}^8$
MHA + V'	$-7.90 \pm 0.20 \pm 1.62$	$+11.69 \pm 0.32 \pm 2.53$
MHA	$-7.89 \pm 0.23 \pm 2.01$	$+11.71 \pm 0.34 \pm 3.08$

	M_7 (NDR)	M_7 (HV)
MHA + V'	$0.12 \pm 0.00 \pm 0.01$	$0.59 \pm 0.01 \pm 0.06$
MHA	$0.12 \pm 0.00 \pm 0.02$	$0.59 \pm 0.01 \pm 0.11$
	M_8 (NDR)	M_8 (HV)
MHA + V'	$2.00 \pm 0.03 \pm 0.20$	$2.15 \pm 0.03 \pm 0.22$
MHA	$1.99 \pm 0.03 \pm 0.36$	$2.15 \pm 0.03 \pm 0.39$

The way of obtention of these results as well as the origin of systematic errors are explained in [1].

We then see that within errors, the two sets of predictions from MHA and from MHA + V' are perfectly consistent with each other.

REFERENCES

1. S. Friot, D. Greynat and E. de Rafael, *J. High Energy Phys.* **10** (2004) 043 [hep-ph/0408281]
2. S. Peris, B. Phily and E. de Rafael, *Phys. Rev. Lett.* **86** (2001) 14.
3. M. Davier, L. Girlanda, A. Hoecker and J. Stern, *Phys. Rev.* **D58** (1998) 096014.
4. B.L. Ioffe and K.N. Zybalyuk, *Nucl. Phys.* **A687** (2001) 437.
5. K.N. Zybalyuk, hep-ph/0404230.
6. J. Bijnens, E. Gamiz and J. Prades, *J. High Energy Phys.* **10** (2001) 009.
7. V. Cirigliano, E. Golowich and K. Maltman, *Phys. Rev.* **D68** (2003) 054013.
8. J. Rojo and J.I. Latorre, *J. High Energy Phys.* **0401** (2004) 055.
9. M. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B147** (1979) 385, 447.
10. M. Knecht and E. de Rafael, *Phys. Lett.* **B424** (1998) 335.
11. K. Hagiwara *et al.*, *Phys. Rev. D* **66** (2002) 010001.
12. J. Bijnens, G. Ecker and J. Gasser *The second DAΦNE Physics Handbook* (1994) 125 [hep-ph/9411232].