



**HAL**  
open science

## The final phase of inspiral of neutron stars: realistic equations of state

Dorota Gondek-Rosinska, Michal Bejger, Tomek Bulik, Eric Gourgoulhon, Pawel Haensel, Francois Limousin, Leszek Zdunik

### ► To cite this version:

Dorota Gondek-Rosinska, Michal Bejger, Tomek Bulik, Eric Gourgoulhon, Pawel Haensel, et al.. The final phase of inspiral of neutron stars: realistic equations of state. 2004. hal-00003463v1

**HAL Id: hal-00003463**

**<https://hal.science/hal-00003463v1>**

Preprint submitted on 2 Dec 2004 (v1), last revised 9 Jan 2007 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The final phase of inspiral of neutron stars: realistic equations of state

D. Gondek-Rosińska<sup>a,b,c</sup>, M. Bejger<sup>a</sup>, T. Bulik<sup>a</sup>,  
E. Gourgoulhon<sup>b</sup>, Paweł Haensel<sup>a,b</sup>, F. Limousin<sup>b</sup> L. Zdunik<sup>a</sup>

<sup>a</sup>*CAMK, Bartycka 18, 00716, Warsaw, Poland*

<sup>b</sup>*LUTH, Observatoire de Paris, Universite Paris 7, Place Jules Janssen, 92195  
Meudon Cedex, France*

<sup>c</sup>*Departament de Fisica Aplicada, Universitat d'Alacant, Apartat de correus 99,  
03080 Alacant, Spain*

---

## Abstract

Coalescing compact star binaries are expected to be among the strongest sources of gravitational radiation to be seen by laser interferometers. We present calculations of the final phase of inspiral of equal mass irrotational neutron star binaries and strange quark star binaries. Six types of equations of state at zero temperature are used - three realistic nuclear equations of state of various softness and three different MIT bag models of strange quark matter. We study the precoalescing stage within the Isenberg-Wilson-Mathews approximation of general relativity using a multidomain spectral method. The gravitational-radiation driven evolution of the binary system is approximated by a sequence of quasi-equilibrium configurations at fixed baryon number and decreasing separation. We find that the innermost stable circular orbit (ISCO) is given by an orbital instability for binary strange quark stars and by the mass-shedding limit for neutron star binaries. The gravitational wave frequency at the ISCO, which marks the end of the inspiral phase, is found to be  $\sim 1100 - 1460$  Hz for two  $1.35 M_{\odot}$  irrotational strange stars described by the MIT bag model and between 800 Hz and 1230 Hz for neutron stars.

*Key words:*

gravitational waves-relativity- stars:binaries- stars: neutron- equation of state

---

---

*Email address:* Dorota.Gondek@obspm.fr (D. Gondek-Rosińska).

## 1 Introduction

Coalescing neutron star binaries are considered among the strongest and most likely sources of gravitational waves to be seen by VIRGO/LIGO interferometers [10, 19, 5]. Due to the emission of gravitational radiation, binary compact stars decrease their orbital separation and finally merge. The evolution of a binary system can be separated into three phases : *point-like inspiral* where orbital separation is much larger than the neutron star radius, *hydrodynamical inspiral* where orbital separation is just a few times larger than the radius of the NS so that hydrodynamics play an important role, and *merger* in which the two stars coalesce dynamically. Gravitational waves emitted during the hydrodynamical inspiral and the merger phase could yield important informations about the equation of state (EOS) of dense matter [13, 27, 24, 4, 20]. Up to now, all calculations (except those of [24, 4, 20]) of the hydrodynamical inspiral and of the merger phase have been done for the simplified EOS of dense matter - the polytropic EOS. In the paper we present the results of our studies on the hydrodynamical phase of inspiraling binary systems containing equal mass compact stars described by different realistic EOS of dense matter. The calculations are performed in the framework of *Isenberg-Wilson-Mathews* approximation to general relativity (see Ref. [3] for a review). We consider binary systems consisting of two identical stars. We choose the gravitational mass of each star to be  $1.35 M_{\odot}$  at infinite separation in order to be consistent with recent population synthesis calculations [8] and with the current set of well-measured neutron star masses in relativistic binary radio pulsars [21, 9].

## 2 Equations of state and stellar models

In Fig. 1 we show gravitational mass versus areal radius for sequences of static compact stars. We limit ourselves to neutron stars consisting of nucleons and hyperons and strange quark stars described by the MIT bag model. Depending on the EOS we obtain the radius of a  $1.35 M_{\odot}$  star in the range 10-14 km. We perform calculations for three nuclear EOS of dense matter based on modern many-body calculations. The differences in the M-R relation for stellar models of neutron stars (solid lines) shown in Fig. 1 reflect the uncertainties in the existing theories of the interactions in nuclear matter. We consider one soft (**BPAL12**, [7]) and one stiff (**AkmalPR**, [2]) EOS of matter composed of nucleons, electrons and muons. We considered also one EOS in which hyperons are present at high densities (**GlendNH3**, [17]). The neutron star crust is described by means of a realistic EOS obtained in the many-body calculations (see [4] for details)

Strange stars are currently considered as a possible alternative to neutron




Fig. 1. Gravitational mass  $M$  versus stellar radius  $R$  for sequences of static neutron stars described by three different nuclear equations of state (solid lines) and strange quark stars described by different MIT bag models (dashed lines).

stars as compact objects (see e.g. [28, 22, 16] and references therein). Typically, they are modeled with an EOS based on the MIT-bag model in which quark confinement is described by an energy term proportional to the volume (e.g. [1, 18]) in which quark confinement is described by an energy term proportional to the volume [14]. There are three physical quantities entering the MIT-bag model: the mass of the strange quarks,  $m_s$ , the bag constant,  $B$ , and the strength of the QCD coupling constant  $\alpha$ . In the framework of this model the quark matter is composed of massless u, d quarks, massive s quarks and electrons. Strange stars are self-bound objects, having high density (in the range  $\sim 3 \times 10^{14} - 6.4 \text{g/cm}^3$  [16]) at the surface. Dashed lines in Fig. 1 correspond to sequences of static strange quark stars described by three different sets of parameters of the MIT-bag model: **SQS56** - the standard MIT bag model:  $m_s c^2 = 200 \text{ MeV}$ ,  $\alpha = 0.2$ ,  $B = 56 \text{ MeV/fm}^3$ ; **SQSB60** - the simplified MIT bag model with  $m_s = 0$ ,  $\alpha = 0$ ;  $B = 60 \text{ MeV/fm}^3$ ; **SQSB40** - the "extreme" MIT bag model (relatively low strange quark mass and  $B$  but high  $\alpha$ ):  $m_s c^2 = 100 \text{ MeV}$ ,  $\alpha = 0.6$ ,  $B = 40 \text{ MeV/fm}^3$ .

### 3 Numerical method

In the hydrodynamical phase, since the timescale of orbital shrinking due to the emission of gravitational waves is longer than the orbital period, one may consider a binary system to be in a quasi-equilibrium state (helical Killing vector approximation). For each EOS, we construct so called an *evolutionary sequence* by calculating a sequence of quasi-equilibrium configurations with fixed baryon mass and decreasing orbital separation. The present computations of




Fig. 2. Binding energy as a function of gravitational wave frequency (twice the orbital frequency) along evolutionary sequences of irrotational binaries. The solid lines denote neutron stars, the dashed one strange quark stars and the dotted line point-mass binaries in the third order Post-Newtonian approximation [6]. Each evolutionary sequence finishes at the innermost stable circular orbit. The diamonds correspond to dynamical orbital instability while circles to the mass-shedding limit.

close binary neutron star or strange star systems rely on the assumption of irrotational flow of the fluid and a conformally flat spatial 3-metric (Isenberg-Wilson-Mathews approximation). In order to calculate the last orbits of inspiral phase of binary compact stars we use a numerical code which solves the five coupled, nonlinear, elliptic equations for the gravitational field, supplemented by an elliptic equation for the velocity potential of irrotational flows (see [20] for a discussion on different boundary conditions in the case of strange stars and neutron stars). The code has been already used successfully for calculating the final phase of the inspiral of compact stars [25, 26, 27, 4, 20]. This code is built upon the C++ library LORENE (<http://www.lorene.obspm.fr>). The complete description of the resulting general relativistic equations, the whole algorithm, as well numerous tests of the code can be found in [25]. Additional tests have been presented in Sect. III of [27].

## 4 Results

In Fig. 2 we show the evolution of equal mass binary neutron stars (solid lines) and strange stars (dashed lines) having total gravitational mass  $2.7M_{\odot}$  at infinity. The binding energy  $E_{\text{bind}}$  is defined as the difference between  $M_{\text{ADM}}$  (see [27]) and the total mass of the system at infinity. This energy is equal to total energy emitted by a binary system in gravitational waves. The frequency

of gravitational waves is twice the orbital frequency. Comparison of our numerical results with 3rd order PN point masses calculations [6] reveals a good agreement for small frequencies (large separations). The deviation from PN curves at higher frequencies (smaller separation) is due to hydrodynamical effects, which are not taken into account in the PN approach. A turning point of  $E_{\text{bind}}$  along an irrotational evolutionary sequence indicates the orbital (dynamical) instability [15]. This instability originates both from relativistic effects and hydrodynamical effects. In the case where no turning point of  $E_{\text{bind}}$  occurs along the sequence, the mass-shedding limit (Roche lobe overflow) marks the end of the inspiral phase of the binary system, since recent dynamical calculations for  $\gamma = 2$  polytrope have shown that the time to coalescence was shorter than one orbital period for configurations at the mass-shedding limit (i.e. see [23]). Thus the physical inspiral of binary compact stars terminates by either the orbital instability (turning point of  $E_{\text{bind}}$ ) or the mass-shedding limit. In both cases, this defines the *innermost stable circular orbit (ISCO)*. The end of inspiral phase strongly depends on EOS - for irrotational neutron star binaries a quasi-equilibrium sequence terminates by mass-shedding limit (circles at the end of each line in Fig 2) and for strange stars by orbital instability (shown as diamonds in Fig. 2). The differences in the evolution of binary strange stars and neutron stars stem from the fact that strange stars are principally bound by additional force than gravitation: the strong interaction between quarks. They are self-bound objects having very high adiabatic index at the stellar surface (see [20]). Although the crust of a  $1.35 M_{\odot}$  neutron star contains only a few percent of the stellar mass, this region is easily deformed under the action of the tidal forces resulting from the gravitational field produced by the companion star. The end of inspiral phase of binary stars strongly depends on the stiffness of matter in this region. *The frequency of gravitational waves at the ISCO is one of potentially observable parameters by the gravitational wave detectors.* The frequency of gravitational waves at the ISCO strongly depends on EOS. For irrotational equal mass (of  $1.35 M_{\odot}$  at infinite separation) binaries this frequency is  $\sim 1100 - 1460$  Hz for strange stars described by the MIT bag model and between 800 Hz and 1230 Hz for neutron stars described by nuclear EOS (see also [24, 4]). The 3rd PN approximations for point masses derived by different authors are giving ISCO at very high frequencies of gravitational waves  $> 2$  kHz [6, 11, 12]

## Acknowledgements

This work was partially supported by the KBN grant 5 P03D 01721 and PBZ-KBN-054/P03/2001, by the “Bourses de recherche 2004 de la Ville de Paris” and by the Associated European Laboratory Astro-PF (Astrophysics Poland-France). DGR is grateful to the COSPAR meeting organizers for support.

## References

- [1] C. Alcock, E. Farhi, and A. Olinto, *Astrophys. J.* **310**, 261 (1986).
- [2] A. Akmal, V. R. Pandharipande and D. G. Ravenhall, 1998, *Phys. Rev. C*, **58**, 1804
- [3] T. W. Baumgarte and S. L. Shapiro, *Phys. Rep.* **376**, 41 (2003)
- [4] M. Bejger, D. Gondek-Rosińska, E. Gourgoulhon et al., *Astron. Astrophys.*, in press [preprint: astro-ph/0406234].
- [5] K. Belczynski, V. Kalogera, and T. Bulik *Astrophys. J.* **572**, 407 (2002).
- [6] L. Blanchet, *Phys. Rev. D* **65**, 124009 (2002).
- [7] Bombaci I. 1995, in *Perspectives on Theoretical Nuclear Physics*, 223
- [8] T. Bulik, D. Gondek-Rosińska, K. Belczynski, *Mon. Not. R. Astron. Soc.* **352**, 1372 (2004).
- [9] M. Burgay et al., *Nature*, **426**, 531 (2003).
- [10] M. Burgay, N. D'Amico, A. Possenti et al. *The Highly Relativistic Binary Pulsar PSR J0737-3039A: Discovery and Implications*, astro-ph/0405194.
- [11] T. Damour, P. Jaranowski, and G. Schäfer, *Phys. Rev. D* **62**, 044024 (2000).
- [12] T. Damour, E. Gourgoulhon, and P. Grandclément, *Phys. Rev. D* **66**, 024007 (2002)
- [13] J. A. Faber, P. Grandclément, F. A. Rasio et al., *Phys. Rev. Lett.* **89**, 231102 (2002).
- [14] E. Fahri, and R. L. Jaffe, *Phys. Rev. D* **30**, 112379 (1984)
- [15] J. L. Friedman, K. Uryu, and M. Shibata, *Phys. Rev. D* **65**, 064035 (2002).
- [16] D. Gondek-Rosińska, E. Gourgoulhon and P. Haensel, *Astron. Astrophys.* **412**, 777 (2003).
- [17] N. K. Glendenning, 1985, *ApJ*, 293, 470
- [18] P. Haensel, J.L. Zdunik, and R. Schaeffer, *Astron. Astrophys.* **160**, 121 (1986).
- [19] V. Kalogera, C. Kim, D. R. Lorimer et al. *Astrophys. J.* **601**, 179 (2004).
- [20] F. Limousin, D. Gondek-Rosińska and E. Gourgoulhon, 2004 submitted to *Phys. Rev. D* [preprint: gr-gc/0411127].
- [21] D. R. Lorimer, *Living Rev. Relativity* **4**, 5 (2001)
- [22] J. Madsen, *Lect. Notes Phys.* **516**, 162 (1999).
- [23] P. Marronetti, M. D. Duez, S. L. Shapiro et al., *Phys. Rev. Lett.* **92**, 141101 (2004).
- [24] R. Oechslin, K. Uryu, G. Poghosyan et al., *Mon. Not. R. Astron. Soc.* **349**, 1469 (2004).
- [25] K. Taniguchi, E. Gourgoulhon, and S. Bonazzola, *Phys. Rev. D* **64**, 064012 (2001)
- [26] K. Taniguchi and E. Gourgoulhon, *Phys. Rev. D* **66**, 104019 (2002).
- [27] K. Taniguchi and E. Gourgoulhon, *Phys. Rev. D* **68**, 124025 (2003).
- [28] F. Weber, *Prog. Part. Nucl. Phys.* **54**, 193 (2005).