

The Gap Function Phi(k,w) for a Two-leg t-J Ladder and the Pairing Interaction

Didier Poilblanc, Douglas J. Scalapino

▶ To cite this version:

Didier Poilblanc, Douglas J. Scalapino. The Gap Function Phi(k,w) for a Two-leg t-J Ladder and the Pairing Interaction. Physical Review B: Condensed Matter and Materials Physics (1998-2015), 2005, 71 (17), pp.174403. 10.1103/PhysRevB.71.174403. hal-00003481

HAL Id: hal-00003481 https://hal.science/hal-00003481

Submitted on 7 Dec 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Gap Function $\phi(k, \omega)$ for a Two-leg t-J Ladder and the Pairing Interaction

Didier Poilblanc*

Laboratoire de Physique Théorique CNRS-UMR5152, Université Paul Sabatier, F-31062 Toulouse, France

D. J. Scalapino[†]

Department of Physics, University of California, Santa Barbara, CA 93106 USA (Dated: December 8, 2004)

Abstract

The gap function $\phi(k, \omega)$, determined from a Lanczos calculation for a doped 2-leg t-J ladder, is used to provide insight into the spatial and temporal structure of the pairing interaction. It implies that this interaction is a local near-neighbor coupling which is retarded. The onset frequency of the interaction is set by the energy of an S = 1 magnon-hole-pair and it is spread out over a frequency region of order the bandwith.

PACS numbers: PACS numbers: 75.10.-b, 75.10.Jm, 75.40.Mg

 $^{{}^*} Electronic \ address: \ didier.poilblanc@irsamc.ups-tlse.fr$

[†]Electronic address: djs@vulcan2.physics.ucsb.edu

The momentum and frequency dependence of the superconducting gap function $\phi(k, \omega)$ provides information on the spatial and dynamic structure of the pairing interaction. Traditionally, electron tunneling has been used to explore the frequency dependence of the gap [1]. Presently, angular resolved photoemission spectroscopy [2] (ARPES) and scanning tunneling microscopy [3, 4] (STM) open the possibility of obtaining both k and ω information about the superconducting gap. However, approaches to extract this information on $\phi(k, \omega)$ from such experiments are still being explored. As has been previously discussed, a k and ω dependent gap function for finite t-J lattices can be obtained using Lanczos exact diagonalization [5, 6]. Here, using results for $\phi(k, \omega)$ becomes available.

The Hamiltonian for a 2-leg t-J ladder can be written as

$$H = J_{\text{rung}} \sum_{i} \left(\mathbf{S}_{i1} \cdot \mathbf{S}_{i2} - \frac{1}{4} n_{i1} n_{i2} \right) + J_{\text{leg}} \sum_{i,\alpha} \left(\mathbf{S}_{i+1,\alpha} \cdot \mathbf{S}_{i,\alpha} - \frac{1}{4} n_{i+1,\alpha} n_{i\alpha} \right)$$
$$- t_{\text{rung}} \sum_{i} \left(c_{i,1}^{\dagger} c_{i,2} + h.c. \right) - t_{\text{leg}} \sum_{i,\alpha} \left(c_{i+1,\alpha}^{\dagger} c_{i\alpha} + h.c. \right)$$
(1)

where $c_{i\alpha}$ are projected hole operators (spin indices are omitted) and $\alpha = (1, 2)$ labels the two legs of the ladder. We will consider the isotropic case in which $J_{\text{rung}} = J_{\text{leg}} = J$ and $t_{\text{rung}} = t_{\text{leg}} = t$ and our calculations will be carried out for a periodic 2 × 12 ladder at 1/8 and 1/6 hole doping.

As previously discussed, a gap function can be extracted by combining Lanczos results for the usual one-electron Green's function $G(k, \omega)$ with the Fourier transform of Gorkov's off-diagonal Green's function

$$F(k,t) = i \left\langle Tc_{-k,-\sigma}(t/2)c_{k,\sigma}(-t/2) \right\rangle$$
(2)

Here, for a finite system, this expectation value is taken between the ground states for N and N-2 particles and we choose the phase of F to be zero. We will take N = 22, N-2 = 20 corresponding to an average filling $n \simeq \frac{21}{24} = 0.875$. The Dyson equations [7] relating G and F are

$$[Z(k,\omega)\omega - (\epsilon_k + X(k,\omega)]G(k,\omega) = 1 - \phi(k,\omega)F(k,\omega)$$
(3)

$$[Z(-k,-\omega)\omega - (\epsilon_k + X(-k,-\omega))]F(k,\omega) = -\phi(k,\omega)G(k,\omega)$$
(4)

with Z and X the usual Nambu self energies and $\phi(k, \omega)$ the gap function. Then, using the even $(k \to -k)$ parity of $Z(k, \omega)$, $X(k, \omega)$, ϵ_k and $\phi(k, \omega)$ along with the even ω dependence of $Z(k, \omega)$, $X(k, \omega)$ and $\phi(k, \omega)$, we have

$$\phi(k,\omega) = \frac{F(k,\omega)}{F^2(k,\omega) + G(k,\omega) G(k,-\omega)}.$$
(5)

Alternatively, $Z(k,\omega)$ can be eliminated to obtain an expression for the superconducting gap

$$\Delta(k,\omega) = \frac{\phi(k,\omega)}{Z(k,\omega)} = \frac{2\omega F(k,\omega)}{G(k,\omega) - G(k,-\omega)}.$$
(6)

From a numerical calculation of $G(k, \omega)$ and $F(k, \omega)$, $\Delta(k, \omega)$ has been obtained for a 32site, t-J cluster [5]. This extended previous work by Ohta *et. al* [8] who fit the spectral weight $Im F(k, \omega)$ to a $d_{x^2-y^2}$ BCS-Bogoliubov quasiparticle form in which the frequency dependence of the gap was neglected.

In contrast to the long range order of the superconducting ground state of a 2D lattice, a 2-leg ladder exhibits power law pair field correlations [10, 11] which decay as $x^{-1/\kappa_{\rho}}$. Here, κ_{ρ} is the Luttinger liquid parameter associated with the massless charge mode. This implies that for a ladder of length L, the off-diagonal Green's function $F(k,\omega)$ decays [12, 13, 14] as $(\xi/L)^{\frac{1}{2\kappa_{\rho}}}$. Here, the coherence length ξ is proportional to the inverse of the gap. For our doped ladder L = 12 is of order this coherence length so that we can probe the internal $k - \omega$ structure of a pair.

For J/t = 0.4 and 1/8 doping, the k-dependence of the zero frequency gap $\phi(k_x, k_y, \omega = 0)$ is plotted in Fig. 1 for the bonding $(k_y = 0)$ and antibonding $(k_y = \pi)$ bands. For comparison, the solid curves correspond to ϕ_0 ($a \cos k_x - \cos k_y$) with $\phi_0 = 0.3$ and a = 0.8. This d-wavelike k-dependence of the gap function is similar to the behavior found in previous studies of both the t-J [8] and Hubbard ladders [9]. It implies that the spatial structure of the pairing interaction is dominantly a near-neighbor interaction.

The frequency dependence of the real and imaginary parts of $\phi(k, \omega)$ are shown in Fig. 2 for k values which are near the fermi surface of the bonding (red) and antibonding (green) bands respectively. In general, the gap function $\phi(k, \omega)$ is a complex frequency-dependent function $\phi_1 + i\phi_2$ with the imaginary part associated with dynamic decay processes. For our doped 2-leg ladder with J/t = 0.4, the magnitudes of the zero frequency gaps $\Delta_0 = \Delta(k_F, 0)$ obtained from eq. (6), for both the bonding and antibonding fermi points, are of order 0.15t. The gap, Δ_0 , is reduced from $\phi(k_F, 0)$ due to the renormalization factor Z. The onset of the imaginary part of $\phi_2(k,\omega)$ seen in Fig. 2(b) appears to occur somewhat below $3\Delta_0$. In addition, the peaking in $\phi_2(k,\omega)$ and the rapid rise in $\phi_1(k,\omega)$ as this onset frequency is approached suggest that a particular excitation mode occurs at a frequency Ω such that $\omega = \Delta_0 + \Omega$ determines the onset of $\phi_2(k,\omega)$.

The occurrence of an onset peak in $\phi_2(k,\omega)$ at threshold and the short-range nature of the interaction involving scattering of pairs from the bonding to antibonding band $(q_y \sim \pi)$, imply that the pairing in this energy regime is mediated by an S = 1 channel. This follows from the form of the coherence factor which varies as $\frac{1}{2} \left(1 \pm \frac{\Delta(p+q,\omega) \Delta(p,\omega)}{E(p+q) E(p)} \right)$, with the plus sign associated with the charge channel and the minus sign with the S = 1 spin channel. For a "d-wave-like" gap with $\mathbf{q} = (k_F \text{ (bonding)} - k_F \text{ (antibonding)}, \pi)$ and ω near threshold, the coherence factor goes to 1 for the spin channel and vanishes for the charge channel.

A plot of the low-energy S = 1 excitations for a doped ladder is shown in Fig. 3(a). The solid diamonds show a collective S = 1 bound magnon-hole-pair [15, 16] mode and the open symbols the S = 1 particle-hole continuum. A measure of the spectral weight of the S = 1channel which couples to $\phi(k, \omega)$ is given by the d-wave projection of the spin fluctuation spectral weight

$$V_d(\omega) = \frac{1}{N^2} \sum_{k,k'} (\cos k_x - \cos k_y) \left(\cos k'_x - \cos k'_y\right) S(\mathbf{k} - \mathbf{k}', \omega)$$
(7)

with

$$S(q,\omega) = -\frac{1}{\pi} Im \left\langle S^{\alpha}_{-q} \frac{1}{\omega + E_0 + i\eta - H} S^{\alpha}_q \right\rangle.$$
(8)

Here S_q^{α} is the Fourier transform of the α spin component. A plot of $V_d(\omega)$ versus ω is shown in Fig. 3(b). The peak at $\Omega \simeq 0.15t$ arises from the bound magnon-hole-pair mode and the high frequency weight comes from the particle-hole continuum. We believe that the onset of $\phi_2(k,\omega)$ seen in Fig. 2 is due to a process in which a single particle excitation with energy $\omega = \Omega_m + \Delta_0$ emits a bound magnon-hole-pair and drops down to the gap edge while the remaining $\phi_2(k,\omega) \neq 0$ region arises from a coupling to the S = 1 continuum.

In order to further characterize the dynamic nature of the pairing interaction, we introduce

$$I(k,\Omega) = \frac{1}{\pi} \int_0^\Omega d\omega' \, \frac{\phi_2(k,\omega')}{\omega'} \,. \tag{9}$$

Then, since the gap function satisfies a dispersion relation, one has

$$\phi_1(k,0) = I(k,\Omega \to \infty) + \phi_{\text{static}}(k) \,. \tag{10}$$

Here, $\phi_{\text{static}}(k)$ represents a non-retarded contribution. For example, if one were to make a mean-field approximation in which

$$J\left(\mathbf{S}_{i}\cdot\mathbf{S}_{j}-\frac{1}{4}n_{i}n_{j}\right)\rightarrow-J\left(\left\langle \Delta_{i\delta}\right\rangle \Delta_{i\delta}^{\dagger}+h.c.\right)$$

one would have an effective attractive pairing interaction which is independent of frequency which would contribute to ϕ_{static} . In Fig. 4(a), we have plotted $I(k,\Omega)/\phi_1(k,0)$ versus Ω and one sees that for J/t = 0.4, $I(k,\Omega)$ saturates at over 80% of $\phi_1(k,0)$. This implies that the dominant part of the pairing interaction comes from dynamic processes. Also shown, Fig. 4(b), are similar results for J/t=0.8. In this, unphysically large, J/t regime only 60% of the $\omega = 0$ gap function is associated with a dynamic pairing interaction. This is similar to earlier results for the spatial structure of a pair [17, 18] in which near-neighbor sites for the two holes making up a pair were favored for large values of J/t. For physical values of J/t, next-near-neighbor (diagonal) hole-hole occupation was favored. The diagonal structure arises dynamically and reflects the retarded nature of the pairing interaction for physical values of J/t.

In summary, knowledge of the k- and ω -dependence of the gap function provides information on the spatial and temporal structure of the pairing interaction. Numerical solutions for the two-leg ladder show that the pairing interaction has a short-range, near-neighbor form, leading to momentum transfer processes which scatter pairs between the bonding and antibonding states. The d-wave-like momentum dependence of the gap function and the ω onset of $\phi_2(k, \omega)$ imply that the pairing interaction in this energy regime is mediated by the S = 1 channel. This channel contains both a bound magnon-pair state and a continuum of particle-hole excitations. We believe that the magnon-pair mode is responsible for the onset behavior seen in $\phi_2(k, \omega)$. The dispersion relation for $\phi_1(k, 0)$ shows that for physical values of J/t, the dominant part of the interaction is dynamic with contributions coming from both the magnon-hole-pair mode and the particle-hole continuum.

Acknowledgments

DJS would like to acknowledge support under NSF Grant #DMR02-11166. DP thanks IDRIS (Orsay, France) for use of supercomputer facilities.

- W.L. McMillan and J.M. Rowell, Superconductivity I, Chap. 11, ed. by R.D. Parks, Marcel Dekker Inc., New York (1969).
- [2] A. Damascelli, Z. Hussain, and Z.X. Shen, Ref. Mod. Phys. 75, 473 (2003).
- [3] J.E. Hoffman, E.W. Hudson, K.M. Lang, V. Modhavan, H. Eisaki, S. Uchida, and J.C. Davis, Science 295, 466 (2002).
- [4] C. Howald, H. Eisaki, N. Kaneko, M Greven, and A. Kapitulnik, Phys. Rev. B 67, 14533 (2003).
- [5] D. Poilblanc and D.J. Scalapino, Phys. Rev. B 66, 052513 (2002).
- [6] D. Poilblanc, D.J. Scalapino, and S. Capponi, Phys. Rev. Lett. 91, 137203 (2004).
- [7] In general, the self-energy is decomposed as $X(k,\omega) + \omega(1 Z(k,\omega))$. Although the 2-leg ladder is a Luttinger liquid, the Green's functions for the finite 2 × 12 lattice that we have studied can be parameterized in terms of $Z(k,\omega)$, $X(k,\omega)$, and $\phi(k,\omega)$.
- [8] Y. Ohta, T. Shimozato, R. Eder, and S. Maekawa, Phys. Rev. Lett. 73, 324 (1994).
- [9] T. Dahm and D.J. Scalapino, *Physica C* 288, 33 (1997).
- [10] E. Dagotto and T.M. Rice, *Science* **271**, 618 (1996).
- [11] C.A. Hayward, D. Poilblanc, R.M. Noack, D.J. Scalapino, W. Hanke, Phys. Rev. Lett. 75, 926 (1995).
- [12] S. Capponi and D. Poilblanc, Phys. Rev.B 66, 180503(R) (2002).
- [13] J. Voit, Eur. Phys. J. B 5, 505 (1998); F.J.L. Essler and A.M. Tsvelik, cond-mat/0205294 (2002).
- [14] A bozonization calculation shows that $\Delta(k,\omega)$ given by eq. (6) factors into an $L^{-\frac{1}{2\kappa_{\rho}}}$ factor times a function of ω and k which is independent of L. See E. Orignac and D. Poilblanc, *Phys. Rev. B* 68, 052504 (2003).
- [15] D. Poilblanc, O. Chiappa, J. Riera, S.R. White, and D.J. Scalapino, *Phys. Rev. B* 62, R14633 (2000). See also W. Zheng and C.J. Hamer, *Phys. Rev. B* 66, 085112 (2002).



FIG. 1: The k_x dependence of $\phi_1(k_x, k_y, \omega = 0)$ for the bonding $k_y = 0$ (red) and antibonding $k_y = \pi$ (green) bands. For comparison, the solid curves show $\phi_0(a \cos k_x - \cos k_y)$ with $\phi_0 = 0.3$ and a = 0.8.

- [16] D. Poilblanc, E. Orignac, S.R. White, and S. Capponi, Phys. Rev. B 69, 220406(R) (2004).
- [17] D. Poilblanc, Phys. Rev. B 49, 1477 (1994).
- [18] S.R. White and D.J. Scalapino, Phys. Rev. B 55, 6504 (1997).



FIG. 2: Real (a) and imaginary (b) parts of the gap function $\phi(k, \omega)$ versus ω for a 2 × 12 ladder at values of k near the bonding (red) and antibonding (green) fermi momenta [$\sim (\frac{2\pi}{3}, 0)$ and $\sim (\frac{\pi}{3}, \pi)$ respectively] for J/t = 0.4. Here, $\Delta_0 \sim \Delta(k_F, \Delta_0)$ and Ω_m is the minimum energy for a bound magnon-hole-pair excitation.



FIG. 3: (a) Low-energy electron-hole S = 1 excitations for a 1/8-doped J/t = 0.4 ladder for $q_y = 0$ (circles) and $q_y = \pi$ (diamonds) momenta. The solid diamonds denote a bound magnon-hole-pair collective mode [6]. (b) "d-wave" projection of the spin fluctuation spectral weight with Ω_m the minimum energy of the bound magnon-hole-pair collective mode (1/6 doping).



FIG. 4: The integrated contribution of $\phi_2(k,\omega)/\omega$ to $\phi_1(k,0)$ for the bonding (red), antibonding (green) k-values shown in Fig. 2. (a) for J/t = 0.4 and (b) for J/t = 0.8. The non-retarded contributions are shown by arrows.