



HAL
open science

Differential approximation results for the traveling salesman problem with distances 1 and 2

Jérôme Monnot, Vangelis Paschos, Sophie Toulouse

► **To cite this version:**

Jérôme Monnot, Vangelis Paschos, Sophie Toulouse. Differential approximation results for the traveling salesman problem with distances 1 and 2. *European Journal of Operational Research*, 2003, 145(3), pp.537-548. hal-00003928

HAL Id: hal-00003928

<https://hal.science/hal-00003928>

Submitted on 18 Jan 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Differential approximation results for the traveling salesman problem with distances 1 and 2

Jérôme Monnot Vangelis Th. Paschos* Sophie Toulouse
{monnot,paschos,toulouse}@lamsade.dauphine.fr

Abstract

We prove that both minimum and maximum traveling salesman problems on complete graphs with edge-distances 1 and 2 (denoted by `min_TSP12` and `max_TSP12`, respectively) are approximable within $3/4$. Based upon this result, we improve the standard approximation ratio known for maximum traveling salesman with distances 1 and 2 from $3/4$ to $7/8$. Finally, we prove that, for any $\epsilon > 0$, it is **NP**-hard to approximate both problems better than within $741/742 + \epsilon$. The same results hold when dealing with a generalization of `min_` and `max_TSP12`, where instead of 1 and 2, edges are valued by a and b .

1 Introduction

Given a complete graph on n vertices, denoted by K_n , with edge distances either 1 or 2 the minimum traveling salesman problem (`min_TSP12`) consists of minimizing the cost of a Hamiltonian cycle, the cost of such a cycle being the sum of the distances on its edges (in other words, in finding a Hamiltonian cycle containing a maximum number of 1-edges). The maximum traveling salesman problem (`max_TSP`) consists of maximizing the cost of a Hamiltonian cycle (in other words, of finding a Hamiltonian cycle containing a maximum number of 2-edges). A generalization of `TSP12`, denoted by `TSPab`, is the one where the edge-distances are either a , or b , $a < b$. Both `min_` and `max_TSP12`, and `TSPab` are **NP**-hard.

Given an instance I of an **NP** optimization (**NPO**) problem Π and a polynomial time approximation algorithm **A** feasibly solving Π , we will denote by $\omega(I)$, $\lambda_{\mathbf{A}}(I)$ and $\beta(I)$ the values of the worst solution of I , of the approximated one (provided by **A** when running on I), and the optimal one for I , respectively. Generally (see [8]), the quality of an approximation algorithm for an **NP**-hard minimization (resp., maximization) problem Π is expressed by the ratio (called standard in what follows) $\rho_{\mathbf{A}}(I) = \lambda(I)/\beta(I)$, and the quantity $\rho_{\mathbf{A}} = \inf\{r : \rho_{\mathbf{A}}(I) < r, I \text{ instance of } \Pi\}$ (resp., $\rho_{\mathbf{A}} = \sup\{r : \rho_{\mathbf{A}}(I) > r, I \text{ instance of } \Pi\}$) constitutes the approximation ratio of **A** for Π . Another approximation-quality criterion used by many researchers ([2, 1, 3, 4, 13, 14]) is what in [6, 5] we call *differential-approximation ratio*. It measures how the value of an approximate solution is placed in the interval between $\omega(I)$ and $\beta(I)$. More formally, the differential-approximation ratio of an algorithm **A** is defined as $\delta_{\mathbf{A}}(I) = |\omega(I) - \lambda(I)|/|\omega(I) - \beta(I)|$. The quantity $\delta_{\mathbf{A}} = \sup\{r : \delta_{\mathbf{A}}(I) > r, I \text{ instance of } \Pi\}$ is the differential approximation ratio of **A** for Π . In [2], the term “trivial solution” is used to denote the solution realizing the worst among the feasible solution-values of an instance. Moreover, all the examples in [2] carry over **NP**-hard problems for which worst solution can be trivially computed. This is for example the case of maximum independent set where, given a graph, the worst solution is the empty set, or of minimum vertex cover, where the worst solution is the vertex-set of the input-graph, or even of the

*LAMSADE, Université Paris-Dauphine, Place du Maréchal De Lattre de Tassigny, 75775 Paris Cedex 16, France

minimum graph-coloring where one can trivially color the vertices of the input-graph using a distinct color per vertex. On the contrary, for TSP things are very different. Let us take for example min_TSP. Here, given a graph K_n , the worst solution for K_n is a maximum total-distance Hamiltonian cycle, i.e., the optimal solution of max_TSP in K_n . The computation of such a solution is very far from being trivial since max_TSP is **NP**-hard. Obviously, the same holds when one considers max_TSP and tries to compute a worst solution for its instance, as well as for optimum satisfiability, for minimum maximal independent set and for many other well-known **NP**-hard problems. In order to remove ambiguities about the concept of the worst-value solution of an instance I of an **NPO** problem Π , we will define it as the optimal solution $\text{opt}(\Pi')$ of an **NPO** problem Π' having the same set of instances and feasibility constraints as Π verifying

$$\text{opt}(\Pi') = \begin{cases} \max & \text{opt}(\Pi) = \min \\ \min & \text{opt}(\Pi) = \max \end{cases}$$

In general, no apparent links exist between standard and differential approximations in the case of minimization problems, in the sense that there is no evident transfer of a positive, or negative, result from one framework to the other. Hence, a “good” differential-approximation result implies nothing for the behavior of the approximation algorithm studied when dealing with the standard framework and vice versa. When dealing with maximization problems, we show in [10] that the approximation of a maximization **NPO** problem Π within differential-approximation ratio δ implies its approximation within standard-approximation ratio δ .

The best known standard-approximation ratio known for min_TSP12 is $7/6$ ([11]), while the best known standard inapproximability bound is $743/742 - \epsilon$, for any $\epsilon > 0$ ([7]). On the other hand, the best known standard-ratio max_TSP is $3/4$ ([12]). To our knowledge, no better result is known in standard approximation for max_TSP12. Furthermore, no special study of TSP ab has been performed until now (a trivial standard-approximation ratio of b/a or a/b is in any case very easily deduced for min_ or max_TSP ab).

Here we show that min_ and max_TSP12, and min_ and max_TSP ab are all equi-approximable within $3/4$ for the differential approximation. We also prove that all these problems cannot be approximated better than within $741/742 + \epsilon$, for any $\epsilon > 0$. By the equi-approximability of min_TSP12, max_TSP12, min_TSP ab and max_TSP ab , the results obtained for the case of min_TSP12 apply to the rest of the problems above. Finally, we improve the *standard*-approximation ratio of max_TSP12 from $3/4$ ([12]) to $7/8$.

In what follows, we will denote by $V = \{v_1, \dots, v_n\}$ the vertex-set of K_n , by E its edge-set, and for $v_i v_j \in E$, we denote by $d(v_i, v_j)$ the distance of the edge $v_i v_j \in E$; we consider that the distance-vector is symmetric and integer. Given a feasible TSP-solution $T(K_n)$ of K_n (both min_ and max_TSP have the same set of feasible solutions), we denote by $d(T(K_n))$ its (objective) value. Given a graph G , we denote by $V(G)$ its vertex-set. Finally, given any set C of edges, we denote by $d(C)$ the total distance of C , i.e., the quantity $\sum_{v_i v_j \in C} d(v_i, v_j)$.

2 Differential-approximation preserving reductions for TSP12

In this section we give a differential-approximation preserving result that will be used later.

Theorem 1. *min_TSP12, max_TSP12, min_TSP ab and max_TSP ab are all equi-approximable for the differential approximation.*

Proof. In order to prove the theorem we will prove the following stronger lemma.

Lemma 1. *Consider any instance $I = (K_n, \vec{d})$ (where \vec{d} denotes the edge-distance vector of K_n). Then, any legal transformation $\vec{d} \mapsto \gamma \cdot \vec{d} + \eta \cdot \vec{1}$ of \vec{d} ($\gamma, \eta \in \mathbb{Q}$) produces differentially equi-approximable TSP-problems.*

Proof of lemma 1. Suppose that TSP can be approximately solved within differential-approximation ratio δ and remark that both the initial and the transformed instance have the same set of feasible solutions. By the transformation considered, the value $d(T(K_n))$ of any feasible tour $T(K_n)$ is affinely transformed into $\gamma d(T(K_n)) + \eta n$. Since differential-approximation ratio is stable under affine transformation, the equi-approximability of the original and of the transformed problem is immediately deduced, concluding so the proof of lemma 1. ■

We are ready now to continue the proof of theorem 1. In order to prove that min_TSP12 and max_TSP12 are equi-approximable, it suffices to apply lemma 1 proved just above with $\gamma = -1$ and $\eta = 3$. On the other hand, in order to prove that min_ or max_TSP12 reduces to min_ or max_TSPab, we apply lemma 1 with $\gamma = 1/(b-a)$ and $\eta = (b-2a)/(b-a)$, while for the converse reduction we apply lemma 1 with $\gamma = b-a$ and $\eta = 2a-b$. Since the reductions presented are transitive and composable, the equi-approximability of the pairs (min_TSP12, min_TSPab) and (TSP12, TSPab) proves the theorem. ■

For reasons of simplicity, we deal, in what follows, with min_TSP12. The differential-approximation results obtained can be immediately transferred, by theorem 1, to max_TSP12, min_TSPab and max_TSPab.

3 Approximating min_TSP12

Let us first recall that, given a graph G , a 2-matching is a set M of edges of G such that if $V(M)$ is the set of the endpoints of M , the vertices of the graph $(V(M), M)$ have degree at most 2; in other words, the graph $(V(M), M)$ is a collection of cycles and simple paths. A 2-matching is optimal if it is the largest over all the 2-matchings of G . It is called perfect if any vertex of the graph $(V(M), M)$ has degree equal to 2, i.e., if it constitutes a partition of $V(M)$ into cycles in G . Remark that determining a maximum 2-matching in a graph G is equivalent to determining a minimum total-distance vertex-partition into cycles into $G \cup \bar{G}$ (the complement of G), where the edges of G are considered of distance 1 and the ones of \bar{G} of distance 2.

As shown in [9], *an optimal triangle-free 2-matching can be computed in polynomial time*. As mentioned above, this amounts to computing a triangle-free minimum-distance collection of cycles in a complete graph K_n with edge-distances 1 and 2. Let us denote by M such a collection. Starting from M , we will progressively patch its cycles in order to finally obtain a unique Hamiltonian cycle in K_n .

3.1 Preprocessing M

We first define two operations, namely the *2-exchange* and the *2-patching*, implying two vertex-disjoint cycles of a 2-matching.

Definition 1. Let C_1 and C_2 be two vertex-disjoint cycles. Then:

- a *2-exchange* is any replacement of two edges $v_1u_1 \in C_1$, $v_2u_2 \in C_2$ by the edges v_1v_2 and u_1u_2 ;
- a *2-patching* of C_1 and C_2 is any cycle C resulting from a 2-exchange on C_1 and C_2 , i.e., $C = (C_1 \cup C_2 \setminus \{v_1u_1, v_2u_2\}) \cup \{v_1v_2, u_1u_2\}$, for any pair $(v_1u_1, v_2u_2) \in C_1 \times C_2$. ■

A 2-matching minimal with respect to the 2-exchange operation will be called *2-minimal*. In particular, if all edges have the same cost, then a 2-minimal matching is a tour.

Definition 2. A 2-matching $M = (C_1, C_2, \dots, C_{|M|})$ is 2-minimal if it verifies, $\forall (C_i, C_j) \in M \times M$, $C_i \neq C_j$, $\forall v_1u_1 \in C_i$, $\forall v_2u_2 \in C_j$, $d(v_1, v_2) + d(u_1, u_2) > d(u_1v_1) + d(u_2v_2)$. ■

In other words, a 2-matching M is 2-minimal if any 2-patching of its cycles produces a 2-matching of total distance strictly greater than the one of M . Starting from a 2-matching \hat{M} transformation of \hat{M} into a 2-minimal one M can be performed in polynomial time by the following procedure.

```

BEGIN *2_MIN*
  Mp ← ∅;
  REPEAT
    pick a new set {Ci, Cj} ⊆  $\hat{M}$ ;
    FOR all vkvl ∈ Ci, vpvq ∈ Cj DO
      take edges vkvl ∈ Ci and vpvq ∈ Cj;
      Cij1 ← Ci ∪ Cj \ {vkvl, vpvq} ∪ {vkvp, vlvq};
      Cij2 ← Ci ∪ Cj \ {vkvl, vpvq} ∪ {vkvq, vlvp};
      Cij ← argmin{d(Cij1), d(Cij2)};
      Mp ←  $\hat{M}$  \ {Ci, Cj} ∪ Cij;
      IF d( $\hat{M}$ ) > d(Mp) THEN  $\hat{M}$  ← Mp FI
    OD
  UNTIL no improvement of d( $\hat{M}$ ) is possible;
  OUTPUT M ←  $\hat{M}$ ;
END. *2_MIN*

```

Moreover, suppose that there exist two distinct cycles C and C' of M (the output of the procedure 2_MIN), both containing 2-edges and denote by $uv \in C$ and $u'v' \in C'$ two such edges. Then, $d(uu') + d(vv') \geq 4$, while $d(uv) + d(u'v') = 4$, a contradiction. So, the following proposition holds.

Proposition 1. *In any 2-minimal 2-matching, at most one of its cycles contains 2-edges.*

Remark 1. If the size of a 2-minimal triangle-free 2-matching M is 1, then, since a Hamiltonian tour is a special case of triangle-free 2-matching, M is an optimal min_TSP12-solution. Hence, in what follows we will suppose 2-matchings of size at least 2. ■

Assume now a 2-minimal triangle-free 2-matching $M = (C_1, \dots, C_p, C_0)$, verifying remark 1, where C_0 is the unique cycle of M (if any) containing 2-edges. Construct a graph $H = (V_H, E_H)$; $V_H = \{w_1, \dots, w_p\}$ contains a vertex per cycle of M and, for $i \neq j$, $w_i w_j \in E_H$ iff $\exists (u, v) \in C_i \times C_j$ such that $d(u, v) = 1$. Consider a maximum matching M_H , $|M_H| = q$, of H . With any edge $w_i w_j$ of M_H we associate the pair (C_{i^s}, C_{j^s}) of the corresponding cycles of M . So, M can be described (up to renaming its cycles) as

$$M = \bigcup_{s=1}^q \{C_1^s, C_2^s\} \bigcup_{t=1}^{r=p-2q} \{C_t\} \bigcup \{C_0\} \quad (1)$$

where for $s = 1, \dots, q$, $\exists e^s \in V(C_1^s) \times V(C_2^s)$ such that $d(e^s) = 1$.

Consider M as expressed in (1), denote by V_s the set of the four vertices of C_1^s and C_2^s adjacent to the endpoints of e^s , and construct the bipartite graph $B = (V_B^1 \cup V_B^2, E_B)$ where $V_B^1 = \{w_1, \dots, w_r\}$ (i.e., we associate a vertex with a cycle C_t , $t = 1, \dots, r$), $V_B^2 = \{w^1, \dots, w^q\}$ (i.e., we associate a vertex with a pair (C_1^s, C_2^s) , $s = 1, \dots, q$) and, $\forall (t, s)$, $w_t w^s \in E_B$ iff $\exists u \in C_t$, $\exists v \in V_s$ such that $d(u, v) = 1$. Compute a maximum matching M_B , $|M_B| = q'$ in B . With any edge $w_t w^s \in M_B$ we associate the triple (C_1^s, C_2^s, C_t) . So, M can be described (up to renaming its cycles) as

$$M = \bigcup_{s=1}^{q'} \{C_1^s, C_2^s, C_3^s\} \bigcup_{s=q'+1}^q \{C_1^s, C_2^s\} \bigcup_{t=1}^{r'=r-q'} \{C_t\} \bigcup \{C_0\} \quad (2)$$

where for $s = 1, \dots, q'$, $\exists f^s \in V_s \times V(C_3^s)$ such that $d(f^s) = 1$. In what follows we will reason with respect to M as it has been expressed in (2).

3.2 Computation and evaluation of the approximate solution and a lower bound for the optimal tour

In the sequel, call s.d.e.p. a set of vertex-disjoint elementary paths, denote by PREPROCESS the procedure that starting from a 2-minimal triangle-free 2-matching M leads to (2) and consider the following algorithm.

```

BEGIN (*TSP12*)
  compute a maximum 2-matching  $\hat{M}$  in  $K_n$ ;
   $M \leftarrow 2\_MIN(\hat{M})$ ;
   $M \leftarrow PREPROCESS(M)$ ;
   $D \leftarrow \emptyset$ ;
(1) FOR  $s \leftarrow 1$  TO  $q'$  DO
  let  $g_1^s$  be the edge of  $C_1^s$  adjacent to both  $e^s$  and  $f^s$ ;
  choose in  $C_2^s$  an edge  $g_2^s$  adjacent to  $e^s$ ;
  choose in  $C_3^s$  an edge  $g_3^s$  adjacent to  $f^s$ ;
   $D \leftarrow (D \cup C_1^s \cup C_2^s \cup C_3^s \setminus \{g_1^s, g_2^s, g_3^s\}) \cup \{e^s, f^s\}$ ;
  OD
(2) FOR  $s \leftarrow q' + 1$  TO  $q$  DO
  choose in  $C_1^s$  an edge  $g_1^s$  adjacent to  $e^s$ ;
  choose in  $C_2^s$  an edge  $g_2^s$  adjacent to  $e^s$ ;
   $D \leftarrow (D \cup C_1^s \cup C_2^s \setminus \{g_1^s, g_2^s\}) \cup \{e^s\}$ ;
  OD
(3) FOR  $t \leftarrow 1$  TO  $r'$  DO
  choose any edge  $g_t$  in  $C_t$ ;
   $D \leftarrow (D \cup C_t) \setminus \{g_t\}$ ;
  OD
(4) IF there exists in  $C_0$  an 1-edge
  THEN choose a 2-edge  $g_0$  of  $C_0$  adjacent to an 1-edge  $e$ ;
  ELSE choose any edge  $g_0$  of  $C_0$ ;
   $D \leftarrow (D \cup C_0) \setminus \{g_0\}$ ;
  FI
(5) complete  $D$  in order to obtain a Hamiltonian tour  $T(K_n)$ ;
  OUTPUT  $T(K_n)$ ;
END (*TSP12*)

```

Clearly, both achievement of a 2-minimal triangle free 2-matching and PREPROCESS can be performed in polynomial time. Moreover, steps (1) to (4) are also executed in polynomial time. Finally, step (5) can be performed by arbitrarily ordering (mod $|D|$) the chains of the s.d.e.p. D and then, for $i = 1, \dots, |D|$, adding in D the edge linking the “last” vertex of chain i to the “first” vertex of chain $i + 1$. Consequently, the whole algorithm TSP12 is polynomial. Finally, remark that C_0 may contain either only 2-edges, or both 1- and 2-edges. In the latter case, edge g_0 (in step 4) can be any 2-edge in C_0 , adjacent to an 1-edge of C_0 .

Lemma 2. $d(T(K_n)) \leq d(M) + q + r'$.

Proof. During steps (1) to (4) of algorithm TSP12, set D remains a s.d.e.p. At the end of step (4), D contains M minus the $3q' + 2(q - q') + r' = q' + 2q + r'$ 1-edges of the set

$\cup_{s=1}^{q'} \{g_1^s, g_2^s, g_3^s\} \cup_{s=q'+1}^q \{g_1^s, g_2^s\} \cup_{s=1}^{r'} \{g_t\}$ minus (if $C_0 \neq \emptyset$) one 2-edge of C_0 plus the $2q' + (q - q') = q' + q$ 1-edges of the set $\cup_{s=1}^{q'} \{e_1^s, f_2^s\} \cup_{s=q'+1}^q \{e^s\}$. So D is a s.d.e.p. of size $n - (q + r') - \mathbf{1}_{C_0 \neq \emptyset}$ and of total distance $d(M) - (q + r') - 2 \cdot \mathbf{1}_{C_0 \neq \emptyset}$. Completion of D in order to obtain a tour in K_n , can be done by adding $q + r' + \mathbf{1}_{C_0 \neq \emptyset}$ new edges. Each of these new edges can be, at worst, of distance 2. We so have $d(T(K_n)) \leq d(M) - (q + r') + 2 \cdot \mathbf{1}_{C_0 \neq \emptyset} + 2(q + r' + \mathbf{1}_{C_0 \neq \emptyset}) = d(M) + q + r'$, q.e.d. ■

On the other hand, the optimal tour being a special triangle-free 2-matching, the following lemma holds immediately.

Lemma 3. $\beta(K_n) \geq d(M)$.

3.3 Evaluation of the worst-value solution

In what follows in this section we will exhibit a s.d.e.p., with all edges of distance 2 (called 2-s.d.e.p.). Given such a s.d.e.p. W , one can proceed as in step (5) of algorithm TSP12 (section 3.2), in order to construct a Hamiltonian tour T_w whose total distance is a lower bound for $\omega(K_n)$.

Denote by $E2$ the set of 2-edges of cycle C_0 . If $q = 0$, i.e., $M_H = \emptyset$, and if $C_0 = E2$, then the tour computed by TSP12 is optimal.

Lemma 4. *If $q = 0$ and $C_0 = E2$, then $\delta_{\text{TSP12}}(K_n) = 1$.*

Proof. Let $k = |V(C_0)| = d(M) - n$ and set $V(C_0) = \{a_1, \dots, a_k\}$. By the fact that M is 2-minimal, all the edges of K_n incident to these vertices have distance 2. On the other hand, between two distinct cycles in the set $\{C_1, \dots, C_{p=r'}\}$ of M , there exist only edges of distance 2. Consider the family $\mathcal{F} = \{\{a_1\}, \dots, \{a_k\}, V(C_1), \dots, V(C_p)\}$. By the above remarks, any edge linking vertices of two distinct sets of \mathcal{F} is a 2-edge. Any feasible tour of K_n (a posteriori an optimal one) integrates the $k + p$ sets of \mathcal{F} by using at least $k + p$ 2-edges pairwise linking these sets. Hence, any tour uses at least $k + p$ 2-edges, so does tour $T(K_n)$ computed by algorithm TSP12, q.e.d. ■

So, we suppose in the sequel that $q = 0 \Rightarrow C_0 \neq E2$. We will now prove the existence of a 2-s.d.e.p. W of size $d(M) + 4(q + r') - n$, where M is as expressed by (2). In the sequel, a path with alternating vertices will denote a path such that no two consecutive vertices lie in the same cycle.

Proposition 2. *Between two cycles C_a and C_b of M of size at least k , there always exists a path with alternating vertices from C_a and C_b , which contains at least k 2-edges.*

Proof. Let $\{a_1, \dots, a_{k+1}\}$ and $\{b_1, \dots, b_{k+1}\}$ be $k+1$ successive vertices of two distinct cycles C_a and C_b of size at least k (possibly $a_1 = a_{k+1}$ if $|V(C_a)| = k$ and $b_1 = b_{k+1}$ if $|V(C_b)| = k$). We will show that there exists a path with alternating vertices from C_a and C_b of size $2k - 1$ and of distance at least $3k - 1$. Consider paths $C = \cup_{i=1}^k \{a_i b_i\} \cup_{i=1}^{k-1} \{a_{i+1} b_i\}$ and $D = \cup_{i=2}^{k+1} \{a_i b_i\} \cup_{i=1}^{k-1} \{a_i b_{i+1}\}$. By the 2-minimality of M we get:

$$\begin{aligned} \forall i = 1, \dots, k \quad \max \{d(a_i, b_i), d(a_{i+1}, b_{i+1})\} = 2 &\Rightarrow d(a_i, b_i) + d(a_{i+1}, b_{i+1}) \geq 3 \\ \forall i = 1, \dots, k-1 \quad \max \{d(a_i, b_{i+1}), d(a_{i+1}, b_i)\} = 2 &\Rightarrow d(a_i, b_{i+1}) + d(a_{i+1}, b_i) \geq 3 \end{aligned}$$

Summing the terms of the expression above member-by-member, one obtains:

$$\begin{aligned} \sum_{i=1}^k (d(a_i, b_i) + d(a_{i+1}, b_{i+1})) + \sum_{i=1}^{k-1} (d(a_{i+1}, b_i) + d(a_i, b_{i+1})) &\geq 6k - 3 \\ \Leftrightarrow d(C) + d(D) \geq 6k - 3 &\Rightarrow \max \{d(C), d(D)\} \geq \left\lceil \frac{6k - 3}{2} \right\rceil = 3k - 1 \quad \blacksquare \end{aligned}$$

Application of proposition 2 to any pair (C_1^s, C_2^s) of M results in the following claim.

Claim 1. $\forall s = 1, \dots, q$, there exists a 2-s.d.e.p. W^s of size 4, alternating vertices of cycles C_1^s and C_2^s , containing a vertex of V_s whose degree with respect to W^s is 1.




Figure 1: An example of claim 1.

In figure 1, we show an application of claim 1. We assume $e^s = a_2b_2$; then $\{a_1, b_1\} \subset V_s$. The 2-s.d.e.p. W^s claimed is $\{b_1a_2, (a_3b_3, b_3a_4), a_5b_5\}$ and the degree of b_1 with respect to W^s is 1.

Consider now the s.d.e.p. $W_t^s = W^s \cup W_t'^s$, where W^s as in claim 1 and $W_t'^s$ is any path of size 4 with alternating vertices from C_3^s and C_t , $s = 1, \dots, q'$, $t = 1, \dots, r'$. By the optimality of M_H , any edge linking vertices of C_3^s to vertices of C_t is a 2-edge. Consequently, W_t^s is a 2-s.d.e.p. and the following claim holds.

Claim 2. $\forall s = 1, \dots, q', \forall t = 1, \dots, r'$, there exists a 2-s.d.e.p. W_t^s of size 8, alternating vertices of the cycles C_1^s and C_2^s , and of the cycles C_3^s and C_t .

For $s = q' + 1, \dots, q$, $t = 1, \dots, r'$, consider the triplet (C_1^s, C_2^s, C_t) . Let $e^s = e_1^s e_2^s$, $V_s = \{u_1^s, v_1^s, u_2^s, v_2^s\}$ and consider any four vertices a_t, b_t, c_t and d_t of C_t . By the optimality of M_B , any vertex of C_t is linked to any vertex of V_s exclusively by 2-edges. Moreover, the 2-minimality of M implies that at least one of $u_1^s e_2^s$ and $e_1^s u_2^s$ is of distance 2. If we suppose $d(u_1^s, e_2^s) = 2$ (figure 2), then the path $\{e_2^s, u_1^s, a_t, v_1^s, b_t, u_2^s, c_t, v_2^s, d_t\}$ is a 2-s.e.d.p. Hence, the following claim holds.

Claim 3. $\forall s = q' + 1, \dots, q, \forall t = 1, \dots, r'$, there exists a 2-s.d.e.p. W_t^s of size 8, alternating vertices of the cycles C_1^s , C_2^s and C_t .




Figure 2: The 2-s.d.e.p. W_t^s of claim 3.

Let $r' \geq 2$ and consider, for $t = 1, \dots, r'$, the (residual) cycles C_t . All edges between these cycles are of distance 2. If we denote by a_t, b_t, c_t and d_t four vertices of C_t , the path $\{a_1, \dots, a_t, \dots, a_{r'}, b_1, \dots, b_t, \dots, b_{r'}, c_1, \dots, c_t, \dots, c_{r'}, d_1, \dots, d_t, \dots, d_{r'}\}$ is a 2-s.d.e.p. of size $4r' - 1$ and the following claim holds.

Claim 4. *If $r' \geq 2$, then there exists a 2-s.d.e.p. $W^{r'}$ of size $4r' - 1$ alternating vertices of cycles $C_t, t = 1, \dots, r'$.*

Lemma 5. *If $q \geq 1$, or if $[(C_0 \neq E_2) \text{ and } (r \neq 1)]$, then $\delta_{\text{TSP12}}(K_n) \geq 3/4$.*

Proof. Consider M as expressed in (2). Let us denote by W' a 2-s.e.d.p. on $V \setminus V(C_0)$. From claims 1, 2, 3 and 4:

$$W' = \begin{cases} \bigcup_{s=1}^{r'} W_s^s \bigcup_{s=r'+1}^q W^s & |W'| = 8r' + 4(q - r') = 4(q + r') & q \geq r' \\ \bigcup_{s=1}^q W^s \cup W^{r'} \cup \{\gamma\} & |W'| = 4q + 4r' - 1 + 1 = 4(q + r') & r' > q > 0 \\ W^{r'} & |W'| = 4r' - 1 = 4(q + r') - 1 & r' \geq 2, q = 0 \end{cases} \quad (3)$$

In (3), γ draws an edge linking a vertex of degree 1 with respect to $W^{r'}$ to a vertex of degree 1 in $V(W')$. This last vertex can belong either to $V(C_{\frac{1}{3}})$ if $q' \geq 1$, or to V_q otherwise. By the optimality of M_H and M_B , γ is a 2-edge. For the first line of (3), remark that if $q = r'$, then W' is given by $\bigcup_{s=1}^{r'} W_s^s$, and claims 2 and 3 conclude $W' = 8r' = 4(r' + q)$; otherwise ($q > r'$), claims 1, 2 and 3 conclude the first line of (3). For the second line ($r' > q > 0$), since r' and q are integers, we have $r' \geq 2$. Then, by claim 4, $|W^{r'}| \geq 4r' - 1$ and the expression of the second line follows from the fact that γ is a 2-edge. For the third line of (3), claim 4 gives immediately the result. Hence, in any case, W' is a 2-s.d.e.p. verifying $|W'| = 4(q + r') - 1$ if $q = 0$, $|W'| = 4(q + r')$ otherwise.

We now construct a 2-s.e.d.p. W_0 on $V(C_0)$. Let $g_0 = u_0v_0$ be the edge removed from C_0 during the execution of step (4) of algorithm TSP12. If $C_0 \neq E_2$, then g_0 has been chosen in such a way that one of its endpoints, say v_0 , is adjacent in C_0 to a 1-edge. Let γ' be an edge linking v_0 to a vertex of degree 1 with respect to W' (such a vertex exists since W' is acyclic). By the 2-minimality of M , $d(\gamma') = 2$. Set

$$W_0 = \begin{cases} E_2 \setminus \{g_0\} \cup \{\gamma'\} & |W_0| = d(M) - n & C_0 = E_2 \\ E_2 \cup \{\gamma'\} & |W_0| = d(M) - n + 1 & \text{otherwise} \end{cases}$$

Setting finally $W = W' \cup W_0$, one obtains the 2-s.d.e.p. claimed at the beginning of section 3.3. We have $d(W) = 4(q + r') - \mathbf{1}_{q=0} + d(M) - n + \mathbf{1}_{C_0 \neq E_2} \geq d(M) - n + 4(q + r')$, the last inequality holding because of the hypothesis $q = 0 \Rightarrow C_0 \neq E_2$ made just before lemma 2. Any completion of W in a Hamiltonian cycle of K_n (as in step (5) of algorithm TSP12) would produce a tour T_w of total distance $d(T_w) \geq d(M) + 4(q + r')$. So, $\omega(K_n) \geq d(M) + 4(q + r')$, and combining this expression together with lemmata 2 and 3, the differential ratio $3/4$ is immediately concluded. ■

Lemma 6. *If $q = 0$ and $r = 1$ and $C_0 \neq E_2$, then $\delta_{\text{TSP12}}(K_n) \geq 3/4$.*

Proof. Recall that, by the optimality of M_H , if $q = 0$ and $r = 1$, all edges linking cycles C_0 and C_1 are of distance 2. Let u_0, u_1 and u_2 be three vertices of C_0 such that $d(u_0, u_1) = 1$ and $d(u_1, u_2) = 2$. We denote by u_2 a neighbor of u_1 with respect to C_0 , and by $u_3 \neq u_1$ the neighbor of u_2 in C_0 . Let a, b, c and d be any four vertices of C_1 , and let W_p and W_m be the sets $\{au_1, u_1b, bu_2, u_2c\}$ and $\{u_1u_2\}$, respectively.




Figure 3:

If u_3 is adjacent to two 1-edges in C_0 , i.e., $d(u_2, u_3) = d(u_3, u_4) = 1$, then set $W_p = W_p \cup \{cu_4\}$ (figure 3(a)).

If only one of u_2u_3 and u_3u_4 is of distance 2, then set $W_p = W_p \cup \{cu_3, u_3d\}$ and $W_m = W_m \cup \{\operatorname{argmax}\{d(u_2, u_3), d(u_3, u_4)\}\}$. In figure 3(b), sets W_p and W_m are shown for the case where $\operatorname{argmax}\{d(u_2, u_3), d(u_3, u_4)\} = u_3u_4$, i.e., $d(u_3, u_4) = 2$ and $d(u_2, u_3) = 1$.

Finally, if u_2u_3 and u_3u_4 are both of distance 2 ($d(u_2, u_3) = d(u_3, u_4) = 2$), then set $W_p = W_p \cup \{cu_3, u_3d, du_4\}$ and $W_m = W_m \cup \{u_2u_3, u_3u_4\}$ (figure 3(c)).

In any of the above cases, we consider the 2-s.d.e.p. $W = E_2 \cup W_p \setminus W_m$ with $|W| = d(M) - n + 4 = d(M) - n + 4r'$. We then have $\omega(K_n) \geq d(M) + 4r'$, and combining it with lemmata 2 and 3, the differential ratio $3/4$ is concluded. ■

In all, combining lemmata 2, 3, 4 and 6, the following theorem can be immediately proved.

Theorem 2. *min-TSP12 is approximable within differential-approximation ratio $3/4$.*

Theorems 1 and 2 induce the following corollary.

Corollary 1. *min-TSP12, max-TSP12, min-TSPab and max-TSPab are approximable within differential-approximation ratio $3/4$.*




Figure 4: Tightness of the TSP12 approximation ratio.

Proposition 3. *Ratio $3/4$ is tight for TSP12*

Proof. Consider two cliques and number their vertices by $\{1, \dots, 4\}$ and by $\{5, 6, \dots, n + 8\}$, respectively. Edges of both cliques have all distance 1. Cross-edges ij , $i = 1, 3$, $j = 5, \dots, n + 8$,

	k	TSP12	The algorithm of [11]
Differential ratio	3	0.931100364	0.846702091
	4	0.9000002	0.833333
	5	0.920289696	0.833333
	6	0.9222222	0.833333
Standard ratio	3	0.923350955	0.87013
	4	0.9094018	0.857143
	5	0.92646313	0.857143
	6	0.928178	0.857143

Table 1: A limited comparison between TSP12 and the algorithm of [11] on some worst-case instances of the latter.

are all of distance 2, while every other cross-edge is of distance 1. Unraveling of TSP12 will produce: $T = \{1, 2, 3, 4, 5, 6, \dots, n+7, n+8, 1\}$ (cycle-patching on edges $(1, 4)$ and $(5, n+8)$), while $T_w = \{1, 5, 2, 6, 3, 7, 4, 8, 9, \dots, n+7, n+8, 1\}$ (using 2-edges $(1, 5)$, $(6, 3)$, $(3, 7)$ and $(n+8, 1)$) and $T^* = \{1, 2, n+8, n+7, \dots, 5, 4, 3, 1\}$ (using 1-edges $(4, 5)$ and $(2, n+8)$). Consequently, $\delta_{\text{TSP12}}(K_{n+8}) = 3/4$, q.e.d. ■

In figure 4, the tours T^* and T_w of proposition 3 are shown for $n = 2$. We assume $T = \{1, \dots, 10, 1\}$.

Let us note that the differential approximation ratio of the 7/6-algorithm of [11], when running on K_{n+8} , is also 3/4. The authors of [11] exhibit a family of worst-case instances for their algorithm: one has k cycles of length 4 arranged around a cycle of length $2k$. We have performed a limited comparative study between their algorithm and ours, for $k = 3, 4, 5, 6$ (on 24 graphs). The average differential and standard approximation ratios for the two algorithms are presented in table 1.

Proposition 4. *min-TSPab is approximable within standard-approximation ratio $\rho \leq (1 + (b-a)/(4a))$. This ratio tends to ∞ when $a = o(b)$.*

Proof. Revisit corollary 1. Differential ratio 3/4 for min-TSPab implies $\lambda(K_n)/\beta(K_n) \leq (3/4) + (\omega(K_n)/(4\beta(K_n)))$. Using $\omega(K_n) \leq bn$ and $\beta(K_n) \geq an$, some easy algebra gives the result claimed. ■

Theorem 3. *min- and max-TSPab and min- and max-TSP12 are inapproximable within differential-ratio of at least $742/743 + \epsilon$, $\forall \epsilon > 0$, unless $P=NP$.*

Proof. Consider min-TSP12. Using $n \leq \beta(K_n) \leq \omega(K_n) \leq 2n$, one can see that approximation of min-TSP12 within $\delta = 1 - \epsilon$ implies its approximation within $\rho = 2 - (1 - \epsilon) = 1 + \epsilon$, $0 \leq \epsilon \leq 1$. Then, the inapproximability bound $(743/742 - \epsilon)$ of [7] for min-TSP12 together with theorem 1 conclude the proof. ■

4 An improvement of the standard ratio for the maximum traveling salesman with distances 1 and 2

We propose in this section a non-trivial improvement of the standard-approximation ratio for max-TSP12, by proving the following theorem.

Theorem 4. *max_TSP12 is polynomially approximable within standard-approximation ratio bounded below by $7/8$.*

Proof. Combining, as in the proof of proposition 4, expressions $\delta_{\max_TSP12} \geq 3/4$, $\omega_{\max}(K_n) \geq an$ and $\beta_{\max}(K_n) \leq bn$, one deduces $\rho_{\max_TSP12} \geq (3/4) + (a/4b)$. Setting $a = 1$ and $b = 2$, the result claimed follows immediately. ■

Remark 2. Consider now the following simple way to use the algorithm of [11] in order to solve max_TSP12 on a graph K_n with edge-distances 1 and 2. The graph \bar{K}_n in the first line of the algorithm MTSPALG just above is such that the distance of an edge e of \bar{K}_n is 1 if e has distance 2 in K_n , and 2 if e has distance 1 in K_n .

```
BEGIN *MTSPALG*
  construct  $\bar{K}_n$ ;
  call the algorithm of [11] to compute a tour  $T_{\min}(\bar{K}_n)$ ;
  OUTPUT  $T_{\max}(K_n) \leftarrow T_{\min}(\bar{K}_n)$ ;
END.
```

Let us denote by **A** the $7/6$ -algorithm of [11] called in the second line of MTSPALG. Then, by theorem 1, $\beta(K_n) = 3n - \beta(\bar{K}_n)$ and $\lambda_{\text{MTSPALG}}(K_n) = 3n - \lambda_{\mathbf{A}}(\bar{K}_n)$, where $\beta(K_n)$ is the optimal value for max_TSP12 on K_n and $\beta(\bar{K}_n)$ is the optimal value for min_TSP12 on \bar{K}_n . Then, $\lambda_{\mathbf{A}}(\bar{K}_n)/\beta(\bar{K}_n) \leq 7/6$, together with $\beta(\bar{K}_n) \geq n$ imply $\lambda_{\text{MTSPALG}}(K_n)/\beta(K_n) \geq 2/3$. ■

Note finally that standard-approximation ratio $7/8$ can be obtained by the following direct method.

```
BEGIN *max_TSP12*
  find a triangle-free 2-matching  $M = \{C_1, C_2, \dots\}$ ;
  FOR all  $C_i$  DO delete a minimum-distance edge from  $C_i$  OD
  properly link the remaining paths to obtain a Hamiltonian cycle  $T$ ;
  OUTPUT  $T$ ;
END. *max_TSP12*
```

Let p be the number of cycles of M where 2-edges have been removed during the FOR-loop of algorithm max_TSP12. Then, $\lambda_{\max_TSP12}(K_n) \geq d(M) - p$, $\beta(K_n) \leq d(M)$, and since M is triangle-free, $d(M) \geq 8p$. Consequently, $\lambda_{\max_TSP12}(K_n)/\beta_{\max}(K_n) \geq 7/8$.

The above result, can be extended to the case of max_TSP ab if we consider that here $\lambda_{\max_TSP12}(K_n) \geq d(M) - p(b - a)$, $\beta(K_n) \leq d(M)$ and $d(M) \geq 4bp$. Hence the following corollary holds and concludes the paper.

Corollary 2. *max_TSP ab is polynomially approximable within standard-approximation ratio bounded below by $(3/4) + (a/(4b))$.*

References

- [1] A. Aiello, E. Burattini, M. Furnari, A. Massarotti, and F. Ventriglia. Computational complexity: the problem of approximation. In C. M. S. J. Bolyai, editor, *Algebra, combinatorics, and logic in computer science*, volume I, pages 51–62, New York, 1986. North-Holland.
- [2] G. Ausiello, A. D’Atri, and M. Protasi. Structure preserving reductions among convex optimization problems. *J. Comput. System Sci.*, 21:136–153, 1980.

- [3] G. Ausiello, A. Marchetti-Spaccamela, and M. Protasi. Towards a unified approach for the classification of NP-complete optimization problems. *Theoret. Comput. Sci.*, 12:83–96, 1980.
- [4] M. Bellare and P. Rogaway. The complexity of approximating a nonlinear program. *Math. Programming*, 69:429–441, 1995.
- [5] M. Demange, P. Grisoni, and V. T. Paschos. Differential approximation algorithms for some combinatorial optimization problems. *Theoret. Comput. Sci.*, 209:107–122, 1998.
- [6] M. Demange and V. T. Paschos. On an approximation measure founded on the links between optimization and polynomial approximation theory. *Theoret. Comput. Sci.*, 158:117–141, 1996.
- [7] L. Engebretsen and M. Karpinski. Approximation hardness of TSP with bounded metrics. Report 89, Electr. Colloq. Computational Comp., 2000.
- [8] M. R. Garey and D. S. Johnson. *Computers and intractability. A guide to the theory of NP-completeness*. W. H. Freeman, San Francisco, 1979.
- [9] D. B. Hartvigsen. *Extensions of matching theory*. PhD thesis, Carnegie-Mellon University, 1984.
- [10] J. Monnot, V. T. Paschos, and S. Toulouse. Differential approximation results for the traveling salesman problem. Cahier du LAMSADE 172, LAMSADE, Universit Paris-Dauphine, 2000.
- [11] C. H. Papadimitriou and M. Yannakakis. The traveling salesman problem with distances one and two. *Math. Oper. Res.*, 18:1–11, 1993.
- [12] A. I. Serdyukov. An algorithm with an estimate for the traveling salesman problem of the maximum. *Upravlyaemye Sistemy*, 25:80–86, 1984.
- [13] S. A. Vavasis. Approximation algorithms for indefinite quadratic programming. *Math. Programming*, 57:279–311, 1992.
- [14] E. Zemel. Measuring the quality of approximate solutions to zero-one programming problems. *Math. Oper. Res.*, 6:319–332, 1981.