



HAL
open science

Limit theorems for iterated random topical operators

Glenn Merlet

► **To cite this version:**

| Glenn Merlet. Limit theorems for iterated random topical operators. 2005. hal-00004594v1

HAL Id: hal-00004594

<https://hal.science/hal-00004594v1>

Preprint submitted on 28 Mar 2005 (v1), last revised 8 Jan 2007 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Limit theorems for iterated random topological operators

Glenn MERLET
glenn.merlet@univ-rennes1.fr

March 28, 2005

Abstract

Let $A(n)$ be a sequence of i.i.d. topical (i.e. isotone and additively homogeneous) operators. Let $x(n, x_0)$ be defined by $x(0, x_0) = x_0$ and $x(n, x_0) = A(n)x(n-1, x_0)$. This can modelize a wide range of systems including, task graphs, train networks, Job-Shop, timed digital circuits or parallel processing systems.

When $A(n)$ has the memory loss property, we use the spectral gap method to prove limit theorems for $x(n, x_0)$. Roughly speaking, we show that $x(n, x_0)$ behaves like a sum of i.i.d. real variables. Precisely, we show that with suitable additional conditions, it satisfies a central limit theorem with rate, a local limit theorem, a renewal theorem and a large deviations principle, and we give an algebraic condition to ensure the positivity of the variance in the CLT. When $A(n)$ are defined by matrices in the $(\max, +)$ semi-ring, we give more effective statements and show that the additional conditions and the positivity of the variance in the CLT are generic.

Introduction

An operator $A : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is called additively homogeneous if it satisfies $A(x + a\mathbf{1}) = A(x) + a\mathbf{1}$ for all $x \in \mathbb{R}^d$ and $a \in \mathbb{R}$, where $\mathbf{1}$ is the vector $(1, \dots, 1)'$ in \mathbb{R}^d . It is called isotone if $x \leq y$ implies $A(x) \leq A(y)$, where the order is the product order on \mathbb{R}^d . It is called topical if it is isotone and homogeneous. The set of topical operators on \mathbb{R}^d will be denoted by Top_d .

We recall that the action of matrices with entries in $\mathbb{R}_{\max} = \mathbb{R} \cup \{-\infty\}$ on \mathbb{R}_{\max}^d is defined by $(Ax)_i = \max_j (A_{ij} + x_j)$. When matrix A has no line of $-\infty$, the restriction of this action to \mathbb{R}^d defines a topical operator, also

denoted by A . Such operators are called $(\max, +)$ operators and composition of operators corresponds to the product of matrices in the $(\max, +)$ semi-ring.

Products of random matrices in the usual sense have been intensively investigated. Let us cite H. Furstenberg [Fur63], Y. Guivarc'h and A. Raugi [GR85] or I. Ya. Gol'dsheid and G. A. Margulis [GdM89]. The interested reader can find a presentation of this theory in [BL85]. We investigate analogous problems to those studied by É. Le Page [LP82], but for matrices in the $(\max, +)$ semi-ring and more generally for iterated topological operators.

Let $(A(n))_{n \in \mathbb{N}}$ be a sequence of random topological operators on \mathbb{R}^d . Let $x(n, x_0)$ be defined by

$$\begin{cases} x(0, x_0) &= x_0 \\ x(n, x_0) &= A(n)x(n-1, x_0). \end{cases} \quad (1)$$

We are interested in the asymptotic behavior of $x(n, \cdot)$. We show that with suitable additional conditions, it satisfies a central limit theorem, a local limit theorem, a renewal theorem and a large deviations principle. When the $A(n)$ are $(\max, +)$ operators we give more explicit results.

This class of system can modelize a wide range of situations. A review of applications can be found in the last section of [BM98]. Among other examples the $(\max, +)$ case has been applied to modelize task graphs, cyclic Jackson networks (J. Mairesse [Mai97]) or Job-Shop (G. Cohen et al. [CDQV85]).

This article is divided in four parts. First we present the model of iterated topological operators, including a short review of known limit theorems. Second we state our results and comment them. Third, we recall the formalism of abstract Markov chains and the limit theorems obtained by the spectral gap method that are the key elements of the proofs. Finally we prove our results.

1 Iterated topological operators

1.1 Memory loss property

Dealing with homogeneous operators it is natural to introduce the quotient space of \mathbb{R}^d by the equivalence relation \sim defined by $x \sim y$ if $x - y$ is proportional to $\mathbf{1}$. This space will be called projective space and denoted by $\mathbb{P}\mathbb{R}_{\max}^d$. Moreover \bar{x} will be the equivalence class of x .

The application $\bar{x} \mapsto (x_i - x_j)_{i < j}$ embeds $\mathbb{P}\mathbb{R}_{\max}^d$ onto a subspace of $\mathbb{R}^{\frac{d(d-1)}{2}}$ with dimension $d - 1$. The supremum norm of $\mathbb{R}^{\frac{d(d-1)}{2}}$ therefore induces a distance on $\mathbb{P}\mathbb{R}_{\max}^d$ which will be denoted by δ . A direct computation shows that $\delta(\bar{x}, \bar{y}) = \max_i(x_i - y_i) + \max_i(y_i - x_i)$. By a slight abuse, we will also write $\delta(x, y)$ for $\delta(\bar{x}, \bar{y})$. The projective norm of x will be $|x|_{\mathcal{P}} = \delta(x, 0)$.

Let us recall two well known facts about topological operators. First a topological operator is non-expanding with respect to the infinity norm. Second the operator it defines from $\mathbb{P}\mathbb{R}_{\max}^d$ to itself is non-expanding for δ .

The key property for our proofs is the following:

Definition 1.1.

1. A topological operator A is said to have rank 1, if it defines a constant operator on $\mathbb{P}\mathbb{R}_{\max}^d$: \overline{Ax} does not depend on $x \in \mathbb{R}^d$.
2. The sequence $(A(n))_{n \in \mathbb{N}}$ of Top_d -valued random variables is said to have the memory loss (MLP) property if there exists an N such that $A(N) \cdots A(1)$ has rank 1 with positive probability.

This notion has been introduced by J. Mairesse [Mai97], the $A(n)$ being $(\max, +)$ operators. The denomination rank 1 is natural for $(\max, +)$ operators.

We proved in [Mer04] that this property is generic for i.i.d. $(\max, +)$ operators: it is fulfilled when the support of the law of $A(1)$ is not included in a finite union of hyperplanes.

Although this result could suggest the opposite, the MLP depends on the law of $A(1)$, and not only on its support : if $(U(n))_{n \in \mathbb{N}}$ is an i.i.d. sequence with the support of $U(1)$ equal to $[0, 1]$, and $A(n)$ are the $(\max, +)$ operators defined by the matrices

$$A(n) = \begin{pmatrix} -U(n) & 0 \\ 0 & -U(n) \end{pmatrix},$$

then $(A(n))_{n \in \mathbb{N}}$ has the MLP property iff $\mathbb{P}(U(n) = 0) > 0$.

The weaker condition that there is an operator with rank 1 in the closed semigroup generated by the support of the law of $A(1)$ has been investigated by J. Mairesse for $(\max, +)$ operators. It ensures the weak convergence of $\overline{x}(n, \cdot)$ but does not seem appropriate for our construction.

1.2 Known results

Before describing our analysis, we give a brief review of published limit theorems about $x(n, X^0)$.

J.M. Vincent has proved a first law of large number:

Theorem 1.1 ([Vin97]). *Let $(A(n))_{n \in \mathbb{N}}$ be a stationary ergodic sequence of topological operators and X^0 an \mathbb{R}^d -valued random variable. If $A(1)$ has rank 1 and X^0*

are integrable, then there exists $\overline{\gamma}$ and $\underline{\gamma}$ in \mathbb{R} such that

$$\begin{aligned}\lim_n \frac{\max_i x_i(n, X^0)}{n} &= \overline{\gamma} \text{ a.s.} \\ \lim_n \frac{\min_i x_i(n, X^0)}{n} &= \underline{\gamma} \text{ a.s.}\end{aligned}$$

F. Baccelli and J. Mairesse give a condition to ensure $\overline{\gamma} = \underline{\gamma}$, hence the convergence of $\frac{x(n, X^0)}{n}$:

Theorem 1.2 ([BM98]). *Let $(A(n))_{n \in \mathbb{N}}$ be a stationary ergodic sequence of topical operators and X^0 an \mathbb{R}^d -valued random variable such that $A(1) \cdot \mathbf{1}$ and X^0 are integrable. If there exists an N , such that $A(N) \cdots A(1)$ has a bounded projective image with positive probability, then there exists γ in \mathbb{R} such that*

$$\lim_n \frac{x(n, X^0)}{n} = \gamma \mathbf{1} \text{ a.s.}$$

In this case γ is called the Lyapunov exponent of the sequence. We notice that the MLP property implies a bounded projective image with positive probability.

The following result has been proved by J. Mairesse when the $A(n)$ are $(\max, +)$ operators, but can be extended to topical operators with the same proof. It will be the key point to ensure the spectral gap.

Theorem 1.3 ([Mai97]). *Let $(A(n))_{n \in \mathbb{N}}$ be a stationary ergodic sequence of topical operators with the MLP property. Then $\overline{x}(n, x_0)$ converges in total variation uniformly in x_0 .*

To end this section we mention two limit theorems, which are close to ours, but obtained by different ways. We will compare those results to ours in section 2.3.

With a martingale method J. Resing et al. [RdVH⁺90] have obtained a central limit theorem for $x(n, X^0)$, when the Markov chain $\overline{x}(n, \cdot)$ is aperiodic and uniformly Φ -recurrent. The theorem has been stated for $(\max, +)$ operators, but it should make no difference to use topical ones.

With a subadditivity method, F. Toomey [Too02] has proved a large deviation principle for $x(n, x_0)$ when the projective image of $A(N) \cdots A(1)$ is bounded.

1.3 Principle of the analysis

From now on, $(A(n))_{n \in \mathbb{N}}$ is an i.i.d sequence of topical operators with the MLP property.

The first step of the proof is to split our Markov chain $x(n, \cdot)$ into another Markov chain and a sum of cocycles over this chain, following what É. Le Page made for products of random matrices. For any topical function ϕ from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$, $\phi(Ax) - \phi(x)$ only depends on A and \bar{x} . Therefore $\phi(x(n, \cdot)) - \phi(x(n-1, \cdot))$ only depends on $A(n)$ and $\bar{x}(n-1, \cdot)$. As $\mathbb{P}\mathbb{R}_{\max}^d$ can be seen as an hyperplane of \mathbb{R}^d , $x(n, \cdot)$ can be replaced by $(\phi(x(n, \cdot)), \bar{x}(n, \cdot))$. (cf. lemma 4.3)

According to theorem 1.3, we know that $\bar{x}(n, \cdot)$ converges. On the other hand, by theorem 1.2 $x(n, X^0)$ goes to infinity (if $\gamma \neq 0$) in the direction of $\mathbf{1}$, so $\phi(x(n, \cdot)) \sim \gamma n$. We investigate the oscillations of $\phi(x(n, \cdot)) - \gamma n$. Interesting ϕ 's are defined by $\phi(x) = x_i$, $\phi(x) = \max_i x_i$, $\phi(x) = \min_i x_i$.

The second step is to prove the spectral gap for the operator defining the Markov chain $(A(n), \bar{x}(n-1, \cdot))_{n \in \mathbb{N}}$ and apply the results described in section 3 that give limit theorems for $\phi(x(n, \cdot)) - \gamma n$. The spectral gap follows from the convergence of $\bar{x}(n, \cdot)$, just as by É. Le Page [LP82].

We use two series of results, which are recalled in section 3. The first series are taken from the book by H. Hennion and L. Hervé [HH01] that sums up the classical spectral gap method developed since Nagaev [Nag57] in a general framework. To apply it we demand integrability conditions on $\sup_x |\phi(A(1)x) - \phi(x)|$ to have a Doeblin operator on the space of bounded functions. The second series are taken from the article [HH04] that is a new refinement of the method in the more precise framework of iterated Lipschitz operators. Since our model enters this framework, we get the same results with integrability conditions on $A(1)0$ that have a Doeblin-Fortet operator on weighted spaces. The comparison between the two series of results will be made in section 2.3.

2 Statement of the limit theorems

2.1 General case

From now on, we state the results that we will deduce from the theorems of section 3 in section 4.

For local limit theorem and for renewal theorem we need non arithmeticity conditions. There are two kind of non arithmeticity, depending if the theorem follows from [HH01] or [HH04]. We will denote them respectively by non arithmeticity and algebraic non arithmeticity. When $d = 1$ they both fall down to the usual non-arithmeticity condition for real i.i.d. variables. Algebraic non arithmeticity will be defined before the statement of LLT, but

other non arithmeticity conditions will be defined in section 3 once we have given the general formalism. Unlike algebraic non arithmeticity, they depend on the 2-uple $((A(n))_{n \in \mathbb{N}}, \phi)$, which will be called "the system".

Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property. The sequence $(x(n, \cdot))_{n \in \mathbb{N}}$ is defined by equation (1) and γ is the Lyapunov exponent defined by theorem 1.2.

As the topology of the uniform convergence over compact subset on Top_d has an enumerable basis of open sets, the support of measures on it is well defined. We denote by S_A the support of the law of $A(1)$ and by T_A the semi-group in Top_d generated by S_A .

Theorem 2.1 (CLT). *Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. Let ϕ be topical from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$. Assume one of the following conditions:*

- i) $\sup_x |\phi(A(1)x) - \phi(x)|$ has a moment of order 2,
- ii) $A(1)0$ has a moment of order $4 + \epsilon$ and X^0 has a moment of order $2 + \epsilon$.

Then there exists $\sigma^2 \geq 0$ such that $\frac{x(n, X^0) - n\gamma \mathbf{1}}{\sqrt{n}}$ converges weakly to $\mathcal{N}(0, \sigma^2) \mathbf{1}$.

In the first case, or if $A(1)0$ has a moment of order $6 + \epsilon$ and X^0 has a moment of order $3 + \epsilon$, then

- $\sigma^2 = \lim \frac{1}{n} \int \phi^2(x(n, X^0)) d\mathbb{P}$
- $\sigma = 0$ iff there is a $\theta \in Top_d$ with rank 1 (or every $\theta \in T_A$ with rank 1) such that for any $A \in S_A$ and any $\theta' \in T_A$, $\theta A \theta' = \theta \theta' + \gamma \mathbf{1}$.

Theorem 2.2 (CLT with rate). *Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. Let ϕ be topical from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$. Assume one of the following conditions:*

- i) $\sup_x |\phi(A(1)x) - \phi(x)|$ has a moment of order $l \geq 3$,
- ii) $A(1)0$ has a moment of order $l > 6$.

If $\sigma^2 > 0$ in theorem 2.1, then there exists $C \geq 0$ such that for every initial condition X^0 with a moment of order l , we have

$$\sup_{u \in \mathbb{R}} |\mathbb{P}[\phi(x(n, X^0)) - n\gamma - \phi(X_i^0) \leq \sigma u \sqrt{n}] - \mathcal{N}(0, 1)(-\infty, u)]| \leq \frac{C \left(1 + \mathbb{E} \left[\max \left(\|X^0\|_\infty^l, 1 \right) \right] \right)}{\sqrt{n}}, \quad (2)$$

$$\begin{aligned} & \sup_{u \in \mathbb{R}^d} \left| \mathbb{P}[x(n, X^0) - n\gamma \mathbf{1} \leq \sigma u \sqrt{n}] - \mathcal{N}(0, 1)(-\infty, \min_i u_i) \right| \\ & \leq \frac{C \left(1 + \mathbb{E} \left[\max \left(\|X^0\|_\infty^l, 1 \right) \right] + \mathbb{E} \left[\|A(1)0\|_\infty^l \right] \right)}{n^{\frac{l}{2(l+1)}}}. \end{aligned}$$

Definition 2.1. We say that the sequence $(A(n))_{n \in \mathbb{N}}$ is algebraically arithmetic if there are $a, b \in \mathbb{R}$ and a $\theta \in \text{Top}_d$ with rank 1 (or every $\theta \in T_A$ with rank 1) such that for any $A \in S_A$ and any $\theta' \in T_A$ with rank 1,

$$(\theta A \theta' - \theta \theta')(\mathbb{R}^d) \subset (a + b\mathbb{Z})\mathbf{1}. \quad (3)$$

Otherwise the sequence is algebraically non arithmetic.

Remark 2.1. For any $\theta, \theta' \in \text{Top}_d$ with rank 1 and any $A \in \text{Top}_d$, the function $\theta A \theta' - \theta \theta'$ is constant with value in $\mathbb{R}\mathbf{1}$.

Theorem 2.3 (LLT). *Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. Let ϕ be topical from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$. Assume one of the following conditions:*

- i) $\sup_x |\phi(A(1)x) - \phi(x)|$ has a moment of order 2, $\sigma > 0$, and the system is non arithmetic
- ii) $A(1)0$ has a moment of order $4+\epsilon$, $X^0 \in \mathbb{L}^\infty$ and the sequence $(A(n))_{n \in \mathbb{N}}$ is algebraically non arithmetic.

Then $\sigma > 0$ and there exists a σ -finite measure α on \mathbb{R}^d , so that for any continuous function h with compact support, we have:

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E}_{X^0} [h(x(n, X^0) - n\gamma \mathbf{1} - u \mathbf{1})] - \mathbb{E}_{X^0} \left[e^{-\frac{(u + \phi(X^0))^2}{2n\sigma^2}} \right] \alpha(h) \right| = 0.$$

Moreover the image of α by the function $x \mapsto (\bar{x}, \phi(x))$ is the product of the invariant probability on $\mathbb{P}\mathbb{R}_{\max}^d$ by the Lebesgue measure.

Remark 2.2. As in the usual LLT, this theorem says that the probability for $x(n, X^0)$ to fall in a box decreases as $\frac{1}{\sqrt{n}}$. To replace the continuous functions by indicator functions of the box, we need to know more about ν . In particular, numerical simulation show that some hyperplanes may have a weight for ν , so those hyperplanes could not intersect the boundary of the box.

The algebraic non arithmeticity is optimal in the following sense:

Proposition 2.1. *If the conclusion of theorem 2.3 is true, then $(A(n))_{n \in \mathbb{N}}$ is algebraically non arithmetic.*

Theorem 2.4 (Renewal theorem). *Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. Assume that there is a topical ϕ from \mathbb{R}^d to \mathbb{R} with $\phi(\mathbf{1}) = 1$ such that $\sup_x |\phi(A(1)x) - \phi(x)|$ has a moment of order 2. We denote by α the same measure as in theorem 2.3. If $\gamma > 0$ and the system is weakly non arithmetic, then for any function h continuous with compact support and any initial condition X^0 , we have:*

$$\lim_{a \rightarrow -\infty} \sum_{n \geq 1} \mathbb{E} [h(x(n, X^0) + a\mathbf{1})] = 0,$$

$$\lim_{a \rightarrow +\infty} \sum_{n \geq 1} \mathbb{E} [h(x(n, X^0) + a\mathbf{1})] = \frac{\alpha(h)}{\gamma}.$$

Remark 2.3. The vector $\mathbf{1}$ gives the average direction in which $x(n, X^0)$ is going to infinity. As in the usual renewal theorem, this theorem says that the average number of $x(n, X^0)$ falling in a box is asymptotically proportional to the length of this box, when the box is going to infinity in that direction. As in the LLT, to replace the continuous functions by indicator functions of the box, we need to know more about ν .

Theorem 2.5 (Large deviations). *Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of topical operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. Let ϕ be topical from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$. Assume that $\sup_x |\phi(A(1)x) - \phi(x)|$ has an exponential moment, and that $\sigma^2 > 0$ in theorem 2.1. Then, there exists a non negative strictly convex function c , defined on a neighborhood of 0 and vanishing only at 0 such that for any bounded initial condition X^0 and any $\epsilon > 0$ small enough we have:*

$$\lim_n \frac{1}{n} \mathbb{P} [\phi(x(n, X^0)) - n\gamma > n\epsilon] = -c(\epsilon),$$

$$\lim_n \frac{1}{n} \mathbb{P} [\phi(x(n, X^0)) - n\gamma < -n\epsilon] = -c(-\epsilon).$$

2.2 Max-plus case

When the $A(n)$ are $(\max, +)$ operators, it is natural to chose $\phi(x) = \max_i x_i$. In this case we get $\min_j \max_i A_{ij} \leq \phi(Ax) - \phi(x) \leq \max_{ij} A_{ij}$, so integrability condition can be checked on the last two quantities.

Theorem 2.6 (CLT). Let $(A(n))_{n \in \mathbb{N}}$ be an i.i.d sequence of $(\max, +)$ operators with the MLP property and X^0 an \mathbb{R}^d -valued random variable independent from $(A(n))_{n \in \mathbb{N}}$. If $\max_{ij} A(1)_{ij}$ and $\min_j \max_i A(1)_{ij}$ have a moment of order 2, then there exists $\sigma^2 \geq 0$ such that for every initial condition X^0 ,

(i) $\frac{x(n, X^0) - n\gamma \mathbf{1}}{\sqrt{n}}$ converges weakly to $\mathcal{N}(0, \sigma^2) \mathbf{1}$,

(ii) $\sigma^2 = \lim \frac{1}{n} \int (\max_{i,j} A(n) \cdots A(1)_{ij})^2 d\mathbb{P}$.

Theorems 2.2 to 2.5 are specialized in the same way.

In this case, we also get another condition to avoid degeneracy in the CLT. To state it, we recall a few definitions and results about $(\max, +)$ matrices:

Definition 2.2. For any $k, l, m \in \mathbb{N}$, the product of two matrices $A \in \mathbb{R}_{\max}^{k \times l}$ and $B \in \mathbb{R}_{\max}^{l \times m}$ is the matrix $A \otimes B \in \mathbb{R}_{\max}^{k \times m}$ defined by :

$$\forall 1 \leq i \leq k, \forall 1 \leq j \leq m, (AB)_{ij} := \bigoplus_{p=1}^l A_{ip} \otimes B_{pj}.$$

We notice that if those matrices have no line of $-\infty$, then the $(\max, +)$ operator defined by AB is the composition of those defined by A and the one defined by B .

Definition 2.3. A cycle on a graph is a closed path on the graph. Let A be a square matrix of size k with entries in \mathbb{R}_{\max} .

- i) The graph of A is the directed labelled graph whose vertices are the elements of I_k and whose edges are the (i, j) such that $A_{ij} > -\infty$. The label on (i, j) is A_{ij} . The graph will be denoted by $\mathcal{G}(A)$ and the set of its minimal cycles by $\mathcal{C}(A)$.
- ii) The average weight of a cycle $c = (i_1, \dots, i_n, i_{n+1}) \in \mathcal{C}(A)$ (where $i_1 = i_{n+1}$) is $aw(A, c) := \frac{1}{n} \sum_{j=1}^n A_{i_j i_{j+1}}$.
- iii) The spectral radius of A is $\rho(A) := \max_{c \in \mathcal{C}(A)} aw(A, c)$.
- iv) The critical graph of A is obtained from $\mathcal{G}(A)$ by keeping only vertices and edges belonging to cycles with average weight $\rho(A)$. It will be denoted by $\mathcal{G}^c(A)$.
- v) The cyclicity of a graph is the greatest common divisor of the length of its cycles if it is strongly connected (that is if any vertex can be reached from any other in both directions). Otherwise it is the least common multiple of the cyclicities of its strongly connected components. The cyclicity of A is that of $\mathcal{G}^c(A)$.

vi) The type of A is **scsN-cycC**, where \mathbf{N} is the number of connected components of $\mathcal{G}^c(A)$ and \mathbf{C} the cyclicity of A .

Remark 2.4. Interpretation of powers with $\mathcal{G}(A)$.

If (i_1, i_2, \dots, i_n) is a path on $\mathcal{G}(A)$, its weight is $\sum_{1 \leq j \leq n-1} A_{i_j i_{j+1}}$, so that $(A^{\otimes n})_{ij}$ is the maximum of the weights of length n paths from i to j .

Theorem 2.7 ([CDQV83]). *Assume $\mathcal{G}(A)$ is strongly connected, $\rho(A) = 0$ and the cyclicity of A is 1. Then for n big enough $A^{\otimes n} = Q$, where Q is defined by*

$$\forall(i, j) \in I_k^2, Q_{ij} := \bigoplus_{l \in \mathcal{G}^c(A)} A_{il}^+ \otimes A_{lj}^+.$$

Theorem 2.8. *Assume the hypothesis of theorem 2.6, with $\gamma = 0$. Then the variance σ^2 in theorem 2.6 is 0 if and only if $\rho_{\max}(T_A) = \{0\}$.*

Theorem 3.2 of [Mer04] gives a condition to ensure the memory loss property. This condition also ensures that there are two matrices in S_A with two distinct spectral radius. This proves the following corollary:

Corollary 2.1. *Let the law of $A(1)$ be a probability on the set of $d \times d$ matrices with moment of order 2 whose support is not included in a finite union of affine hyperplanes of $\mathbb{R}^{k \times k}$. Then $x(n, \cdot)$ satisfies the conclusions of theorem 2.6 with $\sigma > 0$.*

We also give a sufficient condition to ensure the algebraic non arithmeticity:

Theorem 2.9. *Assume the hypothesis of theorem 2.3 ii) except the algebraic non arithmeticity and $A(n)$ are $(\max, +)$ operators. If the system is algebraically arithmetic, then there are $a, b \in \mathbb{R}$ such that*

$$\{\rho_{\max}(A) | A \in S_A, \mathcal{G}(A) \text{ strongly connected}\} \subset a + b\mathbb{Z}.$$

Together with corollary 2.1, this proves that the hypothesis are generic in the following sense:

Corollary 2.2. *If the law of $A(1)$ is a probability on the set of $d \times d$ matrices with moment of order $4 + \epsilon$ whose support is not included in an enumerable union of affine hyperplanes of $\mathbb{R}^{k \times k}$. Then $x(n, \cdot)$ satisfies the conclusions of theorem 2.3.*

2.3 Comments

The following table sums up the limit theorems. In each situation we assume the memory-loss property.

Theorems:	$A(1) 0$	Finite moments of $\max_{ij} A(1)_{ij}$ and $\min_j \max_i A(1)_{ij}$	Additional condition
CLT	$4 + \epsilon$	2	
CLT with rate	$6 + \epsilon$	3	
LLT	$4 + \epsilon$	2	NA
Renewal	–	2	NA
LDP	–	exp	$X^0 \in L^\infty$

NA= non arithmeticity

The results of the second column are stated for $(\max, +)$ operators because the $\sup_x |\phi(A(1)x) - \phi(x)|$ is not bounded for general topical operators. For other subclasses of topical operators, one has to choose ϕ such that $\sup_x |\phi(A(1)x) - \phi(x)|$ is integrable. For instance $\phi(x) = \min_i x_i$ is natural for $(\min, +)$ operators. Actually, it should be possible to derive renewal theorem and large deviation principle with the method of [HH04] but this has not been written down.

The results of the first column require stronger integrability conditions but they are also better for two reasons: they are true for any topical operators and the algebraic non arithmeticity does not depend on ϕ . It is expressed without introducing the Markovian operator Q although the system is algebraically non arithmetic iff Q has an eigenvector with eigenvalue of modulus 1. Moreover for $(\max, +)$ operators the algebraic non arithmeticity can be deduced from theorem 2.9.

An important case for $(\max, +)$ operators is when $A_{ij} \in \mathbb{R}^+ \cup \{-\infty\}$ and $A_{ii} \geq 0$ because it modelizes situations where $x_i(n, \cdot)$ is the date of the n th event of type i and the A_{ij} are delays. In this case the integrability of $\max_{ij} A(1)_{ij}$ and $\min_j \max_i A(1)_{ij}$ is equivalent to the integrability of $A(1) 0$.

We mentioned earlier that J. Resing et al. [RdVH⁺90] obtained a central limit theorem. In a sense our result is weaker because the MLP property implies that $\bar{x}(n, \cdot)$ is uniformly Φ -recurrent and aperiodic. But our integrability conditions are much weaker and the MLP property is easier to check.

F. Toomey's large deviations principle only requires the uniform bound of the projective image that is a very strong integrability condition. It suggests that the MLP property should not be necessary. But his formulation of the

LDP is not equivalent to ours and in the $(\max, +)$ case it needs the fixed structure property, that is $\mathbb{P}(A_{ij}(1) = -\infty) \in \{0, 1\}$.

3 Markov chains and quasi-compact operators

3.1 Spectral gap

We recall the formalism of Markov chains and operators, following the presentation by H. Hennion and L.Hervé [HH01]. Then we state the results obtained by use of the spectral gap method, that we want to apply to our model.

The context is defined by:

- a measurable space (E, \mathcal{E}) ,
- a transition probability Q on (E, \mathcal{E}) ,
- a Markov chain X_n on (E, \mathcal{E}) associated with Q ,
- a measurable real valued function ξ on (E, \mathcal{E}) .

The principle is to investigate the characteristic function ϕ_{S_n} of

$$S_n := \sum_{l=1}^n \xi(X_l)$$

by means of the action of the Fourier kernels denoted by Q_t or $Q(t)$ on a suitable Banach space of measurable functions on E . These kernels are defined for any $t \in \mathbb{C}$ by

$$Q_t(x, dy) = e^{it\xi(y)} Q(x, dy)$$

and we immediately have $\phi_{S_n}(t) = \int_E Q_t^n 1 d\lambda(x)$, where λ is the law of X_0 .

As Q is a transition probability, $Q = Q_0$ admits 1 as an eigenvalue. To control the perturbations Q_t of Q and especially $Q_{t/\sqrt{n}}^n$, the key tool is the quasi-compactness. We say that Q has a spectral gap if it satisfy the following definition:

Definition 3.1. A continuous endomorphism Q on a Banach space \mathcal{B} is said to be **quasi-compact**, if there exists a Q -stable decomposition $\mathcal{B} = F \oplus H$ such that:

1. $\dim F < \infty$ and all eigenvalues of $Q|_F$ have modulus $r(Q)$,
2. $r(Q|_H) < r(Q)$.

In this case $\dim F$ will be denoted by $s(Q, \mathcal{B})$.

3.2 First frame

In this subsection, we state the results of [HH01], under a few additional assumptions that make the statements easier.

Definition 3.2. Let (Q, ξ) be as before, and $(\mathcal{B}, \|\cdot\|)$ be a Banach space of continuous functions on E . We say that (Q, \mathcal{B}, ξ) satisfies condition $\mathcal{H}(m)$ if the following properties hold:

- (H1) (i) $1 \in \mathcal{B}$ and if $f \in \mathcal{B}$, then $\bar{f} \in \mathcal{B}$, $f^+ = \max(f, 0)$,
(ii) The norm $\|\cdot\|$ on \mathcal{B} is greater than the infinity norm,
(iii) if $f, g \in \mathcal{B}$ are bounded, then $fg \in \mathcal{B}$.
- (H2) (i) Q has an invariant probability ν ,
(ii) Q is quasi-compact on \mathcal{B} , with $\sup \|Q^n\| < \infty$,
(iii) $\text{Ker}(1 - Q) = \text{span}(1)$.
- (H3) there exists a neighborhood I_0 of 0 in \mathbb{R} such that $Q(\cdot) \in \mathcal{C}^m(I_0, \mathcal{L}_{\mathcal{B}})$ and for $1 \leq k \leq m$, the operator $Q^{(k)}(0)$ is defined by the kernel $Q(x, dy)(i\xi(y))^k$.

Definition 3.3. We say that the system is non arithmetic if $Q(\cdot)$ is continuous from \mathbb{R} to the space $\mathcal{L}_{\mathcal{B}}$ of continuous linear operators on \mathcal{B} and, for all $t \in \mathbb{R}^*$, $r(Q(t)) < 1$.

We say that the system is weakly non arithmetic if $Q(\cdot)$ is continuous from \mathbb{R} to $\mathcal{L}_{\mathcal{B}}$ and, for all $t \in \mathbb{R}^*$, $1 - Q(t)$ is invertible.

We state the limit theorems of [HH01] when $s(Q, \mathcal{B}) = 1$.

Theorem 3.1 ([HH01]). Assume that (Q, \mathcal{B}, ξ) satisfies $\mathcal{H}(2)$ and $s(Q, \mathcal{B}) = 1$. Then ξ^2 is ν -integrable and if $\nu(\xi) = 0$ there exists $\sigma^2 \geq 0$ such that for any initial condition with law λ ,

- (i) $\lim_n \frac{1}{n} \mathbb{E}_{\lambda}[S_n^2] = \sigma^2$,
- (ii) if $\sigma^2 = 0$, then there exists $\xi_1 \in \mathcal{B}$, such that ξ_1^2 is ν -integrable and $\xi(X_1) = \xi_1(X_0) - \xi_1(X_1) \mathbb{P}_{\nu}$ - a.s.,

(iii) for $f \in \mathcal{B}$ and $g \in \mathcal{C}_c(\mathbb{R})$,

$$\lim_n \mathbb{E}_\lambda \left[f(X_n) g\left(\frac{S_n}{\sqrt{n}}\right) \right] = \nu(f) \mathcal{N}(0, \sigma^2)(g),$$

where the convergence is uniform in the initial condition.

Theorem 3.2 ([HH01]). Assume that (Q, \mathcal{B}, ξ) satisfies $\mathcal{H}(3)$. If $\nu(\xi) = 0$ and $\sigma > 0$, then there exists $C > 0$ such that for any initial condition λ ,

$$\sup_{u \in \mathbb{R}} \left| \mathbb{P}_\lambda[S_n \leq u\sigma\sqrt{n}] - \mathcal{N}(0, 1)(-\infty, u) \right| \leq \frac{C(\|\lambda\| + 1)}{\sqrt{n}}.$$

Theorem 3.3 ([HH01]). Assume that (Q, \mathcal{B}, ξ) satisfies $\mathcal{H}(2)$ and $s(Q, \mathcal{B}) = 1$, that $\nu(\xi) = 0$ and $\sigma > 0$. If moreover the system is non arithmetic, then for $f \in \mathcal{B}$ and $g \in \mathcal{C}_c(\mathbb{R})$, we have:

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma\sqrt{2\pi n} \mathbb{E}_\lambda [f(X_n)g(S_n - u)] - e^{-\frac{u^2}{2n\sigma^2}} \nu(f) \mathcal{L}(g) \right| = 0,$$

where \mathcal{L} is the Lebesgue measure and the convergence is uniform in the initial condition λ .

Theorem 3.4 ([HH01]). Assume that (Q, \mathcal{B}, ξ) satisfies $\mathcal{H}(2)$, and that $\nu(\xi) > 0$. If moreover the system is weakly non arithmetic, then for $f \in \mathcal{B}$ and $g \in \mathcal{C}_c(\mathbb{R})$, we have:

$$\begin{aligned} \lim_{a \rightarrow -\infty} \sum_{n \geq 1} \mathbb{E} [f(X_n)g(S_n - a)] &= 0, \\ \lim_{a \rightarrow +\infty} \sum_{n \geq 1} \mathbb{E} [f(X_n)g(S_n - a)] &= \frac{\nu(f)}{\nu(\xi)} \mathcal{L}(g), \end{aligned}$$

where the convergence is uniform in the initial condition λ .

Theorem 3.5 ([HH01]). Assume that (Q, \mathcal{B}, ξ) satisfies $\mathcal{H}(2)$, that $\nu(\xi) = 0$ and $\sigma > 0$. If moreover there exists $\theta > 0$ for which $Q(\cdot)$ is continuous from $\{z \in \mathbb{C} : |\Re(z)| < \theta\}$ to $\mathcal{L}_\mathcal{B}$ and holomorphic on $\{z \in \mathbb{C} : |z| < \theta\}$, then there exists a non-negative strictly convex function c , defined on a neighborhood of 0 and vanishing only at 0, such that, for $\lambda \in \mathcal{B}'_p$ and sufficiently small $\epsilon > 0$, we have:

$$\lim_n \frac{1}{n} \mathbb{P}_\lambda[S_n > n\epsilon] = -c(\epsilon).$$

3.3 Second frame

In their article [HH04] H. Hennion and L. Hervé have proved limit theorems for sequences $\xi(Y_n, Z_{n-1})$, where $(Y_n)_{n \in \mathbb{N}}$ is an i.i.d. sequence of Lipschitz operators, and Z_n is defined by $Z_{n+1} = Y_{n+1}Z_n$. As explained in section 1.3, this is our situation. Moreover in our situation, the Y_n , which are the projective function defined by $A(n)$, are 1-Lipschitz. Following the same proof as [HH04] with this additional condition, we get the theorems of this section. The integrability conditions are weaker, since the Lipschitz coefficient is uniformly bounded. The only difference in the proof is the Hölder inequality of the 4th part of proposition 7.3 of [HH04]: the exponents in the inequality should be changed to 1 and ∞ .

In the sequel, \mathcal{M} is a non-compact metric space and G is the semi-group of its 1-Lipschitz operators for distance δ . We denote by \mathcal{S} the following data:

- (i) For an i.i.d. sequence $(Y_n)_{n \in \mathbb{N}}$ with values in G and law μ and a real variable Z , with values in \mathcal{M} and independent from $(Y_n)_{n \in \mathbb{N}}$, Z_n is defined by

$$Z_0 = Z \text{ and } Z_{n+1} = Y_{n+1}Z_n.$$

- (ii) For $\xi : G \times \mathcal{M} \rightarrow \mathbb{R}$ 1-Lipschitz in the second variable, S_n is defined by

$$S_n = \sum_{k=1}^n \xi(Y_k, Z_{k-1}).$$

- (iii) For a fixed $x_0 \in \mathcal{M}$, every $\eta \geq 1$ and every $n \in \mathbb{N}$, we set $\mathcal{M}_\eta = \mathbb{E}[\delta^\eta(Y_1 x_0, x_0)]$ and $\mathcal{C}_n = \mathbb{E}[c(Y_n \cdots Y_1)]$, where $c(\cdot)$ is the Lipschitz coefficient.

When there is an $N \in \mathbb{N}$ such that $\mathcal{C}_n < 1$, there is a $\lambda_0 \in]0, 1[$, such that $\int_G c(g) (1 + \lambda_0 \delta(gx_0, x_0))^{2\eta} d\mu^{*n}(g) < 1$. We chose one and set the following notations:

- (i) \mathcal{B}_η is the set of functions f from \mathcal{M} to \mathbb{C} such that $m_\eta(f) < \infty$, with norm $\|f\|_\eta = |f|_\eta + m_\eta(f)$, where

$$\begin{aligned} |f|_\eta &= \sup_x \frac{|f(x)|}{(1 + \lambda_0 \delta(x, x_0))^{1+\eta}}, \\ m_\eta(f) &= \sup_{x \neq y} \frac{|f(x) - f(y)|}{\delta(x, y) (1 + \lambda_0 \delta(x, x_0))^\eta (1 + \lambda_0 \delta(y, x_0))^\eta}. \end{aligned}$$

- (ii) \mathcal{B}'_η is the set of probability measures on E defining a continuous functional on \mathcal{B}_η and for any $r > 0$, we set $\mathcal{B}'_{\eta,r} = \{\lambda \in \mathcal{B}'_\eta : \|\lambda\| < r\}$, where $\|\lambda\|$ is the norm of the linear form on \mathcal{B}_η defined by λ .
- (iii) We say that the system is η -non arithmetic if there is no $t \in \mathbb{R} \setminus \{0\}$, no $\rho \in \mathbb{C}$, and no $w \in \mathcal{B}_\eta$ with non-zero constant modulus on the support S_ν of ν such that $|\rho| = 1$ and for all $n \in \mathbb{N}$, we have

$$e^{itS_n w}(Z_n) = \rho^n w(Z_0) \mathbb{P}_{\nu_0} - \text{ a.s. .} \quad (4)$$

Remark 3.1 (non arithmeticity). In the first frame the non arithmeticity condition is about the spectral radius of Q_t . Here we work with the associated P_t that acts on \mathcal{M} instead of (G, \mathcal{M}) (cf. [HH04]). If P_t is quasi-compact, then $r(P_t) = 1$ iff P_t has an eigenvalue ρ with modulus 1. It is shown in proposition 9.1' of [HH04] that if $r(P_t) = 1$, then P_t is quasi-compact as an operator on \mathcal{B}_η and that an eigenvector w with eigenvalue ρ satisfies equation(4).

We are now ready to state the limit theorems by H. Hennion and L. Hervé.

Theorem 3.6 ([HH04]). *Assuming \mathcal{S} , if there exists $\eta \geq 0$ such that $\mathcal{M}_{\eta+1} < \infty$ and n_0 such that $\mathcal{C}_{n_0} < 1$, then S_n converges in law to an invariant distribution ν , such that $\int_{\mathcal{M}} \delta^{\eta+1}(x, x_0) d\nu(x) < \infty$.*

Theorem 3.7 ([HH04]). *Assuming \mathcal{S} , if there exists $\eta > 1$ and $n_0 \in \mathbb{N}$ such that $\mathcal{C}_{n_0} < 1$ and $\mathcal{M}_{2\eta} < \infty$, then, for $f \in \cup_{\eta' < \eta-1} \mathcal{B}_{\eta'}$ and $g \in \mathcal{C}_c(\mathbb{R})$, we have:*

(i) $\lim_n \mathbb{E}_\lambda \left[f(Z_n) g\left(\frac{S_n}{\sqrt{n}}\right) \right] = \nu(f) \mathcal{N}(0, \sigma^2)(g)$, where the convergence is uniform in $\lambda \in \mathcal{B}'_{\eta,r}$.

(ii) If $\eta > 3$, then for $\lambda \in \mathcal{B}'_\eta$, $\lim_n \frac{1}{n} \mathbb{E}_\lambda[S_n^2] = \sigma^2$.

(iii) If $\eta > 3$ and $\sigma^2 = 0$, then there exists $\xi_1 \in \mathcal{B}_{\eta-1}$, such that ξ_1^2 is ν -integrable and $\xi(X_1) = \xi_1(X_0) - \xi_1(X_1) \mathbb{P}_\nu - \text{ a.s. .}$

Theorem 3.8 ([HH04]). *Assuming \mathcal{S} , if there exists $\eta > 3$ and $n_0 \in \mathbb{N}$ such that $\mathcal{C}_{n_0} < 1$ and $\mathcal{M}_{2\eta} < \infty$, if moreover $\nu(\xi) = 0$ and $\sigma > 0$, then there exists $C > 0$ such that for $\lambda \in \mathcal{B}'_\eta$,*

$$\sup_{u \in \mathbb{R}} \left| \mathbb{P}_\lambda[S_n \leq u\sigma\sqrt{n}] - \mathcal{N}(0, 1)(-\infty, u) \right| \leq \frac{C\|\lambda\|_\eta}{\sqrt{n}}.$$

Theorem 3.9 ([HH04]). *Assuming \mathcal{S} , if there exists $\eta > 2$ and $n_0 \in \mathbb{N}$ such that $\mathcal{C}_{n_0} < 1$ and $\mathcal{M}_{2\eta} < \infty$, if moreover $\nu(\xi) = 0$ and there is $1 < \eta' < \eta - 1$ such that the system is η' -non arithmetic, then for $f \in \mathcal{B}_{\eta'}$ and $g \in \mathcal{C}_c(\mathbb{R})$,*

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E}_\lambda [f(Z_n)g(S_n - u)] - e^{-\frac{u^2}{2n\sigma^2}} \nu(f)\mathcal{L}(g) \right| = 0,$$

where \mathcal{L} is the Lebesgue measure and the convergence is uniform in $\lambda \in \mathcal{B}'_{\eta,r}$.

4 Proofs of the limit theorems

4.1 From iterated topical functions to Markov chains

In this section, we show that the hypothesis of the theorems on our model stated in section 2 imply the hypothesis of the general theorems stated in section 3. To apply the results of [HH01],

- the space E will be $Top_d \times \mathbb{P}\mathbb{R}_{\max}^d$ with the Borel σ -algebra,
- the transition probability Q will be defined by

$$Q((A, \bar{x}), D) = P((A(1), \overline{Ax}) \in D),$$

- the Markov chain X_n will be $(A(n), \bar{x}(n-1, \cdot))$,
- the function ξ will be defined by $\xi(A, \bar{x}) = \phi(Ax) - \phi(x)$, where ϕ is a topical function from \mathbb{R}^d to \mathbb{R} such that $\phi(\mathbf{1}) = 1$.
- $\sigma(A) = \sup_x |\xi(A, x)| = \sup_x |\phi(Ax) - \phi(x)| < \infty$ a.s. .

We first need to define the space \mathcal{B} .

Definition 4.1. Let \mathcal{L}^∞ be the space of complex valued bounded continuous functions on $\mathbb{P}\mathbb{R}_{\max}^d$ with the infinity norm.

Let j be the function from $Top_d \times \mathbb{P}\mathbb{R}_{\max}^d$ to $\mathbb{P}\mathbb{R}_{\max}^d$ such that $j(A, \bar{x}) = \overline{Ax}$ and I the function from $\mathbb{R}^{\mathbb{P}\mathbb{R}_{\max}^d}$ to $\mathbb{R}^{(Top_d \times \mathbb{P}\mathbb{R}_{\max}^d)}$ defined by

$$I(\phi) = \phi \circ j.$$

We call \mathcal{B}^∞ the image of \mathcal{L}^∞ by I .

As $(\mathcal{L}^\infty, \|\cdot\|_\infty)$ is a Banach space, I is an injection, and $\|\phi \circ I\|_\infty = \|\phi\|_\infty$, $(\mathcal{B}^\infty, \|\cdot\|_\infty)$ is also a Banach space.

Proposition 4.1. *If $\sigma(A(1))$ has a moment of order m and if $A(n)$ has the MLP property, then $(Q, \xi, \mathcal{B}^\infty)$ satisfies condition $\mathcal{H}(m)$. Moreover, the interval I_0 in condition (H3) is the whole \mathbb{R} and $s(Q, \mathcal{B}^\infty) = 1$.*

To prove (H2), we will use theorem 6.15 of [Mai97] that we restate in our case:

Theorem 4.1 (Mairesse [Mai97]). *If the sequence $(A(n))_{n \in \mathbb{N}}$ of i.i.d. random variables with values in Top_d has memory loss property, then there exists a random variable Y with values in $\mathbb{P}\mathbb{R}_{\max}^d$ such that $Y_n := A(n) \cdots A(1)Y$ is stationary and*

$$\lim_{n \rightarrow \infty} \mathbb{P}(\exists x_0, Y_n \neq \bar{x}(n, x_0)) = 0.$$

In the sequel, ν_0 will be the law of Y that is the limit law of theorem 1.3.

Proof of proposition 4.1. Condition (H1) is trivial, because of the choice of \mathcal{B}^∞ .

To check condition (H2), we take $\nu := \mu \otimes \nu_0$. It is Q -invariant by definition of ν_0 . This proves (i). To prove (ii) and (iii), we investigate the iterates of Q . For any $\phi \in \mathcal{L}^\infty$, and $x \in \mathbb{R}^d$ we have:

$$\begin{aligned} |Q^n(\phi \circ j)(A, \bar{x}) - \nu(\phi \circ j)| &= |Q^n(\phi)(\overline{Ax}) - \nu_0(\phi)| \\ &= \left| \int \phi(\bar{x}(n, \overline{Ax})) d\mathbb{P} - \int \phi(Y_n) d\mathbb{P} \right| \\ &\leq \|\phi\|_\infty 2\mathbb{P}(\exists x_0, Y_n \neq \bar{x}(n, x_0)), \end{aligned}$$

If we denote $\psi \mapsto \nu(\psi)$ by N , we obtain:

$$\|Q^n - N\| \leq 2\mathbb{P}(\exists x, Y_n \neq \bar{x}(n, x)) \rightarrow 0. \quad (5)$$

This proves that $r(Q|_{\text{Ker}N}) < 1$ and that $\sup_n \|Q^n\| < \infty$, so (ii) is checked. It also proves $\text{Ker}(1 - Q) \subset \text{Im}N$, which implies (iii).

To prove (H3) we set $Q_t^{(k)} := e^{it\xi(y)}(i\xi(y))^k Q(x, dy)$ and

$$\Delta_h^{(k)} := Q_{t+h}^{(k)} - Q_t^{(k)} - hQ_t^{(k+1)}. \quad (6)$$

To prove our proposition, it remains to bound $\|\frac{1}{h}\Delta_h^{(k)}\|$ by a quantity that tends to zero with h .

To this aim, we introduce the following function:

$$\begin{cases} f : \mathbb{R} & \rightarrow \mathbb{C} \\ t & \mapsto e^{it} - 1 - it. \end{cases}$$

The calculus will be based on the following estimations on f : $|f(t)| \leq 2t$, and $|f(t)| \leq t^2$.

Now all follows from

$$\Delta_h^{(k)}(\phi \circ j)(A, \bar{x}) = \int \phi(\overline{BAx}) e^{it\xi(B, \overline{Ax})} (i\xi(B, \overline{Ax}))^k f(h\xi(B, \overline{Ax})) d\mu(B). \quad (7)$$

First it implies that

$$\left\| \Delta_h^{(k)}(\phi \circ j) \right\|_\infty \leq \|\phi\|_\infty \int \sigma^k(B) \|f(h\xi(B, \cdot))\|_\infty d\mu(B). \quad (8)$$

Since $|f(t)| \leq t^2$,

$$\frac{1}{|h|} \sigma(B)^k \|f(h\xi(B, \cdot))\|_\infty \leq h\sigma^{k+2}(B) \rightarrow 0.$$

Since $|f(t)| \leq 2t$,

$$\frac{1}{|h|} \sigma^k(B) \|f(h\xi(B, \cdot))\|_\infty \leq 2\sigma^{k+1}(B).$$

When $k < m$, σ^{k+1} is integrable so the dominate convergence theorem and the two last equations show that

$$\int \sigma(B)^k \|f(h\xi(B, \cdot))\|_\infty d\mu(B) = o(h). \quad (9)$$

Finally for any $k < m$, $\|\frac{1}{h}\Delta_h^{(k)}\|$ tends to zero, so $Q_t^{(k+1)}$ is the derivative of $Q_t^{(k)}$ in t .

To prove that $Q_t^{(m)}$ is continuous, we notice that

$$\left(Q_{t+h}^{(m)} - Q_t^{(m)} \right) (\phi \circ j)(A, \bar{x}) = \int \phi(\overline{BAx}) e^{it\xi(B, \overline{Ax})} (i\xi(B, \overline{Ax}))^m g(h\xi(B, \overline{Ax})) d\mu(B). \quad (10)$$

where $g(t) = e^{it} - 1$. Then we apply the same method as before, replacing the estimates on $f(h\xi(B, \cdot))$ by $\|g(h\xi(B, \cdot))\|_\infty \leq h\sigma(B)$ to prove the convergence, and by $\|g(h\xi(B, \cdot))\|_\infty \leq 2$ to prove the domination.

This proves (H3) and the additional assumption of proposition 4.1. \square

To prove theorems 2.1 and 2.3, we will also apply theorems of section 3.3. We take $\mathcal{M} = \mathbb{P}\mathbb{R}_{\max}^d$, $Y_n = A(n)$ and again $\xi(A, \bar{x}) = \phi(Ax) - \phi(x)$. In this case $Z_n = \bar{x}(n, X^0)$ and $S_n = \phi(x(n, x^0)) - \phi(X^0)$ and we get the following proposition:

Proposition 4.2.

1. If $A(1)0$ has a moment of order $\eta \in \mathbb{R}^+$, then $\mathcal{M}_\eta < \infty$. If the sequence has the MLP property, then there is $n_0 \in \mathbb{N}$ such that $\mathcal{C}_{n_0} < 1$. If $\overline{X^0}$ has a moment of order $\eta \in \mathbb{R}^+$, then $f \mapsto \mathbb{E}[f(\overline{X^0})] \in \mathcal{B}'_\eta$.
2. Algebraic non arithmeticity implies η -non arithmeticity for any $\eta > 0$.

The first part of the proposition is obvious. The second part relies on the next two lemma that will be proved after the proposition:

Lemma 4.1. *The support of the invariant measure ν_0 is*

$$S_{\nu_0} := \overline{\{\theta \mathbf{1} \mid \theta \in T_A, \theta \text{ with rank } 1\}}.$$

Lemma 4.2. *If equation (3) is fulfilled for one θ with rank 1 and any $A \in S_A$ and $\theta' \in T_A$ with rank 1, it is fulfilled for any $\theta \in T_A$ with rank 1.*

Proof of proposition 4.2. Let us assume that the system is η -arithmetic. Then there are $w \in \mathcal{B}_\eta$ and $t, a \in \mathbb{R}$ such that for μ -almost every A and ν_0 almost every \overline{x} , we have:

$$e^{it(\phi(Ax) - \phi(x))} w(\overline{Ax}) = e^{ita} w(\overline{x}). \quad (11)$$

As all function in this equation are continuous, it is true for $\overline{x} \in S_{\nu_0}$ and $A \in T_A$. As S_{ν_0} is T_A invariant, we iterate equation (11) and get

$$e^{it(\phi(Tx) - \phi(x))} w(\overline{Tx}) = e^{itan_T} w(\overline{x}), \quad (12)$$

where $T \in T_A$ and n_T is the number of operators of S_A one has to compose to obtain T .

Because of the MLP property, there is a $\theta \in T_A$ with rank 1. We apply equation (12) for $T = \theta A$ and $T = \theta$ and divide the first equation by the second one. As $n_{\theta A} = n_\theta + 1$ and $\overline{\theta Ax} = \overline{\theta x}$, we get

$$e^{it(\phi(\theta Ax) - \phi(\theta x))} = e^{ita}.$$

Setting $b = \frac{2\pi}{t}$, it means that $\phi(\theta Ax) - \phi(\theta x) \in a + b\mathbb{Z}$. As θ has rank one, $(\theta Ax - \theta x) \in \mathbb{R}\mathbf{1}$, so $\theta Ax - \theta x \in (a + b\mathbb{Z})\mathbf{1}$, and the algebraic arithmeticity follows by lemma 4.1. \square

Proof of lemma 4.1. By theorem 4.1, there is sequence of random variables Y_n with law ν_0 , such that $Y_n = A(n) \cdots A(1)Y$. Let K be a compact subset of \mathbb{R}_{\max} such that $Y \in K$ with positive probability.

For any $\theta \in T_A$ and any $\epsilon > 0$, the set V of topical functions A such that $\delta(\overline{Ax}, \overline{\theta x}) \leq \epsilon$ for all $\overline{x} \in K$ is a neighborhood of θ . Therefore the probability for $A(n_\theta) \cdots A(1)$ to be in V is positive and by independence of Y , we have:

$$\mathbb{P}[Y \in K, A(n_\theta) \cdots A(1) \in V] > 0.$$

As $\overline{\theta \mathbf{1}} = \overline{\theta x}$, this means that with positive probability,

$$\delta(Y_{n_\theta}, \overline{\theta \mathbf{1}}) = \delta(A(n_\theta) \cdots A(1)Y, \overline{\theta Y}) \leq \epsilon,$$

so $\overline{\theta \mathbf{1}} \in S_{\nu_0}$.

This proves that $\overline{\{\overline{\theta \mathbf{1}} | \theta \in T_A, \theta \text{ with rank } 1\}} \subset S_{\nu_0}$.

In [Mai97], ν_0 is obtained as the law of $Z = \overline{\lim_n A(1) \cdots A(n) \mathbf{1}}$. Indeed, the MLP property and the Poincaré recurrence theorem ensure that there are almost surely M and N such that $A(N) \cdots A(N+M)$ has rank 1. Therefore $\overline{A(1) \cdots A(n) \mathbf{1}} = \overline{A(1) \cdots A(N+M) \mathbf{1}}$ for $n \geq N+M$ and $Z = \overline{A(1) \cdots A(N+M) \mathbf{1}}$. But $A(1) \cdots A(N+M) \in T_A$ almost surely, so $Z \in \overline{\{\overline{\theta \mathbf{1}} | \theta \in T_A, \theta \text{ with rank } 1\}}$ almost surely and $S_{\nu_0} \subset \overline{\{\overline{\theta \mathbf{1}} | \theta \in T_A, \theta \text{ with rank } 1\}}$. \square

Proof of lemma 4.2. We assume that equation (3) is fulfilled for $\theta = \theta_1$ and any $A \in S_A$ and $\theta' \in T_A$ with rank 1.

Let $A_1, \dots, A_n \in S_A$, such that $\theta_2 = A_1 \cdots A_n$ has rank 1. For any $i \leq n$, $A_i \cdots A_n \theta'$ has rank 1, so $(\theta_1 A_i \cdots A_n \theta' - \theta_1 A_{i+1} \cdots A_n \theta')(\mathbb{R}^d) \subset (a + b\mathbb{Z})\mathbf{1}$.

Summing these inclusions for $i = 1$ to $i = n$, we get $(\theta_1 \theta_2 \theta' - \theta_1 \theta')(\mathbb{R}^d) \subset (na + b\mathbb{Z})\mathbf{1}$ and

$$((\theta_1 \theta_2 A \theta' - \theta_1 A \theta') - (\theta_1 \theta_2 \theta' - \theta_1 \theta'))(\mathbb{R}^d) \subset b\mathbb{Z}\mathbf{1}. \quad (13)$$

Now we write $\theta_2 \theta'$ as

$$\theta_2 \theta' = \theta_1 \theta' + (\theta_1 \theta_2 \theta' - \theta_1 \theta') - (\theta_1 \theta_2 \theta' - \theta_2 \theta').$$

The last part does not depend on θ' , so replacing θ' by $A\theta'$ and subtracting the first version, we get:

$$\theta_2 A \theta' - \theta_2 \theta' = \theta_1 \theta' - \theta_1 A \theta' + ((\theta_1 \theta_2 A \theta' - \theta_1 A \theta') - (\theta_1 \theta_2 \theta' - \theta_1 \theta')).$$

With equation (13), this proves equation (3) for $\theta = \theta_2$. \square

4.2 From Markov chains to iterated topical functions

Propositions 4.1 and 4.2 prove that under the hypothesis of section 2 the conclusions of the theorems of section 3 are true. This gives results about the convergence of $(\phi(x(n, X^0)) - \phi(X^0) - n\nu(\xi), \bar{x}(n, X^0))$.

When X^0 has law ν_0 , the sequence $(A(n), \bar{x}(n, X^0))_{n \in \mathbb{N}}$ is stationary, so it follows from Birkhoff theorem that $\gamma = \int \xi(A, \bar{x}) d\nu_0(\bar{x}) d\mu(A) = \nu(\xi)$.

The following lemma will be useful to go back to $x(n, \cdot)$.

Lemma 4.3. *If ϕ is a topical function from \mathbb{R}^d to \mathbb{R} , the function $\psi : x \mapsto (\phi(x), \bar{x})$ is a Lipschitz homeomorphism with Lipschitz inverse from \mathbb{R}^d onto $\mathbb{R} \times \mathbb{P}\mathbb{R}_{\max}^d$.*

Proof. Let (t, \bar{x}) be an element of $\mathbb{R} \times \mathbb{P}\mathbb{R}_{\max}^d$. Then $\psi(y) = (t, \bar{x})$ if and only if there is an $a \in \mathbb{R}$ such that $y = x + a\mathbf{1}$ and $\phi(x) + a = t$. So the equation has exactly one solution $y = x + (t - \phi(x))\mathbf{1}$ and ψ is invertible.

It is well known that topical function are Lipschitz, and the projection is linear, so it is Lipschitz and so is ψ .

For any $x, y \in \mathbb{R}^d$, we have $x \leq y + \max_i(x_i - y_i)\mathbf{1}$, so $\phi(x) - \phi(y) \leq \max_i(x_i - y_i)$. Therefore, for any $1 \leq i \leq d$, we have $\phi(x) - \phi(y) - (x_i - y_i) \leq \max_i(x_i - y_i) - \min_i(x_i - y_i) = \delta(\bar{x}, \bar{y})$. Permuting x and y , we see that:

$$|\phi(x) - \phi(y) - (x_i - y_i)| \leq \delta(\bar{x}, \bar{y}). \quad (14)$$

Therefore $|x_i - y_i| \leq |\phi(x) - \phi(y)| + \delta(\bar{x}, \bar{y})$ and ψ^{-1} is Lipschitz. \square

Proof of theorem 2.1. Without loss of generality, we assume that $\gamma = 0$. Theorem 3.1 and proposition 4.1 or theorem 3.7 and proposition 4.2 prove that $\frac{\phi(x(n, X^0)) - \phi(X^0)}{\sqrt{n}}$ converges to $\mathcal{N}(0, \sigma^2)$, which means that $\frac{\phi(x(n, X^0)) - \phi(X^0)}{\sqrt{n}} \mathbf{1}$ converges to $\mathcal{N}(0, \sigma^2) \mathbf{1}$. We just estimate the difference between the converging sequence and the one we want to converge:

$$\Delta_n := \left\| \frac{x(n, X^0)}{\sqrt{n}} - \frac{\phi(x(n, X^0)) - \phi(X^0)}{\sqrt{n}} \mathbf{1} \right\|_{\infty} \leq \frac{|\phi(X^0)|}{\sqrt{n}} + \frac{|\bar{x}(n, X^0)|_{\mathcal{P}}}{\sqrt{n}}. \quad (15)$$

Each term of the last sum is a weakly converging sequence divided by \sqrt{n} so it converges to zero in probability. This proves that Δ_n converges to zero in probability, which ensures the convergence to the Gaussian law.

The expression of σ^2 is the direct consequence of theorems 3.1 and 3.7.

If $\sigma = 0$, then again by theorem 3.1 or 3.7, there is a continuous function ξ on $\mathbb{P}\mathbb{R}_{\max}^d$ such that

$$\phi(Ax) - \phi(x) = \xi(\bar{x}) - \xi(\overline{Ax}) \quad (16)$$

for μ -almost every A and ν_0 -almost every \bar{x} . Since all functions are continuous, (16) is true for every $A \in S_A$ and $\bar{x} \in S_{\nu_0}$. By induction we get it for $A \in T_A$ and if $\theta \in T_A$ has rank 1 and $\bar{x} \in S_{\nu_0}$, $\theta A \bar{x} = \theta \bar{x}$, so $\phi(\theta A \bar{x}) = \phi(\theta \bar{x})$.

As $\theta A \bar{x} - \theta \bar{x} \in \mathbb{R}\mathbf{1}$, this means that $\theta A \bar{x} = \theta \bar{x}$. By lemma 4.1, it proves that $\theta A \theta' = \theta \theta'$ for any $\theta, \theta' \in T_A$ with rank 1 and $A \in S_A$.

On the other side let us assume there is θ with rank one such that for any $\theta' \in T_A$ with rank 1 and $A \in S_A$, we have:

$$\theta A \theta' = \theta \theta'. \quad (17)$$

By lemma 4.2 applied with $a = b = 0$, it is true for any $\theta \in T_A$ with rank 1 and by induction, equation (17) is still true for $A \in T_A$.

Therefore, for any $m \in \mathbb{N}$ and $n \geq m + 1$ and any $\theta' \in T_A$ with rank 1, if $A(n) \cdots A(n - m + 1)$ has rank 1, then $x(n, \theta' \mathbf{1}) = A(n) \cdots A(n - m + 1) \theta' \mathbf{1}$ and for any $N \in \mathbb{N}$

$$\begin{aligned} & \mathbb{P}(\|x(n, \theta' \mathbf{1})\|_\infty \leq N) \\ & \geq \mathbb{P}(A(n) \cdots A(n - m + 1) \text{ has rank 1}, \|A(n) \cdots A(n - m + 1) \theta' \mathbf{1}\|_\infty \leq N) \\ & \geq \mathbb{P}(A(m) \cdots A(1) \text{ has rank 1}, \|A(m) \cdots A(1) \theta' \mathbf{1}\|_\infty \leq N). \end{aligned} \quad (18)$$

We fix a $\theta' \in T_A$ with rank one. The MLP property says there is an m such that $\mathbb{P}(A(m) \cdots A(1) \text{ has rank 1}) > 0$. Therefore, there is an $N \in \mathbb{N}$ such that the right member of 18 is a positive number we denote by β .

Equation 18 now implies that for any $\epsilon > 0$, if $n \geq \max(m, N^2 \epsilon^{-2})$, then $\mathbb{P}(\|\frac{1}{\sqrt{n}} x(n, \theta' \mathbf{1})\|_\infty \leq \epsilon) \geq \beta$, so $\mathcal{N}(0, \sigma^2)[- \epsilon, \epsilon] \geq \beta$. When ϵ tends to zero, we get that $\mathcal{N}(0, \sigma^2)(\{0\}) \geq \beta > 0$, which is true only if $\sigma = 0$. \square

Proof of theorem 2.2. Without loss of generality, we assume that $\gamma = 0$. Equation (2) follows from theorem 3.2 and proposition 4.1 or from theorem 3.8 and proposition 4.2

The only fact to check is that for any $f \in \mathcal{B}_\eta$, we have:

$$\lambda(f) \leq C (1 + \mathbb{E}(\max(\|X^0\|_\infty, 1)^\iota)) \|f\|_\eta.$$

It easily follows from the fact that $|f(x)| \leq \|f\|_\eta (1 + |x|_{\mathcal{P}})^{1+\eta}$ and $1 + \eta \leq m$.

Taking $y = 0$ in 14, we get $|\phi(x) - x_i| \leq |x|_{\mathcal{P}}$. Together with (15) it proves that for any $u \in \mathbb{R}^d$ and any $\epsilon > 0$

$$\begin{aligned} & \mathbb{P}[x(n, X^0) \leq \sigma u \sqrt{n}] \\ & \leq \mathbb{P}\left[\min_i x_i(n, X^0) \leq \sigma \min_i u_i \sqrt{n}\right] \end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{P} \left[\phi(x(n, X^0)) \leq (\sigma \min_i u_i + 2\epsilon)\sqrt{n} \right] + \mathbb{P} \left[\frac{|\phi(X^0)|}{\sqrt{n}} \geq \epsilon \right] + \mathbb{P} \left[\frac{|\bar{x}(n, X^0)|_{\mathcal{P}}}{\sqrt{n}} \geq \epsilon \right] \\
&\leq \mathcal{N}(0, 1)(] - \infty, \min_i u_i + \frac{2\epsilon}{\sigma}]) + \frac{C}{\sqrt{n}} + \frac{\mathbb{E} \left(|\phi(X^0)|^l \right)}{(\epsilon\sqrt{n})^l} + \frac{\mathbb{E} \left(|\bar{x}(n, X^0)|_{\mathcal{P}}^l \right)}{(\epsilon\sqrt{n})^l} \\
&\leq \mathcal{N}(0, 1)(] - \infty, \min_i u_i]) + \frac{C}{\sqrt{n}} + \frac{2\epsilon}{\sigma} + \frac{\mathbb{E} \left(|\phi(X^0)|^l \right)}{(\epsilon\sqrt{n})^l} + \frac{\mathbb{E} \left(|\bar{x}(n, X^0)|_{\mathcal{P}}^l \right)}{(\epsilon\sqrt{n})^l}. \tag{19}
\end{aligned}$$

On the other side

$$\begin{aligned}
&\mathbb{P}[x(n, X^0) \leq \sigma u \sqrt{n}] \\
&\geq \mathbb{P} \left[\phi(x(n, X^0)) \leq \sigma \min_i u_i \sqrt{n} \right] \\
&\geq \mathbb{P} \left[\phi(x(n, X^0)) \leq (\sigma \min_i u_i - 2\epsilon)\sqrt{n} \right] - \mathbb{P} \left[\frac{|\phi(X^0)|}{\sqrt{n}} \geq \epsilon \right] - \mathbb{P} \left[\frac{|\bar{x}(n, X^0)|_{\mathcal{P}}}{\sqrt{n}} \geq \epsilon \right] \\
&\geq \mathcal{N}(0, 1)(] - \infty, \min_i u_i - \frac{2\epsilon}{\sigma}]) - \frac{C}{\sqrt{n}} - \frac{\mathbb{E} \left(|\phi(X^0)|^l \right)}{(\epsilon\sqrt{n})^l} - \frac{\mathbb{E} \left(|\bar{x}(n, X^0)|_{\mathcal{P}}^l \right)}{(\epsilon\sqrt{n})^l} \\
&\geq \mathcal{N}(0, 1)(] - \infty, \min_i u_i]) - \frac{C}{\sqrt{n}} - \frac{2\epsilon}{\sigma} - \frac{\mathbb{E} \left(|\phi(X^0)|^l \right)}{(\epsilon\sqrt{n})^l} - \frac{\mathbb{E} \left(|\bar{x}(n, X^0)|_{\mathcal{P}}^l \right)}{(\epsilon\sqrt{n})^l} \tag{20}
\end{aligned}$$

Taking $\epsilon = n^{-\frac{l}{2(l+1)}}$ in (19) and (20) will conclude the proof of theorem 2.2 if we can show that $\mathbb{E} \left(|\bar{x}(n, X^0)|_{\mathcal{P}}^l \right)$ is bounded uniformly in n and X^0 . Without loss of generality, we assume X^0 .

For $n_0 \in \mathbb{N}$, we take $a \geq (\mathbb{P}[A(n_0) \cdots A(1) \text{ has not rank } 1])^{1/n_0}$. But if $A(n) \cdots A(m)$ has not rank 1, then for any integer less than $\frac{n-m}{n_0}$, the operator $A(1+in_0) \cdots A(1+(i+1)n_0)$ has not rank 1 either. From the Independence of the $A(n)$, we deduce

$$\mathbb{P}(A(n) \cdots A(m) \text{ has not rank } 1) \leq a^{n-m-n_0}.$$

We estimate $\delta(A(n) \cdots A(m)0, A(n) \cdots A(n_0+1+m)0)$: it is 0 when $A(n) \cdots A(n_0+1+m)$ has rank 1, and it is always less than $\delta(A(n_0) \cdots A(1)0, 0)$, that is less than $|A(n_0) \cdots A(1)0|_{\mathcal{P}}$. Therefore, we have for any $n \geq m+n_0$

$$\begin{aligned}
&\mathbb{E} \left[\delta^l(A(n) \cdots A(m)0, A(n) \cdots A(n_0+1+m)0) \right] \\
&\leq \mathbb{E} \left[\mathbf{1}_{A(n) \cdots A(n_0+1) \text{ has not rank } 1} |A(n_0) \cdots A(1)0|_{\mathcal{P}}^l \right] \\
&= a^{n-m-2n_0} \mathbb{E} \left[|A(n_0) \cdots A(1)0|_{\mathcal{P}}^l \right]. \tag{21}
\end{aligned}$$

Let $n = qn_0 + r$ be the euclidean division of n by n_0 . Then we have

$$\begin{aligned} |x(n, 0)|_{\mathcal{P}} &= \delta(A(n) \cdots A(1)0, 0) \\ &\leq \sum_{i=1}^q \delta(A(n) \cdots A(in_0 + 1)0, A(n) \cdots A((i-1)n_0 + 1)0) \\ &\quad + \delta(A(n) \cdots A(n-r+1)0, 0). \end{aligned}$$

Therefore we have:

$$\begin{aligned} \left(\mathbb{E} \left[|x(n, 0)|_{\mathcal{P}}^l \right] \right)^{1/l} &\leq \sum_{i=1}^q \left(a^{n-in_0-n_0} \mathbb{E} \left[|A(n_0) \cdots A(1)0|_{\mathcal{P}}^l \right] \right)^{1/l} \\ &\quad + \left(\mathbb{E} \left[|A(r) \cdots A(1)0, 0|_{\mathcal{P}}^l \right] \right)^{1/l}. \end{aligned} \quad (22)$$

We apply this decomposition again (with $n_0 = 1$ and $a = 1$), to check that

$$\left(\mathbb{E} \left[|A(n) \cdots A(1)0|_{\mathcal{P}}^l \right] \right)^{1/l} \leq n \left(\mathbb{E} \left[|A(1)0|_{\mathcal{P}}^l \right] \right)^{1/l}.$$

It follows from the MLP property, that there is $n_0 \in \mathbb{N}$ such that $a < 1$. Introducing the last equation in equation (22), we see that

$$\left(\mathbb{E} \left[|x(n, 0)|_{\mathcal{P}}^l \right] \right)^{1/l} \leq \left(1 + \frac{a^{-n_0 l}}{1 - a^{n_0 l}} \right) n_0 \left(\mathbb{E} \left[|A(1)0|_{\mathcal{P}}^l \right] \right)^{1/l}.$$

□

To go from the abstract LLT and renewal theorem to ours, we will use the following approximation lemma.

Lemma 4.4. *Let h be a continuous function with compact support from $\mathbb{R}^a \times \mathbb{R}^b$. Then there are two continuous functions f_0 and g_0 with compact support in \mathbb{R}^a and \mathbb{R}^b respectively, so that for any $\epsilon > 0$, there are f_i and g_i continuous functions with compact support satisfying:*

$$\forall x \in \mathbb{R}^a, y \in \mathbb{R}^b, |h(x, y) - \sum_i f_i(x)g_i(y)| \leq \epsilon f_0(x)g_0(y).$$

In the sequel, we denote by \mathcal{L} the Lebesgue measure.

Proof theorem 2.3. By theorem 2.1, the algebraic non arithmeticity ensures that $\sigma > 0$.

We apply proposition 4.1 and theorem 3.3 or proposition 4.2 and theorem 3.9. This proves that, if $g \in \mathcal{C}_c(\mathbb{R})$ and if f is a bounded Lipschitz function on $\mathbb{P}\mathbb{R}_{\max}^d$, then for any $x^0 \in \mathbb{R}^d$,

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E} \left(f(\bar{x}(n, x^0)) g(\phi(x(n, x^0)) - \phi(x^0) - u) \right) - e^{-\frac{u^2}{2n\sigma^2}} \nu(f) \mathcal{L}(g) \right| = 0. \quad (23)$$

Moreover these convergences are uniform in x^0 , because δ_{x^0} is bounded independently of x_0 as a linear form on \mathcal{B}^∞ and is in $\mathcal{B}'_{\eta, \|X^0\|_\infty}$ if $|x^0|_{\mathcal{P}} \leq \|X^0\|_\infty$. The uniformity allows us to take any random initial condition X^0 and get

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E} \left[f(\bar{x}(n, X^0)) g(\phi(x(n, X^0)) - u) \right] - \mathbb{E} \left[e^{-\frac{(u + \phi(X^0))^2}{2n\sigma^2}} \right] \nu(f) \mathcal{L}(g) \right| = 0. \quad (24)$$

But the density of bounded Lipschitz functions in $(\mathcal{C}_c(\mathbb{P}\mathbb{R}_{\max}^d), \|\cdot\|_\infty)$ allows us to take f and g continuous functions with compact support in equation (24). Now, it follows from lemma 4.4, that for any h continuous with compact support

$$\limsup_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E} \left[h(\bar{x}(n, X^0), \phi(x(n, X^0)) - u) \right] - \mathbb{E} \left[e^{-\frac{(u + \phi(X^0))^2}{2n\sigma^2}} \right] \nu \otimes \mathcal{L}(h) \right| = 0. \quad (25)$$

According to lemma 4.3 the function $\Phi : x \mapsto (\phi, \bar{x})$ is Lipschitz with a Lipschitz inverse, h has compact support iff $h \circ \Phi$ does. Since $\overline{x + u\mathbf{1}} = \bar{x}$, this concludes the proof. \square

Proof of proposition 2.1. Assume the sequence of random variables is algebraically arithmetic and the conclusion of theorem 2.3 holds.

There are $a, b \in \mathbb{R}$ and θ with rank 1, such that every $A \in S_A$ and $\theta' \in T_A$ with rank 1 satisfy equation (3). We set $t = \frac{2\pi}{b}$ if $b \neq 0$ and $t = 1$ otherwise, and for any $x \in \mathbb{R}^d$, $w(\bar{x}) = e^{it(\phi(\theta x) - \phi(x))}$, where $\phi(x) = \max_i(x_i)$. Equation (3) implies that, for any $A \in S_A$, $y \in \mathbb{R}^d$, and any $\theta' \in T_A$ with rank 1:

$$e^{it(\phi(A\theta'y) - \phi(\theta'y))} w(\overline{A\theta'y}) = e^{ita} w(\overline{\theta'y}).$$

We chose y such that $\phi(\theta'y) = 0$. By induction, we get

$$e^{it(\phi(x(n, \theta'y)) - \phi(\theta'y))} w(\overline{x(n, \theta'y)}) = e^{itna} w(\overline{\theta'y}). \quad (26)$$

For any $f : \mathbb{R} \mapsto \mathbb{R}$ and $g : \mathbb{P}\mathbb{R}_{\max}^d \mapsto \mathbb{R}$ continuous with compact support, the conclusion of theorem 2.3 for h defined by $h(x) = f(\phi(x))(gw)(\bar{x})$ is that:

$$\sigma \sqrt{2\pi n} \mathbb{E} [f(\phi(x(n, \theta'y))) g(\overline{x(n, \theta'y)}) w(\overline{x(n, \theta'y)})] \rightarrow \mathcal{L}(f) \nu_0(gw).$$

Together with equation (26), it means

$$e^{itna} w(\overline{\theta'y}) \sigma \sqrt{2\pi n} \mathbb{E} [e^{it \cdot} f((\phi(x(n, \theta'y)))g(\overline{x}(n, \theta'y)))] \rightarrow \mathcal{L}(f)\nu_0(gw) \quad (27)$$

But conclusion of theorem 2.3 for h defined by $h(x) = (fe^{it \cdot})(\phi(x))g(\overline{x})$ is that:

$$\sigma \sqrt{2\pi n} \mathbb{E} [e^{it \cdot} f((\phi(x(n, \theta'y)))g(\overline{\theta'y})))] \rightarrow \mathcal{L}(fe^{it \cdot})\nu_0(g) \quad (28)$$

Equations (27) and (28) together imply that $a = 0$ and that

$$w(\overline{\theta'y})\mathcal{L}(fe^{it \cdot})\nu_0(g) = \mathcal{L}(f)\nu_0(gw).$$

The right side of the equation does not depend on θ' so by lemma 4.1 w is constant on S_{ν_0} , this proves $\nu_0(gw) = w(\overline{\theta'y})$, so $\mathcal{L}(fe^{it \cdot}) = \mathcal{L}(f)$ that is $e^{it \cdot} = 1$ or $t = 0$. This is a contradiction, which concludes the proof. \square

Proof of theorem 2.4. Applying proposition 4.1 and theorem 3.4 again, we have that, if $g \in \mathcal{C}_c(\mathbb{R})$ and if f is a bounded Lipschitz function on $\mathbb{P}\mathbb{R}_{\max}^d$, then for any $x^0 \in \mathbb{R}^d$,

$$\lim_{a \rightarrow -\infty} \sum_{n \geq 1} \mathbb{E} [f(\phi(x(n, x^0)) - \phi(x^0) + a)g(\overline{x}(n, x^0))] = 0,$$

$$\lim_{a \rightarrow +\infty} \sum_{n \geq 1} \mathbb{E} [f(\phi(x(n, x^0)) - \phi(x^0) + a)g(\overline{x}(n, x^0))] = \frac{\nu(f)\mathcal{L}(g)}{\gamma}.$$

Moreover these convergences are uniform in x^0 , because δ_{x^0} is bounded as a linear form on \mathcal{B}^∞ . The uniformity allows us to remove the $\phi(x^0)$ in the last equations and take any random initial condition. The result follows by the same successive approximations as in the proof of the LLT. \square

Proof of theorem 2.5. Without lost of generality, we can assume that $\gamma = 0$.

The exponential moment of $\sigma(A)$ means that there is a $\theta > 0$ such that $\int e^{\theta\sigma(A)}d\mu(A) < \infty$. An easy bound of the norm of $\xi^k(y)Q(\cdot, dy)$ inspired by the proof of proposition 4.1 ensures that $z \mapsto Q_z$ is analytic on the open ball with center 0 and radius θ . To prove that it is continuous on the domain $\{|\mathcal{R}z| < \theta/2\}$, we apply the same method.

Now theorem 3.5 gives

$$\lim_n \frac{1}{n} \ln \mathbb{P} [\phi(x(n, X^0)) - \phi(X^0) > n\epsilon] = -c(\epsilon).$$

Let $0 < \eta < \epsilon$. For any $n \geq \phi(X^0)/\eta$, we have

$$\mathbb{P} [\phi(x(n, X^0)) - \phi(X^0) > n\epsilon] \geq \mathbb{P} [\phi(x(n, X^0)) > n(\epsilon + \eta)],$$

which implies that

$$\liminf_n \frac{1}{n} \ln \mathbb{P} [\phi(x(n, X^0)) - n\gamma > n\epsilon] \geq -c(\epsilon + \eta).$$

The same method gives

$$\limsup_n \frac{1}{n} \ln \mathbb{P} [\phi(x(n, X^0)) - n\gamma > n\epsilon] \leq -c(\epsilon - \eta).$$

By continuity of c , the first equality is proved. The second one follows from the same method applied to $-\phi$ instead of ϕ . \square

4.3 Max-plus case

Proof of theorem 2.8. Suppose that $\sigma = 0$. By proposition 4.1 we may apply theorem 3.1. The third point of the theorem says that there exists a bounded Lipschitz function f such that for ν -almost every (A, \bar{x}) :

$$\max_i (Ax)_i - \max_i x_i = f(\bar{x}) - f(\overline{Ax}) \quad (29)$$

Since all functions in that equation are continuous, every $A \in S_A$ and $\bar{x} \in \text{Supp}(\nu_0)$ satisfy equation (29). If $A \in S_A$ and $\bar{x} \in \text{Supp}(\nu_0)$, then $\overline{Ax} \in \text{Supp}(\nu_0)$, so by induction equation (29) is satisfied by A in T_A . As for $A \in T_A$, $A^n \in T_A$, $\max_i A^n x_i$ is bounded. But there exists a k such that $c(A)\rho_{\max}(A) = A^c(A)_{kk}$, so $\max_i (A^{nc(A)}x)_i \geq nc(A)\rho_{\max}(A) + x_k$ and $\rho_{\max}(A) \leq 0$.

As every path on $\mathcal{G}^c(A)$ can be split into a path with length at most d and closed paths whose average length are at most $\rho_{\max}(A)$, we have:

$$A^n x \leq (n - d)\rho_{\max}(A) + d \max_{A_{ij} > -\infty} |A_{ij}| + \max_i x_i,$$

therefore $\rho_{\max}(A) \geq 0$.

If $\rho_{\max}(A) = 0$ for $A \in T_A$, then

$$\begin{aligned} \phi(x(n, 0)) - \phi(0) &= \max_i x_i(n, 0) \\ &= \max_{ij} (A(n) \cdots A(1))_{ij} \\ &\geq \rho_{\max}(A(n) \cdots A(1)) = 0 \text{ a.s.} \end{aligned}$$

Therefore $\mathcal{N}(0, \sigma^2)(\mathbb{R}_+) \geq 1$, and $\sigma = 0$. \square

Proof of theorem 2.9. We assume the system is algebraically arithmetic. Then there are $a, b \in \mathbb{R}$ and $\theta \in T_A$ such that for any $A \in S_A$ and $\theta' \in T_A$ with rank 1, we have: $(\theta A \theta' - \theta \theta')(\mathbb{R}^d) \subset (a + b\mathbb{Z})\mathbf{1}$. Replacing θ' by $A^n \theta'$, we get $(\theta A^{n+1} \theta' - \theta A^n \theta')(\mathbb{R}^d) \subset (a + b\mathbb{Z})\mathbf{1}$ and by induction

$$(\theta A^{n+k} \theta' - \theta A^n \theta')(\mathbb{R}^d) \subset (ka + b\mathbb{Z})\mathbf{1} \quad (30)$$

By theorem 2.7, if $\mathcal{G}(A)$ is strongly connected, for n big enough, $A^{nc(A)} - nc(A)\rho_{\max}(A)\mathbf{1} = (A - \rho_{\max}(A)\mathbf{1})^{nc(A)}$ is constant, because $(A - \rho_{\max}(A)\mathbf{1})^{c(A)}$ has spectral radius 0 and cyclicity 1. Therefore $A^{(n+1)c(A)} = A^{nc(A)} + c(A)\rho_{\max}(A)\mathbf{1}$, and $\theta A^{n+c(A)} \theta' - \theta A^n \theta' = \theta \theta' + c(A)\rho_{\max}(A)\mathbf{1}$. Together with equation (30), this concludes the proof. \square

References

- [BL85] Philippe Bougerol and Jean Lacroix. *Products of random matrices with applications to Schrödinger operators*, volume 8 of *Progress in Probability and Statistics*. Birkhäuser Boston Inc., Boston, MA, 1985.
- [BM98] François Baccelli and Jean Mairesse. Ergodic theorems for stochastic operators and discrete event networks. In *Idempotency (Bristol, 1994)*, volume 11 of *Publ. Newton Inst.*, pages 171–208. Cambridge Univ. Press, Cambridge, 1998.
- [CDQV83] G. Cohen, D. Dubois, J.P. Quadrat, and M. Viot. Analyse du comportement périodique des systèmes de production par la théorie des diodes. Rapport de recherche 191, INRIA, Le Chesnay, France, 1983.
- [CDQV85] G. Cohen, D. Dubois, J.P. Quadrat, and M. Viot. A linear system theoretic view of discrete event processes and its use for performance evaluation in manufacturing. *IEEE Trans. on Automatic Control*, AC-30:210–220, 1985.
- [Fur63] Harry Furstenberg. Noncommuting random products. *Trans. Amer. Math. Soc.*, 108:377–428, 1963.
- [GdM89] I. Ya. Gol'dsheĭ d and G. A. Margulis. Lyapunov exponents of a product of random matrices. *Uspekhi Mat. Nauk*, 44(5(269)):13–60, 1989.

- [GR85] Y. Guivarc’h and A. Raugi. Frontière de Furstenberg, propriétés de contraction et théorèmes de convergence. *Z. Wahrsch. Verw. Gebiete*, 69(2):187–242, 1985.
- [HH01] Hubert Hennion and Loïc Hervé. *Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness*, volume 1766 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [HH04] Hubert Hennion and Loïc Hervé. Central limit theorems for iterated random Lipschitz mappings. *Ann. Probab.*, 32(3A):1934–1984, 2004.
- [LP82] Émile Le Page. Théorèmes limites pour les produits de matrices aléatoires. In *Probability measures on groups (Oberwolfach, 1981)*, volume 928 of *Lecture Notes in Math.*, pages 258–303. Springer, Berlin, 1982.
- [Mai97] J. Mairesse. Products of irreducible random matrices in the $(\max, +)$ algebra. *Adv. in Appl. Probab.*, 29(2):444–477, 1997.
- [Mer04] Glenn Merlet. Memory loss property for products of random matrices in the $(\max, +)$ algebra. Technical report, IRMAR, <http://fr.arxiv.org/abs/math.PR/0405452>, May 2004.
- [Nag57] S. V. Nagaev. Some limit theorems for stationary Markov chains. *Teor. Veroyatnost. i Primenen.*, 2:389–416, 1957.
- [RdVH⁺90] J. A. C. Resing, R. E. de Vries, G. Hooghiemstra, M. S. Keane, and G. J. Olsder. Asymptotic behavior of random discrete event systems. *Stochastic Process. Appl.*, 36(2):195–216, 1990.
- [Too02] Fergal Toomey. Large deviations of products of random topical operators. *Ann. Appl. Probab.*, 12(1):317–333, 2002.
- [Vin97] Jean-Marc Vincent. Some ergodic results on stochastic iterative discrete events systems. *Discrete Event Dynamic Systems*, 7(2):209–232, 1997.