



Loop equations for the semiclassical 2-matrix model with hard edges

Bertrand Eynard

► To cite this version:

Bertrand Eynard. Loop equations for the semiclassical 2-matrix model with hard edges. 2005. hal-00004618

HAL Id: hal-00004618

<https://hal.science/hal-00004618>

Preprint submitted on 1 Apr 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Loop equations for the semiclassical 2-matrix model with hard edges

*B. Eynard*¹

Service de Physique Théorique de Saclay,
F-91191 Gif-sur-Yvette Cedex, France.

Abstract: The 2-matrix model can be defined in a setting more general than polynomial potentials, namely, the semiclassical matrix model. In this case, the potentials are such that their derivatives are rational functions, and the integration paths for eigenvalues are arbitrary homology classes of paths for which the integral is convergent. This choice includes in particular the case where the integration path has fixed endpoints, called hard edges. The hard edges induce boundary contributions in the loop equations. The purpose of this article is to give the loop equations in that semiclassical setting.

1 Introduction

Orthogonal polynomials and biorthogonal polynomials, in the context of random matrices, have been mostly studied for polynomial potentials, on the real axis, or sometimes on homology classes of contours going from ∞ to ∞ [18]. However, it is possible to define matrix models corresponding to a more general context, in particular, the “semi-classical” (called so, because it contains all the classical polynomials). It is defined as follows (see Bertola [1]):

Consider two potentials, (i.e. functions of a complex variable), $V_1(x)$ and $V_2(y)$ whose derivatives $V_1'(x)$ and $V_2'(y)$ are rational functions (notice that ∞ maybe a pole of V_1' (resp. V_2'), this is the case if V_1' (resp. V_2') is a polynomial).

Then consider a generalized integration path $\Gamma = \sum_{i,j} \kappa_{i,j} \gamma_i \times \tilde{\gamma}_i$, such that the following integral is absolutely convergent:

$$\int_{\Gamma} e^{-\frac{N}{t}[V_1(x)+V_2(y)-xy]} dx dy \quad (1-1)$$

¹E-mail: eynard@cea.fr

The possible paths γ (resp. $\tilde{\gamma}$) are described in [1], they can be closed or open:

- If γ (resp. $\tilde{\gamma}$) is a closed contour, the result is non-zero only if it encloses a singularity of $e^{-\frac{N}{t}V_1}$ (resp. $e^{-\frac{N}{t}V_2}$).
- if γ (resp. $\tilde{\gamma}$) is an open contour, its extremities can be:
 - any point in the complex plane, except at the poles of V'_1 (resp. V'_2),
 - a simple pole of $\frac{N}{t}V'_1$ (resp. $\frac{N}{t}V'_2$), with residue $\in \mathbb{R}_-$,
 - a degree ≥ 2 pole of V'_1 (resp. V'_2), if γ (resp. $\tilde{\gamma}$) approaches the pole in a sector where $\Re \frac{N}{t}V_1 > 0$ (resp. $\Re \frac{N}{t}V_2 > 0$).

Some of the extremities are such that $e^{-\frac{N}{t}[V_1(x)+V_2(y)-xy]}$ vanishes, they are the poles of V'_1 (resp. V'_2) of degree at least 2, as well as the simple poles of $\frac{N}{t}V'_1$ (resp. $\frac{N}{t}V'_2$), with residue $\in \mathbb{R}_-$, we will call them “poles”. And some extremities are such that $e^{-\frac{N}{t}[V_1(x)+V_2(y)-xy]}$ does not vanish, they are arbitrary points in the complex plane, we call them “hard edges”, and write them:

$$(X_1, X_2, \dots, X_{K_1}) \quad (\text{resp. } (Y_1, Y_2, \dots, Y_{K_2})) \quad (1-2)$$

Definition 1.1 Consider a path γ in the complex plane. Define the set of “normal matrices” on the path γ :

$$H_N(\gamma) := \{M \in GL_N(\mathbb{C}) \mid \exists U \in U(N), \exists (x_1, \dots, x_N) \in \gamma^N, M = U \text{diag}(x_1, \dots, x_N) U^\dagger\} \quad (1-3)$$

with a measure:

$$dM := \left(\prod_{i>j} (x_i - x_j)^2 \right) dU \prod_{i=1}^N dx_i \quad (1-4)$$

where dU is the Haar measure on $U(N)$, and dx_i are the tangent vectors to the path γ . For a given $M \in H_N(\Gamma)$, the matrix U and the eigenvalues x_i are not uniquely defined, they are defined up to a permutation, and a conjugation by a diagonal unitary matrix, however dM is invariant under these operations, and thus is well defined.

For example, if γ is the real axis we recognize $H_N(\gamma) = H_N$ the set of hermitean matrices.

Then, as in [15], we define the set of normal matrices on a generalized path $\Gamma = \sum_{i,j} \kappa_{i,j} \gamma_i \times \tilde{\gamma}_j$:

Definition 1.2 Let $\Gamma = \sum_{i,j} \kappa_{i,j} \gamma_i \times \tilde{\gamma}_j$ be a linear combination of products of paths. We define:

$$H_N^2(\Gamma) := \{M_1, M_2 \in GL_N(\mathbb{C})^2 \mid \exists U_1, U_2 \in U(N)^2, \exists (x_1, \dots, x_N) \in \mathbb{C}^N, \exists (y_1, \dots, y_N) \in \mathbb{C}^N, \\ M_1 = U_1 \text{diag}(x_1, \dots, x_N) U_1^\dagger, M_2 = U_2 \text{diag}(y_1, \dots, y_N) U_2^\dagger\}$$

$$\forall k = 1, \dots, N, \exists i_k x_k \in \gamma_{i_k}, \forall k = 1, \dots, N, \exists j_k y_k \in \tilde{\gamma}_{j_k} \} \quad (1-5)$$

with a measure $dM_1 dM_2$ such that if $(x_1, \dots, x_N)$ are the eigenvalues of M_1 and $(y_1, \dots, y_N)$ are the eigenvalues of M_2 , and $x_k \in \gamma_{i_k}$ and $y_k \in \tilde{\gamma}_{j_k}$:

$$dM_1 dM_2 := \det_{k,l} (\kappa_{i_k, j_l} e^{x_{i_k} y_{j_l}}) \left(\prod_{i>j} (x_i - x_j) \right) \left(\prod_{i>j} (y_i - y_j) \right) dU_1 dU_2 \prod_{i=1}^N dx_i \prod_{i=1}^N dy_i \quad (1-6)$$

Our goal is to compute the Schwinger-Dyson equations, also called loop equations in the context of random matrices, of the following matrix integral:

$$Z := \int_{H_N^2(\Gamma)} e^{-\frac{N}{t} \text{Tr}[V_1(M_1) + V_2(M_2) - M_1 M_2]} dM_1 dM_2 \quad (1-7)$$

1.1 Notations

Define the following polynomials, which vanish at all the hard edges:

$$s(x) := \prod_{j=1}^{K_1} (x - X_j) = \sum_{r=0}^{K_1} s_r x^r \quad (1-8)$$

$$\tilde{s}(y) := \prod_{j=1}^{K_2} (y - Y_j) = \sum_{r=0}^{K_2} \tilde{s}_r y^r \quad (1-9)$$

The resolvent, (which is a formal series in its large x expansion), is defined by:

$$W(x) := \frac{t}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \right\rangle \quad (1-10)$$

And, up to a shift by the potential, it is more convenient to use $Y(x)$:

$$Y(x) := V_1'(x) - W(x) \quad (1-11)$$

Then, for technical intermediate calculations, define the following expectation values, which are formal series in their large x expansion:

$$U(x, y) := \frac{t}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle \quad (1-12)$$

$$U(x, y, x') := \left\langle \text{Tr} \frac{1}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \text{Tr} \frac{1}{x' - M_1} \right\rangle_c \quad (1-13)$$

$$P(x, y) := \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle \quad (1-14)$$

$$A(x) := \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} \tilde{s}(M_2) V_2'(M_2) \right\rangle \quad (1-15)$$

$$B(x) := \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} \tilde{s}(M_2) \right\rangle \quad (1-16)$$

$$B_k(x) := \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} M_2^k \right\rangle \quad (1-17)$$

$$D(x) := xB(x) + \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle B_j(x) \quad (1-18)$$

Notice that $U(x, y)$ and $U(x, y, x')$ are rational functions of y , and $P(x, y)$, is a rational function of both x and y , whose poles are the same as the poles of V_1' and V_2' .

2 The loop equations

Loop equations, i.e. Schwinger-Dyson equations, merely express the fact that an integral is invariant under a change of variable. Schwinger equations have been extensively studied in the context of field theory, and have been of great importance for the study of matrix models [6]. For the polynomial 2-matrix model, loop equations were first announced to give algebraic equations by M. Staudacher [19], and computed precisely for general potentials, including $1/N^2$ corrections by [12, 13, 14, 10]. They have been solved recently by [8] following the method of [7]. Here, we follow the method of [14].

2.1 Generalities

2.1.1 Changes of variables

Consider a change of variable in $H_n^2(\Gamma)$, of the form:

$$M_1 \rightarrow M_1' = M_1 + \epsilon f(M_1, M_2) + \bar{\epsilon} f^\dagger(M_1, M_2) + O(|\epsilon|^2) \quad (2-1)$$

Remark 2.1 Under such a change of variable, the eigenvalues of M'_1 are no longer on paths γ_{i_k} , but on some paths γ'_{i_k} , which, for ϵ small enough, are small deformations of γ_{i_k} , and thus the integral is indeed unchanged. This is true only if we can deform the contours, i.e. not at the extremities. At poles, the integrand vanishes, so that the integral is still invariant. At hard edges, the integral is invariant only if $M'_1 = M_1$, i.e. $f(M_1, M_2)$ vanishes at hard edges.

To order 1 in $|\epsilon|$, one has:

$$\begin{aligned} & \exp \left[-\frac{N}{t} \text{Tr} (V_1(M'_1) + V_2(M_2) - M'_1 M_2) \right] \\ &= (1 - \epsilon \mathcal{S}(f) - \bar{\epsilon} \bar{\mathcal{S}}(f) + O(|\epsilon|^2)) \exp \left[-\frac{N}{t} \text{Tr} (V_1(M_1) + V_2(M_2) - M_1 M_2) \right] \end{aligned} \quad (2-2)$$

with

$$\mathcal{S}(f) = \frac{N}{t} \text{Tr} (V'_1(M_1) - M_2) f(M_1, M_2) \quad (2-3)$$

The measure dM_1 is multiplied by a Jacobian J , which we expand to order 1 in $|\epsilon|$:

$$dM'_1 = J(f) dM_1 = (1 + \epsilon \mathcal{J}(f) + \bar{\epsilon} \bar{\mathcal{J}}(f) + O(\epsilon^2)) dM_1 \quad (2-4)$$

Loop equations are obtained by writing that the integral is unchanged, i.e.:

$$\begin{aligned} Z &:= \int dM_1 dM_2 \exp \left[-\frac{N}{t} \text{Tr} (V_1(M_1) + V_2(M_2) - M_1 M_2) \right] \\ &= \int dM'_1 dM_2 \exp \left[-\frac{N}{t} \text{Tr} (V_1(M'_1) + V_2(M_2) - M'_1 M_2) \right] \\ &= \int dM_1 dM_2 (1 + \epsilon (\mathcal{J}(f) - \mathcal{S}(f)) + \bar{\epsilon} (\bar{\mathcal{J}}(f) - \bar{\mathcal{S}}(f)) + O(|\epsilon|^2)) \\ &\quad \exp \left[-\frac{N}{t} \text{Tr} (V_1(M_1) + V_2(M_2) - M_1 M_2) \right] \end{aligned} \quad (2-5)$$

Since this must be true for all argument of ϵ , one must have:

$$0 = \int dM_1 dM_2 (\mathcal{J}(f) - \mathcal{S}(f)) \exp \left[-\frac{N}{t} \text{Tr} (V_1(M_1) + V_2(M_2) - M_1 M_2) \right] \quad (2-6)$$

i.e.

$$\langle \mathcal{J}(f) \rangle = \langle \mathcal{S}(f) \rangle \quad (2-7)$$

This equation is called a loop equation.

Let us emphasize again that eq.2-5, and thus eq.2-7 hold only if f vanishes at the hard edges.

2.1.2 Split and merge rules

The rules to compute $\mathcal{J}(f)$ are called split and merge rules and can be found in the litterature, for instance in [14]. Notice that $\mathcal{J}$ is linear, and if f is a product, $\mathcal{J}(f)$ can be computed by the chain rule. Thus, it is useful to determine $\mathcal{J}(f)$ for some particular f . For any two matrices A and B , one has:

- *Split rule:*

$$\mathcal{J} \left(A \frac{1}{x - M_1} B \right) = \text{Tr} A \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} B \quad (2-8)$$

or equivalently:

$$\mathcal{J} (A M_1^k B) = \sum_{j=0}^{k-1} \text{Tr} A M_1^j \text{Tr} M_1^{k-1-j} B \quad (2-9)$$

- *Merge rule:*

$$\mathcal{J} \left(A \text{Tr} \left(\frac{1}{x - M_1} B \right) \right) = \text{Tr} \frac{1}{x - M_1} A \frac{1}{x - M_1} B \quad (2-10)$$

or equivalently:

$$\mathcal{J} (A \text{Tr} (M_1^k B)) = \sum_{j=0}^{k-1} \text{Tr} M_1^j A M_1^{k-j-1} B \quad (2-11)$$

2.2 Equations

In this section, we write the change of - variable, i.e. function f , and the corresponding loop equation of type eq.2-7. We always write the contribution of the Jacobian $\mathcal{J}$ in the LHS and the contribution of the action $\mathcal{S}$ in the RHS.

- computation of $B(x)$: from

$$\begin{aligned} f(M_1, M_2) &= s(M_1) \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \\ &= s(x) \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} - \frac{s(x) - s(M_1)}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \end{aligned} \quad (2-12)$$

(which indeed vanishes at all hard edges) we get:

$$\begin{aligned} &s(x) W(x) \frac{t}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\ &+ s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\ &- \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \end{aligned}$$

$$\begin{aligned}
&= \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} (V'_1(M_1) - M_2) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
(2-13)
\end{aligned}$$

i.e.

$$\begin{aligned}
B(x) &= s(x) \tilde{s}(Y(x)) W(x) \\
&\quad - s(x) \frac{t}{N} \left\langle \text{Tr} \frac{V'_1(x) - V'_1(M_1)}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&\quad - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \tilde{s}(M_2) \right\rangle \\
&\quad - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V'_1(M_1) - M_2) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&\quad + \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&\quad - s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
(2-14)
\end{aligned}$$

• computation of $B_k(x)$: from

$$\begin{aligned}
f(M_1, M_2) &= s(M_1) \frac{1}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \\
&= s(x) \frac{1}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} - \frac{s(x) - s(M_1)}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \\
(2-15)
\end{aligned}$$

(which indeed vanishes at all hard edges) we get:

$$\begin{aligned}
&s(x) W(x) \frac{t}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle \\
&+ s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle_c \\
&- \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle \\
&= \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} (V'_1(M_1) - M_2) \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle \\
(2-16)
\end{aligned}$$

i.e.

$$\begin{aligned}
B_k(x) &= Y(x)^k s(x) W(x) \\
&\quad - s(x) \frac{t}{N} \left\langle \text{Tr} \frac{V'_1(x) - V'_1(M_1)}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle
\end{aligned}$$

$$\begin{aligned}
& -\frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle \\
& -\frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} M_2^k \right\rangle \\
& + \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle \\
& -s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{Y(x)^k - M_2^k}{Y(x) - M_2} \right\rangle_c
\end{aligned}
\tag{2-17}$$

• Computation of $D(x)$:

$$\begin{aligned}
D(x) &= xB(x) \\
&+ \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle B_j(x) \\
&= xs(x) \tilde{s}(Y(x)) W(x) \\
&+ s(x) W(x) \frac{t}{N} \left\langle \text{Tr} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&- xs(x) \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&- x \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \tilde{s}(M_2) \right\rangle \\
&- x \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&+ x \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
&- s(x) \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle \\
&- \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle \\
&- \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_2'(M_2) - M_1) \tilde{s}(M_2) \right\rangle \\
&+ \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle \\
&- xs(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
&- s(x) \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle_c
\end{aligned}$$

$$(2-18) \quad + \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t^2}{N^2} \left\langle \text{Tr } M_2^{s-1-j} \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} M_2^j \right\rangle_c$$

- computation of $A(x)$: By doing a change of variable on M_2 of the form:

$$(2-19) \quad \tilde{f}(M_1, M_2) = s(M_1) \frac{1}{x - M_1} \tilde{s}(M_2)$$

we have:

$$(2-20) \quad \begin{aligned} & \frac{t^2}{N^2} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} M_2^j \text{Tr } M_2^{s-1-j} \right\rangle \\ &= A(x) - xB(x) + \frac{t}{N} \langle \text{Tr } s(M_1) \tilde{s}(M_2) \rangle \end{aligned}$$

i.e.

$$(2-21) \quad \begin{aligned} A(x) &= D(x) \\ &\quad - \frac{t}{N} \langle \text{Tr } s(M_1) \tilde{s}(M_2) \rangle \\ &\quad + \frac{t^2}{N^2} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} M_2^j \text{Tr } M_2^{s-1-j} \right\rangle_c \end{aligned}$$

- Main computation: from

$$(2-22) \quad \begin{aligned} f(M_1, M_2) &= s(M_1) \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \\ &= s(x) \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \\ &\quad - \frac{s(x) - s(M_1)}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \end{aligned}$$

we get:

$$\begin{aligned} & s(x) \frac{t}{N} W(x) \left\langle \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & + s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle_c \\ & - \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr } M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \end{aligned}$$

$$= \frac{t}{N} \left\langle \text{Tr } s(M_1) \frac{1}{x - M_1} (V_1'(M_1) - M_2) \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle$$

(2 - 23)

i.e.

$$\begin{aligned} & s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle_c \\ = & s(x) \tilde{s}(Y(x))V_2'(Y(x))W(x) \\ & - s(x) \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & + \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \tilde{s}(M_2)V_2'(M_2) \right\rangle \\ & - A(x) \end{aligned}$$

(2 - 24)

i.e.

$$\begin{aligned} & s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle_c \\ & + \frac{t^2}{N^2} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \left\langle \text{Tr} \frac{s(M_1)}{x - M_1} M_2^j \text{Tr} M_2^{s-1-j} \right\rangle_c \\ = & s(x) \tilde{s}(Y(x))V_2'(Y(x))W(x) \\ & - s(x) \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & + \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{\tilde{s}(Y(x))V_2'(Y(x)) - \tilde{s}(M_2)V_2'(M_2)}{Y(x) - M_2} \right\rangle \\ & - \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \tilde{s}(M_2)V_2'(M_2) \right\rangle \\ & + \frac{t}{N} \langle \text{Tr } s(M_1) \tilde{s}(M_2) \rangle \\ & - \tilde{D}(x) \end{aligned}$$

(2 - 25)

Now, inserting the value of $D(x)$, and after tedious but straightforward computations, we get:

$$s(x) \tilde{s}(Y(x)) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle_c$$

$$\begin{aligned}
& +s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x-M_1} \text{Tr} \frac{1}{x-M_1} (V_2'(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
& -s(x) \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x-M_1} \text{Tr} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
& +s(x) \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t^2}{N^2} \left\langle \text{Tr} M_2^{s-1-j} \text{Tr} \frac{1}{x-M_1} M_2^j \right\rangle_c \\
& -s(x) \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x-M_1} \text{Tr} \frac{1}{x-M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle_c \\
& + \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x-M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle_c \\
& - \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x-M_1} (V_2'(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
= & s(x) \tilde{s}(Y(x)) ((V_2'(Y(x)) - x)(V_1'(x) - Y(x)) - P(x, y) + t) \\
& -s(x) \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x-M_1} (V_2'(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
& +s(x) \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \langle \text{Tr} M_2^{s-1-j} \rangle \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x-M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle \\
& -\tilde{s}(Y(x)) \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x-M_1} (V_1'(M_1) - M_2) \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle \\
& +\tilde{s}(Y(x)) \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x-M_1} \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle \\
& -\frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x-M_1} \right. \\
& \quad \left. (V_1'(M_1)V_2'(M_2) - M_2V_2'(M_2) - M_1V_1'(M_1) + M_1M_2) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
& -\frac{t^2}{N^2} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x-M_1} \text{Tr} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
& +\frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \langle \text{Tr} M_1^{r-1-i} \rangle \left\langle \text{Tr} \frac{x^i - M_1^i}{x-M_1} (V_2'(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle \\
& +\frac{t^2}{N^2} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \langle \text{Tr} M_2^{s-1-j} \rangle \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x-M_1} (V_1'(M_1) - M_2) \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle \\
& -\frac{t^3}{N^3} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} s_r \tilde{s}_s \langle \text{Tr} M_1^{r-1-i} \rangle \langle \text{Tr} M_2^{s-1-j} \rangle \left\langle \text{Tr} \frac{x^i - M_1^i}{x-M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle
\end{aligned}$$

which we write as follows:

$$\boxed{s(x)\tilde{s}(y) E(x, Y(x)) = \frac{t^2}{N^2} L(x)} \quad (2-27)$$

where $E(x, y)$ is a rational fraction of both x and y with poles at the poles of V'_1 and V'_2 , and at the hard edges:

$$\begin{aligned} & E(x, y) \\ = & \frac{(V'_2(y) - x)(V'_1(x) - y) - P(x, y) + t}{\tilde{s}(y) N} \left\langle \text{Tr} \frac{V'_1(x) - V'_1(M_1)}{x - M_1} (V'_2(M_2) - M_1) \frac{\tilde{s}(y) - \tilde{s}(M_2)}{y - M_2} \right\rangle \\ & + \frac{1}{\tilde{s}(y)} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \left\langle \text{Tr} M_2^{s-1-j} \right\rangle \frac{t}{N} \left\langle \text{Tr} \frac{V'_1(x) - V'_1(M_1)}{x - M_1} \frac{y^j - M_2^j}{y - M_2} \right\rangle \\ & - \frac{1}{s(x)} \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V'_1(M_1) - M_2) \frac{V'_2(y) - V'_2(M_2)}{y - M_2} \right\rangle \\ & + \frac{1}{s(x)} \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{V'_2(y) - V'_2(M_2)}{y - M_2} \right\rangle \\ & - \frac{1}{s(x)\tilde{s}(y)} \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \right. \\ & \quad \left. (V'_1(M_1)V'_2(M_2) - M_2V'_2(M_2) - M_1V'_1(M_1) + M_1M_2) \frac{\tilde{s}(y) - \tilde{s}(M_2)}{y - M_2} \right\rangle \\ & - \frac{1}{s(x)\tilde{s}(y)} \frac{t^2}{N^2} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} \text{Tr} \frac{\tilde{s}(y) - \tilde{s}(M_2)}{y - M_2} \right\rangle \\ & + \frac{1}{s(x)\tilde{s}(y)} \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \right\rangle \left\langle \text{Tr} \frac{x^i - M_1^i}{x - M_1} (V'_2(M_2) - M_1) \frac{\tilde{s}(y) - \tilde{s}(M_2)}{y - M_2} \right\rangle \\ & + \frac{1}{s(x)\tilde{s}(y)} \frac{t^2}{N^2} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \left\langle \text{Tr} M_2^{s-1-j} \right\rangle \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V'_1(M_1) - M_2) \frac{y^j - M_2^j}{y - M_2} \right\rangle \\ & - \frac{1}{s(x)\tilde{s}(y)} \frac{t^3}{N^3} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} s_r \tilde{s}_s \left\langle \text{Tr} M_1^{r-1-i} \right\rangle \left\langle \text{Tr} M_2^{s-1-j} \right\rangle \left\langle \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{y^j - M_2^j}{y - M_2} \right\rangle \end{aligned} \quad (2-28)$$

and in the right hand side of eq.2-27, we have:

$$\begin{aligned} & L(x) \\ = & \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V'_2(Y(x)) - V'_2(M_2)}{Y(x) - M_2} \right\rangle_c \\ & + \frac{1}{\tilde{s}(Y(x))} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} (V'_2(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\tilde{s}(Y(x))} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c \\
& + \frac{1}{\tilde{s}(Y(x))} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \left\langle \text{Tr} M_2^{s-1-j} \text{Tr} \frac{1}{x - M_1} M_2^j \right\rangle_c \\
& - \frac{1}{\tilde{s}(Y(x))} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} \tilde{s}_s \frac{t}{N} \left\langle \text{Tr} M_2^{s-1-j} \right\rangle \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle_c \\
& + \frac{1}{s(x)\tilde{s}(Y(x))} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} \sum_{s=0}^{K_2} \sum_{j=0}^{s-1} s_r \tilde{s}_s \\
& \quad \frac{t}{N} \left\langle \text{Tr} M_2^{s-1-j} \right\rangle \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} \frac{Y(x)^j - M_2^j}{Y(x) - M_2} \right\rangle_c \\
& - \frac{1}{s(x)\tilde{s}(Y(x))} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M_1^{r-1-i} \text{Tr} \frac{x^i - M_1^i}{x - M_1} (V_2'(M_2) - M_1) \frac{\tilde{s}(Y(x)) - \tilde{s}(M_2)}{Y(x) - M_2} \right\rangle_c
\end{aligned}
\tag{2-29}$$

2.3 Examples

Equation eq.2-27 looks rather terrible, but it is actually very simple to use. Let us illustrate it on simple examples.

2.3.1 No hard edges

If there is no hard edges, we have $s(x) = 1$ and $\tilde{s}(y) = 1$, and the loop equation becomes:

$$(V_2'(Y(x)) - x)(V_1'(x) - Y(x)) - P(x, Y(x)) + t = \frac{t^2}{N^2} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle_c
\tag{2-30}$$

which is the well known loop equation of the 2-matrix model with polynomial potentials [14].

2.3.2 1-matrix model

Consider the 1-matrix model with weight $e^{-\frac{N}{t} \text{Tr} V(M)}$. It is equivalent to a 2-matrix model where M_2 is gaussian, i.e. $V_2'(y) = y$, $\tilde{s}(y) = 1$, $V'(x) = V_1'(x) - x$. The equation eq.2-27 becomes:

$$\begin{aligned}
E(x, y) &= (y - x)(V'(x) + x - y) - \frac{t}{N} \left\langle \text{Tr} \frac{V'(x) - V'(M)}{x - M} \right\rangle \\
&\quad - \frac{1}{s(x)} \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M)}{x - M} V'(M) \right\rangle \\
&\quad + \frac{1}{s(x)} \frac{t^2}{N^2} \sum_{r=0}^{K_1} \sum_{i=0}^{r-1} s_r \left\langle \text{Tr} M^{r-1-i} \text{Tr} \frac{x^i - M^i}{x - M} \right\rangle
\end{aligned}$$

(2-31)

and in the right hand side of eq.2-27, we have:

$$L(x) = \left\langle \text{Tr} \frac{1}{x-M} \text{Tr} \frac{1}{x-M} \right\rangle_c \quad (2-32)$$

Since $E(x, y)$ is quadratic in y , this equation defines an hyperelliptical curve.

2.3.3 1-matrix model with only one hard edge

In particular, consider $s(x) = (x - a)$, we have:

$$E(x, y) = (y - x)(V'(x) + x - y) - \frac{t}{N} \left\langle \text{Tr} \frac{V'(x) - V'(M)}{x - M} \right\rangle - \frac{1}{x - a} \frac{t}{N} \langle \text{Tr} V'(M) \rangle \quad (2-33)$$

2.3.4 1-matrix model with only two hard edges

In particular, consider $s(x) = (x - a)(x - b)$, we have:

$$\begin{aligned} E(x, y) &= (y - x)(V'(x) + x - y) - \frac{t}{N} \left\langle \text{Tr} \frac{V'(x) - V'(M)}{x - M} \right\rangle \\ &\quad - \frac{1}{(x - a)(x - b)} \frac{t}{N} \langle \text{Tr} (x + M - a - b) V'(M) \rangle + \frac{t^2}{(x - a)(x - b)} \end{aligned} \quad (2-34)$$

2.3.5 2-matrix model with only one hard edge

Consider $s(x) = (x - a)$ and $\tilde{s}(y) = 1$, we have:

$$\begin{aligned} E(x, y) &= (V'_2(y) - x)(V'_1(x) - y) - P(x, y) + t \\ &\quad - \frac{1}{x - a} \frac{t}{N} \left\langle \text{Tr} (V'_1(M_1) - M_2) \frac{V'_2(y) - V'_2(M_2)}{y - M_2} \right\rangle \end{aligned} \quad (2-35)$$

and in the right hand side of eq.2-27, we have:

$$L(x) = \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V'_2(Y(x)) - V'_2(M_2)}{Y(x) - M_2} \right\rangle_c \quad (2-36)$$

2.3.6 2-matrix model with only two hard edges

Consider $s(x) = (x - a)(x - b)$ and $\tilde{s}(y) = 1$, we have:

$$\begin{aligned}
E(x, y) &= (V_2'(y) - x)(V_1'(x) - y) - P(x, y) + t \\
&\quad - \frac{1}{s(x)} \frac{t}{N} \left\langle \text{Tr} (x + M_1 - a - b) (V_1'(M_1) - M_2) \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle \\
&\quad + \frac{1}{s(x)} \frac{t^2}{N} \left\langle \text{Tr} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle
\end{aligned}
\tag{2-37}$$

and in the right hand side of eq.2-27, we have:

$$L(x) = \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle_c
\tag{2-38}$$

2.3.7 2-matrix model with only one hard edge in x and one hard edge in y

Consider $s(x) = (x - a)$ and $\tilde{s}(y) = (y - b)$, we have:

$$\begin{aligned}
E(x, y) &= (V_2'(y) - x)(V_1'(x) - y) - P(x, y) + t \\
&\quad - \frac{1}{\tilde{s}(y)} \frac{t}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} (V_2'(M_2) - M_1) \right\rangle \\
&\quad - \frac{1}{s(x)} \frac{t}{N} \left\langle \text{Tr} (V_1'(M_1) - M_2) \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle \\
&\quad - \frac{1}{s(x)\tilde{s}(y)} \frac{t}{N} \langle \text{Tr} (V_1'(M_1)V_2'(M_2) - M_2V_2'(M_2) - M_1V_1'(M_1) + M_1M_2) \rangle \\
&\quad - \frac{t^2}{s(x)\tilde{s}(y)}
\end{aligned}
\tag{2-39}$$

and in the right hand side of eq.2-27, we have:

$$\begin{aligned}
L(x) &= \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} \frac{V_2'(Y(x)) - V_2'(M_2)}{Y(x) - M_2} \right\rangle_c \\
&\quad + \frac{1}{\tilde{s}(Y(x))} \left\langle \text{Tr} \frac{1}{x - M_1} \text{Tr} \frac{1}{x - M_1} (V_2'(M_2) - M_1) \right\rangle_c
\end{aligned}
\tag{2-40}$$

3 Large N limit, algebraic curve

In the large N limit, eq.2-27 reduces to an algebraic equation:

$$E(x, Y(x)) = 0
\tag{3-1}$$

One should notice that in the large N limit, there is the factorization [6]: $\langle \text{Tr Tr} \rangle = \langle \text{Tr} \rangle \langle \text{Tr} \rangle$, so that x and y play symmetric roles [17].

We see that $E(x, y)$ has poles only at the poles of V_1' and zeroes of s in x and at the poles of V_2' and the zeroes of $\tilde{s}$ in y .

3.1 Behaviour near hard edges

near an hard edge $x \rightarrow X_i$, such that $s(X_i) = 0$, we have:

$$\begin{aligned} Y^2(x) \sim_{x \rightarrow X_i} & -\frac{1}{s(x)} \frac{t}{N} \left\langle \text{Tr} \frac{s(x) - s(M_1)}{x - M_1} (V_1'(M_1) - M_2) \right\rangle \\ & + \frac{1}{s(x)} \sum_{r=2}^{K_1} \sum_{l=1}^{r-1} s_r \frac{t}{N} \langle \text{Tr} M_1^{r-1-l} \rangle \frac{t}{N} \left\langle \text{Tr} \frac{x^l - M_1^l}{x - M_1} \right\rangle \\ & + \text{finite} \end{aligned} \quad (3-2)$$

Thus:

$$\begin{aligned} \text{Res}_{X_i} Y^2(x) dx &= \frac{1}{s'(X_i)} \frac{t}{N} \left\langle \text{Tr} \frac{s(M_1)}{X_i - M_1} (V_1'(M_1) - M_2) \right\rangle \\ & + \frac{1}{s'(X_i)} \sum_{r=2}^{K_1} \sum_{l=1}^{r-1} s_r \frac{t}{N} \langle \text{Tr} M_1^{r-1-l} \rangle \frac{t}{N} \left\langle \text{Tr} \frac{X_i^l - M_1^l}{X_i - M_1} \right\rangle \end{aligned} \quad (3-3)$$

Hard edges are at the same time poles of $Y(x)$, and zeroes of dx , so that $Y(x)dx$ is finite.

3.2 Behaviour near poles of the potential

Near a finite pole ξ of $V_1'(x)$, we have:

$$Y(x) \underset{x \rightarrow \xi}{\sim} V_1'(x) - W(\xi) + O(x - \xi) \quad (3-4)$$

thus

$$\text{Res}_{\xi} Y(x) dx = \text{Res}_{\xi} V_1'(x) dx \quad (3-5)$$

Near a pole at ∞ , we have:

$$Y(x) \underset{x \rightarrow \infty}{\sim} V_1'(x) - \frac{t}{x} + O(1/x^2) \quad (3-6)$$

thus

$$\text{Res}_{\xi} Y(x) dx = t \quad (3-7)$$

3.3 Determination of the algebraic equation $E(x, y)$

So far, we know that $E(x, y)$ is a rational function of x and y , we know its form, and its behaviour near poles, but most of the coefficients are not determined by the loop equations.

The remaining coefficients of $E(x, y)$, as usual, are determined by extra requirements, which depend on how the matrix model is defined, i.e. on the purpose for which we introduce the matrix model.

The two most frequent definitions of the matrix model are (we mainly follow the presentation of [15]):

3.3.1 Case of the convergent matrix model

In this case, the matrix model is defined by the convergent integral eq.1-7. For generic potentials and hard edges, $\frac{1}{N^2} \ln Z$ has a large N limit F_0 , but has no $1/N$ series expansion. The resolvent, and thus the function $Y(x)$ also, has a large N limit, but no $1/N$ expansion. This fact can be understood from the work of [4].

The large N limit of $Y(x)$ obeys an algebraic equation $E(x, Y(x)) = 0$.

Now, consider an arbitrary $\tilde{E}(x, y)$ satisfying the correct behaviours near poles. It gives a function $\tilde{Y}(x)$, and from it one can compute the free energy F_0 (see the formula in [2]). The $Y(x)$ which is the large N limit of the resolvent is the one for which $\Re(F_0)$ is minimal (see [15, 4]). That implies in particular that for any contour $\mathcal{C}$ on the algebraic curve $E(x, y) = 0$, one has:

$$\forall \mathcal{C} \ , \quad \Re \oint_{\mathcal{C}} Y(x) dx = 0 \quad (3-8)$$

On an algebraic curve of the type $E(x, y) = 0$, there are at most $2 \times \text{genus}$ independent irreducible cycles, (plus contours around poles), and one can check that the number of constraints of type eq.3-8, exactly matches the number of coefficients of $E(x, y)$ not determined by the pole behaviours. Thus, condition eq.3-8, determines $E(x, y)$ completely. In case there are several solutions, one determines a unique one, by choosing the absolute minimum of F_0 .

3.3.2 Case of the formal matrix model

The formal matrix model can be defined in a combinatoric way, as a formal series, generating discrete surfaces. The formal expansion is obtained by expanding the matrix integral eq.1-7 with the Feynman graph technics [5, 16, 6].

In that model, $\ln Z$ has a $1/N$ expansion by definition, as well as the resolvent, and all expectation values. The resolvent of the formal matrix model, is thus obtained by solving the loop equation eq.2-27, order by order in $1/N^2$.

That model, in addition to the potentials and hard edges, depends on a “vacuum” around which the Feynman expansion is performed. This vacuum is characterized by a set of “filling fractions”, as follows (see [15] for more details): The potential $V_1(x) + V_2(y) - xy$ has a certain number of extrema, which are given by the algebraic equation:

$$V_1'(x) = y, \quad V_2'(y) = x \quad (3-9)$$

i.e.

$$V_2'(V_1'(x)) = x \quad (3-10)$$

Let us call K the degree of that algebraic equation, i.e. the number of its solutions:

$$(\bar{x}_1, \bar{y}_1), \dots, (\bar{x}_K, \bar{y}_K) \quad (3-11)$$

The eigenvalues of matrices M_1, M_2 which extremize $\text{Tr}(V_1(M_1) + V_2(M_2) - M_1 M_2)$ must be among the K solutions described above, or can be trapped on contours stopping at hard edges. The filling fractions $(\epsilon_1, \dots, \epsilon_{K+K_1+K_2})$ represent the number of eigenvalues equal to each solutions of eq.3-10, i.e. there are $N\epsilon_1$ eigenvalues of M_1 equal to $\bar{x}_1, \dots$. One must have:

$$\sum_{i=1}^{K+K_1+K_2} \epsilon_i = 1 \quad (3-12)$$

The average number of eigenvalues of M_1 in the vicinity of a point $(\bar{x}, \bar{y})$, is a contour integral of the resolvent, along a contour which surrounds $(\bar{x}, \bar{y})$:

$$2i\pi N\epsilon_i = - \oint_{\mathcal{A}_i} W(x)dx = \oint_{\mathcal{A}_i} Y(x)dx \quad (3-13)$$

and it can be non-zero only on irreducible cycles of the algebraic curve.

Thus, in the Formal matrix model, a set of irreducible cycle contour integrals are fixed. One can verify that the number of filling fractions matches exactly the number of coefficients of $E(x, y)$ not determined by the pole behaviours, Thus, eq.3-13 is sufficient to determine completely the rational function $E(x, y)$.

4 Conclusion

The purpose of this article was to write down the loop-equations for the so-called semiclassical 2-matrix model. We find, that the loop equation becomes an algebraic equation $E(x, y) = 0$ in the large N limit, with poles at the poles of the potentials, and at the hard edges. The hard edges are such that the resolvent has a simple pole, and dx has a zero, so that the differential form ydx is regular.

The loop equation determines the form of the algebraic equation $E(x, y)$, but does not determine its coefficients. The coefficients are determined by extra assumptions, related to which definition of the matrix model is considered. In the formal matrix model, the $\mathcal{A}$ cycle integrals of ydx are fixed parameters of the model, and that determines $E(x, y)$ completely. In the convergent matrix model, the real parts of both $\mathcal{A}$ and $\mathcal{B}$ cycle integrals of ydx must vanish, and that determines $E(x, y)$ completely.

Let us also remark that the function eq.2-27 $E(x, y)$ is unchanged under the exchange $x \leftrightarrow y$, $V_1 \leftrightarrow V_2$, $s \leftrightarrow \tilde{s}$, which is the generalization of Matytsin's duality property [17]:

$$X(Y(x)) = x \tag{4-1}$$

The consequences of that algebraic equation, can then be studied. This is done for instance in [3]. One can also expect to generalize the works of [9], or [7], or [8], i.e. the computation of all correlation functions and their $1/N^2$ expansion, and further, compute the expansion of the free energy [10, 11].

Aknowledgements

The author wants to thank Marco Bertola for numerous discussions, and for the motivation to compute those loop equations. This work was partly supported by the european network ENIGMA (MRTN-CT-2004-5652).

References

- [1] M. Bertola, "Bilinear semi-classical moment functionals and their integral representation", J. App. Th. **121** 71–99 (2003).
- [2] M. Bertola, "Free Energy of the Two-Matrix Model/dToda Tau-Function", preprint CRM-2921 (2003), hep-th/0306184.
- [3] M. Bertola, "Higher Order Observables of the Two-Matrix Model with semiclassical potentials", preprint CRM-???? (2005)
- [4] G. Bonnet, F. David, B. Eynard, "Breakdown of universality in multi-cut matrix models", *J.Phys. A* **33** 6739-6768 (2000).
- [5] E. Brezin, C. Itzykson, G. Parisi, and J. Zuber, *Comm. Math. Phys.* **59**, 35 (1978).

- [6] P. Di Francesco, P. Ginsparg, J. Zinn-Justin, “2D Gravity and Random Matrices”, *Phys. Rep.* **254**, 1 (1995).
- [7] B. Eynard, “Topological expansion for the 1-hermitian matrix model correlation functions”, JHEP/024A/0904, xxx, hep-th/0407261.
- [8] B. Eynard, N. Orantin, “Topological expansion for the 2- matrix model correlation functions”, in preparation.
- [9] B. Eynard, N. Orantin, “Mixed correlation functions in the 2-matrix model, and the Bethe ansatz”, preprint SPHT05/037.
- [10] B. Eynard, “Large N expansion of the 2-matrix model”, *JHEP* **01** (2003) 051, hep-th/0210047.
- [11] B. Eynard, “Large N expansion of the 2-matrix model, multicut case”, preprint SPHT03/106, ccscd-00000521, math-ph/0307052.
- [12] B. Eynard, “Eigenvalue distribution of large random matrices, from one matrix to several coupled matrices” *Nucl. Phys. B* **506**, 633 (1997), cond-mat/9707005.
- [13] B. Eynard, “Correlation functions of eigenvalues of multi-matrix models, and the limit of a time dependent matrix”, *J. Phys. A: Math. Gen.* **31**, 8081 (1998), cond-mat/9801075.
- [14] B. Eynard, “Master loop equations, free energy and correlations for the chain of matrices”, JHEP11(2003)018, xxx, hep-th/0309036, ccscd-00000572.
- [15] B. Eynard, “Polynômes biorthogonaux, problème de Riemann-Hilbert et géométrie algébrique”, Habilitation à diriger les recherches, université Paris VII, (2005).
- [16] V.A. Kazakov, “Ising model on a dynamical planar random lattice: exact solution”, *Phys Lett.* **A119**, 140-144 (1986).
- [17] A. Matytsin, “on the large N limit of the Itzykson Zuber Integral”, *Nuc. Phys.* **B411**, 805 (1994), hep-th/9306077.
- [18] M.L. Mehta, *Random Matrices*, 2nd edition, (Academic Press, New York, 1991).
- [19] M. Staudacher, “ Combinatorial solution of the 2-matrix model”, *Phys. Lett.* **B305** (1993) 332-338.