



HAL
open science

Bidendriform bialgebras, trees, and free quasi-symmetric functions

Loïc Foissy

► **To cite this version:**

Loïc Foissy. Bidendriform bialgebras, trees, and free quasi-symmetric functions. Journal of Pure and Applied Algebra, 2007. hal-00004902

HAL Id: hal-00004902

<https://hal.science/hal-00004902>

Submitted on 11 May 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bidendriform bialgebras, trees, and free quasi-symmetric functions

L. Foissy

*Laboratoire de Mathématiques - UMR6056, Université de Reims
Moulin de la Housse - BP 1039 - 51687 REIMS Cedex 2, France **

ABSTRACT: we introduce bidendriform bialgebras, which are bialgebras such that both product and coproduct can be split into two parts satisfying good compatibilities. For example, the Malvenuto-Reutenauer Hopf algebra and the non-commutative Connes-Kreimer Hopf algebras of planar decorated rooted trees are bidendriform bialgebras. We prove that all connected bidendriform bialgebras are generated by their primitive elements as a dendriform algebra (bidendriform Milnor-Moore theorem) and then is isomorphic to a Connes-Kreimer Hopf algebra. As a corollary, the Hopf algebra of Malvenuto-Reutenauer is isomorphic to the Connes-Kreimer Hopf algebra of planar rooted trees decorated by a certain set. We deduce that the Lie algebra of its primitive elements is free in characteristic zero (G. Duchamp, F. Hivert and J.-Y. Thibon conjecture).

RESUME : nous introduisons les bigèbres bidendriformes, qui sont des bigèbres dont le produit et le coproduit peuvent être scindés en deux avec de bonnes compatibilités. Par exemple, l'algèbre de Hopf de Malvenuto-Reutenauer et les algèbres de Hopf non-commutative de Connes-Kreimer sur les arbres plans enracinés décorés sont des bigèbres bidendriformes. Nous montrons que toute bigèbre bidendriforme connexe est engendrée par ses éléments totalement primitifs comme algèbre dendriforme (version bidendriforme du théorème de Milnor-Moore) et qu'elle est alors isomorphe à une algèbre de Hopf de Connes-Kreimer. En conséquence, l'algèbre de Hopf de Malvenuto-Reutenauer est isomorphe à l'algèbre de Connes-Kreimer des arbres plans enracinés décorés par un certain ensemble. On en déduit que l'algèbre de Lie de ses éléments primitifs est libre en caractéristique zéro (conjecture de G. Duchamp, F. Hivert et J.-Y. Thibon).

Contents

1 Dendriform and codendriform bialgebras	3
1.1 Dendriform algebras and coalgebras	3
1.2 Dendriform and codendriform bialgebras	4
1.3 Free dendriform algebras	5
2 Bidendriform bialgebras	8
2.1 Definition	8
2.2 Totally primitive elements	9
2.3 Connected bidendriform bialgebras	12
2.4 Projections on $Prim_{\leftarrow}(A)$ and $Prim_{tot}(A)$	13

*e-mail : loic.foissy@univ-reims.fr

3	Tensor product and dendriform modules	15
3.1	Tensor product of dendriform algebras	15
3.2	Dendriform modules	17
3.3	Bidendriform structure on $\mathcal{A}^{\mathcal{D}}$	18
3.4	Totally primitive elements of $\mathcal{A}^{\mathcal{D}}$ and universal property	20
4	Application to the Hopf algebra \mathbf{FQSym}	21
4.1	Recalls	21
4.2	Bidendriform structure on \mathbf{FQSym}	21
5	Bidendriform pairings	24
5.1	Definition	24
5.2	Hopf dual of a bidendriform bialgebra	24
5.3	Bidendriform pairing on a connected graded bidendriform bialgebra	27
5.4	Application to $\mathcal{H}^{\mathcal{D}}$	29

Introduction

The Hopf algebra \mathbf{FQSym} of Malvenuto-Reutenauer, also called Hopf algebra of free quasi-symmetric functions ([5, 16, 17, 18]) has certain interesting properties. For example, it is known that it free as an algebra and cofree as a coalgebra; it has a non-degenerate Hopf pairing; it can be given a structure of dendriform algebra. The Hopf algebras $\mathcal{H}^{\mathcal{D}}$ of planar rooted trees, introduced in [7, 8, 9] have a lot of similar properties: they are free as algebras and cofree as coalgebras; they have a non-degenerate Hopf pairing (although not so explicit as the Malvenuto-Reutenauer algebra's); they also are dendriform algebras. So a natural question is: are these two objects isomorphic? More precisely, is there a set \mathcal{D} of decorations such that $\mathcal{H}^{\mathcal{D}}$ is isomorphic to \mathbf{FQSym} ?

In order to answer positively this question, we study more in details dendriform algebras. This notion is introduced by Loday and Ronco in [15] and is studied in [2, 16, 17, 22, 21]. A dendriform algebra A is a (non unitary) associative algebra, such that the product can be split into two parts \prec and \succ (left and right products), with good compatibilities (which mean that (A, \prec, \succ) is a bimodule over itself). The notion of dendriform bialgebra or Hopf dendriform algebra is also given. These are dendriform algebra with a coassociative coproduct $\tilde{\Delta}$, satisfying good relations with \prec and \succ . We introduce here the notion of bidendriform bialgebra (section 2): a bidendriform bialgebra A is a dendriform bialgebra such that the coproduct $\tilde{\Delta}$ can be split into two parts Δ_{\prec} and Δ_{\succ} (left and right coproducts), such that $(A, \Delta_{\succ}, \Delta_{\prec})$ is a bimodule over itself. There are also compatibilities between the left and right coproducts and the left and right products. As this set of axioms is self-dual, the dual of a finite-dimensional bidendriform bialgebra is also a bidendriform bialgebra. An example of bidendriform bialgebra is \mathbf{FQSym} (section 4), or, more precisely, its augmentation ideal: as \mathbf{FQSym} is a dendriform algebra and it is self dual as a Hopf algebra, we can also split the coproduct into two parts. Fortunately, the left and right coproducts defined in this way satisfy the wanted compatibilities with the left and right products.

We would like to give $\mathcal{H}^{\mathcal{D}}$ a structure of bidendriform algebra too. The method used for \mathbf{FQSym} fails here: the left and right coproducts defined by duality, denoted here by Δ'_{\prec} and Δ'_{\succ} , do not satisfy the compatibilities with the left and right products. Hence, we have to proceed in a different way. For this, we consider the category of dendriform algebras and give it a tensor product $\overline{\otimes}$ (section 3). This tensor product is a little bit different of the usual one, as dendriform algebras are not unitary objects: we have to add a copy of both algebras to their tensor products. We also define the notion of dendriform module over a dendriform algebra. Then the notions of dendriform bialgebra and bidendriform bialgebra become more clear: the coproduct of a dendriform bialgebra A has to be a morphism of dendriform algebras from A

to $A\overline{\otimes}A$; the left and right coproducts of a bidendriform bialgebra A have to be morphism of dendriform modules from A to $A\overline{\otimes}A$. Now, as the dendriform algebra $\mathcal{H}^{\mathcal{D}}$ is freely generated by the elements \cdot_d as a dendriform algebra, it is possible to define a unique structure of bidendriform bialgebra over $\mathcal{H}^{\mathcal{D}}$ (or, more exactly, on its augmentation ideal $\mathcal{A}^{\mathcal{D}}$) by $\Delta_{\prec}(\cdot_d) = \Delta_{\succ}(\cdot_d) = 0$ (theorem 31).

Let us now study more precisely the notion of bidendriform bialgebra. For a given bidendriform bialgebra A , we consider totally primitive elements of A , that is to say elements which vanish under both Δ_{\prec} and Δ_{\succ} . We say that A is connected if, for every element a , the various iterated coproducts all vanish on a for a great enough rank. Then, if A is connected, we prove that A is generated as a dendriform algebra by its totally primitive elements (theorem 21). This theorem can be seen as a bidendriform version of the Milnor-Moore theorem ([20]), which says that a cocommutative, connected Hopf algebra is generated by its primitive elements (in characteristic zero).

To precise this result, we consider the bidendriform bialgebra $\mathcal{A}^{\mathcal{D}}$. We prove that its space of totally primitive elements is reduced to the space of its generators: the elements \cdot_d . In other terms, the triple $(Dend, coDend, Vect)$ is a good triple of operads, with the language of [14]. This implies that every connected bidendriform bialgebra is freely generated by its totally primitive elements, so is isomorphic to a $\mathcal{H}^{\mathcal{D}}$ for a well chosen \mathcal{D} (theorem 39). We apply this result to **FQSym**: then, for a certain \mathcal{D} , **FQSym** is isomorphic to $\mathcal{H}^{\mathcal{D}}$ as a bidendriform bialgebra, and hence as a Hopf algebra. This allows us to answer a conjecture of [5]: if the characteristic of the base field is zero, then the Lie algebra of primitive elements of **FQSym** is free (corollary 40), as we already proved this result for $\mathcal{H}^{\mathcal{D}}$ in [8].

As a connected bidendriform bialgebra is isomorphic to a certain $\mathcal{H}^{\mathcal{D}}$, it is self-dual. We define the notion of bidendriform pairing; for example, the pairing of [5] on **FQSym** is a bidendriform pairing, whereas the pairing of [7] on $\mathcal{H}^{\mathcal{D}}$ is not. We construct another pairing on $\mathcal{H}^{\mathcal{D}}$, which is non-degenerate, symmetric and bidendriform. This pairing can be studied in the same way as in [7]: we give an inductive way to compute it and a combinatorial interpretation of this pairing.

Thanks. I am grateful to Ralf Holtkamp for suggestions which greatly improve section 3.

Notation. K is a commutative field of any characteristic.

1 Dendriform and codendriform bialgebras

1.1 Dendriform algebras and coalgebras

Definition 1 (See [2, 15, 16, 17, 21, 22]). A dendriform algebra is a family (A, \prec, \succ) such that:

1. A is a K -vector space and:

$$\prec: \begin{cases} A \otimes A & \longrightarrow & A \\ a \otimes b & \longrightarrow & a \prec b, \end{cases} \quad \Bigg| \quad \succ: \begin{cases} A \otimes A & \longrightarrow & A \\ a \otimes b & \longrightarrow & a \succ b. \end{cases}$$

2. For all $a, b, c \in A$:

$$(a \prec b) \prec c = a \prec (b \prec c + b \succ c), \tag{1}$$

$$(a \succ b) \prec c = a \succ (b \prec c), \tag{2}$$

$$(a \prec b + a \succ b) \succ c = a \succ (b \succ c). \tag{3}$$

Remark. If A is a dendriform algebra, we put:

$$m: \begin{cases} A \otimes A & \longrightarrow & A \\ a \otimes b & \longrightarrow & ab = a \prec b + a \succ b. \end{cases}$$

Then (1) + (2) + (3) is equivalent to the fact that m is associative. Hence, a dendriform algebra is a special (non unitary) associative algebra.

By duality, we obtain the notion of dendriform coalgebra:

Definition 2 A dendriform coalgebra is a family $(C, \Delta_{\prec}, \Delta_{\succ})$ such that:

1. C is a K -vector space and:

$$\Delta_{\prec} : \begin{cases} C & \longrightarrow C \otimes C \\ a & \longrightarrow \Delta_{\prec}(a) = a'_{\prec} \otimes a''_{\prec}, \end{cases} \quad \Bigg| \quad \Delta_{\succ} : \begin{cases} C & \longrightarrow C \otimes C \\ a & \longrightarrow \Delta_{\succ}(a) = a'_{\succ} \otimes a''_{\succ}. \end{cases}$$

2. for all $a \in C$:

$$(\Delta_{\prec} \otimes Id) \circ \Delta_{\prec}(a) = (Id \otimes \Delta_{\prec} + Id \otimes \Delta_{\succ}) \circ \Delta_{\prec}(a), \quad (4)$$

$$(\Delta_{\succ} \otimes Id) \circ \Delta_{\prec}(a) = (Id \otimes \Delta_{\prec}) \circ \Delta_{\succ}(a), \quad (5)$$

$$(\Delta_{\prec} \otimes Id + \Delta_{\succ} \otimes Id) \circ \Delta_{\succ}(a) = (Id \otimes \Delta_{\succ}) \circ \Delta_{\succ}(a). \quad (6)$$

Remarks.

1. If C is a dendriform coalgebra, we put:

$$\tilde{\Delta} : \begin{cases} C & \longrightarrow C \otimes C \\ a & \longrightarrow \tilde{\Delta}(a) = \Delta_{\prec}(a) + \Delta_{\succ}(a) = a' \otimes a''. \end{cases}$$

Then (4) + (5) + (6) is equivalent to the fact that $\tilde{\Delta}$ is coassociative. Hence, a dendriform coalgebra is a special (non counitary) coassociative coalgebra.

2. If $A = \bigoplus A_n$ is a \mathbb{N} -graded dendriform algebra, such that its homogeneous parts are finite-dimensional, then $(A^{*g}, \prec^*, \succ^*)$ is a \mathbb{N} -graded dendriform coalgebra (A^{*g} is the graded dual of A , that is to say $A^{*g} = \bigoplus A_n^* \subseteq A^*$).
3. In the same way, if C is a \mathbb{N} -graded dendriform coalgebra, such that its homogeneous parts are finite-dimensional, then $(C^{*g}, \Delta_{\prec}^*, \Delta_{\succ}^*)$ is a \mathbb{N} -graded dendriform algebra. (In fact, for any dendriform coalgebra C , the whole linear dual C^* is a dendriform algebra).

Definition 3 Let A be a dendriform coalgebra. We put:

$$\begin{aligned} Prim(A) &= \{a \in A / \tilde{\Delta}(a) = 0\}, \\ Prim_{\prec}(A) &= \{a \in A / \Delta_{\prec}(a) = 0\}, \\ Prim_{\succ}(A) &= \{a \in A / \Delta_{\succ}(a) = 0\}, \\ Prim_{tot}(A) &= Prim_{\succ}(A) \cap Prim_{\prec}(A) = \{a \in A / \tilde{\Delta}(a) = \Delta_{\prec}(a) = \Delta_{\succ}(a) = 0\}. \end{aligned}$$

1.2 Dendriform and codendriform bialgebras

Definition 4 (See [2, 16, 17, 21, 22]). A dendriform bialgebra is a family $(A, \prec, \succ, \tilde{\Delta})$ such that:

1. (A, \prec, \succ) is a dendriform algebra.
2. $(A, \tilde{\Delta})$ is a coassociative (non counitary) coalgebra.
3. The following compatibilities are satisfied: for all $a, b \in A$,

$$\tilde{\Delta}(a \prec b) = a'b' \otimes a'' \prec b'' + a' \otimes a'' \prec b + a'b \otimes a'' + b' \otimes a \prec b'' + b \otimes a, \quad (7)$$

$$\tilde{\Delta}(a \succ b) = a'b' \otimes a'' \succ b'' + a' \otimes a'' \succ b + ab' \otimes b'' + b' \otimes a \succ b'' + a \otimes b. \quad (8)$$

Remarks.

- (7) + (8) is equivalent to: for all $a \in A$,

$$\tilde{\Delta}(ab) = a'b' \otimes a''b'' + a' \otimes a''b + ab' \otimes b'' + a'b \otimes a'' + b' \otimes ab'' + a \otimes b + b \otimes a. \quad (9)$$

If A is a dendriform bialgebra, we put $\bar{A} = A \oplus K$, which is given a structure of associative algebra and coassociative coalgebra in the following way: for all $a, b \in A$,

$$\begin{aligned} 1.a = a, \quad a.1 = a, \quad 1.1 = 1, \quad a.b = ab \text{ (product in } A); \\ \Delta(1) = 1 \otimes 1, \quad \Delta(a) = 1 \otimes a + a \otimes 1 + \tilde{\Delta}(a). \end{aligned}$$

Then (9) means that \bar{A} is a bialgebra. A dendriform bialgebra is then the augmentation ideal of a special bialgebra.

- Another interpretation of (7) and (8) will be given in section 3.

By duality, we define the notion of codendriform bialgebra:

Definition 5 A codendriform bialgebra is a family $(A, m, \Delta_{\prec}, \Delta_{\succ})$ such that:

- $(A, \Delta_{\prec}, \Delta_{\succ})$ is a dendriform coalgebra.
- (A, m) is an associative (non unitary) algebra.
- The following compatibilities are satisfied: for all $a, b \in A$,

$$\Delta_{\succ}(ab) = a'b'_{\succ} \otimes a''b''_{\succ} + a' \otimes a''b + ab'_{\prec} \otimes b''_{\prec} + b'_{\prec} \otimes ab''_{\prec} + a \otimes b, \quad (10)$$

$$\Delta_{\prec}(ab) = a'b'_{\prec} \otimes a''b''_{\prec} + a'b \otimes a'' + ab'_{\succ} \otimes b''_{\succ} + b'_{\succ} \otimes ab''_{\succ} + b \otimes a. \quad (11)$$

Remarks.

- (10)+(11) is equivalent to (9). Hence, if A is a codendriform bialgebra, as before $\bar{A} = A \oplus K$ is given a structure of bialgebra. A codendriform bialgebra is then the augmentation ideal of a special bialgebra.
- If A is a \mathbb{N} -graded codendriform bialgebra, such that its homogeneous parts are finite-dimensional, then $(A^{*g}, \Delta_{\prec}^*, \Delta_{\succ}^*, m^*)$ is a \mathbb{N} -graded dendriform bialgebra.
- In the same way, if A is a \mathbb{N} -graded dendriform bialgebra, such that its homogeneous parts are finite-dimensional, then $(A^{*g}, \tilde{\Delta}^*, \prec^*, \succ^*)$ is a \mathbb{N} -graded codendriform bialgebra.

1.3 Free dendriform algebras

Let us recall here the construction of the Connes-Kreimer Hopf algebra of planar decorated rooted trees (See [7, 8, 9] for more details). It is a non-commutative version of the Connes-Kreimer Hopf algebra of rooted trees for Renormalisation ([4, 11, 12, 13]).

Definition 6

- A *rooted tree* t is a finite graph, without loops, with a special vertex called *root* of t . A *planar rooted tree* t is a rooted tree with an imbedding in the plane. The *weight* of t is the number of its vertices. the set of planar rooted trees will be denoted by \mathbb{T} .
- Let \mathcal{D} be a nonempty set. A planar rooted tree decorated by \mathcal{D} is a planar tree with an application from the set of its vertices into \mathcal{D} . The set of planar rooted trees decorated by \mathcal{D} will be denoted by $\mathbb{T}^{\mathcal{D}}$.

Example. If $a, b, c, d, e \in \mathcal{D}$:

$$\begin{aligned} \Delta \left(\cdot_a \mathbb{Y}_b^c \cdot^d \right) &= \cdot_a \mathbb{Y}_b^c \otimes 1 + 1 \otimes \cdot_a \mathbb{Y}_b^c + \cdot_a \otimes \mathbb{Y}_b^c + \mathbb{Y}_b^c \otimes \cdot_a + \cdot_a \cdot e \cdot d \otimes \mathbb{1}_b^c + \cdot e \cdot d \otimes \cdot_a \mathbb{1}_b^c \\ &+ \cdot_a \mathbb{V}_c^d \otimes \cdot_b + \mathbb{V}_c^d \otimes \cdot_a \cdot b + \cdot_a \cdot e \otimes \mathbb{1}_b^d + \cdot_e \otimes \cdot_a \mathbb{1}_b^d + \cdot_a \cdot d \otimes \mathbb{1}_b^e + \cdot_d \otimes \cdot_a \mathbb{1}_b^e. \end{aligned}$$

The counit ε is given by:

$$\varepsilon : \begin{cases} \mathcal{H}^{\mathcal{D}} & \longrightarrow K \\ F \in \mathbb{F}^{\mathcal{D}} & \longrightarrow \delta_{F,1}. \end{cases}$$

We proved in [8] that $\mathcal{H}^{\mathcal{D}}$ is isomorphic to the free dendriform algebra generated by \mathcal{D} , which is described in [16, 21] in terms of planar binary trees. So the augmentation ideal $\mathcal{A}^{\mathcal{D}}$ of $\mathcal{H}^{\mathcal{D}}$ inherits a structure of dendriform algebra, also described in [8] with the help of another basis of $\mathcal{H}^{\mathcal{D}}$, introduced by duality. Hence, $\mathcal{A}^{\mathcal{D}}$ is freely generated by the \cdot_d 's, $d \in \mathcal{D}$, as a dendriform algebra. Here is an example of a computation of a product \prec in $\mathcal{A}^{\mathcal{D}}$. For all $x \in \mathcal{A}^{\mathcal{D}}$,

$$\cdot_d \prec x = B_d^+(x),$$

where $B_d^+ : \mathcal{H}^{\mathcal{D}} \longrightarrow \mathcal{H}^{\mathcal{D}}$ is the linear application which send a forest $t_1 \dots t_n$ to the planar decorated tree obtained by grafting t_1, \dots, t_n on a common root decorated by d . (This comes from the description of \prec in terms of graftings in [8] and proposition 36 of [7]).

As $\mathcal{H}^{\mathcal{D}}$ is self-dual ([7]), $\mathcal{A}^{\mathcal{D}}$ is given a structure of codendriform bialgebra. The description in [8] of the left and right products in the dual basis of forests allows us to describe this structure with the following definition:

Definition 7 Let $F = t_1 \dots t_n \in \mathbb{F}^{\mathcal{D}}$, $F \neq 1$. The set $\mathcal{Adm}'_{\prec}(F)$ is the set of cuts $(c_1, \dots, c_n) \in \mathcal{Adm}(F)$ such that c_n is the total cut of t_n if F is not a single tree, and \emptyset otherwise. The set $\mathcal{Adm}'_{\succ}(F)$ is $\mathcal{Adm}(F) - \mathcal{Adm}'_{\prec}(F)$.

The dendriform coalgebra structure of $\mathcal{A}^{\mathcal{D}}$ is then given in the following way: For all $F \in \mathbb{F}^{\mathcal{D}} - \{1\}$,

$$\Delta'_{\prec}(F) = \sum_{c \in \mathcal{Adm}'_{\prec}(F)} P^c(F) \otimes R^c(F), \quad \Delta'_{\succ}(F) = \sum_{c \in \mathcal{Adm}'_{\succ}(F)} P^c(F) \otimes R^c(F).$$

The product of $\mathcal{A}^{\mathcal{D}}$ is the product induced by the product of $\mathcal{H}^{\mathcal{D}}$.

Examples.

1. If $t \in \mathbb{T}^{\mathcal{D}}$, $\Delta'_{\prec}(t) = 0$.
2. If $a, b, c, d, e \in \mathcal{D}$.

$$\begin{aligned} \Delta'_{\prec} \left(\cdot_a \mathbb{Y}_b^c \cdot^d \right) &= \mathbb{Y}_b^c \otimes \cdot_a, \\ \Delta'_{\succ} \left(\cdot_a \mathbb{Y}_b^c \cdot^d \right) &= \cdot_a \otimes \mathbb{Y}_b^c + \cdot_a \cdot e \cdot d \otimes \mathbb{1}_b^c + \cdot_e \cdot d \otimes \cdot_a \mathbb{1}_b^c + \cdot_a \mathbb{V}_c^d \otimes \cdot_b + \mathbb{V}_c^d \otimes \cdot_a \cdot b \\ &+ \cdot_a \cdot e \otimes \mathbb{1}_b^d + \cdot_e \otimes \cdot_a \mathbb{1}_b^d + \cdot_a \cdot d \otimes \mathbb{1}_b^e + \cdot_d \otimes \cdot_a \mathbb{1}_b^e. \end{aligned}$$

Moreover, $\mathcal{H}^{\mathcal{D}}$ can be graded. A set \mathcal{D} is said to be graded when it is given an application $|\cdot| : \mathcal{D} \longrightarrow \mathbb{N}$. We denote $\mathcal{D}_n = \{d \in \mathcal{D} / |d| = n\}$. We then put, for all $F \in \mathbb{F}^{\mathcal{D}}$:

$$|F| = \sum_{s \in \text{vert}(F)} |\text{decoration of } s|,$$

where $vert(F)$ is the set of vertices of F . Then $\mathcal{H}^{\mathcal{D}}$ is given a graded Hopf algebra structure and $\mathcal{A}^{\mathcal{D}}$ is given a graded (co)dendriform bialgebra structure by putting, for all $F \in \mathbb{F}^{\mathcal{D}}$, F homogeneous of degree $|F|$. When \mathcal{D}_0 is empty and \mathcal{D}_n is finite for all n , then the homogeneous parts of $\mathcal{A}^{\mathcal{D}}$ are finite-dimensional and $(\mathcal{A}^{\mathcal{D}})_0 = (0)$. Moreover, if those conditions occur, we have the following result:

Proposition 8 *We consider the following formal series:*

$$D(X) = \sum_{n=1}^{+\infty} \text{card}(\mathcal{D}_n) X^n, \quad R(X) = \sum_{n=0}^{+\infty} \dim(\mathcal{H}_n^{\mathcal{D}}) X^n.$$

$$\text{Then } R(X) = \frac{1 - \sqrt{1 - 4D(X)}}{2D(X)}.$$

Proof. Similar to the proof of theorem 75 of [6]. \square

2 Bidendriform bialgebras

2.1 Definition

We now introduce the notion of bidendriform bialgebra. A bidendriform bialgebra is both a dendriform bialgebra and a codendriform bialgebra, with some compatibilities.

Definition 9 A bidendriform bialgebra is a family $(A, \prec, \succ, \Delta_{\prec}, \Delta_{\succ})$ such that:

1. (A, \prec, \succ) is a dendriform algebra.
2. $(A, \Delta_{\prec}, \Delta_{\succ})$ is a dendriform coalgebra.
3. The following compatibilities are satisfied: for all $a, b \in A$,

$$\Delta_{\succ}(a \succ b) = a'b'_{\succ} \otimes a'' \succ b''_{\prec} + a' \otimes a'' \succ b + b'_{\prec} \otimes a \succ b''_{\prec} + ab'_{\prec} \otimes b''_{\prec} + a \otimes b, \quad (12)$$

$$\Delta_{\succ}(a \prec b) = a'b'_{\prec} \otimes a'' \prec b''_{\prec} + a' \otimes a'' \prec b + b'_{\prec} \otimes a \prec b''_{\prec}, \quad (13)$$

$$\Delta_{\prec}(a \succ b) = a'b'_{\prec} \otimes a'' \succ b''_{\prec} + ab'_{\prec} \otimes b''_{\prec} + b'_{\prec} \otimes a \succ b''_{\prec}, \quad (14)$$

$$\Delta_{\prec}(a \prec b) = a'b'_{\prec} \otimes a'' \prec b''_{\prec} + a'b \otimes a'' + b'_{\prec} \otimes a \prec b''_{\prec} + b \otimes a. \quad (15)$$

Remarks.

1. (13) + (15) and (12) + (14) are equivalent to (7) and (8), so a bidendriform bialgebra is a special dendriform bialgebra.
2. (12) + (13) and (14) + (15) are equivalent to (10) and (11), so a bidendriform bialgebra is a special codendriform bialgebra.
3. If A is a graded bidendriform bialgebra, such that its homogeneous parts are finite-dimensional, then $(A^{*g}, \Delta_{\prec}^*, \Delta_{\succ}^*, \prec^*, \succ^*)$ is also a graded bidendriform bialgebra, as the transposes of (12) and (15) are themselves, and (13) and (14) are transposes from each other.

2.2 Totally primitive elements

We here give several results which will be useful to prove theorem 21.

First part. Let A be a dendriform coalgebra. We define inductively:

$$\Delta_{\prec}^0 = Id, \quad \Delta_{\prec}^1 = \Delta_{\prec}, \quad \Delta_{\prec}^n = (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-1}) \circ \Delta_{\prec}^{n-1}.$$

For all $n \in \mathbb{N}$, $\Delta_{\prec}^n : A \longrightarrow A^{\otimes(n+1)}$.

Lemma 10 For all $n \in \mathbb{N}^*$, $i \in \{2, \dots, n\}$,

$$\begin{aligned} (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-1}) \circ \Delta_{\prec}^{n-1} &= \Delta_{\prec}^n, \\ (\underbrace{Id \otimes \dots \otimes Id}_{i-1} \otimes \tilde{\Delta} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i}) \circ \Delta_{\prec}^{n-1} &= \Delta_{\prec}^n. \end{aligned}$$

Proof. The first equality is a consequence of the definition of Δ_{\prec} . Let us show the second one. By (4):

$$\begin{aligned} & (\underbrace{Id \otimes \dots \otimes Id}_{i-1} \otimes \tilde{\Delta} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i}) \circ \Delta_{\prec}^{n-1} \\ &= (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-1}) \circ \dots \circ (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i+2}) \\ & \quad \circ (((Id \otimes \tilde{\Delta}) \circ \Delta_{\prec}) \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i}) \circ \Delta_{\prec}^{n-i} \\ &= (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-1}) \circ \dots \circ (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i+2}) \\ & \quad \circ (((\Delta_{\prec} \otimes Id) \circ \Delta_{\prec}) \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i}) \circ \Delta_{\prec}^{n-i} \\ &= (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-1}) \circ \dots \circ (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i+2}) \\ & \quad \circ (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i+1}) \circ (\Delta_{\prec} \otimes \underbrace{Id \otimes \dots \otimes Id}_{n-i}) \circ \Delta_{\prec}^{n-i} \\ &= \Delta_{\prec}^n. \quad \square \end{aligned}$$

Lemma 11 Let $a \in A$, such that $\Delta_{\prec}^n(a) = 0$. Then $\Delta_{\prec}^{n-1}(a) \in Prim_{\prec}(A) \otimes Prim(A)^{\otimes(n-1)}$.

Proof. By lemma 10, $\Delta_{\prec}^{n-1}(a)$ vanishes under $\Delta_{\prec} \otimes Id^{\otimes(n-1)}$, so belongs to $Prim_{\prec}(A) \otimes A^{\otimes(n-1)}$. Moreover, if $i \geq 2$, $\Delta_{\prec}^{n-1}(a)$ vanishes under $Id^{\otimes(i-1)} \otimes \tilde{\Delta} \otimes Id^{\otimes(n-i)}$, so belongs to $A^{\otimes(i-1)} \otimes Prim(A) \otimes A^{\otimes(n-i)}$. \square

Suppose now that A is a bidendriform bialgebra. Let $a_1, \dots, a_n \in A$. We define inductively:

$$\begin{aligned} \omega(a_1) &= a_1, \\ \omega(a_1, a_2) &= a_2 \prec a_1, \\ \omega(a_1, \dots, a_n) &= a_n \prec \omega(a_1, \dots, a_{n-1}). \end{aligned}$$

Lemma 12 Let $a_1 \in Prim_{\prec}(A)$ and $a_2, \dots, a_n \in Prim(A)$. Let $k \in \mathbb{N}$. Then:

$$\Delta_{\prec}^k(\omega(a_1, \dots, a_n)) = \sum_{1 \leq i_1 < i_2 < \dots < i_k < n} \omega(a_1, \dots, a_{i_1}) \otimes \dots \otimes \omega(a_{i_k+1}, \dots, a_n).$$

In particular:

$$\begin{cases} \Delta_{\prec}^{n-1}(\omega(a_1, \dots, a_n)) &= a_1 \otimes \dots \otimes a_n, \\ \Delta_{\prec}^k(\omega(a_1, \dots, a_n)) &= 0 \text{ if } k \geq n. \end{cases}$$

Proof. By induction on k . It is immediate for $k = 0$. Let us show the result for $k = 1$ by induction on n . It is obvious for $n = 1$. We suppose that $n \geq 2$. As $a_n \in \text{Prim}(A)$, we have, by (15):

$$\begin{aligned} \Delta_{\prec}(\omega(a_1, \dots, a_n)) &= \omega(a_1, \dots, a_{n-1})'_{\prec} \otimes a_n \prec \omega(a_1, \dots, a_{n-1})''_{\prec} + \omega(a_1, \dots, a_{n-1}) \otimes a_n \\ &= \sum_{1 \leq i < n-2} \omega(a_1, \dots, a_i) \otimes a_n \prec \omega(a_{i+1}, \dots, a_{n-1}) + \omega(a_1, \dots, a_{n-1}) \otimes a_n \\ &= \sum_{1 \leq i < n-2} \omega(a_1, \dots, a_i) \otimes \omega(a_{i+1}, \dots, a_{n-1}, a_n) + \omega(a_1, \dots, a_{n-1}) \otimes a_n \\ &= \sum_{1 \leq i < n-1} \omega(a_1, \dots, a_i) \otimes \omega(a_{i+1}, \dots, a_n). \end{aligned}$$

We suppose that the result is true at rank k . Then:

$$\begin{aligned} &\Delta_{\prec}^{k+1}(\omega(a_1, \dots, a_n)) \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_k < n} \Delta_{\prec}(\omega(a_1, \dots, a_{i_1})) \otimes \omega(a_{i_1+1}, \dots, a_{i_2}) \otimes \dots \otimes \omega(a_{i_k+1}, \dots, a_n) \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} < n} \omega(a_1, \dots, a_{i_1}) \otimes \dots \otimes \omega(a_{i_{k+1}+1}, \dots, a_n). \quad \square \end{aligned}$$

For all $a \in A$, we put $N_{\prec}(a) = \inf\{n \in \mathbb{N} / \Delta_{\prec}^{n-1}(a) = 0\} \in \mathbb{N} \cup \{+\infty\}$.

Lemma 13 *Let $a \in A$, such that $N_{\prec}(a)$ is finite. Then a can be written as a linear span of terms $\omega(a_1, \dots, a_n)$, $n \leq N_{\prec}(a)$, $a_1 \in \text{Prim}_{\prec}(A)$, $a_2, \dots, a_n \in \text{Prim}(A)$.*

Proof. By induction on $n = N_{\prec}(a)$. If $n = 0$, then $a = 0$. If $n = 1$, then $a \in \text{Prim}_{\prec}(A)$ and the result is obvious. Suppose $n \geq 2$. By lemma 11, we then put:

$$\Delta_{\prec}^{n-1}(a) = \sum a_1 \otimes \dots \otimes a_n \in \text{Prim}_{\prec}(A) \otimes \text{Prim}(A)^{\otimes(n-1)}.$$

By lemma 12, $\Delta_{\prec}^{n-1}(a - \sum \omega(a_1, \dots, a_n)) = 0$. The induction hypothesis applied to $a - \sum \omega(a_1, \dots, a_n)$ gives the result. \square

Second part. Suppose that A is a dendriform coalgebra. We define inductively:

$$\tilde{\Delta}^0 = \text{Id}, \quad \tilde{\Delta}^1 = \tilde{\Delta}, \quad \tilde{\Delta}^n = \underbrace{(\text{Id} \otimes \dots \otimes \text{Id})}_{n-1} \otimes \tilde{\Delta} \circ \tilde{\Delta}^{n-1}.$$

For all $n \in \mathbb{N}$, $\tilde{\Delta}^n : A \rightarrow A^{\otimes(n+1)}$. Note that $\tilde{\Delta}^n$ could be defined for any coassociative coalgebra $(A, \tilde{\Delta})$.

Lemma 14 *Let $a \in \text{Prim}_{\prec}(A)$. Then for all $n \in \mathbb{N}$, $\tilde{\Delta}^n(a) \in A^{\otimes n} \otimes \text{Prim}_{\prec}(A)$.*

Proof. It is obvious if $n = 0$. Suppose $n \geq 1$. By (5):

$$\begin{aligned} (\text{Id} \otimes \Delta_{\prec}) \circ \tilde{\Delta} &= (\text{Id} \otimes \Delta_{\prec}) \circ \Delta_{\prec} + (\text{Id} \otimes \Delta_{\prec}) \circ \Delta_{\succ} \\ &= (\text{Id} \otimes \Delta_{\prec}) \circ \Delta_{\prec} + (\Delta_{\succ} \otimes \text{Id}) \circ \Delta_{\prec} \\ &= (\text{Id} \otimes \Delta_{\prec} + \Delta_{\succ} \otimes \text{Id}) \circ \Delta_{\prec}. \end{aligned}$$

By coassociativity of $\tilde{\Delta}$, we have:

$$\begin{aligned} (Id^{\otimes n} \otimes \Delta_{\prec}) \circ \tilde{\Delta}^n(a) &= (\tilde{\Delta}^{n-1} \otimes Id \otimes Id) \circ (Id \otimes \Delta_{\prec}) \circ \tilde{\Delta}(a) \\ &= (\tilde{\Delta}^{n-1} \otimes Id \otimes Id) \circ (Id \otimes \Delta_{\prec} + \Delta_{\succ} \otimes Id) \circ \Delta_{\prec}(a) \\ &= 0. \end{aligned}$$

Hence, $\tilde{\Delta}^n(a) \in A^{\otimes n} \otimes Prim_{\prec}(A)$. \square

Lemma 15 *Let $a \in A$, such that $\tilde{\Delta}^n(a) = 0$. Then $\tilde{\Delta}^{n-1}(a) \in Prim(A)^{\otimes n}$. Moreover, if $a \in Prim_{\prec}(A)$, then $\tilde{\Delta}^{n-1}(a) \in Prim(A)^{\otimes(n-1)} \otimes Prim_{tot}(A)$.*

Proof. By coassociativity of $\tilde{\Delta}$, for all $i \in \{1, \dots, n\}$:

$$(Id^{\otimes(i-1)} \otimes \tilde{\Delta} \otimes Id^{\otimes(n-i)}) \circ \tilde{\Delta}^{n-1}(a) = \tilde{\Delta}^n(a) = 0,$$

so $\tilde{\Delta}^{n-1}(a) \in Prim(A)^{\otimes n}$. The second assertion comes from lemma 14. \square

Suppose now that A is a bidendriform bialgebra. Let $a_1, \dots, a_n \in A$. We define inductively:

$$\begin{aligned} \omega'(a_1) &= a_1, \\ \omega'(a_1, a_2) &= a_1 \succ a_2, \\ \omega'(a_1, \dots, a_n) &= \omega'(a_1, \dots, a_{n-1}) \succ a_n. \end{aligned}$$

Lemma 16 *Let $a_1, \dots, a_n \in Prim(A)$. Let $k \in \mathbb{N}$. Then:*

$$\tilde{\Delta}^k(\omega'(a_1, \dots, a_n)) = \sum_{1 \leq i_1 < i_2 < \dots < i_k < n} \omega'(a_1, \dots, a_{i_1}) \otimes \dots \otimes \omega'(a_{i_k+1}, \dots, a_n).$$

In particular:

$$\begin{cases} \tilde{\Delta}^{n-1}(\omega'(a_1, \dots, a_n)) &= a_1 \otimes \dots \otimes a_n, \\ \tilde{\Delta}^k(\omega'(a_1, \dots, a_n)) &= 0 \text{ if } k \geq n. \end{cases}$$

Moreover, if $a_n \in Prim_{tot}(A)$, then $\omega'(a_1, \dots, a_n) \in Prim_{\prec}(A)$.

Proof. By induction on k . It is immediate if $k = 0$. Let us show the result for $k = 1$ by induction on n . It is obvious for $n = 1$. Suppose $n \geq 2$. As $a_n \in Prim(A)$, by (8):

$$\begin{aligned} \tilde{\Delta}(\omega'(a_1, \dots, a_n)) &= \omega'(a_1, \dots, a_{n-1})' \otimes \omega'(a_1, \dots, a_{n-1})'' \succ a_n + \omega(a_1, \dots, a_{n-1}) \otimes a_n, \\ &= \sum_{1 \leq i < n-2} \omega'(a_1, \dots, a_i) \otimes \omega'(a_{i+1}, \dots, a_{n-1}) \succ a_n + \omega'(a_1, \dots, a_{n-1}) \otimes a_n \\ &= \sum_{1 \leq i < n-2} \omega'(a_1, \dots, a_i) \otimes \omega'(a_{i+1}, \dots, a_{n-1}, a_n) + \omega'(a_1, \dots, a_{n-1}) \otimes a_n \\ &= \sum_{1 \leq i < n-1} \omega'(a_1, \dots, a_i) \otimes \omega'(a_{i+1}, \dots, a_n). \end{aligned}$$

Suppose that the result is true at rank k . Then:

$$\begin{aligned} &\tilde{\Delta}^{k+1}(\omega'(a_1, \dots, a_n)) \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_k < n} \tilde{\Delta}(\omega'(a_1, \dots, a_{i_1})) \otimes \omega'(a_{i_1+1}, \dots, a_{i_2}) \otimes \dots \otimes \omega'(a_{i_k+1}, \dots, a_n) \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_{k+1} < n} \omega'(a_1, \dots, a_{i_1}) \otimes \dots \otimes \omega'(a_{i_{k+1}+1}, \dots, a_n). \end{aligned}$$

Suppose $a_n \in Prim_{\prec}(A)$. By (14), by putting $x = \omega'(a_1, \dots, a_{n-1})$, $\Delta_{\prec}(\omega'(a_1, \dots, a_n)) = \Delta_{\prec}(x \succ a_n) = 0$. So $\omega'(a_1, \dots, a_n) \in Prim_{\prec}(A)$. \square

For all $a \in A$, we put $N(a) = \inf\{n \in \mathbb{N} / \tilde{\Delta}^{n-1}(a) = 0\} \in \mathbb{N} \cup \{+\infty\}$.

Lemma 17 *Let $a \in \text{Prim}_{\prec}(A)$, such that $N(a)$ is finite. Then a can be written as a linear span of terms $\omega'(a_1, \dots, a_n)$, $n \leq N(a)$, $a_1, \dots, a_{n-1} \in \text{Prim}(A)$, $a_n \in \text{Prim}_{\text{tot}}(A)$.*

Proof. Induction on $n = N(a)$. If $n = 0$, then $a = 0$. If $n = 1$, then $a \in \text{Prim}_{\text{tot}}(A) = \text{Prim}(A) \cap \text{Prim}_{\prec}(A)$. Suppose that $n \geq 2$ and put, by lemma 15:

$$\tilde{\Delta}^{n-1}(a) = \sum a_1 \otimes \dots \otimes a_n \in \text{Prim}(A)^{\otimes(n-1)} \otimes \text{Prim}_{\text{tot}}(A).$$

We put $a' = a - \sum \omega'(a_1, \dots, a_n)$. By lemma 16, $a' \in \text{Prim}_{\prec}(A)$ and $N(a') < n$. Hence, a' satisfies the induction hypothesis, so the result is true for a . \square

2.3 Connected bidendriform bialgebras

We prove in this paragraph that if A is a connected (definition 18) bidendriform bialgebra, it is generated by its totally primitive elements (theorem 21).

Let C be a dendriform coalgebra. We define inductively:

$$\begin{aligned} \mathcal{P}_C(0) &= \{Id_C\} \subseteq \mathcal{L}(C), \\ \mathcal{P}_C(1) &= \{\Delta_{\prec}, \Delta_{\succ}\} \subseteq \mathcal{L}(C, C^{\otimes 2}), \\ \mathcal{P}_C(n) &= \left\{ \left(Id^{\otimes(i-1)} \otimes \Delta_{\prec} \otimes Id^{\otimes(n-i)} \right) \circ P / P \in \mathcal{P}_C(n-1), 1 \leq i \leq n \right\} \\ &\quad \cup \left\{ \left(Id^{\otimes(i-1)} \otimes \Delta_{\succ} \otimes Id^{\otimes(n-i)} \right) \circ P / P \in \mathcal{P}_C(n-1), 1 \leq i \leq n \right\} \subseteq \mathcal{L}(C, C^{\otimes(n+1)}). \end{aligned}$$

Definition 18

1. Let C be a dendriform coalgebra. It is said *connected* if, for all $a \in C$, there exists $n_a \in \mathbb{N}$, such that for all $P \in \mathcal{P}_C(n_a)$, $P(a) = 0$.
2. Let C be a connected dendriform coalgebra. For all $a \in C$, we put:

$$\text{deg}_p(a) = \inf\{n \in \mathbb{N} / \forall P \in \mathcal{P}_C(n), P(a) = 0\} \in \mathbb{N}.$$

3. Let C be a connected dendriform coalgebra. For all $n \in \mathbb{N}$, we put:

$$C^{\leq n} = \{a \in C / \text{deg}_p(a) \leq n\}.$$

Then $C^{\leq 0} = (0)$, $C^{\leq 1} = \text{Prim}_{\text{tot}}(C)$, and $(C^{\leq n})_{n \in \mathbb{N}}$ is a increasing filtration.

Remarks.

1. For all $n \in \mathbb{N}$, $\Delta_{\prec}^n \in \mathcal{P}_C(n)$ et $\tilde{\Delta}^n \in \text{Vect}(\mathcal{P}_C(n))$. Hence, if C is connected, then for all $a \in C$, $N_{\prec}(a)$ and $N(a)$ are finite and smaller than $\text{deg}_p(a)$.
2. Let C be a \mathbb{N} -graded dendriform coalgebra, such that $C_0 = (0)$ (the homogeneous parts of C may not be finite-dimensional). Then C is connected, as, for all $a \in C_n$, for all $P \in \mathcal{P}(n)$:

$$P(a) \in \bigoplus_{k_1 + \dots + k_{n+1} = n} C_{k_1} \otimes \dots \otimes C_{k_{n+1}} = (0).$$

Moreover, for all $a \in C$, $\text{deg}_p(a) \leq |a|$, where $|a|$ is the degree of a for the gradation of C .

Lemma 19 *Let C be a connected dendriform coalgebra. For all $n \geq 1$:*

$$\begin{aligned}\Delta_{\prec}(C^{\leq n}) &\subseteq C^{\leq n-1} \otimes C^{\leq n-1}, \\ \Delta_{\succ}(C^{\leq n}) &\subseteq C^{\leq n-1} \otimes C^{\leq n-1}, \\ \tilde{\Delta}(C^{\leq n}) &\subseteq C^{\leq n-1} \otimes C^{\leq n-1}.\end{aligned}$$

Proof. Let $a \in C^{\leq n}$. Let $P \in \mathcal{P}_C(n-1)$. Then $(P \otimes Id) \circ \Delta_{\prec} \in \mathcal{P}_C(n)$, so:

$$(P \otimes Id) \circ \Delta_{\prec}(a) = 0.$$

Hence, $\Delta_{\prec}(a) \in C^{\leq n-1} \otimes C$. In the same way, we obtain $\Delta_{\prec}(a) \in C \otimes C^{\leq n-1}$ by considering $(Id \otimes P) \circ \Delta_{\prec}$, so $\Delta_{\prec}(a) \in C^{\leq n-1} \otimes C^{\leq n-1}$. The procedure is the same for $\Delta_{\succ}(a)$, and the result for $\tilde{\Delta}(a)$ is obtained by addition. \square

The following lemma is now immediate:

Lemma 20 *Let C be a connected dendriform coalgebra and $k, n \in \mathbb{N}^*$. Then:*

$$\Delta_{\prec}^k(C^{\leq n}) \subseteq (C^{\leq n-1})^{\otimes(k+1)}, \quad \tilde{\Delta}^k(C^{\leq n}) \subseteq (C^{\leq n-1})^{\otimes(k+1)}.$$

Theorem 21 *Let A be a connected (as a dendriform coalgebra) bidendriform bialgebra. Then A is generated (as a dendriform algebra) by $Prim_{tot}(A)$.*

Proof. Let B be the dendriform subalgebra of A generated by $Prim_{tot}(A)$. Let $a \in A$. We denote $deg_p(a) = n$. Let us show that $a \in B$ by induction on n . If $n = 0$, then $a = 0 \in B$. If $n = 1$, then $a \in Prim_{tot}(A) \subseteq B$. Suppose $n \geq 2$. As A is connected, by remark 1 after definition 18, $N_{\prec}(a) = k$ is finite and smaller than n . By lemma 13, we can suppose $a = \omega(a_1, \dots, a_k)$, $k \leq n$, $a_1 \in Prim_{\prec}(A)$, $a_2, \dots, a_k \in Prim(A)$. We have two different cases.

1. If $k \geq 2$, by lemma 20, $\Delta_{\prec}^{k-1}(\omega(a_1, \dots, a_k)) = a_1 \otimes \dots \otimes a_k \in (A^{\leq n-1})^{\otimes k}$, so, for all i , $deg_p(a_i) < n$, so $a_i \in B$. Hence, $a \in B$.
2. If $k = 1$, then $a \in Prim_{\prec}(A)$. As A is connected, by remark 1 after definition 18, $N(a) = l$ is finite and smaller than n . By lemma 17, we can suppose $a = \omega'(b_1, \dots, b_l)$, $l \leq n$, $b_1, \dots, b_{l-1} \in Prim(A)$, $b_l \in Prim_{tot}(A)$. We have two different cases.
 - (a) If $l \geq 2$, by lemma 20, $\tilde{\Delta}^{l-1}(\omega'(b_1, \dots, b_l)) = b_1 \otimes \dots \otimes b_l \in (A^{\leq n-1})^{\otimes l}$, so, for all i , $deg_p(b_i) < n$, so $b_i \in B$. Hence, $a \in B$.
 - (b) If $l = 1$, then $a = b_1 \in Prim_{tot}(A) \subseteq B$. \square

2.4 Projections on $Prim_{\prec}(A)$ and $Prim_{tot}(A)$

We here define an eulerian idempotent for connected bidendriform bialgebras.

Let A be a bidendriform bialgebra. We put:

$$A^{D2} = Vect(x \prec y, x \succ y \mid x, y \in A).$$

We define $m_{\prec}^n : A^{\otimes n} \longrightarrow A$ inductively:

$$\begin{aligned}m_{\prec}^1(a_1) &= a_1, \\ m_{\prec}^2(a_1 \otimes a_2) &= a_2 \prec a_1, \\ m_{\prec}^n(a_1 \otimes \dots \otimes a_n) &= m_{\prec}^{n-1}(a_2 \otimes \dots \otimes a_n) \prec a_1.\end{aligned}$$

Proposition 22 Let A be a bidendriform bialgebra such that for all $a \in A$, $N_{\prec}(a)$ is finite. We consider the following application:

$$T_1 : \begin{cases} A & \longrightarrow A \\ a & \longrightarrow \sum_{k=1}^{+\infty} (-1)^{k+1} m_{\prec}^k \circ \Delta_{\prec}^{k-1}(a). \end{cases}$$

Then T_1 is a projection on $\text{Prim}_{\prec}(A)$. Moreover, for all $a \in A$, $T_1(a) = a + A^{D^2}$.

Proof. Remark first that T_1 is well defined, as for all $a \in A$, $\Delta_{\prec}^{k-1}(a) = 0$ for k great enough. Let us show that $T_1(a) \in \text{Prim}_{\prec}(A)$ for all a . By lemma 13, we can suppose $a = \omega(a_1, \dots, a_n)$, avec $a_1 \in \text{Prim}_{\prec}(A)$, $a_2, \dots, a_n \in \text{Prim}(A)$. If $n = 1$, then $T_1(a) = a_1 \in \text{Prim}_{\prec}(A)$. Suppose $n \geq 2$.

We consider the following binary trees: for all $k \in \mathbb{N}$,

$$t_k^{(d)} = \begin{array}{c} \overbrace{\phantom{\Delta_{\prec}^{k-1}}} \\ \vdots \\ \Delta_{\prec} \end{array}; \quad t_k^{(g)} = \begin{array}{c} \overbrace{\phantom{\Delta_{\prec}^{k-1}}} \\ \vdots \\ \Delta_{\prec} \end{array}.$$

Then m_{\prec}^k and ω can be graphically represented in the following way:

$$\begin{cases} \omega(x_1, \dots, x_k) & = t_k^{(d)}.x_k \otimes \dots \otimes x_1, \\ m_{\prec}^k(x_1 \otimes \dots \otimes x_k) & = t_k^{(g)}.x_k \otimes \dots \otimes x_1. \end{cases}$$


By lemma 12, we have:

$$\Delta_{\prec}^{k-1}(\omega(a_1, \dots, a_n)) = \sum_{n_1 + \dots + n_k = n} t_{n_1}^{(d)}.(a_{n_1} \otimes \dots \otimes a_1) \otimes \dots \otimes t_{n_k}^{(d)}.(a_n \otimes \dots \otimes a_{n_1 + \dots + n_{k-1} + 1}).$$

(The n_i 's are positive, non zero integers). Then:

$$T_1(\omega(a_1, \dots, a_n)) = \sum_{k=1}^{+\infty} \sum_{n_1 + \dots + n_k = n} (-1)^{k+1} t_{n_1, \dots, n_k}.(a_n \otimes \dots \otimes a_1),$$

where the n_i 's are positive, non zero integers and t_{n_1, \dots, n_k} is the following tree:



As $t_{1, n_2, \dots, n_k} = t_{1+n_2, \dots, n_k}$, we obtain:

$$\begin{aligned} T_1(\omega(a_1, \dots, a_n)) &= \sum_{k=1}^{+\infty} \sum_{\substack{n_1 + \dots + n_k = n, \\ n_1 \geq 2}} (-1)^{k+1} t_{n_1, \dots, n_k}.(a_n \otimes \dots \otimes a_1) \\ &\quad + \sum_{k=1}^{+\infty} \sum_{n_1 + \dots + n_k = n-1} (-1)^{k+2} t_{1, n_1, \dots, n_k}.(a_n \otimes \dots \otimes a_1) \\ &= \sum_{k=1}^{+\infty} \sum_{\substack{n_1 + \dots + n_k = n, \\ n_1 \geq 2}} (-1)^{k+1} t_{n_1, \dots, n_k}.(a_n \otimes \dots \otimes a_1) \\ &\quad - \sum_{k=1}^{+\infty} \sum_{\substack{n'_1 + \dots + n_k = n, \\ n'_1 \geq 2}} (-1)^{k+1} t_{n'_1, \dots, n_k}.(a_n \otimes \dots \otimes a_1) \quad (\text{where } n'_1 = 1 + n_1) \\ &= 0. \end{aligned}$$

Hence, $T_1(a) \in \text{Prim}_{\prec}(A)$ for all $a \in A$. Moreover, if $a \in \text{Prim}_{\prec}(A)$, $T_1(a) = a$. So T_1 is a projection on $\text{Prim}_{\prec}(A)$. Finally, for all $a \in A$, we have:

$$T_1(a) = a + \sum_{k=2}^{+\infty} (-1)^{k+1} m_{\prec}^k \circ \Delta_{\prec}^{k-1}(a) = a + A^{D^2}. \square$$

We define $m_{\prec}^n : A^{\otimes n} \rightarrow A$ inductively:

$$\begin{aligned} m_{\prec}^1(a_1) &= a_1, \\ m_{\prec}^2(a_1 \otimes a_2) &= a_1 \succ a_2, \\ m_{\prec}^n(a_1 \otimes \dots \otimes a_n) &= a_1 \succ m_{\prec}^{n-1}(a_2 \otimes \dots \otimes a_n). \end{aligned}$$

With lemmas 14 and 16, we can show the following proposition in the same way as proposition 22:

Proposition 23 *Let A be a bidendriform bialgebra such that for all $a \in \text{Prim}_{\prec}(A)$, $N(a)$ is finite. We consider the following application:*

$$T_2 : \begin{cases} \text{Prim}_{\prec}(A) & \longrightarrow A \\ a & \longrightarrow \sum_{k=1}^{+\infty} (-1)^{k+1} m_{\prec}^k \circ \tilde{\Delta}^{k-1}(a). \end{cases}$$

Then T_2 is a projector from $\text{Prim}_{\prec}(A)$ into $\text{Prim}_{\text{tot}}(A)$. Moreover, for all $a \in A$, $T_2(a) = a + A^{D^2}$.

Corollary 24 *Let A be a bidendriform bialgebra such that for all $a \in A$, $N_{\prec}(a)$ is finite and for all $A \in \text{Prim}_{\prec}(A)$, $N(a)$ is finite (for example, A is connected). We consider the application $T = T_2 \circ T_1 : A \rightarrow A$. Then T is a projection from A into $\text{Prim}_{\text{tot}}(A)$. Moreover, for all $a \in A$, $T(a) = a + A^{D^2}$. So $A = \text{Prim}_{\text{tot}}(A) + A^{D^2}$.*

Proof. Immediate, by composition. \square

Corollary 25 *Let A be a \mathbb{N} -graded bidendriform bialgebra, such that the homogeneous part of A are finite-dimensional and $A_0 = (0)$. Then $A = \text{Prim}_{\text{tot}}(A) \oplus A^{D^2}$.*

Proof. Then A is connected (remark 2 after definition 18). By corollary 24, $A = \text{Prim}_{\text{tot}}(A) + A^{D^2}$. In the same way, the bidendriform bialgebra A^{*g} is connected. We then have $A^{*g} = \text{Prim}_{\text{tot}}(A^{*g}) + (A^{*g})^{D^2}$. By taking the orthogonal:

$$(0) = (A^{*g})^{\perp} = \text{Prim}_{\text{tot}}(A^{*g})^{\perp} \cap ((A^{*g})^{D^2})^{\perp} = A^{D^2} \cap \text{Prim}_{\text{tot}}(A).$$

So $A = \text{Prim}_{\text{tot}}(A) \oplus A^{D^2}$. \square

3 Tensor product and dendriform modules

3.1 Tensor product of dendriform algebras

We here show how the category of dendriform algebras can be given a structure of tensor category. As dendriform algebras are not objects with unit, we have to extend the usual tensor product in order to obtain a copy of A and B in the tensor product of A and B .

Definition 26 Let A, B be two vector spaces. Then:

$$A \overline{\otimes} B = (A \otimes B) \oplus (K \otimes B) \oplus (A \otimes K).$$

Proposition 27 *Let A, B be two dendriform algebras. Then $A\overline{\otimes}B$ is given a structure of dendriform algebra in the following way: for all $a_1, a_2 \in A, b_1, b_2 \in B$,*

$$\left. \begin{array}{l} (a_1 \otimes b_1) \prec (a_2 \otimes b_2) = a_1.a_2 \otimes b_1 \prec b_2, \\ (a_1 \otimes b_1) \prec (a_2 \otimes 1) = a_1.a_2 \otimes b_1, \\ (a_1 \otimes b_1) \prec (1 \otimes b_2) = a_1 \otimes b_1 \prec b_2, \\ (a_1 \otimes 1) \prec (a_2 \otimes b_2) = 0, \\ (a_1 \otimes 1) \prec (a_2 \otimes 1) = a_1 \prec a_2 \otimes 1, \\ (a_1 \otimes 1) \prec (1 \otimes b_2) = 0, \\ (1 \otimes b_1) \prec (a_2 \otimes b_2) = a_2 \otimes b_1 \prec b_2, \\ (1 \otimes b_1) \prec (a_2 \otimes 1) = a_2 \otimes b_1, \\ (1 \otimes b_1) \prec (1 \otimes b_2) = 1 \otimes b_1 \prec b_2; \end{array} \right| \begin{array}{l} (a_1 \otimes b_1) \succ (a_2 \otimes b_2) = a_1.a_2 \otimes b_1 \succ b_2, \\ (a_1 \otimes b_1) \succ (a_2 \otimes 1) = 0, \\ (a_1 \otimes b_1) \succ (1 \otimes b_2) = a_1 \otimes b_1 \succ b_2, \\ (a_1 \otimes 1) \succ (a_2 \otimes b_2) = a_1.a_2 \otimes b_2, \\ (a_1 \otimes 1) \succ (a_2 \otimes 1) = a_1 \succ a_2 \otimes 1, \\ (a_1 \otimes 1) \succ (1 \otimes b_2) = a_1 \otimes b_2, \\ (1 \otimes b_1) \succ (a_2 \otimes b_2) = a_2 \otimes b_1 \succ b_2, \\ (1 \otimes b_1) \succ (a_2 \otimes 1) = 0, \\ (1 \otimes b_1) \succ (1 \otimes b_2) = 1 \otimes b_1 \succ b_2. \end{array}$$

Proof. Direct computations. \square

Remark. $A \otimes K$ is a dendriform subalgebra of $A\overline{\otimes}B$ which is isomorphic to A and $K \otimes B$ is a dendriform subalgebra of $A\overline{\otimes}B$ which is isomorphic to B .

Proposition 28 *1. Let A, B, C be unitary dendriform algebras. Then the following application is an isomorphism of dendriform algebras:*

$$\left\{ \begin{array}{l} (A\overline{\otimes}B)\overline{\otimes}C \longrightarrow A\overline{\otimes}(B\overline{\otimes}C) \\ (a \otimes b) \otimes c \longrightarrow a \otimes (b \otimes c). \end{array} \right.$$

2. Let A, A', B, B' be unitary dendriform algebras and $\phi : A \longrightarrow A', \psi : B \longrightarrow B'$ be morphisms of unitary dendriform algebras. We then define:

$$\phi\overline{\otimes}\psi : \left\{ \begin{array}{l} A\overline{\otimes}B \longrightarrow A'\overline{\otimes}B' \\ a \otimes b \longrightarrow \phi(a) \otimes \psi(b), \\ a \otimes 1 \longrightarrow \phi(a) \otimes 1, \\ 1 \otimes b \longrightarrow 1 \otimes \psi(b), \end{array} \right.$$

for all $a \in A, b \in B$. Then $\phi\overline{\otimes}\psi$ is a morphism of dendriform algebras.

In other terms, the category of unitary dendriform algebras is a tensor category with $\overline{\otimes}$.

Proof. Direct computations. \square

We can now reformulate the axioms of bidendriform bialgebras. Let (A, \prec, \succ) be a dendriform algebra and let $\Delta_{\prec}, \Delta_{\succ} : A \longrightarrow A \otimes A$. We put $\tilde{\Delta} = \Delta_{\prec} + \Delta_{\succ}$. We then define:

$$\begin{aligned} \overline{\Delta}_{\prec} : & \left\{ \begin{array}{l} A \oplus K \longrightarrow A\overline{\otimes}A \\ a \longrightarrow \Delta_{\prec}(a) + a \otimes 1, \\ 1 \longrightarrow 0, \end{array} \right. \\ \overline{\Delta}_{\succ} : & \left\{ \begin{array}{l} A \oplus K \longrightarrow A\overline{\otimes}A \\ a \longrightarrow \Delta_{\succ}(a) + 1 \otimes a, \\ 1 \longrightarrow 0, \end{array} \right. \\ \Delta : & \left\{ \begin{array}{l} A \oplus K \longrightarrow A\overline{\otimes}A \\ a \longrightarrow \tilde{\Delta}(a) + a \otimes 1 + 1 \otimes a, \\ 1 \longrightarrow 0. \end{array} \right. \end{aligned}$$

Note that $\Delta = \overline{\Delta}_{\prec} + \overline{\Delta}_{\succ}$. We can easily prove the following assertions:

1. (4)-(6) for Δ_{\prec} and Δ_{\succ} are equivalent to (4)-(6) for $\overline{\Delta}_{\prec}$ and $\overline{\Delta}_{\succ}$.

2. (12)-(15) is equivalent to: for all $a, b \in A$,

$$\begin{aligned}\overline{\Delta}_{\succ}(a \succ b) &= \Delta(a) \succ \overline{\Delta}_{\succ}(b), & \overline{\Delta}_{\prec}(a \succ b) &= \Delta(a) \succ \overline{\Delta}_{\prec}(b), \\ \overline{\Delta}_{\succ}(a \prec b) &= \Delta(a) \prec \overline{\Delta}_{\succ}(b), & \overline{\Delta}_{\prec}(a \prec b) &= \Delta(a) \prec \overline{\Delta}_{\prec}(b).\end{aligned}$$

3. (8)-(7) is equivalent to: for all $a, b \in A$,

$$\Delta(a \succ b) = \Delta(a) \succ \Delta(b), \quad \Delta(a \prec b) = \Delta(a) \prec \Delta(b).$$

In other terms, (8)-(7) is equivalent to $\Delta : A \longrightarrow A \overline{\otimes} A$ is a morphism of dendriform algebras.

3.2 Dendriform modules

(See for example [19] for complements on operads). We denote by $\mathcal{P}_{dend} = (\mathcal{P}_{dend}(n))_{n \in \mathbb{N}^*}$ the operad of dendriform algebras (we consider here non- Σ -operads, that is to say there is no action of the symmetric groups). In other words, \mathcal{P}_{dend} is the operad generated by \prec and $\succ \in \mathcal{P}_{dend}(2)$ and the following relations:

$$\begin{aligned}\prec \circ (\prec, I) &= \prec \circ (I, \prec + \succ), \\ \prec \circ (\succ, I) &= \succ \circ (I, \prec), \\ \succ \circ (\prec + \succ, I) &= \succ \circ (I, \succ).\end{aligned}$$

Let A be a dendriform algebra. A dendriform module over A is a vector space M together with applications, for all $n \in \mathbb{N}^*$:

$$\begin{cases} \mathcal{P}_{dend}(n) \otimes A^{\otimes(n-1)} \otimes M & \longrightarrow & M \\ p \otimes (a_1 \otimes \dots \otimes a_{n-1}) \otimes m & \longrightarrow & p.(a_1 \otimes \dots \otimes a_{n-1} \otimes m), \end{cases}$$

satisfying the same associativity relations and unit relation as those for \mathcal{P}_{dend} -algebras. In other terms, a dendriform module over A is a vector space M together with two applications:

$$\dashv: \begin{cases} A \otimes M & \longrightarrow & M \\ a \otimes m & \longrightarrow & a \dashv m, \end{cases} \quad \vdash: \begin{cases} A \otimes M & \longrightarrow & M \\ a \otimes m & \longrightarrow & a \vdash m, \end{cases}$$

with the following compatibilities: for all $a, b \in A, m \in M$,

$$(a \prec b) \dashv m = a \dashv (b \dashv m + b \vdash m), \quad (16)$$

$$(a \succ b) \dashv m = a \vdash (b \dashv m), \quad (17)$$

$$(a \prec b + a \succ b) \vdash m = a \vdash (b \vdash m). \quad (18)$$

(We have $a \dashv m = \prec.(a \otimes m)$ and $a \vdash m = \succ.(a \otimes m)$).

Remarks.

1. If M is a dendriform module over A , then $\dashv + \vdash$ gives M a structure of left module over the associative algebra $(A, \prec + \succ)$ by (16)+(17)+(18). This action will be denoted by $a.m = a \dashv m + a \vdash m$.
2. A is a dendriform module over itself with $\dashv = \prec$ and $\vdash = \succ$.
3. Suppose that A is a dendriform bialgebra. Then, as Δ is a morphism of dendriform algebras, $A \overline{\otimes} A$ is given a structure of a dendriform module over A with, for all $a \in A, \sum b \otimes c \in A \overline{\otimes} A$:

$$a \dashv \left(\sum b \otimes c \right) = \sum \Delta(a) \prec (b \otimes c), \quad a \vdash \left(\sum b \otimes c \right) = \sum \Delta(a) \succ (b \otimes c).$$

Moreover, Δ is a morphism of dendriform modules.

4. Let $(A, \prec, \succ, \tilde{\Delta})$ be a dendriform bialgebra and let $\Delta_{\prec}, \Delta_{\succ} : A \longrightarrow A \otimes A$. Observe that (12)-(15) is equivalent to: $\overline{\Delta}_{\prec}, \overline{\Delta}_{\succ}$ are morphisms of dendriform modules.

Proposition 29 *Let A be a free dendriform algebra generated by a subspace V and let M be a dendriform module over A . For any linear application $\phi : V \longrightarrow M$, there exists a unique morphism of dendriform modules $\Phi : A \longrightarrow M$, such that $\Phi|_V = \phi$.*

Proof.

Unicity. Because V generates A as a dendriform algebra, and hence as a dendriform module.

Existence. As A is freely generated by V , we can suppose that $A = \bigoplus_{n \geq 1} \mathcal{P}_{dend}(n) \otimes V^{\otimes n}$. We

then define:

$$\Phi : \begin{cases} A & \longrightarrow M \\ p \otimes (v_1 \otimes \dots \otimes v_n) & \longrightarrow p.(v_1 \otimes \dots \otimes v_{n-1} \otimes \phi(v_n)) \end{cases}$$

for all $p \in \mathcal{P}_{dend}(n)$, $v_1, \dots, v_n \in V$. Then Φ fits the asked conditions. \square

Proposition 30 *Let $(A, \prec, \succ, \tilde{\Delta})$ be a dendriform bialgebra, with applications $\Delta_{\prec}, \Delta_{\succ} : A \longrightarrow A \otimes A$. We put:*

$$\begin{aligned} \overline{\Delta}_{\prec} &: \begin{cases} A & \longrightarrow A \overline{\otimes} A \\ a & \longrightarrow \Delta_{\prec}(a) + a \otimes 1, \end{cases} \\ \overline{\Delta}_{\succ} &: \begin{cases} A & \longrightarrow A \overline{\otimes} A \\ a & \longrightarrow \Delta_{\succ}(a) + 1 \otimes a, \end{cases} \\ \Delta &: \begin{cases} A & \longrightarrow A \overline{\otimes} A \\ a & \longrightarrow \tilde{\Delta}(a) + a \otimes 1 + 1 \otimes a. \end{cases} \end{aligned}$$

We suppose the following conditions:

1. $(A, \Delta_{\prec}, \Delta_{\succ})$ satisfies relations (4), (5) and (6) on a set of generators of the dendriform algebra A . Moreover, $\Delta_{\prec} + \Delta_{\succ} = \tilde{\Delta}$ on this set of generators.
2. Δ_{\prec} and Δ_{\succ} are morphisms of A -dendriform modules.

Then A is a bidendriform bialgebra.

Proof. Let us first show relation (4). We put $X = \{a \in A / a \text{ satisfies (4)}\}$. We have:

$$\begin{aligned} X &= Ker((\Delta_{\prec} \otimes Id) \circ \Delta_{\prec} - (Id \otimes \Delta_{\prec} + Id \otimes \Delta_{\succ}) \circ \Delta_{\prec}) \\ &= Ker((\overline{\Delta}_{\prec} \otimes Id) \circ \overline{\Delta}_{\prec} - (Id \otimes \overline{\Delta}_{\prec} + Id \otimes \overline{\Delta}_{\succ}) \circ \overline{\Delta}_{\prec}), \end{aligned}$$

which is the kernel of a certain morphism of dendriform modules from A into $A \overline{\otimes} A \overline{\otimes} A$ by hypothesis 2. So X is a dendriform submodule of A , hence a dendriform subalgebra of A . As it contains a set of generators by hypothesis 1, it is A . We can prove (5), (6), and the fact that $\tilde{\Delta} = \Delta_{\prec} + \Delta_{\succ}$ on the whole A in the same way. As the hypothesis 2 is a reformulation of axioms (12)-(15), A is a bidendriform bialgebra. \square

3.3 Bidendriform structure on \mathcal{A}^D

Unfortunately, $(\mathcal{A}^D, \prec, \succ, \Delta'_{\prec}, \Delta'_{\succ})$ is not a bidendriform bialgebra. For example, for $a = b = \bullet_d$:

$$a \prec b = \mathbf{1}_d^a, \quad \Delta'_{\prec}(a \prec b) = 0,$$

whereas $a'b'_{\prec} \otimes a'' \prec b''_{\prec} + a'b \otimes a'' + b'_{\prec} \otimes a \prec b''_{\prec} + b \otimes a = \bullet_d \otimes \bullet_d$. So (15) is not satisfied.

Theorem 31 *There is a unique structure of bidendriform bialgebra (with the already known dendriform bialgebra structure) on $\mathcal{A}^{\mathcal{D}}$ such that for all $d \in \mathcal{D}$, $\Delta_{\prec}(\cdot_a) = \Delta_{\succ}(\cdot_a) = 0$. Hence, $(\mathcal{A}^{\mathcal{D}}, \prec, \succ, \Delta_{\prec}, \Delta_{\succ})$ is a bidendriform bialgebra, which induces the structure of Hopf algebra of $\mathcal{H}^{\mathcal{D}}$ already described.*

Proof. We use the notations of proposition 30.

Unicity. As the \cdot_a 's generate $\mathcal{A}^{\mathcal{D}}$, there is at most one way to extend $\overline{\Delta}_{\prec}$ and $\overline{\Delta}_{\succ}$ to A as morphisms of dendriform modules, with notations of proposition 30.

Existence. By proposition 29, we can extend $\overline{\Delta}_{\prec}$ and $\overline{\Delta}_{\succ}$ to A as morphisms of dendriform modules. So the second condition of proposition 30 is satisfied. The first one is trivially satisfied on the set of generators $\{\cdot_a / d \in \mathcal{D}\}$. \square

When \mathcal{D} is a graded set, this structure obviously respects the gradation of $\mathcal{A}^{\mathcal{D}}$:

Corollary 32 *If \mathcal{D} is a graded set, then $\mathcal{A}^{\mathcal{D}}$ is a graded bidendriform bialgebra.*

Proposition 33 *Let $F \in \mathbb{F}^{\mathcal{D}}$, $F \neq 1$. We consider the set $\text{Adm}_{\prec}(F)$ of admissible cuts of F satisfying one of these two conditions:*

1. *c cuts one of the edges which are on the path from the root of the last tree of F to the leave which is at most on the right of F .*
2. *c cuts totally the last tree of F if F is not a single tree.*

Then:

$$\Delta_{\prec}(F) = \sum_{c \in \text{Adm}_{\prec}(F)} P^c(F) \otimes R^c(F). \quad (19)$$

Prove. We denote by $F'_{\ll} \otimes F''_{\ll}$ the second member of (19). Let us prove the result by induction on $n = \text{weight}(F)$. If $n = 1$, then $F = \cdot_a$ and the result is obvious. If $n \geq 2$, we have two possible cases:

1. $F = GH$, with $\text{weight}(G), \text{weight}(H) < n$. By a study of $\text{Adm}_{\prec}(F)$, we easily have:

$$\begin{aligned} F'_{\ll} \otimes F''_{\ll} &= G'H'_{\ll} \otimes G''H''_{\ll} + G'H \otimes G'' + GH'_{\ll} \otimes H''_{\ll} + GH'_{\ll} \otimes H''_{\ll} + H \otimes G \\ &= G'H'_{\prec} \otimes G''H''_{\prec} + G'H \otimes G'' + GH'_{\prec} \otimes H''_{\prec} + GH'_{\prec} \otimes H''_{\prec} + H \otimes G, \end{aligned}$$

by the induction hypothesis on H . By (11), this is equal to $\Delta_{\prec}(F)$.

2. $F = B_d^+(G)$, $\text{weight}(G) = n - 1$. By a study of $\text{Adm}_{\prec}(G)$, we easily have:

$$\begin{aligned} F'_{\ll} \otimes F''_{\ll} &= G'_{\ll} \otimes B_d^+(G''_{\ll}) + G \otimes \cdot_a \\ &= G'_{\prec} \otimes B_d^+(G''_{\prec}) + G \otimes \cdot_a \\ &= G'_{\prec} \otimes \cdot_a \prec G''_{\prec} + G \otimes \cdot_a, \end{aligned}$$

by the induction hypothesis on H . As $F = \cdot_a \prec G$, by (15) for $a = \cdot_a$, this is equal to $\Delta_{\prec}(F)$. \square

Example.

$$\begin{aligned} \Delta_{\prec} \left(\cdot_a \overset{e}{\underset{b}{\text{Y}}}_c^d \right) &= \overset{e}{\underset{b}{\text{Y}}}_c^d \otimes \cdot_a + \cdot_a \cdot e \cdot d \otimes \mathbf{1}_b^e + \cdot_e \cdot d \otimes \cdot_a \mathbf{1}_b^e \\ &\quad + \cdot_a \overset{e}{\underset{c}{\text{V}}}_c^d \otimes \cdot_b + \overset{e}{\underset{c}{\text{V}}}_c^d \otimes \cdot_a \cdot b + \cdot_a \cdot d \otimes \mathbf{1}_b^e + \cdot_d \otimes \cdot_a \mathbf{1}_b^e, \\ \Delta_{\succ} \left(\cdot_a \overset{e}{\underset{b}{\text{Y}}}_c^d \right) &= \cdot_a \otimes \overset{e}{\underset{b}{\text{Y}}}_c^d + \cdot_a \cdot e \otimes \mathbf{1}_b^d + \cdot_e \otimes \cdot_a \mathbf{1}_b^d. \end{aligned}$$

3.4 Totally primitive elements of $\mathcal{A}^{\mathcal{D}}$ and universal property

We here prove that a connected bidendriform bialgebra is isomorphic to $\mathcal{A}^{\mathcal{D}}$ for a certain set \mathcal{D} .

Proposition 34 $Prim_{tot}(\mathcal{A}^{\mathcal{D}}) = Vect(\bullet_d / d \in \mathcal{D})$.

Proof. In an immediate way, $\bullet_d \in Prim_{tot}(\mathcal{A}^{\mathcal{D}})$. Let us show the other inclusion.

First case. Suppose that \mathcal{D} is finite. We graduate \mathcal{D} by putting all its elements in degree 1. Then $\mathcal{A}^{\mathcal{D}}$ is graded, with $\mathcal{A}_0^{\mathcal{D}} = (0)$ and $\mathcal{A}_n^{\mathcal{D}}$ finite-dimensional for all n . By corollary 25, $\mathcal{A}^{\mathcal{D}} = Prim_{tot}(\mathcal{A}^{\mathcal{D}}) \oplus (\mathcal{A}^{\mathcal{D}})^{D^2}$. As $\mathcal{A}^{\mathcal{D}}$ is freely generated by the \bullet_d 's, $Vect(\bullet_d / d \in \mathcal{D}) \oplus (\mathcal{A}^{\mathcal{D}})^{D^2} = \mathcal{A}^{\mathcal{D}}$, and this implies that $Prim_{tot}(\mathcal{A}^{\mathcal{D}}) \subseteq Vect(\bullet_d / d \in \mathcal{D})$.

General case. Let $p \in Prim_{tot}(\mathcal{A}^{\mathcal{D}})$. There exists a finite subset \mathcal{D}' of \mathcal{D} , such that $p \in \mathcal{A}^{\mathcal{D}'}$. Then $p \in Vect(\bullet_d / d \in \mathcal{D}') \subseteq Vect(\bullet_d / d \in \mathcal{D})$. \square

Theorem 35 *Let A be a bidendriform bialgebra. For all $d \in \mathcal{D}$, let $p_d \in Prim_{tot}(A)$. There exists a unique morphism of bidendriform bialgebras:*

$$\Psi : \begin{cases} \mathcal{A}^{\mathcal{D}} & \longrightarrow A \\ \bullet_d & \longrightarrow p_d. \end{cases}$$

Moreover, we have:

1. If the p_d 's are linearly independent, then Ψ is monic.
2. If the dendriform coalgebra A is connected and if the family $(p_d)_{d \in \mathcal{D}}$ linearly generates $Prim_{tot}(A)$, then Ψ is epic.
3. If the dendriform coalgebra A is connected and if the family $(p_d)_{d \in \mathcal{D}}$ is a linear basis of $Prim_{tot}(A)$, then Ψ is an isomorphism.

Proof. As $\mathcal{A}^{\mathcal{D}}$ is freely generated by the \bullet_d 's, Ψ defines a unique morphism of dendriform algebras. As \bullet_d and p_d are both totally primitive, Ψ is a morphism of bidendriform bialgebras. We now prove the two assertions.

1. We graduate \mathcal{D} by putting all its elements of degree 1. Suppose that $Ker(\Psi) \neq (0)$. Let $x \in Ker(\Psi)$, non-zero, of minimal degree n . So Ψ is monic on $\mathcal{A}_{<n}^{\mathcal{D}} = \mathcal{A}_0^{\mathcal{D}} \oplus \dots \oplus \mathcal{A}_{n-1}^{\mathcal{D}}$. Moreover, $\Delta_{<}(x) \in \mathcal{A}_{<n}^{\mathcal{D}} \otimes \mathcal{A}_{<n}^{\mathcal{D}}$ and $(\Psi \otimes \Psi) \circ \Delta_{<}(x) = \Delta_{<}(\Psi(x)) = 0$. By injectivity, $\Delta_{<}(x) = 0$. In the same way, $\Delta_{>}(x) = 0$. So $x \in Prim_{tot}(\mathcal{A}^{\mathcal{D}}) = Vect(\bullet_d / d \in \mathcal{D})$ (proposition 34). As the p_d 's are linearly independent, $Ker(\Psi) \cap Vect(\bullet_d / d \in \mathcal{D}) = (0)$: contradiction. So Ψ is monic.
2. Then $Im(\Psi)$ is a dendriform subalgebra of A which contains $Prim_{tot}(A)$. As A is connected, $Prim_{tot}(A)$ generates A (theorem 21), so Ψ is epic.
3. Comes from 1 and 2. \square

This corollary is immediate:

Corollary 36 *1. Let A be a connected bidendriform bialgebra. Let $(p_d)_{d \in \mathcal{D}}$ be a basis of $Prim_{tot}(A)$. Then the morphism $\Psi : \mathcal{A}^{\mathcal{D}} \longrightarrow A$ described in theorem 35 is an isomorphism of bidendriform bialgebras.*

2. *Let A be a \mathbb{N} -graded bidendriform bialgebra which is connected as a dendriform coalgebra. Let $(p_d)_{d \in \mathcal{D}}$ be a basis of $Prim_{tot}(A)$ made of homogeneous elements. Then \mathcal{D} is given a gradation by putting $|d| = |p_d|$. Then the morphism $\Psi : \mathcal{A}^{\mathcal{D}} \longrightarrow A$ described in theorem 35 is an isomorphism of graded bidendriform bialgebras.*

Corollary 37 *Let A be a \mathbb{N} -graded bidendriform bialgebra such that $A_0 = (0)$ and A_n is finite-dimensional for all $n \in \mathbb{N}$. We consider the following formal series:*

$$P(X) = \sum_{n=1}^{+\infty} \dim(\text{Prim}_{\text{tot}}(A)_n) X^n, \quad R(X) = \sum_{n=1}^{+\infty} \dim(A_n) X^n.$$

$$\text{Then } P(X) = \frac{R(X)}{(1 + R(X))^2}.$$

Proof. Immediate if $A = (0)$. Suppose that $A \neq (0)$ (so $P(X) \neq 0$). By corollary 36, we can suppose $A = \mathcal{A}^{\mathcal{D}}$. By proposition 34, $P(X) = D(X)$ of proposition 8. Hence:

$$\begin{aligned} R(X) + 1 &= \frac{1 - \sqrt{1 - 4P(X)}}{2P(X)} \\ \Rightarrow 2P(X)(R(X) + 1) - 1 &= -\sqrt{1 - 4P(X)} \\ \Rightarrow 4P(X)^2(R(X) + 1)^2 + 1 - 4P(X)(R(X) + 1) &= 1 - 4P(X) \\ \Rightarrow P(X)(R(X) + 1)^2 - R(X) - 1 &= -1 \\ \Rightarrow P(X) &= \frac{R(X)}{(R(X) + 1)^2}. \quad \square \end{aligned}$$

4 Application to the Hopf algebra \mathbf{FQSym}

4.1 Recalls

(See [3, 5, 18]). The algebra \mathbf{FQSym} is the vector space generated by the elements $(\mathbf{F}_u)_{u \in \mathbb{S}}$, where \mathbb{S} is the disjoint union of the symmetric groups S_n ($n \in \mathbb{N}$). Its product and its coproduct are given in the following way: for all $u \in S_n$, $v \in S_m$, by putting $u = (u_1 \dots u_n)$,

$$\begin{aligned} \Delta(\mathbf{F}_u) &= \sum_{i=0}^n \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_n)}, \\ \mathbf{F}_u \cdot \mathbf{F}_v &= \sum_{\zeta \in sh(n, m)} \mathbf{F}_{(u \times v) \cdot \zeta^{-1}}, \end{aligned}$$

where $sh(n, m)$ is the set of (n, m) -shuffles, and st is the standardisation. Its unit is $1 = \mathbf{F}_\emptyset$, where \emptyset is the unique element of S_0 . Moreover, \mathbf{FQSym} is a \mathbb{N} -graded Hopf algebra, by putting $|\mathbf{F}_u| = n$ if $u \in S_n$.

Examples.

$$\begin{aligned} \mathbf{F}_{(12)} \mathbf{F}_{(123)} &= \mathbf{F}_{(12345)} + \mathbf{F}_{(13245)} + \mathbf{F}_{(13425)} + \mathbf{F}_{(13452)} + \mathbf{F}_{(31245)} \\ &\quad + \mathbf{F}_{(31425)} + \mathbf{F}_{(31452)} + \mathbf{F}_{(34125)} + \mathbf{F}_{(34152)} + \mathbf{F}_{(34512)}, \end{aligned}$$

$$\begin{aligned} \Delta(\mathbf{F}_{(12543)}) &= 1 \otimes \mathbf{F}_{(12543)} + \mathbf{F}_{(1)} \otimes \mathbf{F}_{(1432)} + \mathbf{F}_{(12)} \otimes \mathbf{F}_{(321)} \\ &\quad + \mathbf{F}_{(123)} \otimes \mathbf{F}_{(21)} + \mathbf{F}_{(1243)} \otimes \mathbf{F}_{(1)} + \mathbf{F}_{(12543)} \otimes 1. \end{aligned}$$

4.2 Bidendriform structure on \mathbf{FQSym}

Let $(\mathbf{FQSym})_+ = \text{Vect}(\mathbf{F}_u / u \in S_n, n \geq 1)$ be the augmentation ideal of \mathbf{FQSym} . We define $\prec, \succ, \Delta_\prec$ and Δ_\succ on $(\mathbf{FQSym})_+$ in the following way: for all $u \in S_n$, $v \in S_m$, by putting

$$u = (u_1 \dots u_n),$$

$$\begin{aligned} \mathbf{F}_u \prec \mathbf{F}_v &= \sum_{\substack{\zeta \in sh(n,m) \\ \zeta^{-1}(n+m)=n}} \mathbf{F}_{(u \times v) \cdot \zeta^{-1}}, \\ \mathbf{F}_u \succ \mathbf{F}_v &= \sum_{\substack{\zeta \in sh(n,m) \\ \zeta^{-1}(n+m)=n+m}} \mathbf{F}_{(u \times v) \cdot \zeta^{-1}}, \\ \Delta_{\prec}(\mathbf{F}_u) &= \sum_{i=u^{-1}(n)}^{n-1} \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_n)}, \\ \Delta_{\succ}(\mathbf{F}_u) &= \sum_{i=1}^{u^{-1}(n)-1} \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_n)}. \end{aligned}$$

Examples.

$$\begin{aligned} \mathbf{F}_{(12)} \prec \mathbf{F}_{(123)} &= \mathbf{F}_{(13452)} + \mathbf{F}_{(31452)} + \mathbf{F}_{(34152)} + \mathbf{F}_{(34512)}, \\ \mathbf{F}_{(12)} \succ \mathbf{F}_{(123)} &= \mathbf{F}_{(12345)} + \mathbf{F}_{(13245)} + \mathbf{F}_{(13425)} + \mathbf{F}_{(31245)} + \mathbf{F}_{(31425)} + \mathbf{F}_{(34125)}, \end{aligned}$$

$$\begin{aligned} \Delta_{\prec}(\mathbf{F}_{(12543)}) &= \mathbf{F}_{(123)} \otimes \mathbf{F}_{(21)} + \mathbf{F}_{(1243)} \otimes \mathbf{F}_{(1)}, \\ \Delta_{\succ}(\mathbf{F}_{(12543)}) &= \mathbf{F}_{(1)} \otimes \mathbf{F}_{(1432)} + \mathbf{F}_{(12)} \otimes \mathbf{F}_{(321)}. \end{aligned}$$

Theorem 38 $((\mathbf{FQSym})_+, \prec, \succ, \Delta_{\prec}, \Delta_{\succ})$ is a connected bidendriform bialgebra.

Proof. The structure of dendriform bialgebra is already introduced in [22], so we already have (1)-(3), and (12)+(14), (13)+(15). We consider the symmetric non-degenerate pairing on $(\mathbf{FQSym})_+$ defined in [5, 18] by $\langle \mathbf{F}_\sigma, \mathbf{F}_\tau \rangle = \delta_{\sigma, \tau^{-1}}$. Note that $\Delta_{\prec}, \Delta_{\succ}$ are the transposes of \prec, \succ for this pairing. So $((\mathbf{FQSym})_+, \Delta_{\prec}, \Delta_{\succ})$ is a codendriform coalgebra. So we already have (4)-(6), and (12)+(13), (14)+(15). It is then enough to prove (13). We have:

$$\begin{aligned} \Delta_{\succ}(\mathbf{F}_{(u_1 \dots u_n)} \prec \mathbf{F}_{(v_1 \dots v_m)}) &= \Delta_{\succ} \left(\sum_{1 \leq j_1 < \dots < j_m < n+m} \mathbf{F}_{\underbrace{(u_1 \dots v_1 + n \dots v_m + n \dots u_n)}_{v_i + n \text{ in position } j_i}} \right) \\ &= \sum_{1 \leq j_1 < \dots < j_m < n+m} \mathbf{F}_{st(u_1 \dots)} \otimes \mathbf{F}_{st(\dots m+n \dots u_n)} \\ &= \underbrace{(\mathbf{F}_{(u_1 \dots u_n)})' (\mathbf{F}_{(v_1 \dots v_n)})'_{\succ} \otimes (\mathbf{F}_{(u_1 \dots u_n)})'' \prec (\mathbf{F}_{(v_1 \dots v_n)})''_{\succ}}_{\text{terms with } u_i \text{'s and } v_j \text{'s on both sides on the } \otimes} \\ &\quad + \underbrace{(\mathbf{F}_{(u_1 \dots u_n)})' \otimes (\mathbf{F}_{(u_1 \dots u_n)})'' \prec \mathbf{F}_{(v_1 \dots v_n)}}_{\text{terms with all the } v_i \text{'s on the right on the } \otimes} \\ &\quad + \underbrace{(\mathbf{F}_{(v_1 \dots v_n)})'_{\succ} \otimes \mathbf{F}_{(u_1 \dots u_n)} \prec (\mathbf{F}_{(v_1 \dots v_n)})''_{\succ}}_{\text{terms with all the } u_i \text{'s the right of the } \otimes}. \end{aligned}$$

So (13) is satisfied, and $(\mathbf{FQSym})_+$ is a bidendriform bialgebra. Moreover, it is \mathbb{N} -graded, with $((\mathbf{FQSym})_+)_0 = (0)$, so it is connected. \square

Remark. It is of course possible to prove (4)-(6) directly. For example, the two members of (4) for $a = \mathbf{F}_u$, with $u = (u_1 \dots u_n)$, are equal to:

$$\sum_{\sigma^{-1}(n) \leq i < j < n} \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_j)} \otimes \mathbf{F}_{st(u_{j+1} \dots u_n)}.$$

For (5), we obtain:

$$\sum_{1 \leq i < \sigma^{-1}(n) \leq j < n} \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_j)} \otimes \mathbf{F}_{st(u_{j+1} \dots u_n)}.$$

And for (6):

$$\sum_{1 \leq i < j < \sigma^{-1}(n)} \mathbf{F}_{st(u_1 \dots u_i)} \otimes \mathbf{F}_{st(u_{i+1} \dots u_j)} \otimes \mathbf{F}_{st(u_{j+1} \dots u_n)}.$$

Moreover, $(\mathbf{FQSym})_+$ is graded and connected, with:

$$R(X) = \sum_{n=1}^{+\infty} \dim(\mathbf{FQSym}_n) X^n = \sum_{n=1}^{+\infty} n! X^n.$$

By corollary 37:

$$P(X) = \sum_{n=1}^{+\infty} \dim(\text{Prim}_{tot}((\mathbf{FQSym})_+)_n) X^n = \sum_{n=1}^{+\infty} p_n X^n = \frac{R(X)}{(R(X) + 1)^2}.$$

We obtain then:

n	1	2	3	4	5	6	7	8	9	10	11	12
p_n	1	0	1	6	39	284	2 305	20 682	203 651	2 186 744	25 463 925	319 989 030

Remark. Let $\Psi : \mathcal{A} \rightarrow (\mathbf{FQSym})_+$ be the unique morphism of bidendriform bialgebras which send \bullet to $\mathbf{F}_{(1)}$. It coincides with the morphism $\overline{\Phi}^*$ of [17]. We obtain by theorem 35 that it is monic.

By corollary 36:

Theorem 39 *Let \mathcal{D} be a graded set such that $\mathcal{D}_0 = \emptyset$ and for all $n \geq 1$, $\text{card}(\mathcal{D}_n) = p_n$. Then $(\mathbf{FQSym})_+$ and $\mathcal{A}^{\mathcal{D}}$ are isomorphic as graded bidendriform bialgebras. Hence, \mathbf{FQSym} and $\mathcal{H}^{\mathcal{D}}$ are isomorphic as graded Hopf algebras.*

We can now prove the conjecture 3.8 of [5]:

Corollary 40 *If the field K is of characteristic 0, the Lie algebra $\text{Prim}(\mathbf{FQSym})$ is free.*

Proof. By proposition 141 of [6], the Lie algebra $\text{Prim}(\mathcal{H}^{\mathcal{D}})$ is free. \square

Finally, we give a basis of totally primitive elements of $(\mathbf{FQSym})_+$ of degree smaller than 4. With notations of [5]:

$$\begin{aligned} \mathbf{V}_1 &= \mathbf{F}_1, \\ \mathbf{V}_{231} &= \mathbf{F}_{231} - \mathbf{F}_{132}, \\ \mathbf{V}_{3142} &= \mathbf{F}_{3142} - \mathbf{F}_{2143}, \\ \mathbf{V}_{2431} &= \mathbf{F}_{2431} - \mathbf{F}_{1432}, \\ \mathbf{V}_{2341} &= \mathbf{F}_{2341} - \mathbf{F}_{1342}, \\ \mathbf{V}_{3241} &= \mathbf{F}_{1243} - \mathbf{F}_{1342} - \mathbf{F}_{2143} + \mathbf{F}_{3241}, \\ \mathbf{V}_{3412} - \mathbf{V}_{2413} &= \mathbf{F}_{3412} - \mathbf{F}_{2413}, \\ \mathbf{V}_{3421} - \mathbf{V}_{2413} &= \mathbf{F}_{1423} - \mathbf{F}_{2413} - \mathbf{F}_{1432} + \mathbf{F}_{3421}. \end{aligned}$$

5 Bidendriform pairings

5.1 Definition

Definition 41 Let A and B be two bidendriform bialgebras and let $\langle, \rangle: A \times B \longrightarrow K$ be a bilinear form. We will say that \langle, \rangle is a bidendriform pairing if, for all $a, a_1, a_2 \in A$, $b, b_1, b_2 \in B$, the following conditions are satisfied:

$$\begin{array}{l} \langle a_1 \prec a_2, b \rangle = \langle a_1 \otimes a_2, \Delta_{\prec}(b) \rangle, \quad \left| \quad \langle a, b_1 \prec b_2 \rangle = \langle \Delta_{\prec}(a), b_1 \otimes b_2 \rangle, \right. \\ \langle a_1 \succ a_2, b \rangle = \langle a_1 \otimes a_2, \Delta_{\succ}(b) \rangle, \quad \left| \quad \langle a, b_1 \succ b_2 \rangle = \langle \Delta_{\succ}(a), b_1 \otimes b_2 \rangle. \right. \end{array}$$

Suppose that A and B are graded. Then \langle, \rangle will be said homogeneous if for all homogeneous elements $a \in A$, $b \in B$, $\langle a, b \rangle = 0$ if a and b have different degrees.

Remark. If $\langle, \rangle: A \times B \longrightarrow K$ is a bidendriform pairing, then it induces a Hopf pairing between $\overline{A} = A \oplus K$ and $\overline{B} = B \oplus K$ with, for all $a \in A$, $b \in B$, $\langle 1, b \rangle = 0$, $\langle a, 1 \rangle = 0$, and $\langle 1, 1 \rangle = 1$.

Example. The pairing on $(\mathbf{FQSym})_+$ defined by $\langle \mathbf{F}_u, \mathbf{F}_v \rangle = \delta_{u,v^{-1}}$ is a bidendriform pairing.

5.2 Hopf dual of a bidendriform bialgebra

We here extend the notion of Hopf dual of a Hopf algebra (see [1, 10]) to bidendriform bialgebras. Let A be a dendriform algebra. We denote by A^* the following subspace of the linear dual A^* of A :

$$A^* = \{f \in A^* / f \text{ vanishes on a dendriform ideal of finite codimension of } A\}.$$

For all $a \in A$, we consider the following applications:

$$\begin{array}{l} L_a^{\prec} : \left\{ \begin{array}{l} A^* \longrightarrow A^* \\ f \longrightarrow \left\{ \begin{array}{l} A \longrightarrow K \\ b \longrightarrow L_a^{\prec}(f)(b) = f(a \prec b), \end{array} \right. \end{array} \right. \\ L_a^{\succ} : \left\{ \begin{array}{l} A^* \longrightarrow A^* \\ f \longrightarrow \left\{ \begin{array}{l} A \longrightarrow K \\ b \longrightarrow L_a^{\succ}(f)(b) = f(a \succ b), \end{array} \right. \end{array} \right. \\ R_a^{\prec} : \left\{ \begin{array}{l} A^* \longrightarrow A^* \\ f \longrightarrow \left\{ \begin{array}{l} A \longrightarrow K \\ b \longrightarrow R_a^{\prec}(f)(b) = f(b \prec a), \end{array} \right. \end{array} \right. \\ R_a^{\succ} : \left\{ \begin{array}{l} A^* \longrightarrow A^* \\ f \longrightarrow \left\{ \begin{array}{l} A \longrightarrow K \\ b \longrightarrow R_a^{\succ}(f)(b) = f(b \succ a). \end{array} \right. \end{array} \right. \end{array}$$

Lemma 42 Let $f \in A^*$. The following assertions are equivalent:

1. $f \in A^*$.
2. There exists a finite dimensional subspace U of A^* , stable for all $a \in A$ by L_a^{\prec} , L_a^{\succ} , R_a^{\prec} and R_a^{\succ} , which contains f .

Proof.

1 \implies 2. Let I be a dendriform ideal of A , such that $f(I) = (0)$ and f has a finite codimension. Then I^\perp is finite-dimensional and contains f . Let $g \in I^\perp$ and let $a \in A$. For all $b \in I$:

$$L_a^\prec(g)(b) = g(\underbrace{a \prec b}_{\in I}) = 0,$$

as I is a dendriform ideal of A . So $L_a^\prec(g) \in I^\perp$: I^\perp is stable under L_a^\prec . In the same way, I^\perp is stable under L_a^\succ , R_a^\prec and R_a^\succ .

2 \implies 1. Let $I = U^\perp$. Then I has a finite codimension. Moreover, as $f \in U$, $f(I) = (0)$. Let $a \in A$, $b \in I$. For all $g \in U$:

$$g(a \prec b) = \underbrace{L_a^\prec(g)}_{\in U}(b) = 0,$$

so $a \prec b \in I$. In the same way, $a \succ b$, $b \prec a$ and $b \succ a \in I$, so I is a dendriform ideal. \square

Lemma 43 \prec^* , \succ^* send A^* into $A^* \otimes A^*$.

Proof. Let $f \in A^*$ and U as in lemma 42. Let (u_1, \dots, u_n) be a basis of U . By lemma 42, observe that $U \subseteq A^*$. For all $a \in A$, we put:

$$L_a^\prec(f) = \sum_{i=1}^n v_i(a)u_i.$$

Then the v_i 's are elements of A^* . Let us fix $j \in \{1, \dots, n\}$. There exists $b_j \in A$, such that $v_i(b_j) = \delta_{i,j}$. Then:

$$L_a^\prec(f)(b_j) = \sum_{i=1}^n v_i(a)u_i(b_j) = v_j(a) = f(a \prec b_j) = R_{b_j}^\prec(f)(a),$$

so $v_j = R_{b_j}^\prec(f) \in U$. Moreover, for all $a, b \in A$:

$$\prec^*(f)(a \otimes b) = f(a \prec b) = L_a^\prec(f)(b) = \sum_{i=1}^n v_i(a) \otimes u_i(b) = \sum_{i=1}^n v_i \otimes u_i(a \otimes b),$$

so $\prec^*(f) \in U \otimes U \subseteq A^* \otimes A^*$. In the same way, $\succ^*(f) \in A^* \otimes A^*$. \square

Remark. So (A^*, \prec^*, \succ^*) is a dendriform coalgebra.

Suppose now that A is a bidendriform bialgebra. By duality, $(A^*, \Delta_\prec^*, \Delta_\succ^*)$ is given a dendriform algebra structure. More precisely, for all $f, g \in A^*$, we have, for all $a \in A$:

$$(f \prec g)(a) = (f \otimes g) \circ \Delta_\prec(a), \quad (f \succ g)(a) = (f \otimes g) \circ \Delta_\succ(a).$$

Lemma 44 A^* is a dendriform subalgebra of A^* .

Proof. Let $f, g \in A^*$. By lemma 42, there exists a finite-dimensional subspace U of A^* , containing f and g , stable for all $a \in A$ by L_a^\prec , L_a^\succ , R_a^\prec and R_a^\succ . Let $F, G \in U$. Let $a \in A$. For all $b \in A$, by (15):

$$\begin{aligned} L_a^\prec(F \prec G)(b) &= (F \prec G)(a \prec b) \\ &= (F \otimes G) \circ \Delta_\prec(a \prec b) \\ &= (F \otimes G)(a'b'_\prec \otimes a'' \prec b''_\prec + a'b \otimes a'' + b'_\prec \otimes a \prec b''_\prec + b \otimes a) \\ &= ((L_{a'}^\prec + L_{a'}^\succ)(F) \prec L_{a''}^\prec(G))(b) \\ &\quad + G(a'')((L_{a'}^\prec + L_{a'}^\succ)(F))(b) + (F \prec L_a^\prec(G))(b) + F(a)G(b). \end{aligned}$$

Hence:

$$\begin{aligned} L_a^\prec(F \prec G) &= (L_a^\prec + L_{a'}^\prec)(F) \prec L_{a''}^\prec(G) + G(a'')(L_a^\prec + L_{a'}^\prec)(F) + F \prec L_a^\prec(G) + F(a)G \\ &\in U + U \prec U + U \succ U. \end{aligned}$$

In the same way, we can prove that $V = U + U \prec U + U \succ U$ is stable under L_a^\prec , R_a^\prec and R_a^\succ . As U is finite-dimensional, V is finite-dimensional, and contains $f \prec g$ and $f \succ g$, as $f, g \in U$. So, by lemma 42, $f \prec g$ and $f \succ g \in A^*$. \square

By dualising the axioms of bidendriform bialgebras, we obtain:

Proposition 45 *For any bidendriform bialgebra A , $(A^*, \Delta_\prec^*, \Delta_\succ^*, \prec^*, \succ^*)$ is a bidendriform bialgebra. More precisely, for all $f, g \in A^*$, $a, b \in A$:*

$$\begin{aligned} (f \prec g)(a) &= (f \otimes g) \circ \Delta_\prec(a), & (f \succ g)(a) &= (f \otimes g) \circ \Delta_\succ(a), \\ \Delta_\prec(f)(a \otimes b) &= f(a \prec b), & \Delta_\succ(f)(a \otimes b) &= f(a \succ b). \end{aligned}$$

Proposition 46 *Let A, B be bidendriform bialgebras.*

1. *Let $\Phi : A \longrightarrow B^*$ be a morphism of bidendriform bialgebras. Then the following pairing is a bidendriform pairing:*

$$\langle, \rangle : \begin{cases} A \times B & \longrightarrow K \\ (a, b) & \longrightarrow \langle a, b \rangle = \Phi(a)(b) \end{cases}$$

2. *Suppose that A is connected and that $\langle, \rangle : A \times B \longrightarrow K$ is a bidendriform pairing. Then the following application is a morphism of bidendriform bialgebras:*

$$\Phi : \begin{cases} A & \longrightarrow B^* \\ a & \longrightarrow \Phi(a) : \begin{cases} B & \longrightarrow K \\ b & \longrightarrow \langle a, b \rangle \end{cases} \end{cases}.$$

Proof.

1. Let $a_1, a_2 \in A$, $b \in B$.

$$\begin{aligned} \langle a_1 \prec a_2, b \rangle &= \Phi(a_1 \prec a_2)(b) \\ &= (\Phi(a_1) \prec \Phi(a_2))(b) \\ &= (\Phi(a_1) \otimes \Phi(a_2)) \circ \Delta_\prec(b) \\ &= \langle a_1 \otimes a_2, \Delta_\prec(b) \rangle. \end{aligned}$$

Let $a \in A$, $b_1, b_2 \in B$.

$$\begin{aligned} \langle a, b_1 \prec b_2 \rangle &= \Phi(a)(b_1 \prec b_2) \\ &= \Delta_\prec(\Phi(a))(b_1 \otimes b_2) \\ &= ((\Phi \otimes \Phi) \circ \Delta_\prec(a))(b_1 \otimes b_2) \\ &= \langle \Delta_\prec(a), b_1 \otimes b_2 \rangle. \end{aligned}$$

The two other equalities are proved in the same way.

2. We first prove the following lemma:

Lemma 47 *Let C be a connected dendriform coalgebra. For all $a \in C$, there exists a finite-dimensional dendriform subcoalgebra of C which contains a .*

Proof. We use the notations of definition 18. Let $a \in C$. We put:

$$C_a = Vect \left\{ \begin{array}{l} (f_1 \otimes \dots \otimes f_{i-1} \otimes Id \otimes f_{i+1} \otimes \dots \otimes f_n) \circ P(a) / \\ n \in \mathbb{N}^*, i \in \{1, \dots, n\}, f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_n \in C^*, P \in \mathcal{P}_C(n) \end{array} \right\}.$$

Then C_a is the smallest dendriform subcoalgebra of C which contains a . As C is connected, there are only a finite number of P such that $P(a) \neq 0$, so C_a is finite-dimensional. \square

End of the proof of proposition 46. Let us first prove that $\Phi(a) \in B^*$ for all $a \in A$. As A is connected, there exists a finite-dimensional subcoalgebra C of A which contains a (lemma 47). Let I be the orthogonal of C for the pairing \langle, \rangle . As C is finite-dimensional, I has a finite codimension. Let $b_1 \in B, b_2 \in I$. For all $c \in C$, we have:

$$\langle c, b_1 \prec b_2 \rangle = \langle \underbrace{\Delta_{\prec}(c)}_{\in C \otimes C}, b_1 \otimes b_2 \rangle = 0,$$

so $b_1 \prec b_2 \in I$. In the same way, we prove that I is a dendriform ideal of B . As $a \in C$, $\Phi(a)(I) = \langle a, I \rangle = (0)$, so $\Phi(a) \in B^*$.

Let $a_1, a_2 \in A$. For all $b \in B$:

$$\begin{aligned} \Phi(a_1 \prec a_2)(b) &= \langle a_1 \prec a_2, b \rangle \\ &= \langle a_1 \otimes a_2, \Delta_{\prec}(b) \rangle \\ &= (\Phi(a_1) \otimes \Phi(a_2)) \circ \Delta_{\prec}(b) \\ &= (\Phi(a_1) \prec \Phi(a_2))(b), \end{aligned}$$

so $\Phi(a_1 \prec a_2) = \Phi(a_1) \prec \Phi(a_2)$. Let $a \in A$. For all $b_1, b_2 \in B$:

$$\begin{aligned} (\Delta_{\prec}(\Phi(a)))(b_1 \otimes b_2) &= \Phi(a)(b_1 \prec b_2) \\ &= \langle a, b_1 \prec b_2 \rangle \\ &= \langle \Delta_{\prec}(a), b_1 \otimes b_2 \rangle \\ &= ((\Phi \otimes \Phi) \circ \Delta_{\prec}(a))(b_1 \otimes b_2), \end{aligned}$$

so $\Delta_{\prec}(\Phi(a)) = (\Phi \otimes \Phi) \circ \Delta_{\prec}(a)$. The other equalities are proved in the same way. \square

5.3 Bidendriform pairing on a connected graded bidendriform bialgebra

Theorem 48 *Let A, B be connected bidendriform bialgebras. Let $\langle, \rangle_{|Prim_{tot}}: Prim_{tot}(A) \times Prim_{tot}(B) \rightarrow K$. then $\langle, \rangle_{|Prim_{tot}}$ can be uniquely extended into a bidendriform pairing between A and B . Moreover:*

1. \langle, \rangle is non-degenerate $\iff \langle, \rangle_{|Prim_{tot}}$ is non-degenerate.
2. Suppose that $A = B$. Then \langle, \rangle is symmetric $\iff \langle, \rangle_{|Prim_{tot}}$ is symmetric.
3. Suppose that A and B are \mathbb{N} -graded (we do not suppose that the A_n 's or the B_n 's are finite-dimensional). Then \langle, \rangle is homogeneous $\iff \langle, \rangle_{|Prim_{tot}}$ is homogeneous.

Proof.

Existence. By corollary 25, $B = Prim_{tot}(B) \oplus B^{D2}$. We can then define, for all $p \in Prim_{tot}(A)$:

$$f_p : \begin{cases} B & \longrightarrow K \\ q \in Prim_{tot}(B) & \longrightarrow \langle p, q \rangle_{|Prim_{tot}}, \\ b \in B^{D2} & \longrightarrow 0. \end{cases}$$

Then $\text{Ker}(f_p)$ contains B^{D^2} , so is a dendriform ideal of codimension at most 1. Hence, $f_p \in B^*$. Moreover, for all $a_1, a_2 \in A$, $\Delta_{\prec}(f_p)(a_1 \otimes a_2) = f_p(a_2 \prec a_1) = 0$, so $\Delta_{\prec}(f_p) = 0$. In the same way, $\Delta_{\succ}(f_p) = 0$ and then $f_p \in \text{Prim}_{\text{tot}}(B^*)$.

As A is isomorphic to a bidendriform algebra \mathcal{A}^D (corollary 36), by theorem 35, there exists a unique morphism of bidendriform bialgebras:

$$\Phi : \begin{cases} A & \longrightarrow B^* \\ p \in \text{Prim}_{\text{tot}}(A) & \longrightarrow f_p. \end{cases}$$

We then put $\langle a, b \rangle = \Phi(a)(b)$ for all $a \in A$, $b \in B$. This pairing extends $\langle, \rangle|_{\text{Prim}_{\text{tot}}}$: if $p \in \text{Prim}_{\text{tot}}(A)$, $q \in \text{Prim}_{\text{tot}}(B)$, $\langle p, q \rangle = \Phi(p)(q) = f_p(q) = \langle p, q \rangle|_{\text{Prim}_{\text{tot}}}$. Moreover, as Φ is a morphism of bidendriform bialgebras, \langle, \rangle is a bidendriform pairing (proposition 46).

Unicity. Let \langle, \rangle' a second bidendriform pairing which extends $\langle, \rangle|_{\text{Prim}_{\text{tot}}}$. Let us show that it is equal to \langle, \rangle already defined. Consider:

$$A' = \{a \in A / \forall b \in B, \langle a, b \rangle = \langle a, b \rangle'\}.$$

Let $a_1, a_2 \in A'$. Then, for all $b \in B$,

$$\langle a_1 \prec a_2, b \rangle = \langle a_1 \otimes a_2, \Delta_{\prec}(b) \rangle = \langle a_1 \otimes a_2, \Delta_{\prec}(b) \rangle' = \langle a_1 \prec a_2, b \rangle'.$$

So $a_1 \prec a_2 \in A'$. In the same way, $a_1 \succ a_2 \in A'$, so A' is a dendriform subalgebra. As A is generated by $\text{Prim}_{\text{tot}}(A)$, it is enough to prove that $p \in A'$ for all $p \in \text{Prim}_{\text{tot}}(A)$. Let $b \in B$ and let us show that $\langle p, b \rangle = \langle p, b \rangle'$. It is true by hypothesis if $b \in \text{Prim}_{\text{tot}}(A)$. If it is not the case, we can suppose $b = b_1 \prec b_2$ or $b = b_1 \succ b_2$. Hence:

$$\begin{aligned} \langle p, b \rangle' &= \langle p, b_1 \prec b_2 \rangle' = \langle \Delta_{\prec}(p), b_1 \otimes b_2 \rangle' = 0 = \langle p, b \rangle, \\ \text{or } \langle p, b \rangle' &= \langle p, b_1 \succ b_2 \rangle' = \langle \Delta_{\succ}(p), b_1 \otimes b_2 \rangle' = 0 = \langle p, b \rangle. \end{aligned}$$

We now prove the three equivalences.

Non-degeneracy, \implies . Suppose \langle, \rangle non-degenerate and let $a \in \text{Prim}_{\text{tot}}(A) \cap \text{Prim}_{\text{tot}}(B)^\perp$. Then, as $\text{Prim}_{\text{tot}}(A) \subseteq (B^{D^2})^\perp$, we obtain $a \in (B^{D^2} + \text{Prim}_{\text{tot}}(B))^\perp = B^\perp = (0)$. In the same way, $\text{Prim}_{\text{tot}}(B) \cap \text{Prim}_{\text{tot}}(A)^\perp = (0)$, So $\langle, \rangle|_{\text{Prim}_{\text{tot}}}$ is non-degenerate.

Non-degeneracy, \impliedby . Then $\Phi|_{\text{Prim}_{\text{tot}}(A)}$ is injective, so, by theorem 35, Φ is injective: $B^\perp = (0)$. By permuting the role of A and B , which does not change the pairing \langle, \rangle by unicity, we obtain $A^\perp = (0)$. So \langle, \rangle is non-degenerate.

Symmetry, \implies . Obvious.

Symmetry, \impliedby . We consider $A'' = \{a \in A / \forall b \in A, \langle a, b \rangle = \langle b, a \rangle\}$. Let $a_1, a_2 \in A''$. Then, for all $b \in A$:

$$\langle a_1 \prec a_2, b \rangle = \langle a_1 \otimes a_2, \Delta_{\prec}(b) \rangle = \langle \Delta_{\prec}(b), a_1 \otimes a_2 \rangle = \langle b, a_1 \prec a_2 \rangle.$$

So $a_1 \prec a_2 \in A''$. In the same way, $a_1 \succ a_2 \in A''$, so A'' is a dendriform subalgebra. It is then enough to show that $\text{Prim}_{\text{tot}}(A) \subset A''$. Let $a \in \text{Prim}_{\text{tot}}(A)$ and $b \in A$. We can limit ourselves to the three following cases:

1. $b \in \text{Prim}_{\text{tot}}(A)$. Then $\langle a, b \rangle = \langle b, a \rangle$ by hypothesis on $\langle, \rangle|_{\text{Prim}_{\text{tot}}}$.
2. $b = b_1 \prec b_2$. Then $\langle b, a \rangle = \langle b_1 \otimes b_2, \Delta_{\prec}(a) \rangle = 0 = \langle a, b \rangle$.
3. $b = b_1 \succ b_2$. Then $\langle b, a \rangle = \langle b_1 \otimes b_2, \Delta_{\succ}(a) \rangle = 0 = \langle a, b \rangle$.

So $A'' = A$ and \langle, \rangle is symmetric.

Homogeneity, \implies . Obvious.

Homogeneity, \impliedby . Then Φ takes its values in the graded dual of A and is homogeneous of degree 0, so \langle, \rangle is homogeneous. \square

5.4 Application to $\mathcal{H}^{\mathcal{D}}$

We define a pairing on $Prim_{tot}(\mathcal{A}^{\mathcal{D}})$ in the following way: for all $d, d' \in \mathcal{D}$,

$$\langle \cdot_d, \cdot_{d'} \rangle_{Prim_{tot}(\mathcal{A}^{\mathcal{D}})} = \delta_{d,d'}.$$

This pairing is symmetric, homogeneous, and non-degenerate. By theorem 48, it can be extended on a unique bidendriform pairing \langle, \rangle on $\mathcal{A}^{\mathcal{D}}$. This pairing is also symmetric, homogeneous, and non-degenerate.

Remark. This pairing is not the pairing $(,)$ on $\mathcal{H}^{\mathcal{D}}$ introduced in [7], which is a symmetric and non degenerate Hopf pairing but not a bidendriform pairing:

$$(\cdot_d \prec \cdot_d, \mathbf{1}_d^d) = (\mathbf{1}_d^d, \mathbf{1}_d^d) = 0,$$

$$(\cdot_d \otimes \cdot_d, \Delta_{\prec}(\mathbf{1}_d^d)) = (\cdot_d \otimes \cdot_d, \cdot_d \otimes \cdot_d) = 1.$$

The pairing \langle, \rangle admits a similar description as the pairing of [7]. Let $d \in \mathcal{D}$. We consider the following application:

$$\xi_d : \begin{cases} \mathcal{H}^{\mathcal{D}} & \longrightarrow \mathcal{H}^{\mathcal{D}} \\ F \in \mathbb{F}^{\mathcal{D}} & \longrightarrow \begin{cases} 0 \text{ if the leave of } F \text{ at most on the right is decorated by } d; \\ F \text{ without its leave at most on the right if it is decorated by } d. \end{cases} \end{cases}$$

For example, if $a, b, c, d, e, f \in \mathcal{D}$, $\xi_f(\mathbf{1}_a^b \mathbf{V}_c^e) = \delta_{e,f} \mathbf{1}_a^b \mathbf{1}_c^d$.

Lemma 49 *Let $x, y \in \mathcal{A}^{\mathcal{D}}$, $d \in \mathcal{D}$. Then:*

1. $\langle \cdot_d \otimes x, \Delta_{\prec}(y) \rangle = \langle x, \xi_d(y) \rangle$.
2. $\langle B_d^+(x), y \rangle = \langle x, \xi_d(y) \rangle$.

Proof.

1. We can suppose that $y \in \mathbb{F}^{\mathcal{D}}$. We denote by d' the decoration of leave of y at most on the right. The only eventual admissible cut in $Adm_{\prec}(y)$ (defined in proposition 33) such that $P^c(y) = \cdot_d$ is the cut c_y which cut the edge leading to the leave of y at most on the right, if $d = d'$. Hence, by proposition 33:

$$\begin{aligned} \langle \cdot_d \otimes x, y \rangle &= \sum_{c \in Adm_{\prec}(y)} \langle \cdot_d, P^c(y) \rangle \langle x, R^c(y) \rangle \\ &= \sum_{c \in Adm_{\prec}(y)} \delta_{\cdot_d, P^c(y)} \langle x, R^c(y) \rangle \\ &= \delta_{d,d'} \langle x, R^{c_f}(y) \rangle \\ &= \langle x, \xi_d(y) \rangle. \end{aligned}$$

2. We have $\langle B_d^+(x), y \rangle = \langle \cdot_d \prec x, y \rangle = \langle \cdot_d \otimes x, \Delta_{\prec}(y) \rangle = \langle x, \xi_d(y) \rangle$. \square

Hence, the Hopf pairing (also denoted by \langle, \rangle) induced on $\mathcal{H}^{\mathcal{D}}$ satisfies the following properties:

Theorem 50 $\langle, \rangle: \mathcal{H}^{\mathcal{D}} \times \mathcal{H}^{\mathcal{D}} \longrightarrow K$ is the unique pairing such that:

1. $\forall y \in \mathcal{H}^{\mathcal{D}}, \langle 1, y \rangle = \varepsilon(y)$, where ε is the counit of $\mathcal{H}^{\mathcal{D}}$.
2. $\forall x, y \in \mathcal{H}^{\mathcal{D}}, \langle B_d^+(x), y \rangle = \langle x, \xi_d(y) \rangle$.
3. $\forall x, y, z \in \mathcal{H}^{\mathcal{D}}, \langle xy, z \rangle = \langle x \otimes y, \Delta(z) \rangle$.

Moreover, \langle, \rangle is symmetric, homogenous and non-degenerate, and for all $x, y \in \mathcal{H}^{\mathcal{D}}$, we have $\langle S(x), y \rangle = \langle x, S(y) \rangle$.

Remark. The three points of this theorem allow to compute $\langle F, G \rangle$ for $F, G \in \mathbb{F}^{\mathcal{D}}$, by induction on $weight(F)$.

We now give a combinatorial interpretation of this pairing, inspired by our work in [7]. If $F \in \mathbb{F}^{\mathcal{D}}$, We have two partial orders \geq_{high} et \geq_{left} on the set $vert(F)$ of vertices of F (see [7] for more details). if x, y are two vertices of F , then:

1. We will say that $x \geq_{high} y$ if there is a path from y to x .
2. If s and s' are not comparable for \geq_{high} , then one of this two vertices (say for example x) is more on the left than the other: we will denote this situation by $x \geq_{left} y$.

These partial orders induce a total order $\geq_{d,l}$ on $vert(F)$ defined by:

$$x \geq_{d,l} y \text{ if } (x \geq_{left} y) \text{ or } (y \geq_{high} x).$$

Theorem 51 *Let $F, G \in \mathbb{F}^{\mathcal{D}}$. Let $\mathcal{I}(F, G)$ be the set of bijections f from $vert(F)$ to $vert(G)$ such that:*

1. $\forall x, y \in vert(F), x \geq_{high} y \Rightarrow f(x) \geq_{d,l} f(y)$,
2. $\forall x, y \in vert(F), f(x) \geq_{high} f(y) \Rightarrow x \geq_{d,l} y$,
3. $\forall x \in vert(F), x$ and $f(x)$ have the same decoration.

Then $\langle F, G \rangle = card(\mathcal{I}(F, G))$.

Proof. This is true if $weight(F) \neq weight(G)$, as then $\langle F, G \rangle = 0$ and there is no bijection from $vert(F)$ to $vert(G)$. We suppose that $weight(F) = weight(G) = n$, and let us proceed by induction on n . If $n = 1$, then $\langle \cdot_d, \cdot_{d'} \rangle = \delta_{d,d'} = card(\mathcal{I}(\cdot_d, \cdot_{d'}))$ by condition 3. Suppose that the property is satisfied for all $k < n$ and let $F, G \in \mathbb{F}^{\mathcal{D}}$ with n vertices. We put $F = t_1 \dots t_m$.

If $m = 1$: then $F = t_1 = B_d^+(F')$. So $\langle F, G \rangle = \langle F', \xi_d(G) \rangle$. Let $f \in \mathcal{I}(F, G)$. Denote by r the root of F . Remark that for all $x \in vert(F)$, $x \geq_{high} r$. By condition 1, $f(r)$ is the smallest element of $vert(G)$ for $\geq_{d,l}$: it is the leave of G at most on the right. If it is not decorated by d , then $\langle F, G \rangle = 0$ and $\mathcal{I}(F, G) = \emptyset$ by condition 3. Otherwise, there is a bijection $\mathcal{I}(F, G) \rightarrow \mathcal{I}(F', \xi_d(G))$, sending f to its restriction à $vert(F')$. As $\langle F, G \rangle = \langle F', \xi_d(G) \rangle$, we have the result.

If $m > 1$: we put $F' = t_1 \dots t_{m-1}$. Let $f \in \mathcal{I}(F, G)$; consider $f(som(t_m))$. Let y_2 be a vertex of G , such that there exists $x_1 \in vert(t_m)$, $f(x_1) \geq_{high} y_2$. There exists a unique $x_2 \in som(F)$, such that $f(x_2) = y_2$. By condition 2, $x_1 \geq_{d,l} x_2$ and then $x_2 \in som(t_m)$. Hence, there exists a unique admissible cut c_f of G such that $R^{c_f}(G) = f(t_m)$ and $P^{c_f}(F) = f(F')$. We then have a bijection:

$$\begin{cases} \mathcal{I}(F, G) & \longrightarrow \bigcup_{c \in Adm(G)} \mathcal{I}(F', P^c(G)) \times \mathcal{I}(t_m, R^c(G)) \\ f & \longrightarrow (f|_{som(F')}, f|_{som(t_m)}) \in \mathcal{I}(F', P^{c_f}(G)) \times \mathcal{I}(t_m, R^{c_f}(G)). \end{cases}$$

Hence:

$$\begin{aligned} \langle F, G \rangle &= \sum_{c \in Adm(G)} \langle F', P^c(G) \rangle \langle t_m, R^c(G) \rangle \\ &= \sum_{c \in Adm(G)} \langle F', P^c(G) \rangle \langle t_m, R^c(G) \rangle \end{aligned}$$

$$\begin{aligned}
&= \sum_{c \in \text{Adm}(G)} \text{card}(\mathcal{I}(F', P^c(G))) \text{card}(\mathcal{I}(t_m, R^c(G))) \\
&= \text{card} \left(\bigcup_{c \in \text{Adm}(G)} \mathcal{I}(F', P^c(G)) \times \mathcal{I}(t_m, R^c(G)) \right) \\
&= \text{card}(\mathcal{I}(F, G)). \square
\end{aligned}$$

We consider now the case where \mathcal{D} is reduced to a single element. We can identify $\mathcal{H}^{\mathcal{D}}$ and \mathcal{H} . We will denote B^+ instead of B_d^+ , ξ instead of ξ_d . We define $l_k \in \mathbb{F}$ by induction on k in the following way: $l_0 = 1$, $l_{k+1} = B^+(l_k)$. For example:

$$l_1 = \bullet, l_2 = \mathbf{1}, l_3 = \mathbf{1}, l_4 = \mathbf{1}, l_5 = \mathbf{1} \dots$$

Proposition 52 *Let $F \in \mathbb{F}$, of weight n . Then:*

1. $\langle F, l_n \rangle = 1$.
2. $\langle F, l_{n-1} \bullet \rangle$ is the number of roots of F .
3. $\langle F, \bullet l_{n-1} \rangle$ is the number of leaves of F .

Proof.

1. Induction on n . If $n = 0$, then $F = 1$ and $\langle F, l_0 \rangle = \langle 1, 1 \rangle = 1$. Suppose that the result is true for all forest G of weight $n - 1$. Then $\langle F, l_n \rangle = \langle \xi(F), l_{n-1} \rangle = 1$ as $\xi(F) \in \mathbb{F}$.

2. Let X be the set of $F \in \mathbb{F}$, $F \neq 1$, satisfying 2. Let $G \in \mathbb{F}$, with n vertices. We have $\langle B^+(G), l_n \bullet \rangle = \langle G, \xi(l_n \bullet) \rangle = \langle G, l_n \rangle = 1$, so $B^+(G) \in X$: X contents the trees.

Let $F_1, F_2 \in X$, with respectively n_1 and n_2 vertices. We put $n = n_1 + n_2$. We have:

$$\begin{aligned}
\langle F_1 F_2, l_{n-1} \bullet \rangle &= \langle F_1 \otimes F_2, \Delta(l_{n-1} \bullet) \rangle \\
&= \sum_{i+j=n-1} \langle F_1 \otimes F_2, l_i \bullet \otimes l_j + l_i \otimes l_j \bullet \rangle \\
&= \langle F_1 \otimes F_2, l_{n_1-1} \bullet \otimes l_{n_2} + l_{n_1} \otimes l_{n_2-1} \bullet \rangle \text{ by homogeneity,} \\
&= \langle F_1, l_{n_1-1} \bullet \rangle + \langle F_2, l_{n_2-1} \bullet \rangle.
\end{aligned}$$

So $F_1 F_2 \in X$. As X contents the trees, $X = \mathbb{F} - \{1\}$.

3. Let Y be the set of $F \in \mathbb{F}$, $F \neq 1$, satisfying 3. We have $\langle \bullet, \bullet l_0 \rangle = \langle \bullet, \bullet \rangle = 1$, so $\bullet \in Y$.

Let $G \in Y$, with n vertices. We have $\langle B^+(G), \bullet l_n \rangle = \langle G, \xi(\bullet l_n) \rangle = \langle G, \bullet l_{n-1} \rangle$. As G and $B^+(G)$ have the same number of leaves, $B^+(G) \in Y$, so Y is stable under B^+ .

Let $F_1, F_2 \in Y$, with respectively n_1 and n_2 vertices. We put $n = n_1 + n_2$. We have:

$$\begin{aligned}
\langle F_1 F_2, \bullet l_{n-1} \rangle &= \langle F_1 \otimes F_2, \Delta(\bullet l_{n-1}) \rangle \\
&= \sum_{i+j=n-1} \langle F_1 \otimes F_2, \bullet l_i \otimes l_j + l_i \otimes \bullet l_j \rangle \\
&= \langle F_1 \otimes F_2, \bullet l_{n_1-1} \otimes l_{n_2} + l_{n_1} \otimes \bullet l_{n_2-1} \rangle \text{ by homogeneity,} \\
&= \langle F_1, \bullet l_{n_1-1} \rangle + \langle F_2, \bullet l_{n_2-1} \rangle.
\end{aligned}$$

So $F_1 F_2 \in Y$. As Y is stable under product and under B^+ and contents \bullet , $Y = \mathbb{F} - \{1\}$. \square

Examples. Here are the values of the pairing \langle, \rangle of \mathcal{H} on forests of weight smaller than 4:

$\begin{array}{c c} & \cdot \\ \cdot & 1 \end{array}$	$\begin{array}{c cc} & \cdot & \cdot \\ \cdot & 2 & 1 \\ \cdot & 1 & 1 \end{array}$
---	---

...	6	3	3	2	1
\cdot\cdot	3	2	2	2	1
\cdot\cdot	3	2	2	1	1
\vee	2	2	1	1	1
\vdash	1	1	1	1	1

	\cdot\cdot\cdot	\cdot\cdot	\vee\cdot	\vdash\cdot	\cdot\cdot\cdot	\cdot\cdot\cdot	\cdot\vee	\cdot\vdash	\vee\vee	\vee\vdash	\vee\vee	\vee\vee	\vee\vee
....	24	12	12	8	4	12	6	8	4	6	3	3	2	1
\cdot\cdot\cdot	12	7	7	4	3	7	4	4	3	3	2	2	1	1
\cdot\cdot	12	7	7	5	3	7	4	5	3	3	2	2	2	1
\vee\cdot	8	4	5	3	2	6	3	3	3	2	1	2	1	1
\vdash\cdot	4	3	3	2	2	3	2	2	2	1	1	1	1	1
\cdot\cdot\cdot	12	7	7	6	3	7	4	6	3	6	3	3	2	1
\cdot\cdot\cdot	6	4	4	3	2	4	3	3	2	3	2	2	1	1
\cdot\vee	8	4	5	3	2	6	3	5	3	3	2	2	2	1
\cdot\vdash	4	3	3	3	2	3	2	3	2	3	2	2	2	1
\vee\vee	6	3	3	2	1	6	3	3	3	2	1	2	1	1
\vee\vdash	3	2	2	1	1	3	2	2	2	1	1	1	1	1
\vee\vee	3	2	2	2	1	3	2	2	2	2	1	2	1	1
\vee\vee	2	1	2	1	1	2	1	2	2	1	1	1	1	1
\vdash\vdash	1	1	1	1	1	1	1	1	1	1	1	1	1	1

References

- [1] E. Abe, *Hopf algebras*, Cambridge Tracts in Mathematics, 74, Cambridge University Press, Cambridge-New York, 1980.
- [2] M. Aguiar, *Infinitesimal bialgebras, pre-lie and dendriform algebras*, Lecture Notes in Pure and Appl. Math., no. 237, Dekker, New York, 2004, math.QA/02 11074.
- [3] Marcelo Aguiar and Frank Sottile, *Structure of the Malvenuto-Reutenauer Hopf algebra of permutations*, Adv. Math. **191** (2005), no. 2, 225–275, math.CO/02 03282.
- [4] A. Connes and D. Kreimer, *Hopf algebras, Renormalization and Noncommutative geometry*, Comm. Math. Phys **199** (1998), no. 1, 203–242, hep-th/98 08042.
- [5] G. Duchamp, F. Hivert, and J.-Y. Thibon, *Some generalizations of quasi-symmetric functions and noncommutative symmetric functions*, Springer, Berlin, 2000, math.CO/01 05065.
- [6] L. Foissy, *Les algèbres de Hopf des arbres enracinés décorés*, Thèse de doctorat, Université de Reims, 2002.
- [7] ———, *Les algèbres de Hopf des arbres enracinés, I*, Bull. Sci. Math. **126** (2002), 193–239.
- [8] ———, *Les algèbres de Hopf des arbres enracinés, II*, Bull. Sci. Math. **126** (2002), 249–288.
- [9] R. Holtkamp, *Comparison of Hopf Algebras on Trees*, Arch. Math. (Basel) **80** (2003), no. 4, 368–383.

- [10] A. Joseph, *Quantum groups and their primitive ideals*, Springer-Verlag, 1995.
- [11] D. Kreimer, *On the Hopf algebra structure of perturbative quantum field theories*, Adv. Theor. Math. Phys. **2** (1998), no. 2, 303–334, q-alg/97 07029.
- [12] ———, *On Overlapping Divergences*, Comm. Math. Phys. **204** (1999), no. 3, 669–689, hep-th/98 10022.
- [13] ———, *Combinatorics of (perturbative) Quantum Field Theory*, Phys. Rep. **4–6** (2002), 387–424, hep-th/00 10059.
- [14] J.-L. Loday, *Generalized bialgebras and triples of operads*, available at <http://www-irma.strasbg.fr/~loday/>.
- [15] ———, *Dialgebras*, Lecture Notes in Math., no. 1763, Springer, Berlin, 2001, math.QA/01 02053.
- [16] J.-L. Loday and M. Ronco, *Hopf algebra of the planar binary trees*, Adv. Math. **139** (1998), no. 2, 293–309.
- [17] ———, *Order structure on the algebra of permutations and of planar binary trees*, J. Algebraic Combin. **15** (2002), no. 3, 253–270, math.CO/01 02066.
- [18] C. Malvenuto and Ch. Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, J. Algebra **177** (1995), no. 3, 967–982.
- [19] M. Markl, S. Shnider, and J. Stasheff, *Operads in algebra, topology and physics*, Mathematical Surveys and Monographs, 96, American Mathematical Society, Providence, RI, 2002.
- [20] J.W. Milnor and J. C. Moore, *On the structure of Hopf algebras*, Ann. of Math. (2) **81** (1965), 211–264.
- [21] M. Ronco, *Primitive elements of a free dendriform algebra*, Contemp. Math. **267** (2000), 245–263.
- [22] ———, *Eulerian idempotents and Milnor-Moore theorem for certain non-cocommutative Hopf algebras*, J. Algebra **254** (2002), no. 1, 152–172.