



HAL
open science

Transmission through correlated fermion systems and size of the corresponding effective one-body scatterer

Rafael A. Molina, Dietmar Weinmann, Jean-Louis Pichard

► **To cite this version:**

Rafael A. Molina, Dietmar Weinmann, Jean-Louis Pichard. Transmission through correlated fermion systems and size of the corresponding effective one-body scatterer. *The European Physical Journal B: Condensed Matter and Complex Systems*, 2005, 48, pp.243. 10.1140/epjb/e2005-00403-1 . hal-00007669v1

HAL Id: hal-00007669

<https://hal.science/hal-00007669v1>

Submitted on 25 Jul 2005 (v1), last revised 24 Jan 2006 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Transmission through correlated fermion systems and size of the corresponding effective one-body scatterer

RAFAEL A. MOLINA^{1,2}, DIETMAR WEINMANN³ and JEAN-LOUIS PICHARD^{1,4}

¹ *CEA/DSM, Service de Physique de l'Etat Condensé, Centre d'Etudes de Saclay, 91191 Gif-sur-Yvette, France*

² *Max-Planck-Institut für Physik Komplexer Systeme, Nöthnitzer Str. 38, 01187 Dresden, Germany*

³ *Institut de Physique et Chimie des Matériaux de Strasbourg, UMR 7504 (CNRS-ULP), 23 rue du Loess, BP 43, 67034 Strasbourg Cedex 2, France*

⁴ *Laboratoire de Physique Théorique et Modélisation, Université de Cergy-Pontoise, 95031 Cergy-Pontoise Cedex, France*

PACS. 71.27.+a – Strongly correlated electron systems; heavy fermions.

PACS. 71.10.-d – Theory of electronic transport; scattering mechanisms.

PACS. 73.23.-b – Electronic transport in mesoscopic systems.

Abstract. – In order to extend the Landauer formulation of quantum transport to spinless correlated fermions, we consider a system in which charge carriers interact, connected to two reservoirs by non-interacting one-dimensional leads. We study the size of the attached leads necessary for reducing this many-body scatterer to an effective one-body scatterer with interaction-dependent parameters. To obtain this size, we consider two identical correlated systems connected by a non-interacting lead of length L_C . We demonstrate that the effective one-body transmission of the ensemble deviates by an amount A/L_C from the behavior obtained assuming an effective one-body description for each element and the combination law of scatterers in series. A is maximum for the interaction strength U around which the Luttinger liquid becomes a Mott insulator in the used model, and vanishes when $U \rightarrow 0$ and $U \rightarrow \infty$. Analogies with the Kondo problem are pointed out.

In Landauer's formulation of quantum transport [1], the measure of the conductance g of a coherent system is formulated as a scattering problem between incoherent electron reservoirs. In a two-probe geometry, the system is connected to two reservoirs via leads. For large electron densities inside the system and its attached leads, the Coulomb interaction is screened and the Coulomb to kinetic energy ratio r_s is small. One has essentially a non-interacting system of Fermi energy E_F , where the occupation of the one-body levels is given by a Fermi-Dirac distribution at a temperature T . The system acts as a one-body scatterer and its residual conductance $g(T \rightarrow 0)$ is given (in units of $2e^2/h$ for single channel leads and spin degeneracy) by the probability $|t(E_F)|^2$ of an electron of energy E_F to be elastically transmitted through it.

The problem of describing coherent electronic transport becomes much more complex in the case where the carrier density is low inside the scatterer, the screening ceasing to




Fig. 1 – Left: Scheme of the ring pierced by a flux ϕ used for the embedding method. The correlation cloud induced by L_S interacting sites upon the auxiliary lead is sketched in grey. Right: Amplitude $B(U, L_S)$ characterizing the $1/L$ finite size correction to the charge stiffness.

be effective and the electrons becoming correlated. Such situations occur in quantum point contacts of transverse size smaller than the Fermi wavelength, where a $0.7 (2e^2/h)$ structure is observed [2], and can be expected in molecules [3], atomic chains or contacts [4], quantum dots where few electrons might form a correlated solid, a charge density wave, a Mott insulator, etc. In these cases, the electrons are transmitted from one Fermi reservoir to another through a many-body scatterer.

To extend Landauer's approach to such systems, at least for low temperatures and bias voltages, one needs to reduce the bare many-body scatterer to an effective one-body scatterer with interaction-dependent parameters. This task will be hopeless for an isolated system where electrons interact with a large interaction strength U , but becomes possible when leads where electrons do not interact are attached to it. This has been numerically demonstrated in previous works [5,6] using the embedding method, which allows to extract [5–11] the effective coefficient $|t(E_F, U)|^2$ from the persistent current of a large non-interacting ring embedding the many-body scatterer.

Using the same method, we show in this letter that it is not the region where the electrons interact which acts as an effective one-body scatterer with renormalized parameters, but a larger region where the many-body scatterer induces correlations. This problem is somewhat similar to the Kondo problem, which can be solved using Wilson's numerical renormalization group (NRG) [12]. Instead of using the NRG method, we use the density matrix renormalization group (DMRG) method [13, 14] for a non-interacting ring embedding the many-body scatterer. The modulus of the effective one-body transmission amplitude $|t(E_F, U)|$ is obtained from the persistent current of the ring extrapolated to infinite lead length, while its phase α is given by the Friedel sum rule. To study the size of the region which acts as an effective one-body scatterer, we consider two identical many-body scatterers coupled via a non-interacting lead of size L_C . The effective total transmission coefficient $|t_T(E_F, U)|^2$ of this ensemble is directly determined by the embedding method. Comparing this exact value, calculated as a function of L_C , to the value obtained assuming the combination law of one-body scatterers in series, the existence of induced correlations in the attached leads can be detected and their role studied as a function of the interaction strength U .

To study the mapping of a bare many-body scatterer coupled to leads onto an effective one-body scatterer with interaction-dependent coefficients, we take a model of N spinless

fermions in a chain of $L = L_S + L_L$ sites. The Hamiltonian (with even L_L) reads

$$H = -t_h \sum_{i=2}^L (c_i^\dagger c_{i-1} + c_{i-1}^\dagger c_i) + U \sum_{i=L_L/2+2}^{L_L/2+L_S} [n_i - V_+] [n_{i-1} - V_+] . \quad (1)$$

The hopping amplitude $t_h = 1$ between nearest neighbor sites sets the energy scale, c_i (c_i^\dagger) is the annihilation (creation) operator at site i , and $n_i = c_i^\dagger c_i$. The nearest neighbor repulsion U acts upon L_S consecutive sites and gives rise to many-body scattering. We take a half-filled model ($N = L/2$), with a potential $V_+ = 1/2$ being due to a positive background charge which exactly compensates the repulsion U inside the scatterer. Therefore, our model exhibits particle-hole symmetry and a uniform density, without Friedel oscillations around the scattering region where the fermions interact.

The scattering geometry corresponds to two leads of $L_L/2 \rightarrow \infty$ sites connected by an interacting scatterer of L_S sites. The electrons do not interact in the leads, a necessary condition for having appropriate asymptotic scattering channels in one dimension. For the embedding method, we consider the ring geometry sketched in Fig. 1 (left), the scatterer being closed on itself via a non-interacting lead of L_L sites. This is achieved by adding a hopping term $-t_h c_1^\dagger c_L \exp(i\phi) + \text{h.c.}$ to the Hamiltonian (1), the flux ϕ driving a persistent current $J(U)$ in the ring. As the flux dependence of $J(U)$ extrapolated to the limit $L_L \rightarrow \infty$ demonstrates [6], the many-body scatterer behaves as an effective one-body scatterer, but with an interaction-dependent elastic transmission coefficient $|t(E_F, U)|^2$. Instead of using $J(U)$, it is simpler [6] to get $|t(E_F, U)|^2$ from the charge stiffness

$$D(U, L_S, L) = (-1)^N \frac{L}{2} (E_0(U, L_S, L) - E_\pi(U, L_S, L)) , \quad (2)$$

where $E_0(U, L_S, L) - E_\pi(U, L_S, L)$ is the change of the ground-state energy from periodic to antiperiodic boundary conditions. $D(U, L_S, L)$ is obtained by the DMRG implementation for real Hamiltonians, which can be used to study with a great accuracy systems as large as $L = 120$ sites with $N = 60$ particles. In the limit $L_L \rightarrow \infty$, one gets the modulus

$$|t(E_F, U)| = \sin \left(\frac{\pi}{2} \frac{D_\infty(U, L_S)}{D_\infty(U = 0, L_S)} \right) \quad (3)$$

of the transmission amplitude through the scatterer of L_S sites, $D_\infty(U = 0, L_S)$ being the charge stiffness of the same ring for $U = 0$.

To take the limit $L_L \rightarrow \infty$ is one of the key points of the embedding method. This extrapolation is also required for pure one-body scattering, where the finite size corrections to formula (3) can be expanded [6] in powers of $1/L$. For many-body scattering, the DMRG study gives an empirical scaling law [5, 6]

$$D(U, L_S, L) = D_\infty(U, L_S) \exp \left(\frac{C(U, L_S)}{L} \right) \quad (4)$$

obtained for large L_L and small L_S , which allows to determine the asymptotic value $D_\infty(U, L_S)$ necessary to obtain $|t(E_F, U)|^2$. Expanding this scaling law gives $D(U, L_S, L) - D_\infty(U, L_S) \approx B(U, L_S)/L$ when L is large enough, where $B(U, L_S) = C(U, L_S)D_\infty(U, L_S)$. This is a power law decay, and not an exponential decay with a characteristic scale above which the finite size correction can be neglected. The amplitude $B(U, L_S)$ is shown in Fig. 1 (right) for different values of L_S . $B(U, L_S) \rightarrow 0$ when $U \rightarrow 0$ (no scattering, total transmission) and when $U \rightarrow \infty$




Fig. 2 – Scheme of the set-up with two identical many-body scatterers connected by L_C sites where the carriers do not interact. When L_C is small, the two correlation clouds sketched in grey overlap and the effective one-body scatterer of the ensemble is not given by the effective one-body scatterer of each element and the combination law of one-body scatterers in series.

(total reflection, $D_\infty(U \rightarrow \infty, L_S) \rightarrow 0$). It exhibits a first maximum around $U \approx 1$ which is almost independent of L_S , followed by a second one which depends on L_S .

However, since adding one-body potentials in the region of the L_S sites yields a finite size correction to $D(U, L_S, L)$ even when $U \rightarrow 0$, the interpretation of these corrections is not straightforward. They do not depend only on the correlations induced in the attached lead by the interaction acting inside the scatterer, but also on more trivial one-body aspects. This leads us to use a more direct method, where the finite size effects are only due to many-body correlations. This consists in taking two identical scatterers connected by a scattering-free lead of size L_C in which the electrons do not interact, as sketched in Fig. 2. Since the scattering channels begin at the first attached sites of the leads when $U = 0$, there is a simple combination law for one-body scatterers in series. To study how this combination law is broken with increasing U when L_C is small allows to show that the size of the effective elastic scatterer is larger than the region where the carriers interact. When $U \neq 0$, the scattering channels begin only asymptotically far from the many-body scatterer.

Without interaction, a scatterer is described at an energy E_F by a unitary scattering matrix S_S which can be written in terms of its reflection and transmission amplitudes r, r' and t, t' as

$$S_S = \begin{pmatrix} r & t' \\ t & r' \end{pmatrix}. \quad (5)$$

The scatterer being symmetric upon time reversal, one has $t = t'$, while $r = r'$ if the scatterer is symmetric upon space inversion. The transfer matrix M_S (giving the flux amplitudes at the right side in terms of the flux amplitudes at the left side) reads

$$M_S = \begin{pmatrix} 1/t & r/t \\ r^*/t^* & 1/t^* \end{pmatrix}. \quad (6)$$

The total Hamiltonian and the parity operator can be simultaneously diagonalized if one has inversion symmetry, to give even and odd standing-wave solutions which can be written as $\psi_i^0 = \cos(ki + \delta_0)$ and $\psi_i^1 = \sin(ki + \delta_1)$ at the right side of the scatterer, and $\psi_i^0 = \cos(ki - \delta_0)$ and $\psi_i^1 = \sin(ki - \delta_1)$ at its left side. The two phase shifts δ_0 and δ_1 are related [15] to t and r by $t = (\exp(2i\delta_0) + \exp(2i\delta_1))/2$ and $r = (\exp(2i\delta_0) - \exp(2i\delta_1))/2$. Due to symmetries, S_S or M_S have only two free parameters: the modulus $|t| = \cos(\delta_0 - \delta_1)$ and the phase $\alpha = \delta_0 + \delta_1$ of the transmission amplitude t , the unitarity of S_S ($|t|^2 + |r|^2 = 1$ and $r/r^* = -t/t^*$) giving r . We can determine $|t|$ by the embedding method. The Friedel sum rule [16] gives α . If one introduces a scatterer with inversion symmetry in the central region of a scattering free lead, this rule states [17] that

$$\alpha = \delta_0 + \delta_1 = \pi N_f \quad (7)$$




Fig. 3 – Conductance $g(L_C)$ of the set-up sketched in Fig. 2 with $L_S = 2$, for $U = 1$ (left) and $U = 2$ (right). The points are obtained with the embedding method for the ensemble. The dashed lines at $g = 0.717$ and $g = 0.357$ give the approximate values of $g = |t|^4/(|t|^2 - 2)^2$, with $|t|$ obtained by the embedding method for a single scatterer. The solid lines are fits with the forms $g(L_C) = 0.7174 - 0.057/L_C$ for $U = 1$ and $g(L_C) = 0.3572 - 0.077/L_C$ for $U = 2$.

for spinless fermions in one dimension. N_f is the number of displaced fermions when the scatterer is introduced in the central region. For a uniform filling factor $\nu = 1/2$, $N_f = L_S/2$, and the phase α reads

$$\alpha = \pi N_f = \frac{\pi L_S}{2} = k_F L_S, \quad (8)$$

where $k_F = \pi/2$ is the Fermi wave number. For the spinless case in one dimension with a uniform density, this simply means that the transmitted wave has N_f changes of sign when one transfers a fermion through a scatterer containing N_f others. This is obvious for $U = 0$ as well as for $U \neq 0$. Using the same rule, the ideal ballistic lead of L_C sites has a modulus $|t(L_C)| = 1$ and a phase $\alpha(L_C) = k_F L_C$. Its transfer matrix reads

$$M_C = \begin{pmatrix} e^{-ik_F L_C} & 0 \\ 0 & e^{ik_F L_C} \end{pmatrix}. \quad (9)$$

The combination law of one-body scatterers in series being a simple matrix multiplication for the transfer matrices, the total transfer matrix $M_T(E_F)$ of the ensemble is given by $M_T = M_S \cdot M_C \cdot M_S$. For the total transmission coefficient $|t_T|^2$ through the ensemble, expressed in terms of the transmission t of each element and of L_C , this gives

$$|t_T|^2 = \frac{|t|^4}{2(1 - |t|^2)(1 + \cos(2k_F L_C - 2\alpha)) + |t|^4}. \quad (10)$$

Since $|t| = 1$ when L_S is odd [5, 18], we consider only the case where L_S is even. Taking $\alpha = \pi L_S/2$, one gets for the Landauer conductance $g = 1$ if L_C is odd and $g = |t|^4/(|t|^2 - 2)^2$ if L_C is even.

Fig. 3 shows the conductance g for two scatterers of $L_S = 2$ sites with an interaction $U = 1$ (left figure) and $U = 2$ (right figure) in series, as a function of the length L_C of the coupling lead. The data points are directly obtained from the embedding method, without assuming a combination law for scatterers in series. One can see resonances where $g = 1$ for odd $L_C = \{1, 3, 5, \dots\}$. For even values of L_C , the dashed lines correspond to the L_C -independent values $|t|^4/(|t|^2 - 2)^2$ implied by Eq. (10), the coefficient $|t|$ being obtained using the embedding method for a single scatterer. Within the accuracy of the extrapolation procedures involved




Fig. 4 – Left: Error δg made when using the combination law of Eq. (10) for having the conductance of the set-up sketched in Fig. 2 with $L_S = 2$ and different values of U , as a function of an even number L_C of connecting sites. The lines give an $A(U, L_S)/L_C$ fit. Right: Amplitude $A(U, L_S)$ as a function of U for different values of L_S .

in the determination of both, the transmission $|t|$ of an individual scatterer and the total conductance g , the result of (10) gives the correct value when $L_C \rightarrow \infty$, but overestimates g for small even values of L_C . The difference $\delta g(L_C) = g(L_C \rightarrow \infty) - g(L_C)$ is shown in Fig. 4 (left) for even L_C at different values of U . For even L_C , $\delta g(L_C)$ decays as a function of L_C as

$$\delta g(L_C) \approx \frac{A(U, L_S)}{L_C}, \quad (11)$$

with an amplitude $A(U, L_S)$ which is shown in Fig. 4 (right) as a function of the interaction strength U . This $1/L_C$ decay is reminiscent of the $1/L$ decay characterizing $D(U, L_S, L) - D_\infty(U, L_S)$ and of the screening at large distances (larger than the Thomas-Fermi screening length) of the potential of a point charge by non-interacting electrons (Friedel oscillations, RKKY interactions ...) in one dimension. This suggests that the decay could be faster for leads of higher dimensions ($1/L_C^d$ decay in d dimensions). The amplitude $A(U, L_S) \rightarrow 0$ when $U \rightarrow 0$ (one-body scatterers) and when $U \rightarrow \infty$. In this latter limit, the scatterers become decoupled from the leads, the energy for an electron to enter or to leave a scatterer being $\propto U$. $A(U)$ is maximum near $U = 2$, a value where in the thermodynamic limit $L_S \rightarrow \infty$ the Luttinger liquid becomes [19] a Mott insulator for spinless fermions.

For all odd values of L_C , the data for the total transmission coincide with the value $g = 1$ obtained from Eq. (10), and $\delta g(L_C) = 0$. The even-odd dependence on the parity of L_C shows that the convergences of the phase α and the modulus $|t|$ of the effective scatterer are characterized by different scales. One has $\alpha = \pi N_f$ across the scatterer, directly on a scale L_S , independently of U , while $|t|$ reaches its asymptotic value on a much larger scale. This is not surprising since α depends on the mean density, while $|t|$ depends on the correlations of its fluctuations. In our model with a compensating background charge, the mean density does not exhibit Friedel oscillations. The correlation clouds which have to be included with the many-body scatterer to form the effective one-body scatterer should not be confused with the screening clouds characterizing the charge density.

To obtain the effective one-body matrix $S(E_F)$ of a correlated system of spinless fermions is a problem which displays a certain similarity with the Kondo problem of a spin degree of freedom surrounded by a metallic host. In the two cases, it is crucial to couple the many-body system to non-interacting conduction electrons. For the Kondo problem, the original

three-dimensional model can be mapped onto an equivalent non-interacting one-dimensional tight-binding model embedding a Hubbard impurity. In Wilson's renormalization group transformations [12], the embedded many-body Hamiltonian is progressively mapped onto an effective one-body Hamiltonian describing the low energy states. In this transformation, the coupling between different length scales is taken into account progressively, working out from the impurity to the longer length scales and lower energies. The states at sites near the impurity involve conduction states spanning the full band width $2t_h$, while the states located far from the impurity involve conduction states near the Fermi level, with a progressively reduced band width. This NRG method has been used recently [20] to calculate the effective one-body Hamiltonian of a few Hubbard sites embedded in a non-interacting chain, and the corresponding phase shifts.

In our spinless case, the many-body scatterer would give rise to inelastic and elastic transmission and reflection amplitudes if it was taken in the vacuum. It is due to the non-interacting leads which are attached to it, that inelastic processes become progressively blocked by a Fermi vacuum which takes place in the leads, at a large distance from the scatterer. In a spinless case, our DMRG study leads us to a similar conclusion as NRG studies of impurities with spin, and shows that the effective one-body elastic scatterer necessary for extending the Landauer formulation of coherent transport to correlated fermion systems must include slowly decaying correlations in the attached leads.

We thank Y. Asada, G.-L. Ingold, R.A. Jalabert, O. Sushkov and G. Vasseur for stimulating discussions, and P. Schmitteckert for his DMRG code. R.A. Molina acknowledges the financial support provided through the European Community's Human Potential Program under contract HPRN-CT-2000-00144.

REFERENCES

- [1] R. Landauer, IBM J. Res. Dev. **1**, 223 (1957); M. Büttiker, Phys. Rev. Lett. **57**, 1761 (1986); Y. Imry, *Introduction to Mesoscopic Physics*, Oxford University Press (1997).
- [2] K.J. Thomas, *et al.*, Phys. Rev. Lett. **77**, 135 (1996).
- [3] C. Kergueris *et al.*, Phys. Rev. B **59**, 12505 (1999).
- [4] N. Agraït, A.L. Yeyati and J.M. van Ruitenbeek, Phys. Rep. **377**, 81 (2003).
- [5] R.A. Molina *et al.*, Phys. Rev. B **67**, 235306 (2003).
- [6] R.A. Molina *et al.*, Eur. Phys. J. B **39**, 107 (2004).
- [7] A.O. Gogolin and N.V. Prokof'ev, Phys. Rev. B **50**, 4921 (1994).
- [8] J. Favand and F. Mila, Eur. Phys. J. B **2**, 293 (1998).
- [9] O.P. Sushkov, Phys. Rev. B **64**, 155319 (2001).
- [10] V. Meden and U. Schollwöck, Phys. Rev. B **67**, 193303 (2003).
- [11] T. Rejec and A. Ramšak, Phys. Rev. B **68**, 035342 (2003).
- [12] A.C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge University Press (1993).
- [13] S.R. White, Phys. Rev. Lett. **69**, 2863 (1992); Phys. Rev. B **48**, 10345 (1993).
- [14] *Density Matrix Renormalization – A New Numerical Method in Physics*, ed. by I. Peschel *et al.*, Lecture Notes in Physics, Vol. 528 Springer, Berlin (1999).
- [15] H.J. Lipkin, *Quantum Mechanics, New Approaches to Selected Topics*, North Holland (1973).
- [16] J. Friedel, Phil. Mag. **43**, 153 (1952).
- [17] J.S. Langer and V. Ambegaokar, Phys. Rev. **121**, 1090 (1961); D.C. Langreth, Phys. Rev. **150**, 516 (1966).
- [18] R.A. Molina, D. Weinmann and J.-L. Pichard, Europhys. Lett. **67**, 96 (2004).
- [19] T. Giamarchi, *Quantum Physics in One Dimension*, Oxford University Press (2004).
- [20] A. Oguri, Y. Nisikawa and A.C. Hewson, arXiv:cond-mat/0504771 and references therein.