



**HAL**  
open science

## Permutation representations of the braid group commutator subgroup

Abdelouahab Arouche

► **To cite this version:**

Abdelouahab Arouche. Permutation representations of the braid group commutator subgroup. 2005.  
hal-00007972

**HAL Id: hal-00007972**

**<https://hal.science/hal-00007972>**

Preprint submitted on 17 Aug 2005

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# PERMUTATION REPRESENTATIONS OF THE BRAID GROUP COMMUTATOR SUBGROUP.

ABDELOUAHAB AROUCHE

ABSTRACT. We study the representations of the commutator subgroup  $K_n$  of the braid group  $B_n$  into the symmetric group  $S_r$ . Motivated by some experimental results, we conjecture that every such a representation with  $n > r$  must be trivial.

## 1. INTRODUCTION

In [SiWi1], D. Silver and S. Williams exploited the structure of the kernel subgroup  $K$  of an epimorphism  $\chi : G \rightarrow \mathbb{Z}$ , where  $G$  is a finitely presented group, to show that the set  $Hom(K, \Sigma)$  of representations of  $K$  into a finite group  $\Sigma$  has a structure of a subshift of finite type (SFT), a symbolic dynamical system described by a graph  $\Gamma$ ; namely, there is a one to one correspondence between representations  $\rho : K \rightarrow \Sigma$  and bi-infinite paths in  $\Gamma$ .

We apply this method to the group  $B_n$  of braids with  $n$ -strands, with  $\chi$  being the abelianization homomorphism and  $\Sigma$  the symmetric group of degree  $r$ . The subgroup  $K_n = \ker \chi$  is then the commutator subgroup of  $B_n$ .

It is a well known fact that for a given group  $K$ , there is a finite to one correspondence between its subgroups of index no greater than  $r$  and representations  $\rho : K \rightarrow S_r$ . This correspondence can be described by

$$\rho \longmapsto \{g \in K : \rho(g)(1) = 1\}.$$

The pre-image of a subgroup of index exactly  $r$  consists of  $(r-1)!$  transitive representations  $\rho$ . ( $\rho$  is said to be transitive if  $\rho(K)$  operates transitively on  $\{1, 2, \dots, r\}$ ). This will allow us to draw some conclusions about the subgroups of finite index of  $K_n$ .

We give an algorithm to compute  $Hom(K_n, S_r)$ , for  $n \geq 5$  and  $r \geq n$ . Motivated by some experimental results, we conjecture that  $Hom(K_n, S_r)$  is trivial for  $n \geq 5$  and  $r < n$ . Since every representation in  $Hom(B_n, S_r)$  restricts to an element of  $Hom(K_n, S_r)$ , we enhance the given algorithm in order to compute  $Hom(B_n, S_r)$ .

## 2. GENERALITIES

Let  $B_n$  be the braid group given by the presentation:

$$\langle \sigma_1, \dots, \sigma_{n-1} \mid \begin{array}{l} \sigma_i \sigma_j = \sigma_j \sigma_i; \quad |i-j| \geq 2 \\ \sigma_i \sigma_j \sigma_i = \sigma_j \sigma_i \sigma_j; \quad |i-j| = 1 \end{array} \rangle,$$

*Date:* July 30, 2005.

*2000 Mathematics Subject Classification.* Primary 20F36, 20C40; Secondary 20E07.

*Key words and phrases.* Braid group, commutator subgroup, representation, symmetric group.

(see [BuZi] for additional background). Let  $\beta \in B_n$  be a braid. Then  $\beta$  can be written as :

$$\beta = \sigma_{i_1}^{\varepsilon_1} \cdots \sigma_{i_k}^{\varepsilon_k},$$

with  $i_1, \dots, i_k \in \{1, \dots, n-1\}$  and  $\varepsilon_i = \pm 1$ . Define the exponent sum of  $\beta$  (in terms of the  $\sigma_i$ 's) denoted by  $\exp(\beta)$ , as:

$$\exp(\beta) = \varepsilon_1 + \cdots + \varepsilon_k.$$

Then  $\exp(\beta)$  is an invariant of the braid group, that is, it doesn't depend on the writing of  $\beta$ . Moreover,  $\exp(\beta) : B_n \rightarrow \mathbb{Z}$  is a homomorphism. Let  $H_n$  denote its kernel. Then the Reidemeister-Schreier theorem [LySc] enables us to find a presentation for  $H_n$ . We choose the set

$$\{\cdots, \sigma_1^{-m}, \sigma_1^{-m+1}, \cdots, \sigma_1^{-1}, 1, \sigma_1, \sigma_1^2, \cdots, \sigma_1^m, \cdots\}$$

as a Schreier system of right coset representatives of  $H_n$  in  $B_n$ . Putting  $z_m = \sigma_1^m (\sigma_2 \sigma_1^{-1}) \sigma_1^{-m}$  for  $m \in \mathbb{Z}$ , and  $x_i = \sigma_i \sigma_1^{-1}$  for  $i = 3, \dots, n-1$ , we get the following presentation of  $H_n$  :

$$H_n = \left\langle \begin{array}{l} z_m, \quad m \in \mathbb{Z} \\ x_i, \quad i = 3, \dots, n-1 \end{array} \middle| \begin{array}{l} x_i x_j = x_j x_i, \quad |i-j| \geq 2; \\ x_i x_j x_i = x_j x_i x_j, \quad |i-j| = 1; \\ z_m z_{m+2} = z_{m+1}, \quad m \in \mathbb{Z}; \\ z_m x_3 z_{m+2} = x_3 z_{m+1} x_3, \quad i = 4, \dots, n-1; \\ z_m x_i = x_i z_{m+1}, \quad m \in \mathbb{Z}. \end{array} \right\rangle$$

**Example 1.** For  $n = 3, 4$ , we have:

$$H_3 = \langle z_m \mid z_m z_{m+2} = z_{m+1}; \forall m \in \mathbb{Z} \rangle$$

is a free group on two generators  $z_0 = \sigma_2 \sigma_1^{-1}$  and  $z_{-1} = \sigma_1^{-1} \sigma_2$ , and

$$H_4 = \langle z_m, t \mid \begin{array}{l} z_m z_{m+2} = z_{m+1}, \\ z_m t z_{m+2} = t z_{m+1} t, \end{array} \quad m \in \mathbb{Z} \rangle.$$

Note that  $H_2 = \{1\}$ , since  $B_2 = \langle \sigma_1 \mid \rangle \cong \mathbb{Z}$ .

Now, every commutator in  $B_n$  has exponent sum zero. Conversely, every generator of  $H_n$  is a product of commutators. Hence, we have  $H_n = K_n$ , and  $\exp$  is the abelianization homomorphism.

Each  $K_n$  fits into a split exact sequence:

$$1 \rightarrow K_n \rightarrow B_n \rightarrow \mathbb{Z} \rightarrow 0,$$

and there are "natural" inclusions  $B_n \subset B_{n+1}$  and  $K_n \subset K_{n+1}$ .

### 3. THE REPRESENTATION SHIFT

Our goal is to study representations of  $K_n$  into the symmetric group  $S_r$ . Note that there is a natural homomorphism:

$$\pi : B_n \rightarrow S_n,$$

for all  $n \geq 2$ , given by  $\sigma_i \mapsto (ii+1)$ . This restricts to  $K_n$  to give a non trivial homomorphism:

$$\begin{array}{l} x_i \mapsto (12)(ii+1) \quad i = 3, \dots, n-1 \\ z_m \mapsto \begin{cases} (132), & m \text{ is even} \\ (123), & m \text{ is odd} \end{cases} \end{array}$$

We start with  $n = 3$  and describe  $\text{Hom}(K_3, S_r)$  by means of a graph  $\Gamma$  that we will construct in a step by step fashion [SiWi2]. Note that  $\text{Hom}(K_3, S_r)$  contains  $\pi|_{K_3}$ , for  $r \geq 3$ . Later we will see that  $\text{Hom}(K_3, S_2)$  is not trivial.

A representation  $\rho : K_n \rightarrow S_r$  is a function  $\rho$  from the set of generators  $z_m$  to  $S_r$  such that for each  $m \in \mathbb{Z}$ , the relation:

$$\rho(z_m)\rho(z_{m+2}) = \rho(z_{m+1})$$

holds in  $S_r$ . Any such function can be constructed as follows, beginning with step 0 and proceeding to steps  $\pm 1, \pm 2, \dots$

.

.

.

(step -1) Choose  $\rho(z_{-1})$  if possible such that  $\rho(z_{-1})\rho(z_1) = \rho(z_0)$ .

(step 0) Choose values  $\rho(z_0)$  and  $\rho(z_1)$ .

(step 1) Choose  $\rho(z_2)$  if possible such that  $\rho(z_0)\rho(z_2) = \rho(z_1)$ .

(step 2) Choose  $\rho(z_3)$  if possible such that  $\rho(z_1)\rho(z_3) = \rho(z_2)$ .

.

.

.

This process leads to a bi-infinite graph whose vertices are the maps  $\rho : \{z_0, z_1\} \rightarrow S_r$ , each of which can be regarded as an ordered pair  $(\rho(z_0), \rho(z_1))$ . There is a directed edge from  $\rho$  to  $\rho'$  if and only if  $\rho(z_1) = \rho'(z_0)$  and  $\rho(z_0)\rho'(z_1) = \rho(z_1)$ . In such a case, we can extend  $\rho : \{z_0, z_1\} \rightarrow S_r$  by defining  $\rho(z_2)$  to be equal to  $\rho'(z_1)$ . Now if there is an edge from  $\rho'$  to  $\rho''$ , we can likewise extend  $\rho$  by defining  $\rho(z_3)$  to be  $\rho''(z_1)$ . We implement this process by starting with an ordered pair  $(a_0, a_1)$  of elements of  $S_r$ , and computing at each step a new ordered pair from the old one, so that every edge in the graph looks like:

$$(a_m, a_{m+1}) \rightarrow (a_{m+1}, a_{m+2}),$$

with

$$a_{m+2} = a_m^{-1}a_{m+1}.$$

In our case, the graph  $\Gamma$  we obtain consists necessarily of disjoint cycles. This gives an algorithm for finding  $\text{Hom}(K_3, S_r)$ . Observe that  $\text{Hom}(K_3, S_r)$  is endowed with a shift map

$$\sigma : \rho \mapsto \sigma(\rho)$$

defined by

$$\sigma(\rho) : x \mapsto \rho(\sigma_1 x \sigma_1^{-1}).$$

If we regard  $\rho$  as a bi-infinite path in the graph  $\Gamma$ , then  $\sigma$  correspond to the shift map  $(a_m, a_{m+1}) \mapsto (a_{m+1}, a_{m+2})$ , since  $a_{m+1} = \sigma_1 a_m \sigma_1^{-1}$ . Any cycle in the graph  $\Gamma$  with length  $p$  corresponds to  $p$  representations having least period  $p$ . These are the iterates of some representation  $\rho \in \text{Hom}(K_3, S_r)$  satisfying  $\rho(z_m) = a_m$  and  $\sigma^p(\rho) = \rho$ , since  $a_{m+p} = a_m$ .

In order to minimize calculations, we extract some foreseeable behaviour for various choices of the ordered pair  $(a_0, a_1)$  in the previous algorithm.

First, the dynamics of ordered pairs  $(a_0, a_1)$  such that  $a_0 = 1$  or  $a_1 = 1$  or  $a_0 = a_1$  is entirely known. To be precise, we get a cycle of length 6 unless  $a = a^{-1}$ ,

in which case it is of length 3 (or 1 if and only if  $a = 1$ ).

$$\begin{aligned} (1, a) &\rightarrow (a, a) \rightarrow (a, 1) \rightarrow (1, a^{-1}) \\ &\rightarrow (a^{-1}, a^{-1}) \rightarrow (a^{-1}, 1) \rightarrow (1, a). \end{aligned}$$

Second, when we proceed to a new step, we do not need to take a pair we have already got in a previous cycle, since we would get indeed the same cycle. The following dichotomy will prove useful in the sequel:

**Definition 1.** *If a vertex of a cycle in  $\Gamma$  has equal components, then the cycle is said of type I. Otherwise, it is of type II.*

Note that a cycle is determined by any of its vertices. Furthermore, the type I cycles are determined by elements of  $S_r$ . Recall that a non trivial element  $a \in S_r$  has order two if and only if it is a product of disjoint transpositions. Let  $n_r$  be the number of such elements. This gives us a means to compute the number of type I cycles to be  $\frac{1}{2}(1 + n_r + r!)$  and of representations coming from them to be  $3r! - 2$ .

**Example 2.** *Since  $S_2 = \{1, (12)\}$ ,  $\text{Hom}(K_3, S_2)$  consists only of the following type I cycle:*

$$(1, (12)) \rightarrow ((12), (12)) \rightarrow ((12), 1) \rightarrow (1, (12)),$$

*along with the trivial representation. So  $|\text{Hom}(K_3, S_2)| = 4$ . As for  $\text{Hom}(K_3, S_3)$ , there are three type I cycles of length 3 corresponding to transpositions and one type I cycle of length 6 corresponding to the 3-cycle (123) (and its inverse). Looking at type II cycles, we find two cycles of length 9 corresponding to the pairs  $((23), (12))$  and  $((23), (123))$  and one cycle of length 2 corresponding to the pair  $((123), (132))$ . This last one is exactly the orbit (under the shift map  $\sigma$ ) of  $\pi|_{K_3}$ . All by all, we have  $|\text{Hom}(K_3, S_3)| = 36$ .*

*In the last section we present, among other things, the results of computer calculations of type II cycles in the graphs of  $\text{Hom}(K_3, S_4)$  using Maple.*

Now let us proceed to compute  $\text{Hom}(K_4, S_r)$ . Since  $K_3 \subset K_4$ , every representation  $\rho \in \text{Hom}(K_4, S_r)$  restricts to a representation  $\rho|_{K_3} \in \text{Hom}(K_3, S_r)$ , the latter being described by a cycle. All we have to do is then to check which representation in  $\text{Hom}(K_3, S_r)$  does extend to  $K_4$ . To this end, observe that  $K_4$  is gotten from  $K_3$  by adjunction of a generator  $x_3$  subject to the relations

$$z_m x_3 z_{m+2} = x_3 z_{m+1} x_3; m \in \mathbb{Z}.$$

Hence we may proceed as follows. Take a cycle in  $\text{Hom}(K_3, S_r)$  (by abuse of language, i.e. identify each representation with its orbit, since a representation in  $\text{Hom}(K_3, S_r)$  extends to  $K_4$  if and only if every element in its orbit does), and choose if possible a value  $b_3 \in S_r$  for  $\rho(x_3)$ . This value must satisfy the relations

$$a_m b_3 a_{m+2} = b_3 a_{m+1} b_3;$$

for  $m = 0, \dots, p-1$ , where  $p$  is the cycle's length and the indexation is *mod* $p$ . Observe that the choice  $b_3 = 1$  is convenient, so all cycles extend to  $K_4$ . However, this is the only possibility for type I cycles to extend, for if  $b_3$  commute with some  $a_m$ , then  $b_3 = 1$ . For type II cycles, we find for example that no cycle in  $\text{Hom}(K_3, S_3)$  extends to  $K_4$  with non trivial  $b_3$  and that out of 71 cycles in  $\text{Hom}(K_3, S_4)$  only ten do extend to  $K_4$ , each with three possibilities for  $b_3$  (the same for all; see the last section).

Before giving the general procedure, let us proceed one further step to show that all type I cycles will vanish for  $n \geq 5$ . Take a cycle in  $Hom(K_3, S_r)$ ,  $r \geq 2$  along with a convenient value  $b_3$  of  $\rho(x_3)$ . We look for an element  $b_4 \in S_r$  satisfying :

$$\begin{aligned} a_m b_4 &= b_4 a_{m+1}, & m = 0, \dots, p; \\ b_3 b_4 b_3 &= b_4 b_3 b_4 \end{aligned}$$

Hence, if

$$b_3 b_4 = b_4 b_3$$

then using

$$a_m a_{m+2} = a_{m+1},$$

and

$$a_m b_3 a_{m+2} = b_3 a_{m+1} b_3,$$

we get

$$b_3 = b_4 = 1,$$

and

$$a_m = a_{m+1}, \forall m = 0, \dots, p-1,$$

so that the representation is trivial. So only type II cycles, with non trivial  $b_3$  possibly extend to  $K_5$  (beside the trivial one). So no (type II) cycle in  $Hom(K_3, S_3)$  extends to  $K_5$ . It turns out that no type II cycle in  $Hom(K_3, S_4)$  extends to  $K_5$ .

**Algorithm 1.** *The general procedure for  $Hom(K_n, S_r)$ ,  $n \geq 5$  is to consider only type II cycles along with convenient non trivial values  $b_3, \dots, b_{n-2}$ , which correspond to representations in  $Hom(K_{n-1}, S_r)$  and find a non trivial element  $b_{n-1} \in S_r$  such that the following relations are satisfied:*

$$\begin{aligned} a_m b_{n-1} &= b_{n-1} a_{m+1}, & m = 0, \dots, p; \\ b_{n-1} b_i &= b_i b_{n-1} & i = 3, \dots, n-2; \\ b_{n-1} b_{n-2} b_{n-1} &= b_{n-2} b_{n-1} b_{n-2} \end{aligned}$$

The element  $b_{n-1}$  has to be non trivial, otherwise the representation is trivial. Experimental results lead us to conjecture that the process will stop at step  $n = r$ . That is:

**Conjecture 1.**  *$Hom(K_n, S_r)$  is trivial for  $n \geq r + 1$ .*

It is obvious that a cycle (of any type) can not extend to  $K_n$  if it doesn't extend to  $K_{n-1}$  it is enough for the conjecture to be true that  $Hom(K_{r+1}, S_r)$  be trivial. Recall that for  $n \leq r$ ,  $Hom(K_n, S_r)$  is not trivial since it contains the homomorphism  $\pi|_{K_n} : K_n \rightarrow S_n$ .

#### 4. EXTENSION TO THE BRAID GROUP

In this section, we address the question of extending representations

$$\rho \in Hom(K_n, S_r)$$

to representations

$$\tilde{\rho} \in Hom(B_n, S_r).$$

Applying [SiWi1 (3.5)], the extension is possible if and only if there is an element  $c \in S_r$  such that

$$\begin{aligned} a_m c &= c a_{m+1}, & m = 0, \dots, p-1; \\ c b_i &= b_i c & i = 3, \dots, n-1; \end{aligned}$$

Observe that a necessary condition for a representation  $\rho \in \text{Hom}(K_n, S_r)$  to extend to  $\tilde{\rho} \in \text{Hom}(B_n, S_r)$  is that  $\rho \in \text{Hom}(K_n, A_r)$ , since the alternating group  $A_r$  is the commutator subgroup of  $S_r$ . A sufficient condition is that  $\rho$  be the restriction of some representation  $\hat{\rho} \in \text{Hom}(K_{n+2}, S_r)$  for if this is the case, the choice  $c = b_{n+1}$  will do. In this case, since  $\hat{\rho}$  maps  $K_n$  into  $A_r$ , it also maps  $K_{n+2}$  into  $A_r$ , for the values  $b_i$  are conjugate and  $A_r$  is normal in  $S_r$ . As a result, we get the following

**Proposition 1.** *for  $n \geq 5$ ,  $\text{Hom}(K_n, S_r) = \text{Hom}(K_n, A_r)$ .*

Actually we can enhance our algorithm to one which gives for fixed  $n \geq 5$  and  $r \geq n$  the sets  $\text{Hom}(K_n, S_r)$  and  $\text{Hom}(B_n, S_r)$ .

Step one: find all cycles of both types. This gives  $\text{Hom}(K_3, S_r)$ .

Step two: For the trivial cycle, take any  $c$ . For a type II cycle  $C$ , find  $c \neq 1$  such that  $a_m c = c a_{m+1}$ , for  $m = 0, \dots, p-1$ . This gives  $\text{Hom}(B_3, S_r)$ .

Step three: For a cycle of any type, take  $b_3 = 1$ . For a type II cycle  $C$ , find  $b_3 \neq 1$  such that  $a_m b_3 a_{m+2} = b_3 a_{m+1} b_3$ , for  $m = 0, \dots, p-1$ . This gives  $\text{Hom}(K_4, S_r)$ .

Step four: take a type II cycle  $C$ , along with a convenient  $b_3$ . If this cycle occurs in  $\text{Hom}(B_3, S_r)$  with some convenient  $c$  then :

if  $cb_3 = b_3c$ , then the representation  $[C, b_3]$  moves up to a representation  $[C, b_3, c]$  in  $\text{Hom}(B_4, S_r)$ ;

if  $cb_3c = b_3cb_3$ , then the representation  $[C, b_3]$  moves up to a representation  $[C, b_3, b_4]$  in  $\text{Hom}(K_5, S_r)$  by taking  $b_4 = c$ .

⋮  
⋮  
⋮

Step i: take a representation  $\rho$  in  $\text{Hom}(K_i, S_r)$ , encoded by a type II cycle  $C$  along with convenient values  $b_3, \dots, b_{i-1}$ . if  $[C, b_3, b_4, \dots, b_{i-2}, c]$  occurs in  $\text{Hom}(B_{i-1}, S_r)$  with some convenient  $c$  then:

if  $cb_{i-1} = b_{i-1}c$ , then the representation  $[C, b_3, b_4, \dots, b_{i-1}]$  moves up to a representation  $[C, b_3, b_4, \dots, b_{i-1}, c]$  in  $\text{Hom}(B_i, S_r)$ ;

if  $cb_{i-1}c = b_{i-1}cb_{i-1}$ , then the representation  $[C, b_3, b_4, \dots, b_{i-1}]$  moves up to a representation  $[C, b_3, b_4, \dots, b_{i-1}, b_i]$  in  $\text{Hom}(K_{i+1}, S_r)$  by taking  $b_i = c$ .

⋮  
⋮  
⋮

Note that if the conjecture is true, then for  $n \geq r+1$ , every representation  $\tilde{\rho} : B_n \rightarrow S_r$  factorizes through the abelianized group  $(B_n)_{ab}$ , and has a cyclic image. Hence, there are  $r!$  possible choices for  $\tilde{\rho}$ .

## 5. CONSEQUENCES

Regarding the correspondence between subgroups of finite index of a group  $K$  and its representations into symmetric groups, we investigate the subgroups of index  $r$  of  $K_n$  for low degrees  $r$ . The general principle is to compute the number of transitive representations of  $K$  into  $S_r$  to deduce the number of subgroups of  $K$  with index exactly  $r$ . We start with  $K_3$  as usual. Note that since  $K_3$  is freely generated by  $z_0$  and  $z_{-1}$ , it maps onto any symmetric group, and hence, has subgroups of every index. Now, if a representation in  $\text{Hom}(K_3, S_r)$  is transitive, then so are the representations in its orbit. Consider a type I cycle in  $\text{Hom}(K_3, S_r)$ . Then the representations it defines are transitive if and only if the defining element

$a$  is (with respect to the action of  $S_r$  on  $\{1, \dots, r\}$ ). This exactly means that  $a$  is an  $r$ -cycle. If  $r > 2$  then  $a^2 \neq 1$  and the cycle has length 6.

**Claim 1.** *The number of transitive representations  $\rho \in \text{Hom}(K_3, S_r)$ ,  $r \geq 2$  coming from type I cycles is  $3(r-1)!$ .*

For  $r = 2$ , there are only type I cycles and there is only one 2-cycle, which has length 3; Hence, The number of transitive representations  $\rho \in \text{Hom}(K_3, S_2)$  is 3. The kernels of these representations give rise to subgroups of  $K_3$  with index 2.

**Claim 2.** *There are three subgroups of  $K_3$  with index 2.*

Now we compute the number of subgroups of  $K_3$  with index 3. Among all representations we have seen in example 2, there are six transitive representations coming from the only type I cycle and all representations coming from type II cycles are transitive. Hence:

**Claim 3.** *The number of transitive representations in  $\text{Hom}(K_3, S_3)$  is 26, consequently there are thirteen subgroups of  $K_3$  with index 3.*

We can proceed in this way for every degree  $r$ . To compute the number of transitive representations of  $K_3$  into  $S_r$ , we need only consider those coming from type II cycles, since we already know the number of those coming from type I cycles. This can be done using a computer algebra system, by taking any cycle  $C = (a_0, \dots, a_{p-1})$  and checking if the subgroup  $\langle a_0, \dots, a_{p-1} \rangle$  of  $S_r$  acts transitively on  $\{1, \dots, r\}$ . If so, this gives rise to  $p$  transitive representations in  $\text{Hom}(K_3, S_3)$ . Then we divide the total number by  $(r-1)!$  to find the number of subgroups of  $K_3$  of index  $r$ .

Now let us consider  $\text{Hom}(K_4, S_r)$ . For  $r = 2$  we have, as previously:

**Claim 4.** *There are three subgroups of  $K_4$  with index 2.*

As for transitive representations in  $\text{Hom}(K_4, S_3)$ , since all cycles in  $\text{Hom}(K_3, S_3)$  extend to  $K_4$  with only  $b_3 = 1$ , we have:

**Claim 5.** *There are twenty six transitive representations in  $\text{Hom}(K_4, S_3)$ , hence thirteen subgroups of  $K_3$  with index 3.*

For  $r \geq 4$ , we have  $3(r-1)!$  transitive representations coming from type I cycles, and we must check which representation coming from a type II cycle is transitive. For a cycle  $C = (a_0, \dots, a_{p-1})$  such that  $\langle a_0, \dots, a_{p-1} \rangle$  failed to be transitive, we check if  $\langle a_0, \dots, a_{p-1}, b_3 \rangle$  (with  $b_3$  non trivial) is transitive. Indeed, if  $\langle a_0, \dots, a_{p-1} \rangle$  is transitive, then so is  $\langle a_0, \dots, a_{p-1}, b_3 \rangle$ . Finally, we divide the total number by  $(r-1)!$  to find the number of subgroups of  $K_3$  of index  $r$ .

Now, we consider  $n \geq 5$ , where we get rid of type I cycles. Suppose we have found the transitive representations in  $\text{Hom}(K_{n-1}, S_r)$ . We then take, for fixed  $r$ , a type II cycle  $C = (a_0, \dots, a_{p-1})$  along with values  $b_3, \dots, b_{n-1}$ , such that  $\langle a_0, \dots, a_{p-1}, b_3, \dots, b_{n-2} \rangle$  failed to be transitive and check if  $\langle a_0, \dots, a_{p-1}, b_3, \dots, b_{n-1} \rangle$  is transitive. We may enhance algorithm 1 by checking, each time we get a new type II cycle, if it is transitive, and if not, we re-check at each time the cycle extends from  $K_i$  to  $K_{i+1}$ ,  $i = 3, \dots, n-1$ , after having augmented it with  $b_i$ . Dividing by  $(r-1)!$  the number of transitive representations in  $\text{Hom}(K_n, S_r)$  we find the number of subgroups of  $K_n$  with index  $r$ . As a consequence of conjecture 1, we get the following:



**Conjecture 2.** For  $n \geq 5$  and  $2 \leq r \leq n - 1$ , there are no subgroups of  $K_n$  with index  $r$ . Moreover, every nontrivial representation  $\rho$  of  $K_n$  into  $S_n$  is transitive.

**Remark 1.** We can likewise investigate the number of subgroups of  $B_n$  with a given index  $r$  by looking at transitive representations of  $B_n$  into  $S_r$ . Namely, if conjecture 1 is true, then there is exactly one subgroup of index  $r$  in  $B_n$ , for  $1 \leq r \leq n - 1$ . Moreover, if  $\rho : B_n \rightarrow S_n$  is a representation, then  $\rho|_{K_n}$  is either trivial or transitive, according to conjecture 2. In the first case,  $\rho$  has a cyclic image and we know when it is transitive. In the second case,  $\rho$  is transitive.

## 6. EXPERIMENTAL FACTS

In what follows, we list the type II cycles of  $Hom(K_n, S_r)$  for various (small)  $n$  and  $r$ . A word about the notation: each cycle  $B[a_0, a_1] = [a_2, a_3, \dots, a_{p-1}, a_0, a_1]$  is indexed by its first vertex  $(a_0, a_1)$  and is followed by its length  $p$ . Elements  $\tau \in S_r$  are ordered from 1 to  $r!$  with respect to the lexicographic order on the vectors  $(\tau(1), \dots, \tau(r))$ . It would have taken too much space to list the cycles for  $r = 5$ . We found that there were no (type II) cycles in  $Hom(K_5, S_4)$  nor in  $Hom(K_6, S_5)$ . Furthermore,  $Hom(K_4, S_3)$  contains no type II cycles with non trivial  $b_3$ . This motivated our conjecture 1.

$n = 3; r = 3 :$

B[2, 3] = [5, 6, 2, 5, 3, 6, 5, 2, 3]

9

B[2, 4] = [6, 3, 4, 2, 6, 4, 3, 2, 4]

9

B[4, 5] = [4, 5]

2

$n = 3; r = 4 :$

B[2, 3] = [5, 6, 2, 5, 3, 6, 5, 2, 3]

9

B[2, 4] = [6, 3, 4, 2, 6, 4, 3, 2, 4]

9

B[2, 7] = [8, 2, 7]

3

B[2, 8] = [7, 2, 8]

3

B[2, 9] = [11, 6, 16, 18, 3, 20, 19, 2, 9]

9

B[2, 10] = [12, 3, 23, 21, 6, 14, 13, 2, 10]

9

B[2, 11] = [9, 6, 18, 16, 3, 19, 20, 2, 11]

9

B[2, 12] = [10, 3, 21, 23, 6, 13, 14, 2, 12]

9

B[2, 13] = [19, 22, 4, 23, 15, 12, 11, 2, 13]

9

B[2, 14] = [20, 15, 18, 5, 22, 10, 9, 2, 14]

9

$$B[2, 15] = [21, 22, 2, 21, 15, 22, 21, 2, 15]$$

9

$$B[2, 16] = [22, 15, 16, 2, 22, 16, 15, 2, 16]$$

9

$$B[2, 17] = [23, 7, 24, 23, 2, 17]$$

6

$$B[2, 18] = [24, 7, 18, 17, 2, 18]$$

6

$$B[2, 19] = [13, 22, 23, 4, 15, 11, 12, 2, 19]$$

9

$$B[2, 20] = [14, 15, 5, 18, 22, 9, 10, 2, 20]$$

9

$$B[2, 23] = [17, 7, 23, 24, 2, 23]$$

6

$$B[2, 24] = [18, 7, 17, 18, 2, 24]$$

6

$$B[3, 7] = [13, 15, 3, 13, 7, 15, 13, 3, 7]$$

9

$$B[3, 8] = [14, 22, 17, 14, 3, 8]$$

6

$$B[3, 9] = [15, 7, 9, 3, 15, 9, 7, 3, 9]$$

9

$$B[3, 10] = [16, 7, 11, 5, 15, 23, 20, 3, 10]$$

9

$$B[3, 11] = [17, 22, 11, 8, 3, 11]$$

6

$$B[3, 12] = [18, 15, 4, 14, 7, 21, 19, 3, 12]$$

9

$$B[3, 14] = [8, 22, 14, 17, 3, 14]$$

6

$$B[3, 16] = [10, 7, 5, 11, 15, 20, 23, 3, 16]$$

9

$$B[3, 17] = [11, 22, 8, 11, 3, 17]$$

6

$$B[3, 18] = [12, 15, 14, 4, 7, 19, 21, 3, 18]$$

9

$$B[3, 22] = [24, 3, 22]$$

3

$$B[3, 24] = [22, 3, 24]$$

3

$$B[4, 5] = [4, 5]$$

2

$$B[4, 8] = [20, 16, 17, 4, 13, 8, 16, 20, 17, 13, 4, 8]$$

12

$$B[4, 9] = [21, 20, 4, 9]$$

4

$$B[4, 10] = [22, 13, 18, 6, 21, 11, 7, 4, 10]$$

9

$B[4, 11] = [23, 12, 19, 11, 13, 23, 19, 4, 11]$   
 9  
 $B[4, 12] = [24, 9, 16, 8, 12, 4, 24, 16, 9, 8, 4, 12]$   
 12  
 $B[4, 16] = [12, 9, 4, 16]$   
 4  
 $B[4, 17] = [9, 20, 24, 4, 21, 17, 20, 9, 24, 21, 4, 17]$   
 12  
 $B[4, 18] = [10, 13, 11, 18, 21, 10, 11, 4, 18]$   
 9  
 $B[4, 19] = [14, 21, 18, 19, 12, 14, 18, 4, 19]$   
 9  
 $B[4, 20] = [13, 16, 4, 20]$   
 4  
 $B[4, 22] = [18, 13, 6, 11, 21, 7, 10, 4, 22]$   
 9  
 $B[5, 7] = [14, 16, 6, 23, 9, 22, 19, 5, 7]$   
 9  
 $B[5, 8] = [13, 21, 24, 5, 20, 8, 21, 13, 24, 20, 5, 8]$   
 12  
 $B[5, 9] = [17, 12, 21, 8, 9, 5, 17, 21, 12, 8, 5, 9]$   
 12  
 $B[5, 10] = [18, 9, 14, 10, 20, 18, 14, 5, 10]$   
 9  
 $B[5, 12] = [16, 13, 5, 12]$   
 4  
 $B[5, 13] = [20, 21, 5, 13]$   
 4  
 $B[5, 14] = [19, 16, 23, 14, 9, 19, 23, 5, 14]$   
 9  
 $B[5, 16] = [24, 13, 12, 17, 16, 5, 24, 12, 13, 17, 5, 16]$   
 12  
 $B[5, 19] = [7, 16, 14, 6, 9, 23, 22, 5, 19]$   
 9  
 $B[5, 21] = [9, 12, 5, 21]$   
 4  
 $B[5, 23] = [11, 20, 10, 23, 16, 11, 10, 5, 23]$   
 9  
 $B[6, 7] = [20, 22, 6, 20, 7, 22, 20, 6, 7]$   
 9  
 $B[6, 8] = [19, 15, 24, 19, 6, 8]$   
 6  
 $B[6, 10] = [24, 15, 10, 8, 6, 10]$   
 6  
 $B[6, 12] = [22, 7, 12, 6, 22, 12, 7, 6, 12]$   
 9  
 $B[6, 15] = [17, 6, 15]$   
 3

$B[6, 17] = [15, 6, 17]$   
 3  
 $B[6, 19] = [8, 15, 19, 24, 6, 19]$   
 6  
 $B[6, 24] = [10, 15, 8, 10, 6, 24]$   
 6  
 $B[8, 17] = [24, 8, 17]$   
 3  
 $B[8, 18] = [23, 8, 23, 18, 8, 18]$   
 6  
 $B[8, 24] = [17, 8, 24]$   
 3  
 $B[9, 13] = [9, 13]$   
 2  
 $B[9, 18] = [11, 16, 19, 18, 20, 11, 19, 9, 18]$   
 9  
 $B[10, 14] = [12, 23, 10, 21, 14, 23, 13, 10, 14]$   
 9  
 $B[10, 17] = [10, 19, 17, 19, 10, 17]$   
 6  
 $B[11, 14] = [24, 14, 11, 24, 11, 14]$   
 6  
 $B[12, 20] = [12, 20]$   
 2  
 $B[16, 21] = [16, 21]$   
 2

$n = 4; r = 4 :$

ten cycles of length 2 along with three values  $b_3 = 8, 17, 24$ .

$[8, [4, 5], [4, 9], [4, 16], [4, 20], [5, 12], [5, 13], [5, 21], [9, 13], [12, 20], [16, 21]]$   
 $[17, [4, 5], [4, 9], [4, 16], [4, 20], [5, 12], [5, 13], [5, 21], [9, 13], [12, 20], [16, 21]]$   
 $[24, [4, 5], [4, 9], [4, 16], [4, 20], [5, 12], [5, 13], [5, 21], [9, 13], [12, 20], [16, 21]]$

**Acknowledgement 1.** *I am grateful to Susan G. Williams for many helpful e-mail discussions. I also wish to thank the students M. Menouer and Z. Ziadi for their help in computer search.*

#### REFERENCES

- [BuZi] G. Burde and H. Zieschang, Knots, De Gruyter Studies in Mathematics **5**, De Gruyter, Berlin, 1985.
- [LySc] R.C. Lyndon and P.E. Schupp, Combinatorial group theory, Springer-Verlag, Berlin, 1977.
- [SiWi1] D.S. Silver and S.G. Williams, "Augmented group systems and shifts of finite type," Israel J. Math. **95** (1996) , 213-251.
- [SiWi2] D.S. Silver and S.G. Williams, "Generalized  $n$ -colorings of links," Knot theory, Banach Center Publication, vol **42**, Inst. of Math., Polish Acad. of Sc., Warsaw 1998, 381-394.

USTHB, FAC. MATH. P.O.BOX 32 EL ALIA 16111 BAB EZZOUAR ALGIERS, ALGERIA  
*E-mail address:* [abdarouche@hotmail.com](mailto:abdarouche@hotmail.com)