



HAL
open science

ON THE CLASSIFICATION OF VECTOR BUNDLES WITH PERIODIC MAPS

Abdelouahab Arouche

► **To cite this version:**

Abdelouahab Arouche. ON THE CLASSIFICATION OF VECTOR BUNDLES WITH PERIODIC MAPS. 2005. hal-00007974

HAL Id: hal-00007974

<https://hal.science/hal-00007974>

Preprint submitted on 17 Aug 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE CLASSIFICATION OF VECTOR BUNDLES WITH PERIODIC MAPS

ABDELOUAHAB AROUCHE

ABSTRACT. We give an explicit description for universal principal $U(r)$ -bundles with periodic map by means of equivariant Stiefel manifolds. We then show that the associated equivariant vector bundle is equivalent to the canonical one given by G. Segal. Finally, we investigate some ideals involved in the equivariant K-theory of this classifying space.

1. INTRODUCTION

For a compact Lie group Γ and a topological group G , we call a $(\Gamma; G)$ -bundle any principal G -bundle on which Γ acts by bundle maps [D], [L]. It has been shown in [D] and [L] that a universal $(\Gamma; G)$ -bundle is given by $E(\Gamma; G) = \underset{H \in F}{*} \frac{\Gamma \times G}{H}$, where $F = F(\Gamma; G)$ is the family of subgroups $H \leq \Gamma \times G$ such that $H \cap G = 1$ (see also [AHJM]). This space is characterized by the fact that the fixed points set $E(\Gamma; G)^H$ is contractible if $H \in F$ and empty otherwise [J].

2. TWO LEMMAS

In the sequel, we shall need the following

Lemma 1. *if $G = G_1 \times G_2$, then there is a $(\Gamma \times G)$ -homotopy equivalence :*

$$E(\Gamma; G) \simeq E(\Gamma; G_1) \times E(\Gamma; G_2).$$

Lemma 2. *If $\Gamma_1 \leq \Gamma_2$ and $G_1 \leq G_2$, then there exists a $(\Gamma_1 \times G_1)$ -homotopy equivalence :*

$$E(\Gamma_1; G_1) \simeq \text{res}_{(\Gamma_1 \times G_1)} E(\Gamma_2; G_2).$$

3. THE MAIN THEOREM

In what follows, we restrict our study to the case $\Gamma = \mathbf{Z}_n$, the cyclic group of order n . Recall that the Stiefel manifolds are defined by

$$V_{r,s} = \{(v_1, \dots, v_r), v_i \in \mathbf{C}^s, v_i \cdot v_j = \delta_{ij}, i, j = 1, \dots, r\}$$

and

$$V_{r,\infty} = \varinjlim_s V_{r,s}.$$

Let $\mathbf{Z}_n \times U(r)$ act on the Stiefel manifold $V_{r,\infty}$ by :

$$(\gamma, a) \cdot (v_1, \dots, v_r) = (w_1, \dots, w_r)$$

Date: March 10, 2005.

2000 Mathematics Subject Classification. 19L47, 55R91.

Key words and phrases. Stiefel manifolds, equivariant bundles, completion.

□

4. THE ASSOCIATED \mathbf{Z}_n -VECTOR BUNDLE.

Let \mathbf{M} be the \mathbf{Z}_n -module defined by $\gamma : \mathbf{M} = \mathbf{C}^\infty$ and

$$\gamma \cdot (z_1, z_2, \dots) = (\gamma \cdot z_1, \gamma^2 \cdot z_2, \dots)$$

. Let $G_{r,\infty}$ be the Grassmannian manifold of r -dimensional subspaces of \mathbf{M} . Then the right $U(r)$ -action on $V_{r,\infty}$ defined by

$$(v_1, \dots, v_r) \cdot a = (1, a^{-1}) \cdot (v_1, \dots, v_r)$$

has as a quotient space $V_{r,\infty}/U(r) = G_{r,\infty}$. There is a canonical \mathbf{Z}_n -vector bundle on $G_{r,\infty}$ whose totalspace is

$$E_{\mathbf{M}} = \{(N, z) : z \in N\} \subseteq G_{r,\infty} \times \mathbf{M}$$

and which is universal [S].

Theorem 2. *The \mathbf{Z}_n -vector bundle associated with the principal $(\mathbf{Z}_n; U(r))$ -bundle $V_{r,\infty} \rightarrow G_{r,\infty}$ is equivalent to $E_{\mathbf{M}} \rightarrow G_{r,\infty}$*

Proof. The total space of the associated \mathbf{Z}_n -vector bundle is $V_{r,\infty} \times_{U(r)} \mathbf{C}^r$, with the \mathbf{Z}_n -action :

$$\gamma \cdot [(v_1, \dots, v_r), y] = [(\gamma, \mathbf{I}_r) \cdot (v_1, \dots, v_r), y].$$

Following [H, ch. 7, (7.1)], we define a map

$$f : V_{r,\infty} \times_{U(r)} \mathbf{C}^r \rightarrow E_{\mathbf{M}},$$

by

$$f [(v_1, \dots, v_r), y] = \left(\langle v_1, \dots, v_r \rangle, \sum_{i=1}^r y_i v_i \right)$$

. According to [H], f is a vector bundle isomorphism. It is easy to check its \mathbf{Z}_n -equivariance. □

5. CHARACTERISTIC CLASSES

Let Γ be a compact Lie group and G a topological group. Then $E(\Gamma; G)$ is a right G -space by $e.g = (1, g) \cdot e$; we denote by $B(\Gamma; G)$ the quotient space $E(\Gamma; G)/G$. According to [AHJM], the equivariant K-theory ring $K_\Gamma^*(B(\Gamma; G))$ is isomorphic to the completion $R(\Gamma \times G)_{\hat{F}}$ of the complex representation ring with respect to the F -adic topology. Recall that the F -adic topology is defined on $R(\Gamma \times G)$ by the set of ideals $I_H = \ker \{R(\Gamma \times G) \rightarrow R(H)\}$, for $H \in F$. In the case of $\Gamma = \mathbf{Z}_2$ and $G = \mathbf{T}^r$, we have :

$$F = H_k : k = -1, 0, \dots, r,$$

where

$$H_{-1} = 1,$$

$$H_0 = \{1, (-1, \mathbf{I}_r)\}, H_r = \{1, (-1, -\mathbf{I}_r)\}$$

and for $k = 1, \dots, r-1$:

$$H_k = \left\{ 1, \left(-1, \begin{pmatrix} -\mathbf{I}_k & 0 \\ 0 & \mathbf{I}_{r-k} \end{pmatrix} \right) \right\}.$$

Putting

$$I_k = \ker \{R(\mathbf{Z}_2 \times \mathbf{T}^r) \rightarrow R(H_k)\},$$

the F -adic topology is defined on $R(\mathbf{Z}_2 \times \mathbf{T}^r)$ by the ideal :

$$I = \bigcap_{k=-1}^r I_k.$$

Proposition 1. *The the F -adic topology is defined on $R(\mathbf{Z}_2 \times \mathbf{T}^r)$ by the ideal :*

$$I = I_0 \cong R(\mathbf{Z}_2) \otimes I_{\mathbf{T}^r},$$

where $I_{\mathbf{T}^r}$ denotes the augmentation ideal, and that on $R(\mathbf{Z}_2 \times \mathbf{U}(r))$ is defined by :

$$J = I \cap R(\mathbf{Z}_2 \times \mathbf{U}(r)).$$

In general, the F -adic topology on $R(\mathbf{Z}_n \times \mathbf{U}(r))$ is defined by the ideals $I_{d,j}$ coming from the subgroups :

$$K_{d,j} = \left\langle \lambda, \begin{pmatrix} \lambda^j & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} \right\rangle; \quad d \mid n; \quad j = 1, \dots, d; \quad (\lambda^d = 1).$$

Proof. Easy. The general case follows from the description of $F(\mathbf{Z}_2; U(r))$ given in thm (3.1).

Example 2. $K_{d,d} \cong \mathbf{Z}_d$, $I_{d,d} \cong R(\mathbf{Z}_d) \otimes I_{U(r)}$ and $I_{d,d-1} \cong I_{(\mathbf{Z}_d \times \mathbf{T}^r)} \cap R(\mathbf{Z}_d \times U(r))$.

□

6. HOMOTOPIES IN STIEFEL MANIFOLDS

Let \mathbf{C}^∞ be the telescope $\bigcup_{n \geq 1} \mathbf{C}^n$, and let

$$\begin{aligned} (\mathbf{C}^\infty)^{even} &= \{v \in \mathbf{C}^\infty : v_{2n+1} = 0; \quad \forall n = 1, \dots, \infty.\}, \\ (\mathbf{C}^\infty)^{odd} &= \{v \in \mathbf{C}^\infty : v_{2n} = 0; \quad \forall n = 1, \dots, \infty.\}. \end{aligned}$$

Two maps :

$$g^{even}, g^{odd} : \mathbf{C}^n \times \mathbf{I} \rightarrow \mathbf{C}^{2n}$$

are defined in the following way :

$$g_t^{even}(v_1, \dots, v_n) = (1-t)(v_1, \dots, v_n) + t(0, v_1, \dots, 0, v_n)$$

and

$$g_t^{odd}(v_1, \dots, v_n) = (1-t)(v_1, \dots, v_n) + t(v_1, 0, \dots, v_n, 0).$$

These maps extend to :

$$g^{even}, g^{odd} : \mathbf{C}^\infty \times \mathbf{I} \rightarrow \mathbf{C}^\infty,$$

and satisfy :

$$\begin{aligned} g_0^{even} &= g_0^{odd} = Id_{\mathbf{C}^\infty}; \\ g_1^{even}(\mathbf{C}^\infty) &\subseteq (\mathbf{C}^\infty)^{even}; \quad g_1^{odd}(\mathbf{C}^\infty) \subseteq (\mathbf{C}^\infty)^{odd}. \end{aligned}$$

[H, ch. 3, (6.1)].

Similarly, we define

$$(\mathbf{V}_{r,\infty})^{even} = \{(v_1, \dots, v_r) \in \mathbf{V}_{r,\infty} : v_j^{2n+1} = 0, \quad \forall j = 1, \dots, r; \quad \forall n = 1, \dots, \infty\},$$

and

$$(\mathbf{V}_{r,\infty})^{odd} = \{(v_1, \dots, v_r) \in \mathbf{V}_{r,\infty} : v_j^{2n} = 0, \quad \forall j = 1, \dots, r; \quad \forall n = 1, \dots, \infty\}.$$

