



3-colorability and contacts of segments

Patrice Ossona de Mendez

► To cite this version:

Patrice Ossona de Mendez. 3-colorability and contacts of segments. Série du CAMS, 1999, 171, pp.1-2.
hal-00008547

HAL Id: hal-00008547

<https://hal.science/hal-00008547>

Submitted on 8 Sep 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

3-colorability and contacts of segments [★]

P. Ossona de Mendez

CNRS UMR 0017, EHESS, 54 Boulevard Raspail, 75006, Paris, France.

Abstract. Using a result on planar graph representation [1], we shall prove that vertex 3-colorability is NP-complete for simple contact graphs of segments in 4 directions.

Let V be a finite set of segments in the plane. Associate a graph $G(V)$ called the *intersection graph* of V as follows : V is the vertex set of G and there is an edge connecting v_1 and v_2 if and only if the segments v_1 and v_2 intersect.

G. Ehrlich, S. Even and R. Tarjano [3] proved that vertex 3-colorability of intersection graphs of segments is NP-complete.

It has been conjectured that any planar graph is an intersection graph of segments and this has been proved for 3-colorable graphs [2], what gives another proof of the previous theorem.

In case where the segments in V never cross, but may only touch, the intersection graph of V is a *contact graph of segments*.

Contact graphs of segments are precisely those graphs G that satisfy :

- G is planar
- Any subgraph H of G , induced by $n_H \geq 2$ vertices, has a number of edges at most equal to $2.n_H - 3$

This class includes the planar graphs that are triangle free (3-colorable, according to Grötsch theorem), the outerplanar graphs (3 colorable, as proved by König), 3-regular graphs different from K_4 (3 colorable, according to Brooks' theorem in the cubic case) but also 4 chromatic graphs.

If the segments of V belongs to k directions, $G(V)$ is a *k-dir contact graph of segments*. Obviously, a 3-dir contact graph of segments is 3-colorable.

One may ask if the problem of finding the chromatic number is still NP-complete for contact graphs of segments, or even for 4-dir contact graphs of segments.

Theorem 1. *Vertex 3-colorability is NP-complete for 4-dir contact graphs of segments.*

Proof. As notice in [1], any planar graph G is the contact graph of T-shape. Moreover, the vertex 3-colorability of G is obviously equivalent to the vertex 3-colorability of the 4-dir contact graph of segments G' obtained by transforming each T-shape as shown in the figure. \square

[★] This work was partially supported by the **Esprit LTR Project no 20244-ALCOM IT**.




Fig. 1. Transformation of a T-shape contact graph into a 4-dir contact graph of segments, while preserving vertex 3-colorability

References

1. H. de Fraysseix, P. Ossona de Mendez, and P. Rosenstiehl. On triangle contact graphs. *Combinatorics, Probability and Computing*, 3:233–246, 1994.
2. H. de Fraysseix, P. Ossona de Mendez, and C. Thomassen. On graphs representable by contact graphs. in preparation.
3. G. Ehrlich, S. Even, and R.E. Tarjan. Intersection graphs of curves in the plane. *Journal of Combinatorial Theories, Series B*, 21:8–20, 1986.