

# Life-cycle assessment of straw use in bio-ethanol production: a case-study based on biophysical modelling

Benoit Gabrielle, Nathalie Gagnaire

#### ▶ To cite this version:

Benoit Gabrielle, Nathalie Gagnaire. Life-cycle assessment of straw use in bio-ethanol production: a case-study based on biophysical modelling. Biomass and Bioenergy, 2007, 32 (5), pp.431-441. 10.1016/j.biombioe.2007.10.017 . hal-00008565

## HAL Id: hal-00008565 https://hal.science/hal-00008565

Submitted on 9 Sep 2005

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Life-cycle assessment of straw use in bio-ethanol production: a case-study based on biophysical modelling. Benoît Gabrielle\* and Nathalie Gagnaire Environment and Arable Crops Research Unit, Institut National de la Recherche Agronomique, 78850 Thiverval-Grignon, France. \*: corresponding author. UMR INRA INA P-G Environnement et Grandes Cultures, 78850 Thiverval-Grignon, France. Phone: (+33) 1 30 81 55 51 Fax: (+33) 1 30 81 55 63 E-mail address: Benoit.Gabrielle@grignon.inra.fr

#### ABSTRACT

2	Cereal straw, a by-product in the production of agricultural crops, is
3	considered as a potentially large source of energy supply with an estimated value of
4	$47 \times 10^{18}$ J worldwide. However, there is some debate regarding the actual amounts
5	of straw which could be removed from arable soils without jeopardizing their
6	quality, as well as the potential trade-offs in the overall straw-to-energy chain
7	compared to the use of fossil energy sources.
8	Here, we used a deterministic model of C and N dynamics in soil-crop
9	systems to simulate the effect of straw removal under various sets of soil, climate
10	and crop management conditions in northeastern France. Model results in terms of
11	nitrate leaching, soil C variations, nitrous oxide and ammonia emissions were
12	subsequently inputted into the life cycle assessment (LCA) of a particular bio-
13	energy chain in which straw was used to generate heat and power in a plant
14	producing bio-ethanol from wheat grains.
15	Straw removal had little influence on simulated environmental emissions in
16	the field, and straw incorporation in soil resulted in a sequestration of only 5 to
17	10% of its C in the long-term (30 years). The LCA concluded to significant benefits
18	of straw use for energy in terms of global warming and use of non-renewable
19	energy. Only the eutrophication and atmospheric acidification impact categories
20	were slightly unfavourable to straw use in some cases, with a difference of 8% at
21	most relative to straw incorporation. These results based on a novel methodology
22	thereby confirm the environmental benefits of substituting fossil energy with straw.

- **Key-words**: C-N dynamics cereal straw modelling life cycle
- 2 assessment combined heat and power generation

#### 1 **1INTRODUCTION**

2 Crop residues have recently regained attention as a potentially considerable source of renewable energy. Available residues are estimated at  $10 \times 10^9$  Mg 3 worldwide, corresponding to an energy value of  $47 \times 10^{18}$  J [1]. Among them, cereal 4 5 residues are the largest source, making up two thirds of the total available amount. 6 However, there is an on-going debate on the actual possibilities of straw removal 7 from agricultural cropping systems [2]. As reviewed by the latter authors, current 8 experimental evidences on the effects of straw removal on processes like soil 9 organic matter (SOM) turnover, soil erosion, or crop yields are not consistent 10 because of the strong influence of local conditions (climate, soil type, and crop 11 management). Besides, other types of environmental impacts should be taken into 12 account in order to obtain a complete picture of the advantages and drawbacks of 13 using straw for energy purposes. These include the leaching of nitrate, and the 14 emissions of N trace gases such as ammonia  $(NH_3)$ , nitrogen oxides  $(NO_x)$ , and nitrous oxide (N<sub>2</sub>O), the latter being critical since it is a major contributor to the 15 16 global warming impact of agricultural systems, compared to soil C sequestration 17 [3]. Except for nitrate leaching, there are few references on these effects in the 18 literature, and the patterns are again not consistent across references, and for the 19 same reasons. The time-frame over which the effects of straw removal are 20 investigated is also an issue. For instance, nitrate leaching was shown to decrease in 21 the winter following the first incorporation of wheat straw in a cropping system, 22 compared to a control with no added straw [4]. However the same tendency was 23 reversed after a few years of continued straw incorporation in another trial [5].

1	Deterministic models of C-N dynamics in soil-crop systems provide a
2	unique means of addressing the afore-mentioned issues dealing with straw removal
3	effects. They simulate the major processes governing the impacts cited [6], and
4	make it possible to single out soil, climate, and management factors through
5	scenario analysis [7]. Also, they have the potential to take local context into
6	account, which is important when assessing the environmental impacts of setting
7	up a bio-energy chain in a particular area. Secondly, the environmental assessment
8	should encompass the whole chain to address potential trade-offs along the chain.
9	Life cycle assessment (LCA) provides a comprehensive, standardised framework to
10	deal with such issues, and was already applied to straw [8, 9]. Although the results
11	are generally favourable to cereal straw compared to various fossil feedstock (coal
12	or natural gas), the methodology employed in these studies did not tackle the
13	problem of ecosystem context. For instance, they all used the average emission
14	factor of 1.25% recommended by the IPCC [10] to estimate $N_2O$ emissions,
15	although these are known to be highly variable in time and space [11].
16	In the framework of a case-study on the potential benefits of substituting
17	natural gas with wheat straw in a bio-ethanol production plant, , we therefore set
18	out to predict the effect of wheat straw removal on the dynamics of C and N in
19	arable fields, including N losses (gaseous and leaching). The model is based on
20	CERES [12], as modified to suit French conditions [13]. The second objective of
21	this work was to use the resulting data in a LCA, and to analyze the contribution of
22	field emissions in the overall performance of the straw-based system, compared to
23	the reference system using solely natural gas.

#### 1 2 MATERIALS AND METHODS

#### 2 2.1 System definition and simulation scenarios

3 Our case-study involved the utilization of straw to supply heat and power to 4 an industrial plant producing ethanol from wheat grains via biological conversion. 5 The plant is currently in operation and located in the Picardie region, 200 kms North of Paris, France. It has a production capacity of  $3 \ 10^7 \ 1 \ vr^{-1}$ , and requires 202 6  $10^{6}$  MJ of primary energy per annum. In the reference system (called S1 in the 7 8 following), the plant is powered only by natural gas. In the straw-to-energy system 9 (S2), half of the energy is supplied by a straw-fueled combined heat and power (CHP) unit. The annual straw requirement thus amounts to 96  $10^3$  Mg (dry matter 10 11 basis). Estimates of wheat straw availability around the plant resulted in a collection area of 6 000 km<sup>2</sup> (C. Jacquin, Arvalis, personal communication). 12 However, the studied area was extended to a wider area of ca. 22 000 km<sup>2</sup>, 13 14 encompassing 4 administrative "departments" to investigate the impact of the plant 15 location itself relative to the spatial availability of straw. The study area comprises mostly cropland, of which 45% are planted with 16 cereals. Cereals are mostly rotated with winter oilseed rape and sugar-beet, and 17 potatoes to a little extent. The major soil types occurring in this area are luvisols, 18 19 cambisols, and rendzinas (Soil Survey, INRA Orléans, France). The climate is 20 continental, with influence from the sea in the western end of the zone. To capture 21 the variability in environmental emissions resulting from the differences in climatic 22 and soil conditions across the zone under study, we selected three soils 23 representative of the major types occurring in this zone. Likewise, we selected three

weather stations along a 250 km southeast-northwest transect across the study area.
Table 1 gives the weather statistics for these three stations, while Table 2 lists
selected characteristics of the three soils. The latter comprise an orthic luvisol, a
redoxic luvisol, and a rendzina (FAO classification, [14]). In previous work, the
CERES model was tested in details against experimental data for all three soils
(Table 2).

7 CERES was run on a combination of soil types, weather stations, and crop management scenarios. Management included only two variants: the type of crop 8 9 rotations in which the wheat crops were grown, and the frequency at which wheat 10 straw was removed from the arable field. This frequency, expressed as an average 11 of removal events per year, varied from 0% (no removal) to 100% (straw removed 12 every year). The rotations comprised the following crops: winter wheat (WW), 13 winter barley (WB), silage maize (SM) and grain maize (GM), sugar-beet (SB), and 14 winter oilseed rape (WOR). The following rotations were simulated: WOR-WW-15 WB; WOR-WW-WW; SB-WW-WB; SB-WW-WW; SM-WW; GM-WW; WW in 16 mono-culture. For each WW crop appearing in the rotations, two types were of 17 straw management were simulated upon harvest: removal or soil return. Each 18 rotation thus involved a number of variants according to the occurrence of WW in 19 them. For instance, the WW monoculture involved four years, and thus five 20 variants: straw removal every year; removal every two, three of four years; and no 21 removal in any of the four years.

The rotations and their variants were run on historical series of weather data spanning the 1980-2000 time period. Crop management variables other than straw removal (fertilizer application, sowing date, etc..) were set according to current

recommended practices in the area. A simple N balance was implemented in the
 model to calculate fertilizer N application for winter crops.

3 2.2 The CERES model

4 CERES is a mechanistic model simulating the dynamics of water, carbon 5 and nitrogen in soil-crop systems. It runs on a daily time step and is available for a 6 large range of arable species [12]. It runs from standard weather data incuding: 7 solar radiation, rainfall, air temperature and potential evapo-transpiration. 8 CERES comprises three main sub-models. First, a physical module 9 simulates the transfer of heat, water and nitrate down the soil profile, as well as soil 10 evaporation, plant water uptake and transpiration in relation to climatic demand. 11 Water infiltrates down the soil profile following a tipping-bucket approach, and 12 may be redistributed upwards after evapo-transpiration has dried some soil layers. 13 In both of these equations, we introduced the generalized Darcy's law in order to 14 better simulate water dynamics in fine-textured soils [15]. Next, a micro-biological 15 module simulates the turnover of organic matter in the plough layer, involving both 16 mineralization and immobilisation of inorganic N. It comprises three endogenous 17 soil OM pools: microbial biomass, active humus ('humads'), and passive humus, 18 which decompose according to first-order kinetics, and partly recycle into the 19 microbial biomass. A module for predicting the emissions of N<sub>2</sub>O via the soil 20 nitrification and denitrification pathway was recently incorporated [16]. Also, an 21 ammonia (NH<sub>3</sub>) volatilization module was included in CERES [17]. Lastly, crop 22 net photosynthesis is a linear function of intercepted radiation according to the 23 Monteith approach, with interception depending on leaf are index based on Beer's

law of diffusion in turbid media. Photosynthates are partitioned on a daily basis to
 currently growing organs (roots, leaves, stems, fruits) according to crop
 development stage. The latter is driven by the accumulation of growing degree
 days, as well as cold temperature and day-length for crops sensitive to vernalization
 and photoperiod. Crop N uptake is computed through a supply/demand scheme,
 with soil supply depending on soil nitrate and ammonium concentrations and root
 length density.

#### 8 2.3 Life cycle assessment

9 Life cycle assessment (LCA) was conducted within the framework provided 10 by the BioFit project [8], dedicated to the LCA of bio-fuels in european context. 11 The analysis comprised the bio-ethanol from wheat grains chain, along with the 12 straw for heating and triticale for CHP chains. It follows the ISO norms 14040 and 13 14041 regarding the various stages of LCA: goal and scope definition, inventory 14 analysis, impact assessment, and interpretation [9]. In our case, the functional unit 15 is one litre of ethanol, produced either using natural gas (reference system, S1) or a 16 50%-50% mix of gas and straw (alternative system, S2). The systems are depicted 17 in Figure 1, which highlights the main differences between them. In the S2 system, 18 the removal of straw results in the loss of SOM as well as nutrients, which are 19 supplemented in mineral form. The loss of nutrients is based on the chemical 20 composition of straw, while the SOM loss is a estimated with the CERES 21 simulations. Since straw is considered a by-product of wheat grain production, all 22 the emissions resulting from the cultivation of wheat are allocated to the grains [9]. 23 The wheat straw is pressed into bales, transported by tractor to a temporary storage

on the farm, prior to being collected by trucks and transported to the bio-ethanol
plant. The bales are stored there for a short time and fed directly into a dedicated
boiler for CHP generation. The inventory of environmental outputs was based on
BioFit data, which was supplemented it with data from a more recent LCA based
on the same plant under study here [18]. Following the latter study, we used a
weight-based allocation ratio to partition impacts between the product of interest
(ethanol) and its main by-product (wheat meal for animal feed).

8 The following impact categories were analysed: depletion of natural 9 resources, global warming, atmospheric acidification, eutrophication, and potential 10 for ozone formation. The impacts were expressed in equivalent substances: carbon 11 dioxide (CO<sub>2</sub>) for global warming, sulphur dioxide (SO<sub>2</sub>) for acidification, nitrate 12 (NO<sub>3</sub><sup>-</sup>) for eutrophication, and ethene (C<sub>2</sub>H<sub>4</sub>) for potential ozone formation.

#### 13 3 RESULTS AND DISCUSSION

#### 143.1 Field emissions and effect of straw removal

## 15 3.1.1 Effects of soil type, geographical location, climatic year and crop 16 rotation

Prior to analysing the effects of straw removal *per se*, it is interesting to try
and rank the effects of the other factors included in the simulation scenarios,

including soil type, geographical location, climatic year and crop rotation. Among them, climatic year appeared as the most sensitive factor, the other factors having a similar but smaller influence on model outputs. This is illustrated on Figure 1 in the case of wheat grain yields, which shows that inter-annual variability resulted in

1 standard deviations up to five times higher than those resulting from the variations 2 in crop rotation types. However, comparison with yield census data over the region 3 shows that inter-annual variation might be over-estimated by CERES. The standard 4 deviation calculated across the years from these data is indeed much lower than that 5 estimated by the crop model (Figure 1), even if the census data are likely to smooth 6 out variability because they represent regional averages as opposed to the field-7 scale simulations provided by CERES. The latter also over-estimated the average harvested yield by 10% to 35%, which is not surprising since the model does not 8 9 include the effect of pests and diseases, as well as grain losses upon harvest. 10 Over the four combinations of soil type and climate tested in Figure 1, the time-averaged grain yields varied within a 2 Mg ha<sup>-1</sup> range, the two extremes 11 occurring with the rendzina in the drier climatic location (Fagnières), and the deep 12 13 loam in the wetter location (Abbeville). We considered the latter two combinations 14 as regional extremes, given that the third soil type (redoxic luvisol) represented a 15 medium situation in terms of simulated grain yields, whatever the climatic location. 16 This also applied to the other model outputs, with the exception of ammonia emissions, which were lower by 10 to 15 kg N ha<sup>-1</sup> yr<sup>-1</sup> with the redoxic luvisol 17 18 than with the other two soils. 19 Regarding outputs other than crop yields, the effect of climatic location and

19 Regarding outputs other than crop yields, the effect of chinatic location and
20 soil type may be analysed by comparing the four situations resulting from the
21 combination of the above two extreme soils (rendzina and deep loam) and climatic
22 locations (Abbeville and Fagnières). Switching from the wetter location (Abbeville)
23 to the drier one (Fagnières) with the deep loam resulted in a grain yield decrease of
24 1 Mg DM ha<sup>-1</sup>, on average. Deep drainage also decreased 132 mm, which is very

1 close to the 138 mm difference in annual rainfall across the two locations (Table 1). As a consequence, nitrate leaching decreased 14.6 kg N ha<sup>-1</sup> yr<sup>-1</sup> in the drier climate 2 3 compared to the wetter one. Conversely, with the rendzina soil, switching to a wetter climate resulted in a slight increase in grain yields (0.3 Mg DM ha<sup>-1</sup>), two-4 fold higher amounts of deep drainage, and an 8 kg N ha<sup>-1</sup> yr<sup>-1</sup> increase in nitrate 5 6 leaching on average. In both soils, nitrous oxide emissions were relatively low, ranging from 0.15 to 0.60 kg N-N<sub>2</sub>O ha<sup>-1</sup> yr<sup>-1</sup>. This was especially true for the 7 rendzina soil due to its alkaline pH, which is known to reduce N<sub>2</sub>O production from 8 9 denitrification [19]. The emissions were little affected by climate, decreasing at most 0.20 kg N ha<sup>-1</sup> yr<sup>-1</sup> when switching to the drier climate with the deep loam. 10 11 Ammonia volatilisation increased 80% in the drier climate relative to the wetter 12 one, whether with the deep loam or the rendzina soil.

13 CERES predicted a net increase in soil organic matter all soils, even when 14 straw was removed most of the years. The variations in SOM were very similar 15 across climates, and ranged from 0.15 to 0.75 Mg C ha<sup>-1</sup> yr<sup>-1</sup>, with the exception of 16 the rotation involving grain maize for which the increase was much higher, ranging 17 from 1.0 to 1.3 Mg C ha<sup>-1</sup> yr<sup>1</sup>. These values fall in the higher part of the -2.0 to 2.0 18 Mg C ha<sup>-1</sup> yr<sup>-1</sup> range reported for arable soils in the area over the 1970-1998 time 19 period [20], regardless of crop rotations and straw management.

#### 20 **3.1.2** Effect of straw removal frequency on model outputs

Figures 3 and 4 show the effects of crop rotation and straw removal frequency on various model outputs for the two extreme combinations of soil and climate: the rendzina at Fagnières and the deep loam at Abbeville. The differences

across rotations are generally more important than those related to straw
management for a given rotation. The removal of straw thus bore little effect on
field emissions. The highest effects were noted on crop productivity, with grain
yield being negatively affected by straw removal because of a lower net
mineralisation of N in soils. The yield losses ranged between 0.05 to 0.15 tons DM
ha <sup>-1</sup> for each ton of straw removed, which is in line with the 0.13 Mg grain DM ha <sup>-1</sup>
loss reported for various crops in the US [2]. This corresponds to a straw fertiliser
value ranging from 1.5 to 4.5 kg N $Mg^{-1}$ straw, which is lower than the total N
content of the straw, estimated at 6 kg N Mg <sup>-1</sup> straw DM [21]. Harvested straw
yields paralleled grain yields, and varied between 1.0 and 4.6 Mg straw DM ha <sup>-1</sup>
when averaged over the whole rotation (Figure 3, middle).
Regarding the environmental emissions, nitrate losses tended to decrease
with increasing straw removal in all cases (Figure 4, middle). This represents the
balance between two opposite effects: in the short-term, straw incorporation was
shown to decrease nitrate leaching because of a temporary immobilisation of
mineral N by soil microflora [4, 22]. This effect is not always noted [23], for
instance if straw residues are relatively rich in N [24]. In the longer term on the
other hand, after several years of continued incorporation, the increase in SOM may
lead to higher, uncontrolled N mineralization, and to higher nitrate losses as a

20 consequence [5]. Such was the case in the CERES simulations, which spanned 30

21 years.

Similarly to nitrate leaching, N<sub>2</sub>O emissions decreased slightly with
increasing straw removal, with a rate of 0.1 to 0.25 kg N Mg<sup>-1</sup> straw DM (Figure 4,
top). There is little information on such effect in the literature, but the available

1	results generally emphasize the links with N fertilization: straw return to soil
2	increases to soil's denitrification potential, and to a some extent its capacity to
3	produce $N_2O$ [25], but decrease it temporarily in case the incorporation of straw is
4	followed by an application of mineral of organic fertilizers [26, 27]. This was
5	however not the case here. Nitrous oxide emissions were converted in $CO_2$
6	equivalent using a global warming potential of 270, which corresponds to a 100-yr
7	projection [28]. They were subsequently aggregated with the C balance of the soil
8	to estimate the net impact of the cropping system on global warming. Straw
9	removal contributed to increase the global warming impact in both locations
10	(Figure 3, top), by one Mg equivalent $CO_2$ ha <sup>-1</sup> on average, due to the contribution
11	of incorporated straw to SOM in the long-term. Over the 30-yr simulation period,
12	the model simulated a sequestration potential ranging from 0.05 to 0.1 Mg C $Mg^{-1}$
13	of added straw DM, being lowest for the SB-WW-WW rotation in Fagnières. These
14	figures are in the higher range of the values reported in a recent review [29],
15	namely a range of 0.01 to 0.04 Mg C Mg <sup>-1</sup> straw DM. Wilhem et al. [2] cited a
16	much broader range (0.0 to 0.1 Mg C Mg <sup>-1</sup> straw DM), but noted a major
17	interaction with tillage. Here, the type of rotation was also decisive as regards the
18	sequestration capacity of the cropping system, whether the wheat straw was
19	removed or not. The rotation involving grain maize has the highest sequestration
20	potential because of the high level of residue return from the maize crops.
21	

Ammonia volatilisation correlated negatively with the rate of straw removal in Fagnières, and positively in Abbeville (Figure 4, bottom). These two opposite cases reflect different balances between the two factors which could explain the

1	effect of straw management on NH <sub>3</sub> volatilization: on the one hand, incorporating
2	straw in soil increases its capacity to immobilize mineral fertilizer N, and thus
3	decreases the availability of ammonium to volatilization; on the other hand, straw
4	incorporation increases the net mineralization of nitrogen, being produced in the
5	ammonium form, which is likely to increase NH <sub>3</sub> volatilisation. However, as
6	observed after application of urea [30], straw management had little effect on
7	volatilization. The maximum differences between the treatments with total straw
8	return to soil and straw removal ranged from 1 to 2 kg N-NH <sub>3</sub> ha <sup>-1</sup> yr <sup>-1</sup> in Fagnières,
9	representing only 5 à 10% of the absolute emissions, and these deviations were
10	similar in Abbeville except for two rotations.

Lastly, we tested the influence of the timing of straw removal within the rotations, and found it negligible. There was also very little effect of straw management on the water balance, with straw removal either slightly increasing (Fagnières) or decreasing (Abbeville) deep drainage. Straw removal also decreased annual evapo-transpiration by a few percents, because of its slightly decreasing crop yields.

#### 17 **3.2 Life cycle assessment**

For the purposes of the LCA, we selected one crop rotation which was judged representative of the other rotations, namely the WOR-WW-WW rotation. Table 3 summarises the direct emissions data simulated by CERES for the two extreme soil and climate combinations (the rendzina soil in Fagnières and the deep loam in Abbeville), averaged over time. The main differences between the two soils may be summarised as follows: the deep loam emitted more nitrate than the

rendzina by an order of magnitude, three times as much nitrous oxide, and similar
levels of ammonia, while achieving 25% higher yields (Table 3). Originally, there
were two variants for that rotation, since wheat straw could be removed once or
twice per rotation. Since the simulated emissions differed only by a few percents
between the two variants, we decided to consider only the first one (straw removed
once per rotation).

7 These emissions were inputted to the LCA of the two systems (S1 and S2), whose results are shown in Table 4, while Figure 5 shows their breakdown among 8 9 the various phases (agricultural production, transport of grains and straw to the 10 ethanol plant, conversion to ethanol, combustion of straw and ash disposal). As 11 could be expected [8, 9], the substitution of natural gas with straw resulted in a 12 significant reduction of in the global warming impact, along with non-renewable 13 energy consumption. For each litre of ethanol produced, the relative differences 14 between the reference and straw-based systems amounted to 20% for these two 15 impact categories. When the differences in primary energy consumption between 16 S1 and S2 were expressed relative to the amount of straw used in S2 (Table 4), it is 17 interesting to note that they corresponded to the lower heating value (LHV) of straw, which is of 15 MJ kg<sup>-1</sup> DM. Likewise, the CO<sub>2</sub> savings correspond to 100% 18 19 of the theoretical substitution potential for natural gas, since the latter contains 53 g 20 CO<sub>2</sub> per MJ of LHV [31]. This represents 50% of the straw's total C content. The 21 S2 system was thus very efficient at substituting fossil energy and carbon with non-22 renewable feedstock. This stems from the fact that, compared to natural gas, the 23 extra energy required in the S2 system to collect the straw and manufacture and

operate the straw boiler was 20 times lower than the natural gas savings incurred by
 the use of straw (Figure 5).

3 Regarding the other three categories within the scope of the present LCA, 4 the differences were either nil (ozone creation potential), or dependent on the location considered. Compared to the reference system S1, acidification was 8% 5 6 higher in the S2 system in Fagnières and 5% lower in Abbeville, whereas 7 eutrophication was 3% lower in Fagnières and 0.2% higher in Abbeville. These 8 variations resulted from differences in the two sites in terms of nitrate leaching and 9 ammonia emissions response to straw removal, as discussed in the previous 10 section. They emphasize the influence of local ecosystem context on the overall 11 results of the LCA, which also appeared in the energy balances: the latter were less 12 favourable in Fagnières compared to Abbeville because the yields were lower. The 13 efficiency of the agricultural production phase was thus decreased. Lastly, it should 14 be noted that the straw boiler emitted more compounds involved in eutrophication (in the form of NO<sub>x</sub> and NH<sub>3</sub>) than the natural gas boiler, due to the higher N 15 16 content of straw. It may thus be argued that crop management could be adjusted to 17 decrease the N content of straw, however it is likely to be antagonistic with grain 18 quality targets in terms of protein content, if the wheat grain is marketed for food 19 purposes.

#### 20 3.3 Conclusions

Using the framework of life cycle assessment, we evaluated the interest of substituting cereal straw for natural gas for combined heat and power generation in a bio-ethanol plant. As already shown in previous studies, the main benefits lied in

1 the saving of non-renewable resources and the reduction of greenhouse gas 2 emissions, which proved very efficient. The picture was mitigated regarding other 3 impacts such as eutrophication or acidification, whose outcome actually depended 4 on local ecosystem context (ie, soil type and climatic zone). The use of a 5 biophysical model made it possible to take such factors into account, substantiating the idea that impacts occurring on a local scale should be addressed based on local 6 7 characteristics rather than on national or global averages. This will ultimately mean that some biomass production zones will emerge as performing better than others, 8 9 from the point of view of environmental impacts, and thus induce some kind of 10 spatial differentiation with respect to the implementation of biomass chains. 11 Although this idea is rather intuitive, it had not been implemented yet in life cycle 12 assessment. The use of biophysical models may therefore be expected to play a 13 crucial role in the future development of this methodology. 14 The impacts related to human toxicity and eco-toxicity were disregarded in

this study, although they might play a significant role, especially during the straw burning phase. We had decided such impacts lied beyond the scope of the present study, since it was focused on the agricultural production phase and the use of crop residues, which do not directly involve the use of agrochemicals. Also, the data available for toxicity assessment are limited and the methodology is still under development [32]. Future work along this line is therefore essential to provide a more complete picture of straw to energy chains.

22

#### 1 4 ACKNOWLEDGEMENTS

The authors would like to thank J. Da Silveira for her help in the CERES simulations, and Jean-Claude Sourie for coordinating the research project within which this work was carried out. Financial support was provided by the AGRICE grant programme (ADEME, France).

6

#### 7 5 REFERENCES

8 [1] Lal R. World crop residues production and implications of its use as a biofuel.

9 Environ Internat 2005; 31:575-584.

- 10 [2] Wilhem WW, Johnson JMF, Hatfield JL, Voorhees WB, Linden DR. Crop and
- soil productivity response to corn residue removal: a literature review. Agron J
  2004; 96:1-17.
- 13 [3] Robertson GP, Paul EA, Harwood RR. Greenhouse gases in intensive
- agriculture: Contributions of individual gases to the radiative forcing of theatmosphere. Science 2000; 289:1922-1925.
- 16 [4] Garnier P, Neel C, Aita C, Recous S, Lafolie F, Mary B. Modelling carbon and
- nitrogen dynamics in a bare soil with and without straw incorporation. Eur J
  Soil Sci 2003; 54:555-568.
- 19 [5] Catt JA, Howse KR, Christian DG, Lane PW, Harris GL, Goss MJ. Strategies to
- 20 decrease nitrate leaching in the Brimstone Farm Experiment, Oxfordshire, UK,
- 21 1988-93: the effect of straw incorporation. J Agric Sci (Camb) 1998; 131:309-
- 22 319.

1	[6] Diekkrüger B, Söngerath D, Kersebaum KC, McVoy CW. Validity of
2	agroecosystem models. A comparison of results of different models applied to
3	the same data set. Ecol Model 1995; 81:3-29.
4	[7] Ramanarayanan TS, Storm DE, Smolen MD. Analysis of nitrogen management
5	strategies using EPIC. J Am Water Resour Ass 1998; 34:5.
6	[8] Reinhardt GA. Bio-energy for Europe. Which ones fit best? Rapport final projet
7	Biofit. Heidelberg: IFEU; 2000.
8	[9] Hartmann D, Kaltschmitt M. Electricity generation from solid biomass via co-
9	combustion with coal. Energy and emission balances from a German case
10	study. Biomass and Bioenergy 1999; 16:397-406.
11	[10] IPCC. Revised 1996 IPCC Guidelines for National Greenhouse Gas
12	Inventories Reporting Instructions (Volume 1). wwwipccch 1997.
13	[11] Li C, Zhuang YH, Cao M, Crill PM, Dai ZH, Frolking SE et al. Comparing a
14	process-based agro-ecosystem model to the IPCC methodology for developing
15	a national inventory of N2O emissions from arable lands in China. Nutr
16	Cycling Agroecosys 2001; 60:159-175.
17	[12] Jones CA, Kiniry JR. Ceres-N Maize: a simulation model of maize growth and
18	development. Temple, TX: Texas A&M University Press, College Station;
19	1986.
20	[13] Gabrielle B, Kengni L. Analysis and field-evaluation of the CERES models'
21	soil components: Nitrogen transfer and transformation. Soil Sci Soc Am J
22	1996; 60:142-149.
23	[14] Fao Unesco ISRIC. Carte mondiale des sols, légende révisée. Rapport sur les
24	ressources en sols du monde, Rome: Food and Agriculture Organization; 1989.

1	[15] Gabrielle B, Menasseri S, Houot S. Analysis and field-evaluation of the
2	CERES models' water balance component. Soil Sci Soc Am J 1995; 59:1402-
3	1411.
4	[16] Hénault C, Bizouard F, Laville P, Gabrielle B, Nicoullaud B, Germon JC et al.
5	Predicting in situ soil N2O emissions using NOE algorithm and soil data base.
6	Global Change Biol 2005; 11:115-127.
7	[17] Génermont S, Cellier P. A mechanistic model for estimating ammonia
8	volatilization from slurry applied to bare soil. Agric For Meteorol 1997;
9	88:145-167.
10	[18] ADEME. Non-technical barriers to the development of liquid bio-fuels in
11	Europe. Final project report, EEC Altener Programme. Altener; 1995.
12	[19] Hénault C, Cheneby D, Heurlier K, Garrido F, Perez S, Germon JC.
13	Laboratory kinetics of soil denitrification are useful to discriminate soils with
14	potentially high levels of N2O emission on the field scale. Agronomie 2001;
15	21:713-723.
16	[20] Wylleman R, Mary B, Machet JM, Guérif J, Degrendel M. Evolution des
17	stocks de matière organique dans les sols de grande culture: analyse et
18	modélisation. Persp Agricoles 2001; 270:8-14.
19	[21] SOLAGRO. Faisabilité écologique de la valorisation énergétique de la paille
20	en Ile-de-France. Toulouse: ARENE Ile de France / ADEME; 2001.
21	[22] Silgram M, Chambers BJ. Effects of long-term straw management and
22	fertilizer nitrogen additions on soil nitrogen supply and crop yields at two sites
23	in eastern England. J Agric Sci (Camb) 2002; 139:115-127.

1	[23] Thomsen IK, Christensen BT. Cropping system and residue management
2	effects on nitrate leaching and crop yields. Agric Ecosys Environ 1998; 68:73-
3	84.
4	[24] Mitchell RDJ, Webb J, Harrison R. Crop residues can affect N leaching over
5	at least two winters. Eur J Agron 2001; 15:17-29.
6	[25] Avalakki UK, Strong WM, Saffigna PG. Measurements of gaseous emissions
7	from denitrification of applied 15-N. 2. Effects of temperature and added
8	straw. Austr J Soil Res 1995; 33:89-99.
9	[26] Cai Z, Laughlin RJ, Stevens RJ. Nitrous oxide and dinitrogen emissions from
10	soil under different water regimes and straw amendment. Chemosphere 2001;
11	42:113-121.
12	[27] Hao X, Chang C, Carefoot JM, Janzen HH, Ellert BH. Nitrous oxide
13	emissions from an irrigated soil as affected by fertilizer and straw
14	management. Nutr Cycling Agroecosys 2001; 60:1-8.
15	[28] IPCC. Revised IPCC [Intergovernmental Panel on Climate Change]
16	Guidelines for National Greenhouse Gas Inventories: Bracknell, UK; 1996.
17	[29] Soussana JF, Arrouays D, Balesdent J, Chenu C, Ciais P, Gabrielle B et al.
18	Usages du sol, stockage de carbone et effets connexes: état de l'art. In:Arrouays
19	D, Balesdent J, Germon JC, Jayet PA, Soussana JF, Stengel P, Arrouays D,
20	Balesdent J, Germon JC, Jayet PA, Soussana JF, Stengel Ps. Stocker du
21	carbone dans les sols agricoles de France ?, Paris: INRA Expertise collective;
22	2002, p. 131-133.
23	[30] Tian GM, Cai ZC, Cao JL, Li XP. Factors affecting ammonia volatilisation
24	from a rice-wheat rotation system. Chemosphere 2001; 42:123-129.

1	[31] Riedacker A. Effet de serre et politiques de lutte contre le changement
2	climatique. Mondes en développement 2003; 31:47-70.
3	[32] Pant R, Hoof Gv, Schowanek D, Feijtel TCJ, Koning Ad, Hauschild M et al.
4	Comparison between three different LCIA methods for aquatic ecotoxicity and
5	a product environmental risk assessment: insights from a detergent case study
6	within OMNIITOX. Int J LCA 2004; 9:296-306.
7	[33] Gabrielle B, Da-Silveira J, Houot S, Francou C. Simulating urban waste
8	compost impact on C-N dynamics using a biochemical index. J Envion Qual
9	2004; 33:2333-2342.
10	[34] Gabrielle B, Roche, R., Angas, P., Cantero-Martinez, C., Cosentino, L.,
11	Mantineo, M., Langensiepen, M., Hénault, C., Laville, P., Nicoullaud, B.,
12	Gosse G. A priori parameterisation of the CERES soil-crop models and tests
13	against several European data sets. Agronomie 2002; 22:119-132.
14	

- **Table 1.** Weather statistics for three sites along an East-West transect across the
- 2 studied area.

Site name	Cumulative Cumulative		Mean air	Mean global
	rainfall	potential evapo-	temperature	radiation
	transpiration			
	mm yr <sup>-1</sup>		°C	$MJ m^{-2} d^{-1}$
Abbeville	747	637	10.3	10.65
Fagnières	636	694	10.4	11.55
Mons-en-	627	665	10.6	10.73
Chaussées				

- **Table 2.** Selected properties of the representative soils used in the field
- 2 simulations. The references describe the test of CERES against experimental data
- 3 for the various soils.

Soil classification	Selected properties				Reference
(FAO)					
	Topsoil texture	pН	%C	%CaCO3	
	(UDSA triangle)	1			
Orthic Luvisol	loam	7.8	1.2	1	[13]
Redoxic Luvisol	silt-loam	6.9	1.1	< 1	[33]
Rendzina on chalk	silt-loam	8.7	1.9	75	[34]

**Table 3.** Field emissions used in the LCA for the selected agronomic scenarios: the rendzina soil at Fagnières, and the deep loam at Abbeville, for the oilseed rape-wheat-wheat rotation. Wheat straw is either returned to soil, which corresponds to the reference system (S1), or removed once per rotation, which corresponds the straw-based system S2. The emissions are averaged over the various climatic years simulated. The global warming impact is calculated as the sum of C sequestration in soil organic matter (negative) and the emissions of N<sub>2</sub>O, converted to CO<sub>2</sub> based on a global warming power of 270. The contribution of N<sub>2</sub>O is singled out.

Location Global warming		Ammonia	Nitrate	Crop yield	1	
	impact			leaching		
		N <sub>2</sub> O			Grains	Harvested
		part				straw
	kg eq. C-	CO2 ha <sup>-1</sup>	kg N-NH <sub>3</sub>	kg N-NO3	Mg l	DM yr <sup>-1</sup>
			ha <sup>-1</sup>	ha <sup>-1</sup>		
			Reference s	ystem (S1)		
Fagnières	-800	78	16.2	5.5	7.12	0
Abbeville	-860	210	19.2	48.4	9.52	0
Straw-based system (S2)						
Fagnières	-680	78	17.0	5.0	7.14	1.05
Abbeville	-660	200	17.0	44.0	9.25	1.34

**Table 4.** LCA results for the reference (S1) and straw-based (S2) systems, for the selected agronomic scenarios involving the rendzina soil at Fagnières and the deep loam at Abbeville. For system S2, fluxes are expressed as a difference relative to S1 (S2 - S1). A positive value thus indicates a reduction in environmental impact. The functional unit is one litre of bio-ethanol in the S1 and (S2 - S1) columns, whereas in the rightmost column the fluxes are expressed relative to one kg of straw dry matter used in the bio-ethanol production process.

				(S2-S1)/kg of
Impact category	Unit	<b>S1</b>	(S2 - S1)	straw
Non-renewable	energy	consumption		
Fagnières	MJ	13.0	3.2	15.1
Abbeville	MJ	12.3	3.2	14.9
<b>Global warming</b>	(100 years)			
Fagnières	g eq. CO <sub>2</sub>	977.1	165.1	- 780
Abbeville	g eq. CO <sub>2</sub>	918.4	160.1	- 762
Acidification				
Fagnières	g eq. SO <sub>2</sub>	9.6	- 0.7	3.12
Abbeville	g eq. SO <sub>2</sub>	7.9	0.3	1.21
Eutrophication				
Fagnières	g eq. NO <sub>3</sub>	24.0	- 0.7	3.06
Abbeville	g eq. NO <sub>3</sub>	40.8	0.8	- 4.11
<b>Ozone creation</b>	potential			
Fagnières	g eq. Ethen	0.3	0.01	- 0.02
Abbeville	g eq. Ethen	0.3	0.00	- 0.02

## 1 Figure captions

2	Figure 1. Schematic of the reference system (S1, solid line) and the straw-
3	based system (S2, dashed line). In the S2 system, the CHP in the ethanol
4	conversion plant is operated from a 50%-50% mix of straw and natural gas.
5	Figure 2. Effect of soil type, geographical location (Fagnieres or Abbeville
6	- Abbe.) and climatic year on the wheat yields simulated by CERES. The bars
7	correspond to the mean yields averaged over the various climatic years and crop
8	rotations simulated for each combination of soil type and geographical location.
9	Two series of error bars are shown: one corresponding to the standard deviation
10	across the various crop rotation and straw removal scenarios (n=20), and the other
11	to the average standard deviation across the various climatic years (n=31). The last
12	bar shows the average yield, as estimated from the census data over the study area,
13	and the standard deviation over the 1980-1995 period.
14	Figure 3. Effect of straw removal frequency on the emissions of $N_2O$ (top),
15	nitrate (middle), and ammonia (bottom) in the rendzina and deep loam soils, for the
16	various rotations (WOR-WW-WW: ●; WOR-WW-WB: ○; SM-WW: ▲; GM-
17	WW: $\Delta$ ; SB-WW-WW: <b>•</b> ; SB-WW-WB: $\Box$ ; wheat mono-culture: $\Diamond$ ). The solid lines
18	are linear regressions against straw removal frequency for each rotation. Note the
19	differences in scales for the <i>y</i> -axis between the two soils.
20	Figure 4. Effect of straw removal frequency on cropping systems'
21	greenhouse gas (GHG) balance (top), straw output (middle), and wheat grain yields
22	(bottom) in the rendzina and deep loam soils, for the various rotations (WOR-WW-
23	WW: ●; WOR-WW-WB: ○; SM-WW: ▲; GM-WW:Δ; SB-WW-WW: ■; SB-

1	WW-WB: $\Box$ ; wheat mono-culture: $\Diamond$ ). The solid lines are linear regressions against
2	straw removal frequency for each rotation. The GHG balance, expressed in
3	equivalent $CO_2$ net emissions, compounds two items: $N_2O$ emissions, and soil
4	carbon content variations.
5	Figure 5. Breakdown of the bio-ethanol LCA results among the various
6	production phases: agricultural production (AGRICULT), transport of grains and
7	straw to ethanol plant (TRANSPORT), conversion of wheat grains to ethanol
8	(EtOH), combustion of straw and ash disposal (COMBUSTION). S1 is the
9	reference system while S2 is the straw-based system, and the agronomic scenario
10	corresponds to the deep loam soil.
11	
12	





 $\omega 4 \omega$ 





