



**HAL**  
open science

# Estimation of the Memory Parameter of the Infinite Source Poisson Process

Gilles Fay, François Roueff, Philippe Soulier

► **To cite this version:**

Gilles Fay, François Roueff, Philippe Soulier. Estimation of the Memory Parameter of the Infinite Source Poisson Process. 2007. hal-00008797v3

**HAL Id: hal-00008797**

**<https://hal.science/hal-00008797v3>**

Preprint submitted on 21 Mar 2007 (v3), last revised 3 Sep 2007 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ESTIMATION OF THE MEMORY PARAMETER OF THE INFINITE SOURCE POISSON PROCESS

GILLES FAÏ, FRANÇOIS ROUEFF, AND PHILIPPE SOULIER

ABSTRACT. Long range dependence induced by heavy tails is a widely reported feature of internet traffic. Long range dependence can be defined as the regular variation of the variance of the integrated process, and half the index of regular variation is then referred to as the *Hurst index*. The infinite source Poisson process (a particular case of which is the  $M/G/\infty$  queue) is a simple and popular model with this property, when the tail of the service time distribution is regularly varying. The Hurst index of the infinite source Poisson process is then related to the index of regular variation of the service times. In this paper, we present a wavelet based estimator of the Hurst index of this process, when it is observed either continuously or discretely over an increasing time interval. Our estimator is shown to be consistent and robust to some form of nonstationarity. Its rate of convergence is investigated.

*Keywords: heavy tails, internet traffic, long range dependence, Poisson point processes, semiparametric estimation, wavelets.*

## 1. INTRODUCTION

We consider the infinite source Poisson process with random transmission rate defined by

$$(1.1) \quad X(t) = \sum_{\ell \in \mathbb{N}} U_{\ell} \mathbb{1}_{\{t_{\ell} \leq t < t_{\ell} + \eta_{\ell}\}}, \quad t \geq 0,$$

where

- the arrival times  $\{t_{\ell}\}_{\ell \geq 0}$  are the points of a unit rate homogeneous Poisson process on the positive half-line, independent of the initial conditions;
- the durations and transmission rates  $\{(\eta_{\ell}, U_{\ell})\}$  are independent and identically distributed random variables with values in  $(0, \infty) \times \mathbb{R}$  and independent of the Poisson process and of the initial conditions.

This process was considered by Resnick and Rootzén (2000), Mikosch et al. (2002) among others. The  $M/G/\infty$  queue is a special case, namely, for  $U_{\ell} \equiv 1$ . An important motivation for the infinite source Poisson process is to model the instantaneous rate of the workload going through an Internet link. Although too simple models are generally not relevant for the Internet traffic at the packet level, it is generally admitted that rather simple

models can be used for higher level (the so called *flow level*) traffic such as TCP or HTTP sessions, one of them being the infinite source Poisson process (see Barakat et al., 2002). One way to empirically analyze the Internet traffic at the flow level using the infinite source Poisson process would consist in retrieving all the variables  $\{t_\ell, \eta_\ell, U_\ell\}$  involved in the observed traffic during a given period of time, but this would require to collect all the relevant information in the packets headers (such as source and destination addresses) for separating the aggregated workload into transmission rates at a pertinent level (see Duffield et al., 2002, for many insights about this problem).

It is well known that heavy tails in the durations  $\{\eta_k\}$  result in long range dependence of the process  $X(t)$ . Long range dependence can be defined by the regular variation of the autocovariance of the process or more generally by the regular variation of the variance of the integrated process:

$$\text{var} \left( \int_0^t X(s) ds \right) = L(t)t^{2H},$$

where  $L$  is a slowly varying function at infinity and  $H > 1/2$  is often referred to as the *Hurst index* of the process. For the infinite source Poisson process, the Hurst index  $H$  is related to the tail index  $\alpha$  of the durations by the relation  $H = (3 - \alpha)/2$ . The long range dependence property has motivated many empirical studies of Internet traffic and theoretical ones concerning its impact on queuing (these questions are studied in the  $M/G/\infty$  case in Parulekar and Makowski, 1997).

However, up to the best of our knowledge, no statistical procedure to estimate  $H$  has been rigorously justified. It is the aim of this paper to propose an estimator of the Hurst index of the infinite source Poisson process, and to derive its statistical properties. We propose to estimate  $H$  (or equivalently  $\alpha$ ) from a path of the process  $X(t)$  over a finite interval  $[0, T]$ , observed either continuously or discretely. In practice this can be done by counting all the packets going through some point of the network and then collect local traffic rate measurements. Our estimator is based on the so-called wavelet coefficients of a path. There is a wide literature on this methodology to estimate long range dependence, starting as early as Wornell and Oppenheim (1992) for instance, but we are not aware of rigorous results for non Gaussian or non Stable processes. The main contribution of this paper is thus the proof of the consistency of our estimator. We also investigate the rate of convergence of the estimator in the case  $\alpha > 1$ . If the process is observed continuously, the rate of convergence is good. In the case of discrete observations, the rate is much smaller. Also, the choice of the tuning parameters of the estimators is much more restricted in the latter case, and practitioners should perhaps be aware of this; see Section 4.3 for details.

Outline of the paper. The process  $X$  is formally defined in Section 2. We state our assumptions and, using a point process representation of  $X$ , we establish some of its main properties. The wavelet coefficients are defined and the scaling property of their variances

is obtained in Section 3. The estimator is defined and its properties are established in section 4. The Appendix contains technical Lemmas.

## 2. BASIC PROPERTIES OF THE MODEL

**2.1. Assumptions.** We now introduce the complete assumption on the joint distribution of the transmissions rates and durations.

*Assumption 1.* (i) The random vectors  $\{(\eta, U), (\eta_\ell, U_\ell), \ell \in \mathbb{Z}\}$  are independent with common distribution  $\nu$  on  $(0, \infty) \times \mathbb{R}$  and independent of the homogeneous Poisson point process on the real line with points  $\{t_\ell\}_{\ell \in \mathbb{Z}}$  such that  $t_\ell < t_{\ell+1}$  for all  $\ell$  and  $t_{-1} < 0 \leq t_0$ ;

(ii) there exists a positive integer  $p^*$  such that  $\mathbb{E}[|U|^{p^*}] < \infty$ ;

(iii) there exist a real number  $\alpha \in (0, 2)$  and positive functions  $L_0, \dots, L_{p^*}$  slowly varying at infinity such that for all  $t > 0$  and  $p = 0, \dots, p^*$ ,

$$(2.1) \quad H_p(t) := \mathbb{E} [ |U|^p \mathbb{1}_{\{\eta > t\}} ] = L_p(t) t^{-\alpha} .$$

Since  $\eta > 0$ , the functions  $H_p$  are continuous at zero and  $H_p(0) = \mathbb{E}[|U|^p]$ . Condition (2.1) is equivalent to saying that the functions  $H_p, p = 0, 1, \dots, p^*$ , are regularly varying with index  $-\alpha$ . If  $\alpha > 1$  and  $p^* \geq 2$ , Assumption 1 and Karamata's theorem imply the following asymptotic equivalence:

$$(2.2) \quad \mathbb{E} [ U^2 \{\eta - t\}_+ ] = \mathbb{E} \left[ U^2 \int_{v=t}^{\infty} \mathbb{1}_{\{v < \eta\}} dv \right] = \int_{v=t}^{\infty} H_2(v) dv \sim \frac{1}{\alpha - 1} L_2(t) t^{1-\alpha} .$$

*Remark 2.1.* Assumption 1 will be used with  $p^* = 2$  to prove the regular variation of the autocovariance function of the process  $X$  and with  $p^* = 4$  to prove consistency of our estimators. It can be related to the theory of multivariate regular variation (see for instance Maulik et al., 2002). But the definitions of multivariate regular variation involve vague convergence and do not necessarily ensure the convergence of moments required here.

*Remark 2.2.* We do not assume that  $U$  is nonnegative. This allows to consider applications other than teletraffic modeling. For instance, the process  $X$  could be used to model the volatility of some financial time series.

*Remark 2.3.* In the sequel, we will often have to separate the cases  $\mathbb{E}[\eta] = \infty$  and  $\mathbb{E}[\eta] < \infty$ . These cases are respectively implied by  $\alpha < 1$  and  $\alpha > 1$ . If  $\alpha = 1$ , the finiteness of  $\mathbb{E}[\eta]$  depends on the precise behavior of  $L_0$  at infinity.

*Example 2.1.* Assumption 1 implies in particular that the tail of the distribution of  $\eta$  is regularly varying with index  $\alpha$ . This in turns implies Assumption 1 if  $U$  and  $\eta$  are independent and  $\mathbb{E}[|U|^{p^*}] < \infty$ , in which case the functions  $L_p$  differ by a multiplicative constant.

*Example 2.2.* Assumption 1 also holds in the following case which is of interest in teletraffic modeling. In a TCP/IP traffic context,  $\eta$  and  $U$  represent respectively the duration of a download session and its intensity (bit rate). Then  $W := U\eta$  represents the amount of transmitted data. We assume that, for some  $u_0 > 0$ , there exist two regimes,  $U \geq u_0$  (xDSL/LAN/Cable connection), and  $U \in (0, u_0)$  (RTC connection) such that:

- The distribution of  $W$  given  $U = u \geq u_0$  is heavy tailed and independent of  $u$ :  $\mathbb{P}(W \geq w|U = u) = L(w)w^{-\alpha}$  ;
- The distribution of  $W$  given  $U = u \in (0, u_0)$  is light-tailed uniformly of  $u$ ; for instance, we assume exponentially decaying tails:

$$\mathbb{P}(W \geq w|U = u) \leq \exp(-\beta w^{-\gamma}),$$

for some  $\beta > 0$  and  $\gamma > 0$ .

An explicit example for two such regimes is obtained when the conditional density of  $W$  given  $U = u$  is equal to  $\alpha w^{-\alpha-1} \mathbb{1}_{\{w \geq 1\}}$  if  $u \geq u_0$  and  $\exp(-w)$  if  $u < u_0$ .

Concerning the distribution of  $U$  we only assume that:

- $\mathbb{P}(U \geq u_0) > 0$ ,  $\mathbb{E}[|U|^{-\alpha-\epsilon}] < \infty$  for some  $\epsilon > 0$  and  $\mathbb{E}[|U|^{p^*}] < \infty$ .

Then (2.1) holds for  $p \leq p^*$ . Indeed,

$$\begin{aligned} \mathbb{E}[U^p \mathbb{1}_{\{\eta > t\}} \mathbb{1}_{\{U \geq u_0\}}] &= \mathbb{E}[U^p \mathbb{1}_{\{W > Ut\}} \mathbb{1}_{\{U \geq u_0\}}] \\ &= \mathbb{E}[U^p L(Ut)(Ut)^{-\alpha} \mathbb{1}_{\{U \geq u_0\}}] \\ (2.3) \quad &= L(t)t^{-\alpha} \mathbb{E}[U^{p-\alpha} L(Ut)/L(t) \mathbb{1}_{\{U \geq u_0\}}]. \end{aligned}$$

Since  $L$  is slowly varying at infinity,  $\lim_{t \rightarrow \infty} L(ut)/L(t) = 1$ , uniformly with respect to  $u$  in compact sets of  $(0, +\infty)$  and there exists  $t_0 > 0$  such that for  $u \geq u_0$ ,  $t \geq t_0$

$$\frac{L(ut)}{L(t)} \leq (1 + \alpha)u^{\alpha/2}.$$

See *e.g.* Resnick 1987, Proposition 0.8. Then by the Dominated Convergence Theorem,

$$(2.4) \quad \lim_{t \rightarrow \infty} \mathbb{E}[U^{p-\alpha} L(Ut)/L(t) \mathbb{1}_{\{U \geq u_0\}}] = \mathbb{E}[U^{p-\alpha} \mathbb{1}_{\{U \geq u_0\}}]$$

Consider now the low-bitrate regime. We have, since, for all  $x > 0$ ,  $\exp\{-\beta x^\gamma\} \leq Cx^{-\alpha-\epsilon}$  for some positive constant  $C$ ,

$$\mathbb{E}[U^p \mathbb{1}_{\{\eta > t\}} \mathbb{1}_{\{U < u_0\}}] \leq \mathbb{E}[U^p \exp\{-\beta(Ut)^\gamma\} \mathbb{1}_{\{U < u_0\}}] \leq Ct^{-\alpha-\epsilon} \mathbb{E}[U^{p-\alpha-\epsilon} \mathbb{1}_{\{U < u_0\}}].$$

Using the assumption on  $U$ , since  $p \geq 0$ , the rightmost expectation in the previous display is finite and we get that

$$\lim_{t \rightarrow \infty} t^\alpha L^{-1}(t) \mathbb{E}[U^p \mathbb{1}_{\{\eta > t\}} \mathbb{1}_{\{U < u_0\}}] = 0.$$

This, (2.3) and (2.4) imply that, as  $t \rightarrow \infty$ ,  $\mathbb{E}[U^p \mathbb{1}_{\{\eta > t\}}] t^\alpha \sim L(t) \mathbb{E}[U^{p-\alpha} \mathbb{1}_{\{U \geq u_0\}}]$  hence is slowly varying and Assumption 1 holds.

**2.2. Point Process representation and stationary version.** Let  $\mathcal{N}$  denote a Poisson point process on a set  $E$  endowed with a  $\sigma$ -field  $\mathcal{E}$  with intensity measure  $\mu$ , that is a random measure such that for any disjoint  $A_1, \dots, A_p$  in  $\mathcal{E}$ ,  $\mathcal{N}(A_1), \dots, \mathcal{N}(A_p)$  are independent random variables with Poisson law with respective parameters  $\mu(A_i)$ ,  $i = 1, \dots, p$ . The main property of Poisson point processes that we will use is the following cumulant formula (see for instance Resnick, 1987, Chapter 3). For any positive integer  $p$  and functions  $f_1, \dots, f_p$  such that  $\int |f_i| d\mu < \infty$  and  $\int |f_i|^p d\mu < \infty$  for all  $i = 1, \dots, p$ , the  $p$ -th order joint cumulant of  $\mathcal{N}(f_1), \dots, \mathcal{N}(f_p)$  exists and is given by

$$(2.5) \quad \text{cum}(\mathcal{N}(f_1), \dots, \mathcal{N}(f_p)) = \int f_1 \cdots f_p d\mu .$$

Let  $N_S$  be the point processes on  $\mathbb{R} \times (0, \infty) \times \mathbb{R}$  with points  $(t_\ell, \eta_\ell, U_\ell)_{\ell \in \mathbb{Z}}$ , *i.e.*  $N_S = \sum_{\ell \in \mathbb{Z}} \delta_{t_\ell, \eta_\ell, U_\ell}$ . Under Assumption 1(i), it is a Poisson point process with intensity measure  $\text{Leb} \otimes \nu$ , where  $\text{Leb}$  is the Lebesgue measure on  $\mathbb{R}$ . For  $t, u \in \mathbb{R}$ , define

$$\begin{aligned} A_t &= \{(s, v) \in \mathbb{R} \times \mathbb{R}_+ \mid s \leq t < s + v\} ; \\ B_u &= \{\lambda u \mid \lambda \in [1, \infty)\} \quad . \end{aligned}$$




FIGURE 1. The set  $A_t$

We can now show that if  $\mathbb{E}[\eta] < \infty$ , one can define a stationary version for  $X$  and provide its second order properties.

**Proposition 2.1.** *If Assumption 1(i) holds and  $\mathbb{E}[\eta] < \infty$ , then the process*

$$(2.6) \quad X_S(t) = \sum_{\ell \in \mathbb{Z}} U_\ell \mathbb{1}_{\{t_\ell \leq t < t_\ell + \eta_\ell\}}$$

*is well defined and strictly stationary. It has the following point process representation*

$$(2.7) \quad X_S(t) = \int_0^\infty N_S(A_t \times B_u) du - \int_{-\infty}^0 N_S(A_t \times B_u) du .$$

*Let  $K_0 = \sup\{\ell > 0 \mid t_{-\ell} + \eta_{-\ell} > 0\}$ ,  $\tilde{U}_\ell = U_{-\ell}$  and  $\tilde{\eta}_\ell = \eta_{-\ell} + t_{-\ell}$ , then, for all  $t \geq 0$ ,*

$$(2.8) \quad X_S(t) = \sum_{\ell=1}^{K_0} \tilde{U}_\ell \mathbb{1}_{\{t < \tilde{\eta}_\ell\}} + X(t).$$

If moreover  $p^* \geq 2$ , then  $X_S$  has finite variance and

$$\begin{aligned} \mathbb{E}[X_S(t)] &= \mathbb{E}[U\eta] , \\ \text{cov}(X_S(0), X_S(t)) &= \mathbb{E}[U^2(\eta - t)_+] = \int_t^\infty H_2(v)dv . \end{aligned}$$

*Remark 2.4.* Note that if  $\alpha > 1$ , then  $\mathbb{E}[\eta] < \infty$  and by Karamata's theorem

$$\text{cov}(X_S(0), X_S(t)) \sim \frac{1}{\alpha - 1} L_2(t) t^{1-\alpha} , \quad t \rightarrow +\infty .$$

*Proof.* The number of non vanishing terms in the sum (2.6) is  $N_S(A_t \times \mathbb{R})$  and has a Poisson distribution with mean  $\mathbb{E} \int_{\mathbb{R}} \mathbb{1}_{A_t}(s, \eta) ds = \mathbb{E}[\eta]$ . Thus  $X_S$  is well defined and stationary since  $N_S$  is stationary. The number of indices  $\ell > 0$  such that  $t_{-\ell} + \eta_{-\ell} > 0$  is  $N_S(A_0 \times \mathbb{R})$ , hence if  $K_0$  is the largest of those  $\ell$ 's, it is almost surely finite and

$$\sum_{t_\ell < 0} U_\ell \mathbb{1}_{\{t_\ell \leq t < t_\ell + \eta_\ell\}} = \sum_{\ell=1}^{K_0} \tilde{U}_\ell \mathbb{1}_{\{t < \tilde{\eta}_\ell\}} .$$

Hence (2.8).

The point process representation (2.7), Formulas (2.5) and (2.2) finally yield the given expressions for the mean and the covariance.  $\square$

Relation (2.8) shows that the stationary version  $X_S$  can be defined by changing the initial condition of the system. More generally one could consider *any* initial conditions, that is, any process defined as in the right-hand side of (2.8) with  $K_0$  and  $\tilde{\eta}_\ell, \ell > 0$  finite. Since the initial conditions almost surely vanish after a finite period, they have a negligible impact on the estimation procedure. Thus, our result on  $X$  easily generalizes to any such initial conditions, and, in particular, to the stationary version  $X_S$ , when it exists.

Applying similar arguments as those used for showing Proposition 2.1, we obtain

**Proposition 2.2.** *The process  $X$  admits a point process representation:*

$$(2.9) \quad X(t) = \int_0^\infty N_S(A_t^+ \times B_u) du - \int_{-\infty}^0 N_S(A_t^+ \times B_u) du ,$$

where  $A_t^+ = A_t \cap \mathbb{R}_+^2$ .

If Assumption 1 hold with  $p^* \geq 2$ , then the process  $X$  is nonstationary with expectation and autocovariance function given, for  $s \leq t$  by

$$\begin{aligned} \mathbb{E}[X(t)] &= \mathbb{E}[U(\eta \wedge t)] , \\ \text{cov}(X(s), X(t)) &= \mathbb{E}[U^2\{s - (t - \eta)_+\}_+] = \int_{t-s}^t H_2(v)dv . \end{aligned}$$

By the Uniform Convergence Theorem for slowly varying functions, the following asymptotic equivalence of the covariance holds. For any  $\alpha \in (0, 2)$  and all  $t > s > 0$ , as  $T \rightarrow \infty$ ,

$$(2.10) \quad \text{cov}(X(Tt), X(Ts)) \sim CL_2(T) T^{1-\alpha}$$

with  $C = \int_{t-s}^s v^{-\alpha} dv$ .

In accordance with the notation in use in the context of long memory processes, we can define the Hurst index of the process  $X$  as  $H = (3 - \alpha)/2$ , because the variance of the process integrated between 0 and  $T$  increases as  $T^{2H}$ . If  $\alpha < 1$ , then  $H > 1$ . This case has been considered for instance by Resnick and Rootzén (2000)

### 3. WAVELET COEFFICIENTS

**3.1. Continuous observation.** Let  $\psi$  be a bounded real valued function with compact support in  $[0, M]$  and such that

$$(3.1) \quad \int_0^M \psi(s) ds = 0 .$$

For integers  $j \geq 0$  and  $k \in \mathbb{Z}$ , define

$$(3.2) \quad \psi_{j,k}(s) = 2^{-j/2} \psi(2^{-j}s - k).$$

The wavelet coefficients of the path are defined as (see e.g. Cohen, 2003)

$$(3.3) \quad d_{j,k} = \int_0^\infty \psi_{j,k}(s) X(s) ds .$$

Assume that a path of the process  $X$  is observed continuously between times 0 and  $T$ . Since  $\psi_{j,k}$  has support in  $[k2^j, (k+M)2^j]$ , the coefficients  $d_{j,k}$  can be computed for all  $(j, k)$  such that  $T2^{-j} \geq M$  and  $k = 0, 1, \dots, T2^{-j} - M$ .

According to Lemma 5.1, one may define, for all  $j$  and  $k$ ,

$$(3.4) \quad d_{j,k}^S = \sum_{\ell \in \mathbb{Z}} U_\ell \int_{t_\ell}^{t_\ell + \eta} \psi_{j,k}(s) ds .$$

As precised in Lemma 5.1, if  $\mathbb{E}[\eta] < \infty$ , we have  $d_{j,k}^S = \int_0^\infty \psi_{j,k}(s) X_S(s) ds$ . Nevertheless, even if  $\mathbb{E}[\eta] = \infty$ , the sequence of coefficients at a given scale  $j$ ,  $\{d_{j,k}^S, k \in \mathbb{Z}\}$  is stationary. Moreover, the definition (3.4) yields

**Lemma 3.1.** *Let Assumption 1 hold with  $p^* \geq 2$ . We have*

$$(3.5) \quad \mathbb{E}[d_{j,k}^S] = 0, \quad \text{var}(d_{j,k}^S) = \mathcal{L}(2^j) 2^{(2-\alpha)j},$$

where

$$(3.6) \quad \mathcal{L}(z) := z^\alpha \int_0^\infty \int_{-\infty}^\infty \left( \int_{-\infty}^\infty \left\{ \int_t^{t+uz^{-1}} \psi(s) ds \right\}^2 dt \right) w^2 \nu(dw, dv)$$

is slowly varying as  $z \rightarrow \infty$ . More precisely, we have the asymptotic equivalence

$$(3.7) \quad \mathcal{L}(z) \sim C_{\mathcal{L}} L_2(z) \quad \text{as } z \rightarrow \infty,$$

with  $C_{\mathcal{L}} = \alpha \int_0^\infty \int_{-\infty}^\infty \left\{ \int_x^{x+y} \psi(s) ds \right\}^2 dx t^{-\alpha-1} dt > 0$ .

*Proof.* See Appendix 5. □

**3.2. Wavelet coefficients in discrete time.** Let  $\phi$  be a bounded  $\mathbb{R} \rightarrow \mathbb{R}$  function with compact support included in  $[-M + 1, 1]$  and such that

$$(3.8) \quad \sum_{k \in \mathbb{Z}} \phi(t - k) = 1, \quad t \in \mathbb{R}.$$

Let  $I_\phi$  denote the operator defined on the set of functions  $x : \mathbb{R} \rightarrow \mathbb{R}$  by

$$(3.9) \quad I_\phi[x](t) = \sum_{k \in \mathbb{Z}} x(k) \phi(t - k).$$

The wavelet coefficients of  $x$  are then defined as the wavelet coefficients of  $I_\phi[x]$ .

From a computational point of view, it is convenient to chose  $\phi$  and  $\psi$  to be the so-called father and mother wavelets of a multiresolution analysis, see for instance Meyer (1992). The simplest choice is to take  $\phi$  and  $\psi$  associated to the Haar system, in which case  $M = 1$ ,  $\phi = \mathbb{1}_{[0,1]}$  and  $\psi = \mathbb{1}_{[0,1/2]} - \mathbb{1}_{[1/2,1]}$ .

If the process  $X$  is observed discretely, we still denote  $d_{j,k}^D$  its wavelet coefficients :

$$(3.10) \quad d_{j,k}^D = \int \psi_{j,k}(s) I_\phi[X](s) ds .$$

If we observe  $X(0), X(1), \dots, X(T-1)$  for some positive integer  $T$ , we can compute  $d_{j,k}^D$  for all  $j, k$  such that  $0 \leq k \leq 2^{-j}(T - M + 1) - M$ . Roughly, for  $2^j \geq T/M$ , no coefficients can be computed and if  $2^j < T/M$  the number of computable wavelet coefficients at scale  $2^{-j}$  is of order  $T2^{-j} + 1 - M$  for  $j$  and  $T$  large.

*Remark 3.1.* Observe that the choice of time units is unimportant here. Indeed, in Assumption 1, changing the time units simply amounts to adapt the slowly varying functions  $L_k$ 's and the rate of the arrival process  $\{t_k\}$ . Clearly both adaptations do not modify our results as far as precise multiplicative constants are not considered.

**3.3. Averaged observations.** We describe now a third observation scheme for which our results can easily be extended. Suppose that  $T$  is a positive integer and that we observe local averages of the trajectory

$$\bar{X}(k) := \int_k^{k+1} X(t) dt = \int X(t) \phi_H(t-k) dt, \quad k = 0, 1, \dots, T-1,$$

where  $\phi_H := \mathbb{1}_{[0,1]}$  is the Haar wavelet. Let  $\bar{\Gamma}_\phi$  denote the operator on locally integrable functions  $x$  defined by

$$\bar{\Gamma}_\phi[x](t) = \sum_{k \in \mathbb{Z}} \left( \int x(s) \phi_H(s-k) ds \right) \phi(t-k)$$

For this observation scheme, as in Section 3.2, one may compute the wavelet coefficients of the function  $\bar{\Gamma}_\phi[X]$  at all scale and location indices  $(j, k)$  such that  $0 \leq k \leq 2^{-j}(T - M + 1) - M$ . If  $\phi = \phi_H$  and  $\psi$  is the Haar mother wavelet,  $\psi = \mathbb{1}_{[0,1/2)} - \mathbb{1}_{[1/2,1)}$ , then, the wavelet coefficients of  $\bar{\Gamma}_\phi[X]$  are precisely the continuous wavelet coefficients defined in (3.3). For any other choice of  $\phi$  and  $\psi$ , this is no longer true. We will not treat this case but all our results can be extended at the cost of more technicalities.

#### 4. ESTIMATION

Tail index estimation methods do not seem appropriate here for estimating the parameter  $\alpha$ . Indeed, the parameter  $\alpha$  is the tail index of the unobserved durations  $\{\eta_k\}$ , whereas the observed process  $X(t)$  always has finite variance ( $\mathbb{E}[|X(t)|^p] < \infty$  if and only if  $\mathbb{E}[|U^p|] < \infty$  and the marginal distribution of  $X(t)$  is Poisson if  $U = 1$  almost surely). But as shown by Proposition 2.2,  $\alpha$  is related to the second order properties of the process: the coefficient  $H = (3 - \alpha)/2$  can be viewed as its Hurst index, *i.e.*  $H$  governs the rate of decay of the autocovariance function of the process. Therefore it seems natural to use an estimator of the Hurst index.

**4.1. The estimator.** Lemma 3.1 provides the rationale for the following minimum contrast estimator of  $\alpha$  which is related to the local Whittle estimator, cf. Künsch (1987). Let

- (i)  $\mathbf{d}_{j,k}$  denote the wavelet coefficients which are actually available; these may be obtained from continuous time ( $\mathbf{d}_{j,k} = d_{j,k}$ ) or discrete time ( $\mathbf{d}_{j,k} = d_{j,k}^D$ ) observations;
- (ii)  $\Delta$  be a set of indices  $(j, k)$  of available wavelet coefficients;

Denote the mean scale index over  $\Delta$

$$\delta := \frac{1}{\#\Delta} \sum_{(j,k) \in \Delta} j.$$

The reduced local Whittle contrast function is

$$(4.1) \quad \hat{W}(\alpha') = \log \left( \sum_{(j,k) \in \Delta} \frac{\mathbf{d}_{j,k}^2}{2^{(2-\alpha')j}} \right) + \delta \log(2)(2 - \alpha') .$$

The local Whittle estimator of  $\alpha$  is then defined as

$$(4.2) \quad \hat{\alpha} := \arg \min_{\alpha' \in (0,2)} \hat{W}(\alpha')$$

In order to simplify the proof of our result, from now on, we take  $\Delta$  of the form

$$\Delta = \{(j, k) ; J_0 < j \leq J_1, \quad 0 \leq k \leq n_j - 1\} ,$$

with

$$J = \max\{j; 2^j \leq (T - M + 1)/(M + 1)\} , \\ n_j = 2^{J-j}$$

and integers  $J_0$  and  $J_1$  such that

$$(4.3) \quad 0 < J_0 < J_1 \leq J .$$

The sequence of integers  $J$  depends on  $T$  in such a way that  $2^J \asymp T$ . Note that the dependence of the sequences  $J, J_0, J_1, n_j$  etc. with respect to  $T$  is omitted in our notation.

**4.2. Consistency.** Our estimator is consistent in the potentially unstable case, that is when  $\alpha$  is not assumed to be in  $(1, 2)$ , provided that the assumptions on the functions  $\phi$  and  $\psi$  are strengthened. We assume that

$$(4.4) \quad \int_{-\infty}^{\infty} s \psi(s) ds = 0 ,$$

and there exist constants  $a$  and  $b$  such that for all  $t \in \mathbb{R}$ ,

$$(4.5) \quad \sum_{k \in \mathbb{Z}} k \phi(t - k) = a + bt .$$

These conditions are not satisfied by the Haar wavelet, but hold for any Daubechies wavelets; see Cohen (2003).

**Theorem 4.1.** *Let Assumption 1 hold with  $p^* \geq 4$ . Assume that  $J_0$  and  $J_1$  depend on  $T$  in such a way that*

$$(4.6) \quad \lim_{T \rightarrow \infty} J_0 = \lim_{T \rightarrow \infty} (J_1 - J_0) = \infty ,$$

$$(4.7) \quad \limsup_{T \rightarrow \infty} J_0/J < 1/\alpha ,$$

$$(4.8) \quad \limsup_{T \rightarrow \infty} J_1/J < 1/(2 - \alpha) .$$

*Then  $\hat{\alpha}$  is a consistent estimator of  $\alpha$ . Moreover if  $\alpha \in (1, 2)$ , Conditions (4.4), (4.5) and (4.8) are not necessary for the same result to hold.*

*Remark 4.1.* Conditions (4.6), (4.7) and (4.8) are satisfied by the choice  $J_0 = \lfloor J/2 \rfloor$  and  $J_1 = \lfloor J/2 + \log(J) \rfloor$ .

*Proof of Theorem 4.1.* For clarity of notation, we denote  $\sum_j = \sum_{j=J_0+1}^{J_1}$ ,  $\Delta_j := \{k : (j, k) \in \Delta\}$  and  $\#\Delta_j = n_j$ . Elementary computations give

$$(4.9) \quad \delta = J_0 + 2 + (J_0 - J_1)/(2^{J_1 - J_0} - 1)$$

so that  $\delta - (J_0 + 2) \rightarrow 0$  under (4.6). By Karamata's representation theorem, the slowly varying function  $\mathcal{L}$  defined in (3.6) can be written as

$$\mathcal{L}(z) = c(1 + r(z)) \exp \left\{ \int_1^z \frac{\ell(s)}{s} ds \right\},$$

with  $c > 0$  and  $\lim_{z \rightarrow \infty} \ell(z) = \lim_{z \rightarrow \infty} r(z) = 0$ . Define  $\mathcal{L}_0(z) = c \exp \left\{ \int_1^z \frac{\ell(s)}{s} ds \right\}$ ,  $r^*(z) = \sup_{z' \geq z} |r(z')|$  and  $\ell^*(z) = \sup_{z' \geq z} |\ell(z')|$ . The functions  $r^*$  and  $\ell^*$  are non increasing and tend to zero at infinity. Introduce some notation that will be used throughout the proof.

$$W(\alpha') = \log \left( \sum_j 2^{(\alpha' - \alpha)j} n_j \mathcal{L}(2^j) \right) + \delta \log(2)(2 - \alpha'),$$

$$W_0(\alpha') = \log \left( \sum_j 2^{(\alpha' - \alpha)j} n_j \mathcal{L}_0(2^j) \right) + \delta \log(2)(2 - \alpha'),$$

$$w_{j,0}(\alpha') := \frac{2^{(\alpha' - \alpha)j} n_j \mathcal{L}_0(2^j)}{\sum_{j'} 2^{(\alpha' - \alpha)j'} n_{j'} \mathcal{L}_0(2^{j'})}, \quad w_j(\alpha') := \frac{2^{(\alpha' - \alpha)j} n_j \mathcal{L}(2^j)}{\sum_{j'} 2^{(\alpha' - \alpha)j'} n_{j'} \mathcal{L}(2^{j'})},$$

$$v_j = \mathcal{L}(2^j) 2^{(2-\alpha)j}, \quad \Lambda_j = n_j^{-1} \sum_{k=0}^{n_j-1} \{v_j^{-1}(\mathbf{d}_{j,k})^2 - 1\},$$

$$E(\alpha') := \sum_j w_j(\alpha') \Lambda_j.$$

We have

$$W(\alpha') - W_0(\alpha') = \log \left( 1 + \frac{\sum_j 2^{(\alpha' - \alpha)j} n_j \mathcal{L}(2^j) r(2^j)}{\sum_j 2^{(\alpha' - \alpha)j} n_j \mathcal{L}_0(2^j)} \right).$$

The fraction inside the logarithm in the last display is bounded by  $r^*(2^{J_0})$ , thus for  $J$  large enough

$$\sup_{\alpha'} |W(\alpha') - W_0(\alpha')| \leq Cr^*(2^{J_0}).$$

Standard algebra yields

$$W'_0(\alpha') = \log 2 \sum_j w_{j,0}(\alpha')(j - \delta)$$

$$W''_0(\alpha') = \log^2(2) \sum_j w_{j,0}(\alpha')(j - \sum_{j'} w_{j',0}(\alpha')j')^2.$$

By Lemma 5.13, under (4.6),

$$\lim_{T \rightarrow \infty} W'_0(\alpha) = 0, \quad \lim_{T \rightarrow \infty} W''_0(\alpha) = 2.$$

Thus, there exist  $\eta > 0$  and  $\zeta > 0$  such that

$$\liminf_{T \rightarrow \infty} \inf_{\alpha' \in (\alpha - \eta, \alpha + \eta)} W''_0(\alpha') > \zeta.$$

This implies that for large  $T$ , and some positive constant  $c$

$$(4.10) \quad W(\hat{\alpha}) - W(\alpha) \geq W'_0(\alpha) \log(2)(\hat{\alpha} - \alpha) + c(\hat{\alpha} - \alpha)^2 - 2r^*(2^{J_0})$$

Since  $W'_0(\alpha) \rightarrow 0$  and  $|\hat{\alpha} - \alpha| \leq 2$ , this implies that for all  $\epsilon > 0$ ,

$$(4.11) \quad \limsup_{T \rightarrow \infty} \mathbb{P}((\hat{\alpha} - \alpha)^2 > \epsilon) \leq \limsup_{T \rightarrow \infty} \mathbb{P}(W(\hat{\alpha}) - W(\alpha) \geq c\epsilon).$$

Write

$$(4.12) \quad \begin{aligned} \hat{W}(\alpha') &= W(\alpha') + \log \{1 + E(\alpha')\}, \\ W(\hat{\alpha}) - W(\alpha) &= \hat{W}(\hat{\alpha}) - \hat{W}(\alpha) - \log \{1 + E(\hat{\alpha})\} + \log \{1 + E(\alpha)\} \\ &\leq 2 \sup_{\alpha' \in (0,2)} |\log \{1 + E(\alpha')\}|. \end{aligned}$$

Consistency will follow from (4.11) and (4.12) provided that we can prove that  $\sup_{\alpha' \in (0,2)} |E(\alpha')| = o_P(1)$ . If  $\alpha > 1$ , take  $\varepsilon \in (0, (\alpha - 1)/2)$  such that  $\limsup J_0/J < 1/(\alpha + \varepsilon)$  which is possible by assumption (4.7). Define

$$(4.13) \quad J_2 = \begin{cases} J_1 & \text{if } \alpha \leq 1, \\ J_1 \wedge [J/(\alpha + \varepsilon)] & \text{if } \alpha > 1. \end{cases}$$

so that, for  $T$  large enough,  $J_0 < J_2 \leq J_1$ . Write

$$E(\alpha') = \sum_{j=J_0+1}^{J_2} w_j(\alpha') \Lambda_j + \sum_{j=J_2+1}^{J_1} w_j(\alpha') \Lambda_j =: E_1(\alpha') + E_2(\alpha'),$$

with the convention  $\sum_{j=J_2+1}^{J_1} = 0$  if  $J_2 = J_1$ . By Lemma 5.11,

$$(4.14) \quad \sup_{\alpha' \in (0,2)} |E_1(\alpha')| = O_P(2^{-\xi_1 J})$$

for some positive  $\xi_1$ . Treat now  $E_2$  for  $\alpha > 1$  and  $J_2 = [J/(\alpha + \varepsilon)] > J_1$ . For all  $\alpha' \in (0, 2)$ , we have  $\alpha' - \alpha - 1 < -2\varepsilon$ . Since  $\mathcal{L}$  is slowly varying, we obtain, for some positive constant  $C$ , for all  $j = J_2 + 1, J_2 + 2, \dots, J_1$ ,  $w_j(\alpha') \leq C 2^{-\varepsilon(J_2 - J_0)}$ . Using Lemma 5.10, it follows that

$$(4.15) \quad \mathbb{E} \left[ \sup_{\alpha' \in (0,2)} |E_2(\alpha')| \right] \leq C (J_1 - J_2) 2^{-\varepsilon(J_2 - J_0)} = O(2^{-\xi_2 J}),$$

for some  $\xi_2 > 0$  because  $\limsup J_0/J < 1/(\alpha + \varepsilon)$ . This concludes the proof.  $\square$

### 4.3. Rate of Convergence in the stable case.

**Theorem 4.2.** *Let Assumption 1 hold with  $\alpha \in (1, 2)$  and  $p^* = 4$ . Assume moreover that  $L_4$  is bounded and that  $\mathcal{L}(z) = c + O(z^{-\beta})$  with  $c > 0$  and  $\beta > 0$ .*

*If  $X$  is observed continuously on  $[0, T]$ , that is,  $\mathbf{d}_{j,k} = d_{j,k}$ , then the rate of convergence in probability of  $\hat{\alpha}$  is  $T^{-\beta/(2\beta+\alpha)}$ , obtained for  $J_0 = \lfloor J/\{2\beta + \alpha\} \rfloor$  and  $J_1 = J$ .*

*If  $X$  is observed at discrete time points  $1, 2, \dots, T$ , that is,  $\mathbf{d}_{j,k} = d_{j,k}^D$ , then the rate of convergence in probability of  $\hat{\alpha}$  is  $T^{-\gamma/(2\gamma+\alpha)}$  with  $\gamma = \beta \wedge (2 - \alpha)$ , obtained for  $J_0 = \lfloor J/\{2\gamma + \alpha\} \rfloor$  and  $J_1 = J$ .*

*Remark 4.2.* Observe that the choice of  $J_0$  corresponding to the best rate for  $\hat{\alpha}$  depends both on the unknown smoothness parameter  $\beta$  and on the parameter  $\alpha$  itself. The case of discrete observations is similar to the continuous time observations but with the smoothness parameter  $\beta$  replaced by  $\gamma = \beta \wedge (2 - \alpha)$ , resulting in a slower rate of convergence. This can be explained by the aliasing induced by the interpolation step (3.9). It is clear that these rates of convergence are the best possible for our estimator under the assumption on  $\mathcal{L}$ , since this choice of  $J_0$  makes the squared bias and the variance of the same order of magnitude. However, to our knowledge, the best possible rate of convergence for the estimation of  $\alpha$  under these observations schemes is an open question. In other words, it is still to be found out whether our estimator is rate optimal.

The rate of convergence of our estimator is derived under assumptions on the function  $\mathcal{L}$ . The following lemma allows to check them through conditions on the joint distribution of  $(U, \eta)$ .

**Lemma 4.3.** *Let Assumption 1 hold.*

(i) *If there exists positive constants  $c$  and  $\beta$  such that, as  $t \rightarrow \infty$ ,*

$$L_2(t) = c + O(t^{-\beta}),$$

*then there exists a constant  $c'$  such that, as  $z \rightarrow \infty$ ,*

$$(4.16) \quad \mathcal{L}(z) = c' + \begin{cases} O(z^{-\beta}) & \text{if } \beta < 2 - \alpha, \\ O(z^{\alpha-2} \log z) & \text{if } \beta = 2 - \alpha, \\ O(z^{\alpha-2}) & \text{if } \beta > 2 - \alpha. \end{cases}$$

(ii) *If there exists positive constants  $c$  and  $\beta$  such that, as  $t \rightarrow 0$ ,*

$$\mathbb{E}[U^2 \{1 - \cos(\eta t)\}] = c |t|^{-\alpha} \{1 + O(|t|^\beta)\},$$

*then there exists a constant  $c'$  such that, as  $z \rightarrow \infty$ ,*

$$(4.17) \quad \mathcal{L}(z) = c' + O(z^{-\beta}),$$

provided that  $\psi$  belongs to the Sobolev space  $W^{(\alpha+\beta)/2-1}$ , that is,

$$(4.18) \quad \int_{-\infty}^{\infty} (1 + |\xi|)^{(\alpha+\beta)-2} |\psi^*(\xi)|^2 d\xi < \infty ,$$

where  $\psi^*$  denotes the Fourier transform of  $\psi$ ,

$$(4.19) \quad \psi^*(\xi) = \int_0^M \psi(t) e^{-i\xi t} dt .$$

*Example 4.1.* Assume that  $\eta$  has a Pareto distribution, i.e.  $\mathbb{P}(\eta > t) = (1 \vee t)^{-\alpha}$ , and is independent of  $U$ . This corresponds to Lemma 4.3(i) with  $\beta = \infty$ , and we can easily compute an exact expression for the  $O(z^{\alpha-2})$  term:

$$\mathcal{L}(z) = c' + \frac{\alpha \mathbb{E}[U^2]}{2 - \alpha} z^{\alpha-2} + o(z^{\alpha-2}) .$$

The best possible rate of convergence of  $\hat{\alpha}$  is thus  $T^{-(2-\alpha)/(4-\alpha)}$ , irrelevant of the observation scheme.

*Example 4.2.* Let  $\alpha \in (1, 2)$  and suppose that  $\eta$  is the absolute value of a symmetric  $\alpha$ -stable random variable. Then Assumption 1 holds, say, if  $U$  is independent of  $\eta$  and has sufficiently many finite moments, and

$$\mathbb{E}[\cos(\eta t)] = \exp(-\sigma |t|^\alpha) = 1 - \sigma |t|^\alpha + O(|t|^{2\alpha}).$$

By Lemma 4.3, the best possible rate of convergence of  $\hat{\alpha}$  is thus  $T^{-\gamma/(2\gamma+\alpha)}$  with  $\gamma = \alpha$  for continuous time observations and  $\gamma = 2 - \alpha$  for discrete time observations.

*Proof of Theorem 4.2.* In the following, we give a decomposition of the error valid under the assumption

$$0 < \liminf_{T \rightarrow \infty} \frac{J_0}{J} \leq \limsup_{T \rightarrow \infty} \frac{J_0}{J} < 1.$$

Optimizing  $J_0$  in this decomposition will then give the result. We use the same notation as in the proof of Theorem 4.1 with  $J_1 = J$ . We first give a first rough rate of convergence for  $\hat{\alpha}$  by adapting the proof of Theorem 4.1. Under the present assumptions,  $\mathcal{L}_0(z) = c$ , which implies  $W'_0(\alpha) = 0$ , and  $r^*(z) = O(z^{-\beta})$  as  $z \rightarrow \infty$ . Then, (4.10), (4.12), (4.14) and (4.15) yield:

$$(4.20) \quad (\hat{\alpha} - \alpha)^2 = O_P(2^{-\xi J} + 2^{-\beta J_0}).$$

Since  $\hat{\alpha}$  is consistent and  $\alpha$  is an interior point of the parameter set, the first derivative of the contrast function vanishes at  $\hat{\alpha}$  with probability tending to one. Hence

$$0 = \frac{\sum_{(j,k) \in \Delta} j 2^{(\hat{\alpha}-2)j} \mathbf{d}_{j,k}^2}{\sum_{(j,k) \in \Delta} 2^{(\hat{\alpha}-2)j} \mathbf{d}_{j,k}^2} - \delta \log(2) .$$

By definition of  $\delta$ , this yields

$$\begin{aligned} 0 &= \sum_{(j,k) \in \Delta} (j - \delta) 2^{(\hat{\alpha}-2)j} \mathbf{d}_{j,k}^2 \\ &= \sum_{(j,k) \in \Delta} (j - \delta) 2^{(\alpha-2)j} \mathbf{d}_{j,k}^2 + \log(2)(\hat{\alpha} - \alpha) \sum_{(j,k) \in \Delta} j(j - \delta) 2^{(\hat{\alpha}-2)j} \mathbf{d}_{j,k}^2 \end{aligned}$$

for a random  $\tilde{\alpha}$  between  $\alpha$  and  $\hat{\alpha}$ . By definition of  $\Lambda_j$ , the last display implies

$$\hat{\alpha} - \alpha = - \frac{\sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) (1 + \Lambda_j)}{\log 2 \sum_j j(j - \delta) 2^{(\tilde{\alpha}-\alpha-1)j} \mathcal{L}(2^j) (1 + \Lambda_j)} .$$

Denote the sum in the denominator of the last expression by  $D$ , and write

$$\begin{aligned} D &= \sum_j j(j - \delta) 2^{-j} \mathcal{L}(2^j) + \sum_j j(j - \delta) \mathcal{L}(2^j) 2^{-j} (2^{(\tilde{\alpha}-\alpha)j} - 1) (1 + \Lambda_j) \\ &\quad + \sum_j j(j - \delta) 2^{-j} \mathcal{L}(2^j) \Lambda_j =: S + R_1 + R_2 . \end{aligned}$$

Using Lemma 5.13 and (4.9), one easily gets that  $S \sim 2^{1-J_0}$  as  $J \rightarrow \infty$ .

Using Lemma 5.10, and the fact that  $|\tilde{\alpha} - \alpha| \leq |\hat{\alpha} - \alpha| = o_P(J^{-2})$ , one gets similarly  $R_1 = o_P(2^{-J_0})$ . To bound  $R_2$ , we proceed as for bounding  $E(\alpha')$  in the proof of Theorem 4.1 (here with  $\alpha' = \alpha > 1$ ): we write  $\sum_j = \sum_{j=J_0+1}^{J_2} + \sum_{j=J_2+1}^J$  and apply Lemmas 5.10 and 5.11 to obtain  $R_2 = o_P(2^{-J_0})$ . Hence, we finally get

$$(4.21) \quad \hat{\alpha} - \alpha = \frac{2^{J_0}}{2 \log 2} \left\{ \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) + \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) \Lambda_j \right\} \{1 + o_p(1)\} .$$

In (4.21), the terms inside the curly brackets are interpreted as a deterministic bias term and a stochastic fluctuation term. The bias is bounded as follows:

$$(4.22) \quad 2^{J_0} \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) = 2^{J_0} \sum_j (j - \delta) 2^{-j} (\mathcal{L}(2^j) - c) = O(2^{-\beta J_0}) .$$

In the case of continuous time observations, *i.e.*  $\mathbf{d}_{j,k} = d_{j,k}^S$  or  $\mathbf{d}_{j,k} = d_{j,k}$ , we have

$$(4.23) \quad \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) \Lambda_j = O_P(2^{-J/2 + (\alpha/2 - 1)J_0}) .$$

Gathering this bound with (4.21) and (4.22), and setting  $J_0 = J/(2\beta + \alpha)$  yields the first claim of Theorem 4.2, *i.e.*  $\hat{\alpha} - \alpha = O_P(2^{-\beta/(2\beta + \alpha)})$ . We now prove (4.23). Define  $\beta_j = n_j^{-1} \sum_{k=0}^{n_j-1} \{v_j^{-1} (d_{j,k}^S)^2 - 1\}$ . Then  $\beta_j = \Lambda_j$  if  $\mathbf{d}_{j,k} = d_{j,k}^S$ . Since  $\alpha > 1$ , Lemmas 3.1 and 5.2 yield, for some positive constant  $C$ ,

$$(4.24) \quad \mathbb{E}[\beta_j^2] = \text{var}(\beta_j) \leq C \frac{L_4(2^j)}{\mathcal{L}^2(2^j)} 2^{\alpha j - J} .$$

Since  $\mathcal{L}$  is bounded away from zero and  $L_4$  is bounded by assumption, the ratio  $L_4/\mathcal{L}^2$  is also bounded. The Minkowski inequality then yields, for some constant  $C > 0$ ,

$$\mathbb{E} \left[ \left( \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) \beta_j \right)^2 \right]^{1/2} \leq C 2^{-J/2} \sum_j |j - \delta| 2^{(\alpha/2-1)j} = O(2^{-J/2+(\alpha/2-1)J_0}).$$

If  $\mathbf{d}_{j,k} = d_{j,k}$ , we use (5.40) in Lemma 5.10, and obtain  $\mathbb{E}[|\Lambda_j - \beta_j|] \leq C n_j^{-1/2}$  for some constant  $C > 0$ . Hence, in this case, since  $-1/2 < \alpha/2 - 1$ ,

$$\mathbb{E} \left[ \left| \sum_j (j - \delta) 2^{-j} \mathcal{L}(2^j) (\Lambda_j - \beta_j) \right| \right] \leq C 2^{-J/2} \sum_j |j - \delta| \mathcal{L}(2^j) 2^{-j/2} = o(2^{-J/2+(\alpha/2-1)J_0}).$$

The last two displays imply (4.23).

We now briefly adapt the previous proof to the case of discrete observations. Define  $v_j^D = \mathbb{E}[(d_{j,k}^{SD})^2]$ . Lemma 5.6 [iii] implies  $v_j^D = v_j + O(1)$ . Thus we have  $v_j^D = \mathcal{L}_D(2^j) 2^{(2-\alpha)j}$  and

$$\mathcal{L}_D(z) = \mathcal{L}(z) + O(z^{\alpha-2}) = c + O(z^{-\gamma}),$$

with  $\gamma = \beta \wedge (2 - \alpha)$ . Then, defining

$$\Lambda_j^D = n_j^{-1} \sum_{k=1}^{n_j} \{(v_j^D)^{-1} \mathbf{d}_{j,k}^2 - 1\},$$

we obtain that (4.21) still holds with  $\mathcal{L}_D$  and  $\Lambda^D$  replacing  $\mathcal{L}$  and  $\Lambda$ , respectively. Lemma 5.12 implies that  $\Lambda_j^D$  has the same order of magnitude as  $\Lambda_j$ , so that the stochastic fluctuation term has the same order of magnitude as in the previous case. The difference comes from the bias term, which is  $O(2^{-\gamma J_0})$ . Thus,  $\hat{\alpha} - \alpha = O_P(2^{-\gamma J_0} + 2^{-\gamma J_0})$ , and setting  $J_0 = J/(2\gamma + \alpha)$  yields the second claim of Theorem 4.2. □

## 5. CONCLUDING REMARKS

In this work, we have proved the validity of a wavelet method for the estimation of the long memory parameter of an infinite source Poisson traffic model, both in a stable or in an unstable state, that is, when it does or does not converge to a stationary process. We have shown that a suitable choice of the scales in the estimator (see Remark 4.1) yields a consistent estimator in both situations and we checked that the estimator is robust to discrete data sampling.

However the study of the rates raises some questions concerning the optimality of this estimator. To draw a comparison, suppose that one *directly* observes the durations  $\eta_1, \dots, \eta_n$  of the clients arrived at times  $t_1, \dots, t_n$  in  $[0, T]$ . Then one can use the Hill estimator for estimating the tail index  $\alpha$ . Since  $T$  and  $n$  are asymptotically proportional and  $\eta_1, \dots, \eta_n$  are iid, the rates of this estimator are those derived in Hall and Welsh (1984). In particular, if  $\eta$  has a Pareto distribution, then a parametric rate  $\sqrt{T}$  can be

obtained. On the other hand, in the same situation, our wavelet estimator defined on the observations  $\{X(t), t \in [0, T]\}$  has a dramatically deteriorating rate for  $\alpha$  close to 2. It remains to find out whether this discrepancy comes from the choice of the estimator or from the fact that the durations  $\eta_k$  are not directly observed.

Finally, let us draw a practical conclusion from our study. Some precaution should be made on the choice of the scales used in the estimation as shown by the conditions on  $J_0$  and  $J_1$ . In particular, if only discrete observations are available, the best possible rate of convergence is obtained for a much bigger value of  $J_0$  than if continuous observations are available. Too small a value of  $J_0$  will induce an important bias for finite samples. Practitioners should be aware of this restriction and be careful in the interpretation of the results. Those questions will be tackled numerically in a future work.

#### APPENDIX: TECHNICAL RESULTS

**Lemma 5.1.** *Let Assumption 1 hold. Let  $f$  be a bounded measurable compactly supported function such that  $\int f(s) ds = 0$ . Define*

$$\tilde{f}(t, v, w) = w \int_t^{t+v} f(s) ds .$$

Then  $\int |\tilde{f}(t, v, w)|^p dt \nu(dv, dw) < \infty$ ,

$\mathbb{E}[N_S(\tilde{f})] = 0$  and  $\int_0^\infty X(s) f(s) ds = N_S(\tilde{f} \mathbb{1}_{\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}})$ . If moreover  $\mathbb{E}[\eta] < \infty$ , then  $N_S(\tilde{f}) = \int X_S(s) f(s) ds$ .

*Proof.* Assume that  $f$  has compact support in  $[0, M]$ . Then  $\int_t^{t+v} f(s) ds = 0$  if  $t + v < 0$  or  $t > M$ . If moreover  $\int_0^M f(s) ds = 0$ , then the former integral also vanishes if  $t < 0$  and  $t + v > M$ . Then, denoting  $\|f\|_1 = \int |f(s)| ds$ , we have

$$\int_{-\infty}^{\infty} \left| \int_t^{t+v} f(s) ds \right|^p dt \leq \|f\|_1^p \int_{-\infty}^{\infty} \{\mathbb{1}_{[0, M]}(t) + \mathbb{1}_{[0, M]}(t + v)\}^p dt = 2^p M \|f\|_1^p .$$

Thus  $\int |\tilde{f}(t, v, w)|^p dt \nu(dv, dw) \leq 2^p M \|f\|_1^p \mathbb{E}[|U|^p]$ . The fact that  $\mathbb{E}[N_S(\tilde{f})] = 0$  is then a straightforward application of Fubini's Theorem. The other properties follow directly from the definition of  $N_S$ .  $\square$

For  $p = 1, \dots, p^*$ , define the signed measures

$$(5.1) \quad \nu_p(dv) := \int u^p \nu(dv, du) .$$

*Proof of Lemma 3.1.* Using the above notation, we have  $d_{j,k}^S = N_S(\tilde{\psi}_{j,k})$ , since (3.1) implies that  $\int \psi_{j,k}(s) ds = 0$  for all  $j, k$ , hence the wavelet coefficients have zero mean by

Lemma 5.1. To compute the variance, we apply the cumulant formula (2.5) and a change of variable.

$$\begin{aligned} \text{var}(d_{j,k}^S) &= \int_0^\infty \nu_2(dv) \int_{-\infty}^\infty \left\{ \int_t^{t+v} \psi_{j,k}(s) ds \right\}^2 dt \\ &= 2^{2j} \int_0^\infty \nu_2(dv) \int_{-\infty}^\infty \left\{ \int_t^{t+2^{-j}v} \psi(s) ds \right\}^2 dt = 2^{(2-\alpha)j} \mathcal{L}(2^j), \end{aligned}$$

by definition of  $\mathcal{L}$ , see (3.6). To conclude the proof, we show (3.7). For  $p = 2, \dots$ , define

$$(5.2) \quad K_p(y) := \int_{-\infty}^\infty \left\{ \int_t^{t+y} \psi(u) du \right\}^p dt = \int_{-\infty}^\infty (\Psi(t+y) - \Psi(t))^p dt,$$

where  $\Psi(u) := \int_0^u \psi(y) dy$ . By (3.1),  $\Psi$  has compact support in  $[0, M]$  and,

$$(5.3) \quad |K_p(y)| \leq \int_{-M}^M |\Psi(t+y) - \Psi(t)|^p dt \leq 2M \|\psi\|_\infty^p |y|^p, \quad y \in \mathbb{R}.$$

If  $|y| \geq M$ , either  $\Psi(t+y)$  or  $\Psi(t)$  vanishes for any  $t$ , hence

$$(5.4) \quad K_p(y) = (1 + (-1)^p) \|\Psi\|_p^p, \quad |y| \geq M.$$

From (5.2), the two first derivatives of  $K_p$  can be expressed as

$$(5.5) \quad K_p'(y) = p \int_0^M \psi(u) (\Psi(u) - \Psi(u-y))^{p-1} du,$$

$$(5.6) \quad K_p''(y) = p(p-1) \int_0^M \psi(u) \psi(u-y) (\Psi(u) - \Psi(u-y))^{p-2} du,$$

which are continuous by continuity of  $\Psi$ , and with support in  $[-M, M]$  by (5.4). Inserting  $K_2(v/z) = \int_0^{v/z} K_2'(t) dt$  in (3.6) and inverting the integrals yield

$$(5.7) \quad \mathcal{L}(z) = z^\alpha \int_0^M K_2'(t) H_2(z t) dt = L_2(z) \int_0^M \frac{L_2(z t)}{L_2(z)} K_2'(t) t^{-\alpha} dt \sim_{z \rightarrow \infty} C_{\mathcal{L}} L_2(z),$$

Indeed, for any  $\epsilon \in (0, M)$

$$L_2(z) \int_\epsilon^M \frac{L_2(z t)}{L_2(z)} K_2'(t) t^{-\alpha} dt \sim L_2(z) \int_\epsilon^M K_2'(t) t^{-\alpha} dt$$

as  $z \rightarrow \infty$ , by the Uniform Convergence Theorem for slowly varying functions. Since  $|K_2'(y)| \leq c|y|$ , we have for  $z \geq 1/\epsilon$

$$\begin{aligned} \int_0^\epsilon L_2(z t) K_2'(t) t^{-\alpha} dt &= z^{\alpha-1} \int_0^{z\epsilon} L_2(u) K_2'(u/z) u^{-\alpha} du \\ &\leq C z^{\alpha-2} \left\{ \int_0^1 L_2(u) u^{-\alpha+1} du + \int_1^{z\epsilon} L_2(u) u^{-\alpha+1} du \right\} \end{aligned}$$

Using  $u^{-\alpha}L_2(u) = H_2(u) \rightarrow \mathbb{E}|U|^2$  as  $u \rightarrow 0$ , Karamata's Theorem and Potter's bound (see Resnick, 1987), we obtain the bound

$$L_2(z)^{-1} \int_0^\epsilon L_2(zt)K_2'(t)t^{-\alpha} dt \leq C\{L_2(z)^{-1}z^{\alpha-2} + \epsilon^{2-\alpha}\frac{L_2(z\epsilon)}{L_2(z)}\} \leq C\epsilon^{(2-\alpha)/2}.$$

(5.7) follows by letting  $\epsilon$  go to zero and noting that since  $K_2'(0) = 0$ ,  $C_{\mathcal{L}} := \int_0^M K_2'(t)t^{-\alpha} dt = \alpha \int_0^M K_2(t)t^{-\alpha-1} dt$  is a positive constant.  $\square$

*Proof of Lemma 4.3.* From (5.7), we have

$$\mathcal{L}(z) = \int_0^M K_2'(t)t^{-\alpha} \{c + r(zt)\} dt = c' + \int_0^M K_2'(t)t^{-\alpha} r(zt) dt,$$

where  $L_2(t) = c + r(t)$ . Since  $r(t) = O(t^{-\beta})$  as  $t \rightarrow \infty$  and  $r(t) = H_2(t)t^\alpha - c$ , there exists  $C > 0$  such that  $|r(t)| \leq Ct^{-\beta}$ . By (5.5) and (5.6), we have  $K_2'(t) \sim \|K_2\|_2 t$  as  $t \rightarrow 0$ . If  $\beta < 2 - \alpha$ , Eq (4.16) follows immediately. For  $\beta \geq 2 - \alpha$ , let us write

$$\begin{aligned} \left| \int_0^M K_2'(t)t^{-\alpha} r(zt) dt \right| &= \left| \int_0^{z^{-1}} K_2'(t)t^{-\alpha} r(zt) dt + \int_{z^{-1}}^M K_2'(t)t^{-\alpha} r(zt) dt \right| \\ &\leq C_1 \int_0^{z^{-1}} t^{1-\alpha} dt + C_2 z^{-\beta} \int_{z^{-1}}^M t^{1-\alpha-\beta} dt, \end{aligned}$$

where we used that  $r(zt) = H_2(zt)(zt)^\alpha - c$  is bounded for  $zt \leq 1$  for the first integral, and  $|r(t)| \leq Ct^{-\beta}$  for the second one. This yields (4.16).

Let us now show (4.17). Recall that  $\psi^*$  denotes the Fourier transform of  $\psi$ , see (4.19). The Sobolev condition (4.18) on  $\psi$  and (3.1) imply that

$$\int |\xi|^{\alpha+\beta-2} |\psi^*(\xi)|^2 d\xi < \infty$$

so that the lemma is a direct consequence of the following formula; for all  $z > 0$ ,

$$(5.8) \quad \mathcal{L}(z) = \frac{z^\alpha}{\pi} \int_{-\infty}^{\infty} \xi^{-2} |\psi^*(\xi)|^2 \mathbb{E}[U^2 \{1 - \cos(\eta\xi/z)\}] d\xi.$$

We now prove (5.8). For  $p = 2$ , (5.5) gives, since  $\psi\Psi$  integrates to  $\Psi^2/2$  and  $\Psi(0) = \Psi(M) = 0$ ,

$$K_2'(y) = 2 \int_0^M \psi(u)(\Psi(u) - \Psi(u-y)) du = -2 \int_{-\infty}^{\infty} \psi(u) \Psi(u-y) du.$$

By integration by parts one gets that  $\psi^*(\xi) = \int_{-\infty}^{\infty} \psi(t)e^{-i\xi t} dt = i\xi \int_{-\infty}^{\infty} \Psi(t)e^{-i\xi t} dt$ . Then, by Fubini's theorem, we obtain

$$\int_{-\infty}^{\infty} K_2'(y) e^{-iy\xi} dy = -2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi(u)e^{-iu\xi} \Psi(u-y) e^{i(u-y)\xi} dudy = \frac{2}{i\xi} |\psi^*(\xi)|^2.$$

The function  $K'_2$  is bounded and compactly supported (see the proof of Lemma 3.1), hence square integrable. By Parseval's formula, we find, for all  $y \geq 0$ ,

$$K_2(y) = \int_{-\infty}^{\infty} \mathbb{1}_{[0,y]}(u) K'_2(u) du = \pi^{-1} \int_{\xi=-\infty}^{\infty} \xi^{-2} |\psi^*(\xi)|^2 (1 - e^{i\xi y}) d\xi .$$

From (5.7), we have

$$\mathcal{L}(z) = z^\alpha \int_0^\infty K'_2(t) H_2(z t) dt = z^\alpha \mathbb{E}[U^2 K_2(\eta/z)] ,$$

Since  $\xi^{-2} |\psi^*(\xi)|^2$  is integrable, we may replace  $K_2$  by the above formula and exchange expectation and integration signs, giving

$$\mathcal{L}(z) = \frac{z^\alpha}{\pi} \int_{\xi=-\infty}^{\infty} \xi^{-2} |\psi^*(\xi)|^2 \mathbb{E}[U^2 (1 - e^{i\xi t})] d\xi .$$

Observing that  $\xi^{-2} |\psi^*(\xi)|^2$  is an even function of  $\xi$  and that the imaginary part of the expectation is an odd function of  $\xi$ , we get (5.8).  $\square$

**Lemma 5.2.** *Let Assumption 1 hold with  $p^* \geq 4$ . Then, there exists a positive constant  $C > 0$  such that*

$$\text{var} \left( \sum_{k=0}^{n-1} (d_{j,k}^S)^2 \right) \leq Cn \{ L_2^2(2^j) 2^{(4-2\alpha)j} + L_4(2^j) 2^{(3-\alpha)j} \} .$$

*Note that the first term dominates for  $\alpha < 1$  and the second one dominates for  $\alpha > 1$ .*

*Proof.* Write

$$(5.9) \quad \text{var} \left( \sum_{k=0}^{n-1} (d_{j,k}^S)^2 \right) = 2 \sum_{k,k'=0}^{n-1} \text{cov}^2(d_{j,k}^S, d_{j,k'}^S) + \sum_{k,k'=0}^{n-1} \text{cum}(d_{j,k}^S, d_{j,k}^S, d_{j,k'}^S, d_{j,k'}^S) .$$

For any positive integer  $p$ ,  $k \in \mathbb{Z}$  and  $v \geq 0$ , define

$$K_p(v, k) := \int_{-\infty}^{\infty} \left\{ \int_t^{t+v} \psi(s-k) ds \right\}^p \left\{ \int_t^{t+v} \psi(s') ds' \right\}^p dt .$$

We have the following properties:  $K_p(v, -k) = K_p(v, k)$ , by Cauchy Schwarz Inequality, for all  $k \in \mathbb{Z}$  and  $v > 0$ ,  $|K_p(v, k)| \leq K_p(v, 0) = K_{2p}(v)$ , where we use the notation (5.2); if  $k > v + M$ , then the support  $[0, M]$  of  $\psi$  cannot intersect both  $[t-k, t-k+v]$  and  $[t, t+M]$  so that  $K_p(v, k) = 0$ ; if  $k > M$  and  $v > k + M$ , then the support of  $\psi$  cannot intersect both  $[t-k, t-k+v]$  and  $[t, t+M]$  without being included in one of them, so that, by (3.1),  $K_p(v, k) = 0$ . Gathering the above facts gives, for all  $k \in \mathbb{Z}$  and  $v > 0$ ,

$$(5.10) \quad |K_p(v, k)| \leq K_{2p}(v) \{ \mathbb{1}(|k| \leq M) + \mathbb{1}(v \in [|k| - M, |k| + M]) \} .$$

Recall that  $d_{j,k}^S = N_S(\tilde{\psi}_{j,k})$ . Thus Formula (2.5) implies

$$(5.11) \quad \text{cov}(d_{j,k}^S, d_{j,k'}^S) = 2^{2j} \int_0^\infty K_1(v, |k - k'|) \nu_2(2^j dv) ,$$

$$(5.12) \quad \text{cum}(d_{j,k}^S, d_{j,k}^S, d_{j,k'}^S, d_{j,k'}^S) = 2^{3j} \int_0^\infty K_2(v, |k - k'|) \nu_4(2^j dv) .$$

Thus, separating the cases  $|k - k'| \leq M$  and  $|k - k'| > M$  in the sums of the RHS in (5.9) and using (5.10), we get

$$(5.13) \quad n^{-1} \text{var} \left( \sum_{k=0}^{n-1} (d_{j,k}^S)^2 \right) \leq 4(2M+1) 2^{4j} \left( \int_0^\infty K_2(v) \nu_2(2^j dv) \right)^2$$

$$(5.14) \quad + 4 2^{4j} \sum_{l=M+1}^{n-1} \left( \int_{l-M}^{l+M} K_2(v) \nu_2(2^j dv) \right)^2$$

$$(5.15) \quad + 2(2M+1) 2^{3j} \int_0^\infty K_4(v) \nu_4(2^j dv)$$

$$(5.16) \quad + 2 2^{3j} \sum_{l=M+1}^{n-1} \int_{l-M}^{l+M} K_4(v) \nu_4(2^j dv)$$

The RHS in (5.13) is  $4(2M+1) (\text{var}(d_{j,k}^S))$ , cf. the proof of Lemma 3.1. In this proof, the properties of  $K_4$  are similar to those of  $K_2$  (see (5.3) and (5.5)) and this yields similarly that (5.15) is asymptotically equivalent to  $2(2M+1)L_4(2^j)2^{(3-\alpha)j}$ . By elementary computations, the term (5.16) is less than

$$2 2^{3j} (2M+1) \int_1^{n+M-1} K_4(v) \nu_4(2^j dv)$$

which is less than (5.15). The term (5.14) is treated similarly by noting that

$$\begin{aligned} \sum_{l=M+1}^{n-1} \left( \int_{l-M}^{l+M} K_2(v) \nu_2(2^j dv) \right)^2 &\leq \int_0^\infty K_2(v) \nu_2(2^j dv) \sum_{l=M+1}^{n-1} \int_{l-M}^{l+M} K_2(v) \nu_2(2^j dv) \\ &\leq (2M+1) \left( \int_0^\infty K_2(v) \nu_2(2^j dv) \right)^2 \end{aligned}$$

so that it is of the same order as (5.13). This concludes the proof.  $\square$

**Lemma 5.3.** *Let Assumption 1 hold with  $p^* \geq 2$ . Then the following assertions hold.*

(i) *If  $\alpha \in (1/2, 2)$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,*

$$(5.17) \quad \sum_{k=0}^{n-1} (\mathbb{E}[d_{j,k}])^2 \leq C 2^{(3-2\alpha)j} L_1^2(2^j) .$$

(ii) If  $\alpha \in (1, 2)$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.18) \quad \sum_{k=0}^{n-1} \text{var}(d_{j,k} - d_{j,k}^S) \leq C 2^{(2-\alpha)j} L_2(2^j) .$$

(iii) If  $\alpha \in (0, 1]$ , for all  $\epsilon > 0$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.19) \quad \sum_{k=0}^{n-1} \text{var}(d_{j,k} - d_{j,k}^S) \leq C 2^{(2-\alpha)j} L_2(2^j) n^{1-\alpha+\epsilon} .$$

(iv) If (4.4) hold then, there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.20) \quad \sum_{k=0}^{n-1} (\mathbb{E}[d_{j,k}])^2 \leq C 2^{(3-2\alpha)j} L_1^2(2^j) .$$

*Proof.* By Lemma 5.1 we have that  $d_{j,k} = N_S(\tilde{\psi}_{j,k} \mathbb{1}_{\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}})$ . Thus

$$(5.21) \quad \begin{aligned} \mathbb{E}[d_{j,k}] &= \int_0^\infty \nu_1(dv) \int_0^\infty du \int_u^{u+v} \psi_{j,k}(w) dw \\ &= - \int_0^\infty \nu_1(dv) \int_{-\infty}^0 du \int_u^{u+v} \psi_{j,k}(w) dw , \end{aligned}$$

since (3.1) implies that  $\int_{-\infty}^\infty du \int_u^{u+v} \psi_{j,k}(w) dw = 0$ . Define  $\Psi(t) = \int_0^t \psi(s) ds$ . Again, (3.1) implies that  $\Psi$  also has compact support in  $[0, M]$ . Thus

$$(5.22) \quad \begin{aligned} \int_{-\infty}^0 du \int_u^{u+v} \psi_{j,k}(w) dw &= 2^{3j/2} \int_{-\infty}^{-k} (\Psi(x + 2^{-j}v) - \Psi(x)) dx \\ &= 2^{3j/2} \int_0^{(2^{-j}v-k) \wedge M} \Psi(x) dx , \end{aligned}$$

Inserting (5.22) in (5.21) and exchanging the two integrals give that

$$\begin{aligned} |\mathbb{E}[d_{j,k}]| &\leq 2^{3j/2} \int_0^M |\Psi(x)| H_1(2^j(x+k)) dx \\ &= 2^{(3/2-\alpha)j} L_1(2^j) \int_0^M |\Psi(x)| (x+k)^{-\alpha} \frac{L_1(2^j(x+k))}{L_1(2^j)} dx . \end{aligned}$$

Along the lines that lead to (5.7), we get  $|\mathbb{E}[d_{j,0}]| \leq C 2^{(3/2-\alpha)j} L_1(2^j) \int_0^M |\Psi(x)| x^{-\alpha} dx$ . The integral in this RHS is finite for since  $\Psi(x) = O(x)$  at  $x = 0$ . Note now that for any  $\epsilon > 0$ , there exists some constant  $C_\epsilon$  such that  $\forall j, t \geq 1, L_1(tx)/L_1(t) \leq C_\epsilon x^\epsilon$ . Then, for  $k \geq 0$

$$(5.23) \quad |\mathbb{E}[d_{j,k}]| \leq C 2^{(3/2-\alpha)j} L_1(2^j) \int_0^M |\Psi(x)| (x+k)^{\epsilon-\alpha} dx \leq C 2^{(3/2-\alpha)j} L_1(2^j) (k \vee 1)^{\epsilon-\alpha} .$$

Hence (5.17) for  $\alpha > 1/2$  by choosing  $\epsilon$  small enough.

We now prove (5.18) and (5.19). By Lemma 5.1,  $d_{j,k}^S - d_{j,k} = N_S(\tilde{\psi}_{j,k} \mathbb{1}_{\mathbb{R}_- \times \mathbb{R}_+ \times \mathbb{R}})$ . Applying (2.5) yields

$$(5.24) \quad \text{var}(d_{j,k}^S - d_{j,k}) = \int_0^\infty \nu_2(dv) \int_{-\infty}^0 du \left( \int_u^{u+v} \psi_{j,k}(w) dw \right)^2 .$$

Proceeding as for (5.22), one obtains

$$(5.25) \quad \begin{aligned} \int_{-\infty}^0 du \left( \int_u^{u+v} \psi_{j,k}(w) dw \right)^2 &= 2^{2j} \int_{-\infty}^{-k} (\Psi(x + 2^{-j}v) - \Psi(x))^2 dx \\ &= 2^{2j} \int_0^{(2^{-j}v-k)_+ \wedge M} \Psi^2(y) dy . \end{aligned}$$

The case  $\alpha > 1$  is then similar as for establishing (5.17) from (5.22) and (5.21). The case  $\alpha \in (0, 1]$  is also similar except that the sum  $\sum_{k=0}^{n-1}$  is no longer bounded independently of  $n$ .

We now show (5.20). If the function  $\psi$  satisfies (4.4) then  $\int_0^M \Psi(s) ds = 0$ . In (5.22), it gives

$$(5.26) \quad \left| \int_{-\infty}^0 du \int_u^{u+v} \psi_{j,k}(w) dw \right| \leq M \|\Psi\|_\infty \mathbb{1}(v \in [k2^j, (k+M)2^j]) .$$

Hence, by inserting this in (5.21), for  $k \geq 1$

$$(5.27) \quad |\mathbb{E}[d_{j,k}]| \leq C 2^{(3/2-\alpha)j} L_1(2^j) \{ \tilde{H}_1(k, j) - \tilde{H}_1(k+M, j) \} ,$$

where

$$\tilde{H}_1(k, j) := \frac{2^{\alpha j} H_1(k2^j)}{L_1(2^j)} = k^{-\alpha} \frac{L_1(k2^j)}{L_1(2^j)} .$$

Since  $L_1(k2^j)/L_1(2^j)$  is less than any positive power of  $k$ , there exists  $K > 0$  such that, for all  $k \geq K$  and  $j \geq 0$ ,  $\tilde{H}_1(k, j) \leq 1$ . Note also that by definition of  $\tilde{H}_1(k, j)$ , it is nonincreasing with respect to  $k$ . Thus, for all  $n > K + M$ ,

$$\begin{aligned} \sum_{k=K}^{n-1} \{ \tilde{H}_1(k, j) - \tilde{H}_1(k+M, j) \}^2 &\leq \sum_{k=K}^{n-1} \{ \tilde{H}_1(k, j) - \tilde{H}_1(k+M, j) \} \\ &= \sum_{k=K}^{K+M-1} \tilde{H}_1(k, j) - \sum_{k=n}^{n+M-1} \tilde{H}_1(k, j) \leq M . \end{aligned}$$

Inserting this in (5.27) and using (5.23) for bounding  $\sum_{k=0}^{M-1} (\mathbb{E}[d_{j,k}])^2$  yield (5.20).  $\square$

**Lemma 5.4.** *Define, for all  $t \in \mathbb{R}$  and  $v > 0$ ,*

$$(5.28) \quad g_{t,v} = \mathbf{I}_\phi[\mathbb{1}_{[t,t+v]}] \quad \text{and} \quad h_{t,v} = \mathbb{1}_{[t,t+v]} - g_{t,v} .$$

*We have the following properties for all  $t \in \mathbb{R}$  and all  $v > 0$ .*

- (i)  $\int g_{s,v}(t) ds = v$ ,
- (ii)  $\|g_{t,v}\|_\infty \leq M \|\phi\|_\infty$  and the support of  $g_{t,v}$  is included in  $[\lceil t \rceil - M + 1, \lceil t + v \rceil]$ ,
- (iii) the support of  $1 - g_{t,v}$  is included in  $(-\infty, \lceil t \rceil] \cup [\lceil t + v \rceil - M + 1, \infty)$ ,
- (iv)  $\|h_{t,v}\|_\infty \leq 1 + M \|\phi\|_\infty$  and the support of  $h_{t,v}$  is included in  $[t - M + 1, \lceil t \rceil] \cup [t + v - M + 1, \lceil t + v \rceil]$ .

*Proof.* By definition of  $g_{s,v}$  in (5.28), we have, using that  $\phi$  has a compact support and then (3.8),

$$\int g_{s,v}(t) ds = \int \sum_{k \in \mathbb{Z}} \mathbb{1}_{(k-v, k]}(s) \phi(t - k) ds = v \sum_{k \in \mathbb{Z}} \phi(t_k) = v.$$

We have, for all non-integer  $s \in \mathbb{R}$ ,

$$|g_{t,v}(s)| \leq \|\phi\|_\infty \sum_{k \in \mathbb{Z}} \mathbb{1}_{(-M+1, 1)}(s) \leq M \|\phi\|_\infty.$$

Hence the bound  $\|h_{t,v}\|_\infty \leq 1 + \|g_{t,v}\|_\infty$ . Denote by  $\lceil s \rceil$  the smallest integer larger than or equal to  $s$ . We have

$$(5.29) \quad g_{t,v}(s) = \sum_{k=\lceil t \rceil}^{\lceil t+v \rceil-1} \phi(s - k), \quad s \in \mathbb{R},$$

so that  $g_{t,v}$  has its support included in the segment  $[\lceil t \rceil - M + 1, \lceil t + v \rceil]$ . It follows that the support of  $h_{t,v}$  is included in  $[t - M + 1, \lceil t + v \rceil]$ . Now, by (3.8), we have

$$\begin{aligned} 1 - g_{t,v} &= \mathbb{I}_\phi[1 - \mathbb{1}_{[t, t+v]}] = -\mathbb{I}_\phi[\mathbb{1}_{[t, t+v]^c}], \\ h_{t,v} &= \mathbb{1}_{[t, t+v]^c} - \mathbb{I}_\phi[\mathbb{1}_{[t, t+v]^c}], \end{aligned}$$

where  $A^c$  denotes the complementary set of  $A$  in  $\mathbb{R}$ . The support of  $\mathbb{I}_\phi[\mathbb{1}_{[t, t+v]^c}]$  is included in  $(-\infty, \lceil t \rceil] \cup [\lceil t + v \rceil - M + 1, \infty)$ ; thus  $h_{t,v}$ 's one is in  $(-\infty, \lceil t \rceil] \cup [t + v - M + 1, \infty)$ . Hence the result.  $\square$

**Lemma 5.5.** *Let Assumption 1 hold. Let  $f$  be a bounded measurable compactly supported function such that  $\int f(s) ds = 0$ . Define*

$$\hat{f}(t, v, w) = w \int_{-\infty}^{\infty} g_{t,v}(s) f(s) ds, \quad \check{f}(t, v, w) = w \int_{-\infty}^{\infty} h_{t,v}(s) f(s) ds.$$

*Then, for  $p = 1, \dots, p^*$ ,  $\int |\hat{f}(t, v, w)|^p dt \nu(dv, dw) < \infty$ ,  $\int |\check{f}(t, v, w)|^p dt \nu(dv, dw) < \infty$ ,  $\int \mathbb{I}_\phi[X](s) f(s) ds = N_S(\hat{f} \mathbb{1}_{\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}})$ , and  $\mathbb{E}[N_S(\hat{f})] = \mathbb{E}[N_S(\check{f})] = 0$ . If moreover  $\mathbb{E}[\eta] < \infty$ , then  $N_S(\hat{f}) = \int \mathbb{I}_\phi[X_S](s) f(s) ds$ .*

*Proof.* Assume that  $f$  has compact support in  $[0, M]$ . Applying Lemma 5.4(ii) and proceeding as in the proof of Lemma 5.1, we have, for some numerical constant  $C$  whose face

value may change upon each appearance,

$$\left| \int_{-\infty}^{\infty} g_{t,v}(s) f(s) ds \right| \leq C \|f\|_1 \{ \mathbb{1}_{[M-1, 2M-1]}(\lceil t \rceil) + \mathbb{1}_{[0, M]}(\lceil t+v \rceil) \},$$

$$\int |\hat{f}(t, v, w)|^p dt \nu(dv, dw) \leq C \|f\|_1^p \mathbb{E}[|U|^p].$$

Applying Lemma 5.4(iv), we obtain a similar bound for  $\int \check{f}(t, v, w) dt \nu(dv, dw)$ . Thus we can apply Fubini's Theorem and Lemma 5.4(i).

$$\mathbb{E}[N_S(\hat{f})] = \int_{\mathbb{R}_+ \times \mathbb{R}} \left( \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} g_{t,v}(s) f(s) ds \right\} dt \right) w \nu(dv, dw),$$

$$\int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} g_{t,v}(s) f(s) ds \right\} dt = \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} g_{t,v}(s) dt \right\} f(s) ds = v \int_{-\infty}^{\infty} f(s) ds = 0.$$

Finally,  $\check{f} = \tilde{f} - \hat{f}$ , thus  $\mathbb{E}[N_S(\check{f})] = \mathbb{E}[N_S(\tilde{f})] - \mathbb{E}[N_S(\hat{f})] = 0$ .  $\square$

Applying Lemma 5.5, we can extend the definition of  $d_{j,k}^{SD}$  in (3.10) to the case  $\mathbb{E}[\eta] = \infty$  by

$$(5.30) \quad d_{j,k}^{SD} = N_S(\hat{\psi}_{j,k}).$$

- Lemma 5.6.** (i) Let Assumption 1 hold with  $p^* \geq 1$  and  $\alpha \in (0, 2)$ . Then  $\mathbb{E}[d_{j,k}^{SD}] = 0$  for all  $j \geq 0$  and  $k \in \mathbb{Z}$ .
- (ii) Let Assumption 1 hold with  $p^* \geq 2$  and  $\alpha \in (0, 2)$ . Then  $\text{var}(d_{j,k}^S - d_{j,k}^{SD})$  is bounded uniformly for  $j \in \mathbb{N}$  and  $k \in \mathbb{Z}$ .
- (iii) Let Assumption 1 hold with  $p^* \geq 2$  and  $\alpha \in (1, 2)$ . Then  $|\text{var}(d_{j,k}^S) - \text{var}(d_{j,k}^{SD})|$  is bounded uniformly for  $j \in \mathbb{N}$  and  $k \in \mathbb{Z}$ .

*Proof.* The first claim is given by Lemma 5.5. By Lemma 5.5, we have  $d_{j,k}^S - d_{j,k}^{SD} = N_S(\check{\psi}_{j,k})$ . Hence, applying (2.5),

$$\text{var}(d_{j,k}^S - d_{j,k}^{SD}) = \int_0^\infty \nu_2(dv) \int_{-\infty}^{\infty} \left( \int_{-\infty}^{\infty} h_{t,v}(s) \psi_{j,k}(s) ds \right)^2 dt.$$

By Lemma 5.4 (iv) and since the support of  $\psi$  is in  $[0, M]$ , we have, for all  $t \in \mathbb{R}$  and  $v \geq 0$ ,

$$(5.31) \quad \left| \int_{-\infty}^{\infty} h_{t,v}(s) \psi_{j,k}(s) ds \right| \leq 2^{-j/2} (1 + M \|\phi\|_\infty) \|\psi\|_\infty (B(t+1) + B(t+v+1)),$$

where, for all  $u \in \mathbb{R}$ ,

$$(5.32) \quad B(u) = \text{Leb}([u-M, u] \cap 2^j[k, k+M]) \leq M \mathbb{1}_{\{2^j k \leq u \leq 2^j k + (2^j + 1)M\}}.$$

This implies that  $\int B(u)^2 du$  is bounded by  $2^j$  up to a multiplicative constant and it follows that  $\text{var}(d_{j,k}^S - d_{j,k}^{SD}) = O(1)$ . We now suppose that  $\alpha \in (1, 2)$ . Using that

$\text{var}(d_{j,k}^{SD}) = \text{var}(d_{j,k}^S) + \text{var}(d_{j,k}^{SD} - d_{j,k}^S) + 2\text{cov}(d_{j,k}^S, d_{j,k}^{SD} - d_{j,k}^S)$ , it only remains to show that  $\text{cov}(d_{j,k}^S, d_{j,k}^{SD} - d_{j,k}^S) = O(1)$ . Applying (2.5), we have

$$\begin{aligned} \text{cov}(d_{j,k}^S, d_{j,k}^S - d_{j,k}^{SD}) &= \text{cov}(N_S(\psi_{j,k}), N_S(\check{\psi}_{j,k})) \\ &= \int_k^\infty \nu_2(dv) \int_{-\infty}^\infty \left( \int_{-\infty}^\infty h_{t,v}(s) \psi_{j,k}(s) ds \right) \left( \int_t^{t+v} \psi_{j,k}(s) ds \right) dt . \end{aligned}$$

By (5.31), (5.32) and

$$\left| \int_t^{t+v} \psi_{j,k}(s) ds \right| \leq \|\psi\|_\infty 2^{-j/2} v ;$$

we get  $\text{cov}(d_{j,k}^S, d_{j,k}^{SD} - d_{j,k}^S) = O(1)$ .  $\square$

**Lemma 5.7.** *For all  $\alpha \in (1/2, 2)$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,*

$$(5.33) \quad \sum_{k=0}^{n-1} (\mathbb{E}[d_{j,k}^D])^2 \leq C 2^{(3-2\alpha)j} L_1^2(2^j) .$$

For all  $\alpha \in (1, 2)$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.34) \quad \sum_{k=0}^{n-1} \text{var}(d_{j,k}^D - d_{j,k}^{SD}) \leq C 2^{(2-\alpha)j} L_2(2^j) .$$

For all  $\alpha \in (0, 1]$  and all  $\epsilon > 0$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.35) \quad \sum_{k=0}^{n-1} \text{var}(d_{j,k}^D - d_{j,k}^{SD}) \leq C 2^{(2-\alpha)j} L_2(2^j) n^{1-\alpha+\epsilon} .$$

If moreover (3.1), (3.8), (4.4) and (4.5) hold, then, for all  $\alpha \in (0, 2)$ , there exists  $C > 0$  such that, for all  $j \geq 0$  and  $n \geq 1$ ,

$$(5.36) \quad \sum_{k=0}^{n-1} (\mathbb{E}[d_{j,k}^D])^2 \leq C 2^{(3-2\alpha)j} L_1^2(2^j) .$$

*Proof.* Using the previous notation, we can write  $d_{j,k}^D = N_S(\hat{\psi}_{j,k} \mathbb{1}_{\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}})$ . Thus  $d_{j,k}^{SD} - d_{j,k}^D = N_S(\hat{\psi}_{j,k} \mathbb{1}_{\mathbb{R}_- \times \mathbb{R}_+ \times \mathbb{R}})$  and

$$\begin{aligned} \mathbb{E}[d_{j,k}^D] &= - \int_0^\infty \nu_1(dv) \int_{-\infty}^0 \left\{ \int_{-\infty}^\infty g_{t,v}(s) \psi_{j,k}(s) ds \right\} dt , \\ \text{var}(d_{j,k}^{SD} - d_{j,k}^D) &= \int_0^\infty \nu_2(dv) \int_{-\infty}^0 \left( \int_{-\infty}^\infty g_{t,v}(s) \psi_{j,k}(s) ds \right)^2 dt . \end{aligned}$$

By (3.1)  $\int g_{t,v}(s) \psi_{j,k}(s) ds = \int \{1 - g_{t,v}(s)\} \psi_{j,k}(s) ds$ . Hence this integral vanishes when the support of  $\psi_{j,k}$  does not intersect the support of  $g_{t,v}$  or when it does not intersect the

support of  $1 - g_{t,v}$ . The support of  $\psi_{j,k}$  is included in  $2^j[k, k + M]$  and Lemma 5.4 gives the supports of  $g_{t,v}$  and  $1 - g_{t,v}$ . We thus obtain

$$2^j k > t + v \text{ or } 2^j(k + M) < t - M + 1 \Rightarrow \int g_{t,v}(s) \psi_{j,k}(s) ds = 0$$

$$2^j k > t \text{ and } 2^j(k + M) < t + v - M + 1 \Rightarrow \int g_{t,v}(s) \psi_{j,k}(s) ds = 0 .$$

From Lemma 5.4, we know that  $\|g_{t,v}\|_\infty \leq M\|\phi\|_\infty$ ; hence

$$\left| \int g_{t,v}(s) \psi_{j,k}(s) ds \right| \leq M \|\phi\|_\infty \|\psi_{j,k}\|_1 = M \|\phi\|_\infty \|\psi\|_1 2^{j/2} .$$

The three last displays give that, for all  $t \leq 0$ ,

$$\left| \int g_{t,v}(s) \psi_{j,k}(s) ds \right| \leq C 2^{j/2} \mathbb{1}(0 < t - 2^j k + v < 2^j M + M - 1)$$

and, finally, for all  $k, j \geq 0$ ,

$$(5.37) \quad \left| \int_{-\infty}^0 \left\{ \int g_{t,v}(s) \psi_{j,k}(s) ds \right\} dt \right| \leq C 2^{3j/2} \{(k - 2^{-j}v)_+ \wedge (2M - 1)\}$$

$$\int_{-\infty}^0 \left\{ \int g_{t,v}(s) \psi_{j,k}(s) ds \right\}^2 dt \leq C 2^{2j} \{(k - 2^{-j}v)_+ \wedge (2M - 1)\} ,$$

which are similar bounds as those given by (5.22) and (5.25). The proofs of (5.33), (5.34) and (5.35) then follow as in the proof of Lemma 5.3. It only remains to show (5.36). Again the proof is similar to the proof of (5.20) in Lemma 5.3. One only needs to provide a bound similar as (5.26), namely,

$$\left| \int_{-\infty}^0 \left\{ \int g_{t,v}(s) \psi_{j,k}(s) ds \right\} dt \right| \leq C 2^{3j/2} \mathbb{1}(v \in [2^j k, 2^j(k + M')]) ,$$

for some constant  $M'$ . As we already have (5.37) at hand, it is now sufficient to show that, for all  $v \geq 2^j(k + M) + M - 1$ ,

$$(5.38) \quad \int_{-\infty}^0 \left\{ \int g_{t,v}(s) \psi_{j,k}(s) ds \right\} dt = - \int_0^\infty \left\{ \int g_{t,v}(s) \psi_{j,k}(s) ds \right\} dt = 0 .$$

Since by Lemma 5.4  $g_{t+v,\infty}$  has support in  $[t + v - M + 1, \infty)$ , we have, for all  $v \geq 2^j(k + M) + M - 1$  and  $t > 0$ ,

$$\int g_{t+v,\infty}(s) \psi_{j,k}(s) ds = 0 .$$

By (5.29), we have  $g_{t,v} + g_{t+v,\infty} = g_{t,\infty}$ , hence we only need to show that (5.38) holds for  $v = \infty$ . Observe that, by (5.29) and (4.5),

$$\int_0^\infty g_{t,\infty}(s) dt = \int_0^\infty \sum_{k \in \mathbb{Z}} \mathbb{1}(k \geq t) \phi(t - k) dt = \sum_{k \in \mathbb{Z}} k \phi(t - k)$$

is linear in  $t$ . By Fubini's theorem and using (3.1) and (4.4), we obtain (5.38) with  $v = \infty$ , which concludes the proof.  $\square$

**Lemma 5.8.** *Let Assumption 1 hold with  $p^* \geq 2$  and  $\alpha > 1$ . Then, there exists  $C > 0$  such that, for all  $n \geq 1$  and  $j \geq 0$ ,*

$$\begin{aligned} \sum_{k=0}^{n-1} \mathbb{E}[(d_{j,k} - d_{j,k}^S)^2] &\leq C v_j ; \\ \sum_{k=0}^{n-1} \mathbb{E}[(\mathbf{d}_{j,k} - d_{j,k}^S)^2] &\leq C(v_j + n) . \end{aligned}$$

**Lemma 5.9.** *Let Assumption 1 hold with  $p^* \geq 2$  and  $\alpha \in (0, 1]$ . Assume that  $\phi$  and  $\psi$  satisfy (4.4) and (4.5). Then, for all  $\epsilon > 0$ , then there exists  $C > 0$  such that, for all  $n \geq 1$  and  $j \geq 0$ ,*

$$\sum_{k=0}^{n-1} \mathbb{E}[(\mathbf{d}_{j,k} - d_{j,k}^S)^2] \leq C(L_1^2(2^j) 2^{(3-2\alpha)j} + v_j n^{1-\alpha+\epsilon} + n) ,$$

*Proof of Lemmas 5.8 and 5.9.* Write  $\mathbb{E}[(\mathbf{d}_{j,k} - d_{j,k}^S)^2] = (\mathbb{E}[\mathbf{d}_{j,k}])^2 + \text{var}(\mathbf{d}_{j,k} - d_{j,k}^S)$  and apply Lemmas 3.1, 5.3, 5.6 and 5.7.  $\square$

**Lemma 5.10.** *Let Assumption 1 hold with  $\alpha \in (1, 2)$  and  $p^* \geq 2$ . Then*

$$(5.39) \quad \sup_{0 \leq j \leq J} \mathbb{E}|\Lambda_j| = O(1) ;$$

$$(5.40) \quad \sup_{n \geq 1, j \geq 0} n^{-1/2} \mathbb{E} \left[ \left| v_j^{-1} \sum_{k=0}^{n-1} \{(d_{j,k}^S)^2 - d_{j,k}^2\} \right| \right] < \infty .$$

**Lemma 5.11.** *Let Assumption 1 hold with  $\alpha \in (0, 2)$  and  $p^* \geq 4$ . If  $\alpha \leq 1/2$ , assume moreover (4.4) and (4.5).*

*Let  $J^*$  be a sequence depending on  $J$  such that  $\limsup J^*/J < (1/\alpha) \wedge (1/(2-\alpha))$ . Then, there exists  $\epsilon > 0$  such that*

$$(5.41) \quad \sup_{u \in \mathcal{S}} \left| \sum_{j=J_0+1}^{J^*} u_j \Lambda_j \right| = O_P(2^{-\epsilon J})$$

where  $\mathcal{S}$  is the set of sequences  $u = (u_0, \dots)$  satisfying  $\sum_{j \in \mathbb{N}} |u_j| \leq 1$ .

*Proof of Lemmas 5.10 and 5.11.* Recall that  $d_{j,k}^S$  is always well defined, though unobservable in the case  $\mathbb{E}[\eta] = \infty$ . As previously, we denote  $\beta_j = n_j^{-1} \sum_{k=0}^{n_j-1} \{v_j^{-1}(d_{j,k}^S)^2 - 1\}$  and

$r_j = \Lambda_j - \beta_j$ . By Jensen's inequality, we have

$$\sup_{u \in \mathcal{S}} \left| \sum_{j=J_0+1}^{J^*} u_j \beta_j \right|^2 \leq \sum_{j=J_0+1}^{J^*} \beta_j^2,$$

hence, applying Lemma 5.2, we obtain, for some slowly varying function  $\tilde{\mathcal{L}}$ ,

$$\mathbb{E} \left[ \sup_{u \in \mathcal{S}} \left| \sum_{j=J_0+1}^{J^*} u_j \beta_j \right|^2 \right] \leq 2^{-J} \sum_{j=J_0+1}^{J^*} \tilde{\mathcal{L}}(2^j) 2^{(\alpha \vee 1)j} \leq C \tilde{\mathcal{L}}(2^{J^*}) 2^{(\alpha \vee 1)J^* - J} = o(1).$$

To bound the remainder terms, the cases  $\alpha > 1$  and  $\alpha \leq 1$  must be considered separately.

$$\begin{aligned} \mathbb{E}[|r_j|] &\leq (n_j v_j)^{-1} \sum_{k=0}^{n_j-1} \mathbb{E}[|(d_{j,k}^S)^2 - \mathbf{d}_{j,k}^2|] \\ &\leq (n_j v_j)^{-1} \sum_{k=0}^{n_j-1} \mathbb{E}[(d_{j,k}^S - \mathbf{d}_{j,k})^2] + 2(n_j v_j)^{-1/2} \left\{ \sum_{k=0}^{n_j-1} \mathbb{E}[(d_{j,k}^S - \mathbf{d}_{j,k})^2] \right\}^{1/2}. \end{aligned}$$

If  $\alpha > 1$ , we can apply Lemma 5.8 and obtain

$$(5.42) \quad \mathbb{E}[|r_j|] = O(1/\sqrt{n_j} + 1/\sqrt{v_j}).$$

Since  $\mathbb{E}[\beta_j] = 0$ , this proves (5.39). In Lemma 5.8, for  $\mathbf{d}_{j,k} = d_{j,k}$ , (5.42) can be made more precise, giving (5.40). Now, noting that  $\alpha > 1$  implies  $J^* < J/\alpha$ , we obtain

$$\mathbb{E} \left[ \sup_{u \in \mathcal{S}} \left| \sum_{j=J_0+1}^{J^*} u_j r_j \right| \right] \leq \sum_{j=J_0+1}^{J^*} \mathbb{E}[|r_j|] = O(2^{(\alpha/2-1)J_0}).$$

If  $\alpha \leq 1$ , we apply Lemma 5.9 and obtain, for some arbitrarily small  $\epsilon > 0$ ,

$$(5.43) \quad \begin{aligned} \mathbb{E}[|r_j|] &\leq C \left( 2^{(1-\alpha/2)j-J/2} + n_j^{(\epsilon-\alpha)/2} + 1/\sqrt{v_j} \right), \\ \mathbb{E} \left[ \sup_{u \in \mathcal{S}_J} \left| \sum_j u_j r_j \right| \right] &= O \left( 2^{\{(2-\alpha)J^*-J\}/2} + 2^{(J^*-J)(\epsilon-\alpha)/2} + 2^{(\alpha/2-1)J_0} \right). \end{aligned}$$

□

**Lemma 5.12.** *Let Assumption 1 hold with  $p^* \geq 4$  and  $\alpha \in (1, 2)$ . Then, there exists a positive constant  $C > 0$ ,*

$$(5.44) \quad \text{var} \left( \sum_{k=0}^{n-1} (d_{j,k}^{SD})^2 \right) \leq CL_4(2^j) n 2^{(3-\alpha)j}.$$

*Proof.* The proof is along the same line as the proof of Lemma 5.2 except that (5.11) and (5.12) are replaced by

$$(5.45) \quad \text{cov}(d_{j,k}^{SD}, d_{j,k'}^{SD}) = \int_0^\infty K_1(v, j, |k - k'|) \nu_2(dv),$$

$$(5.46) \quad \text{cum}(d_{j,k}^{SD}, d_{j,k}^{SD}, d_{j,k'}^{SD}, d_{j,k'}^{SD}) = \int_0^\infty K_2(v, j, |k - k'|) \nu_4(dv).$$

where

$$K_p^D(v, j, k) := 2^{-jp} \int_{-\infty}^\infty \left\{ \int_{-\infty}^\infty g_{t,v}(s) \psi(2^{-j}s - k) ds \right\}^p \left\{ \int_{-\infty}^\infty g_{t,v}(s') \psi(2^{-j}s') ds' \right\}^p dt$$

Denote

$$K_p^D(v, j) := 2^{-jp/2} \int_{-\infty}^\infty \left\{ \int_{-\infty}^\infty g_{t,v}(u) \psi(2^{-j}u) du \right\}^p dt$$

It first check that  $K_p^D(v, j, k)$  and  $K_p^D(v, j)$  satisfy similar properties as the ones needed for  $K_p(v, k)$  and  $K_p(v)$  in the proof of Lemma 5.2. Since, for any integer  $k$ ,  $g_{t,v}(s + k2^j) = g_{t-k2^j,v}(s)$ , we have  $K_p^D(v, j, k) = K_p^D(v, j, -k)$  and, by Cauchy Schwartz Inequality,  $|K_p^D(v, j, k)| \leq |K_{2p}^D(v, j)|$ . Recall that Lemma 5.4 implies that the support of  $g_{t,v}$  is included in  $[t - M + 1, t + v + 1]$ . We thus have that, if  $2^j k > v + M(1 + 2^j)$ , the support of  $\psi(2^{-j}\cdot)$  being included in  $[0, 2^j M]$  cannot intersect the one of  $g_{t-2^j k, v}$  and the one of  $g_{t,v}$  at the same time, implying  $K_p^D(v, j, k) = 0$ . If  $k > M$  and  $v > 2^j k + M(2^j - 1)$ , the support of  $\psi$  cannot intersect both the one of  $g_{t-2^j k, v}$  and the one of  $g_{t,v}$  without being included in one of them, so that, by (3.1),  $K_p^D(v, j, k) = 0$ . Hence we obtain the property corresponding to (5.10)

$$(5.47) \quad |K_p^D(v, j, k)| \leq K_{2p}^D(v, j) \{ \mathbb{1}(|k| \leq M) + \mathbb{1}(v \in [2^j |k| - M(1 + 2^j), 2^j |k| + M(2^j - 1)]) \}.$$

Proceeding as in the proof of Lemma 5.2, this bound yields

$$(5.48) \quad n^{-1} \text{var} \left( \sum_{k=0}^{n-1} (d_{j,k}^{SD})^2 \right) \leq C \left\{ \left( \int_0^\infty K_2^D(v, j) \nu_2(dv) \right)^2 + \int_0^\infty K_4^D(v, j) \nu_4(dv) \right\}.$$

From (5.28), we have

$$\int_{-\infty}^\infty g_{t,v}(u) \psi(2^{-j}u) du = \int_t^{t+v} \psi(2^{-j}u) du - R(t, v, j),$$

where

$$R(t, v, j) := \int_{-\infty}^\infty h_{t,v}(u) \psi(2^{-j}u) du = 2^{j/2} \int_{-\infty}^\infty h_{t,v}(u) \psi_{j,0}(u) du.$$

Hence, since  $\int_t^{t+v} \psi(2^{-j}u) du \leq M (v \wedge 2^j) \|\psi\|_\infty$ , using that, for any real numbers  $a$  and  $b$ ,  $|(a+b)^p - a^p| \leq p|b|(|b|^{p-1} \vee |a|^{p-1})$ , we find

$$K_p^D(v, j) = 2^{j(1+p/2)} K_p(2^{-j}v) + O\left(2^{-jp/2} \int_{t=-\infty}^{\infty} |R(t, v, j)| (|R(t, v, j)|^{p-1} \vee v^{(p-1)}) dt\right).$$

Using (5.31) and (5.32) gives that  $R(t, v, j) = O(1)$  and that  $\int R(t, v, j) dt = O(2^j)$ ; hence the above  $O$ -term is bounded by  $2^{j(1-p/2)}(v \wedge 2^j)^{p-1}$  up to a multiplicative constant independent of  $v$ . This in turn gives

$$\int_0^\infty K_p^D(v, j) \nu_p(dv) = 2^{j(1+p/2)} \int_0^\infty K_p(2^{-j}v) \nu_p(dv) + O\left(2^{j(1-p/2)} \int_0^\infty (v \wedge 2^j)^{p-1} \nu_p(dv)\right).$$

We know that (5.3) and (5.5) yield  $\int_0^\infty K_p(v) \nu_2(2^j dv) \sim L_p(2^j) 2^{-\alpha j}$ ; hence, since  $|K_p^D(v) - K_p(v)|$  is bounded independently of  $v$ ,

$$\int_0^\infty K_p^D(v) \nu_2(2^j dv) \sim L_p(2^j) 2^{-\alpha j}$$

Applying the two last displays to the cases  $p = 2$  and  $p = 4$  with  $\alpha > 1$  successively yield

$$\begin{aligned} \int_0^\infty K_2^D(v, j) \nu_2(dv) &\leq C (L_2(2^j) 2^{(2-\alpha)j} + 1), \\ \int_0^\infty K_4^D(v, j) \nu_4(dv) &\leq C (L_4(2^j) 2^{(3-\alpha)j} + L_4(2^j) 2^{(2-\alpha)j}). \end{aligned}$$

This, with (5.48) concludes the result.  $\square$

**Lemma 5.13.** *Let  $\rho$  be a positive real and  $\rho' := (2^\rho - 1)^{-1}$ . Let  $\ell^*$  be a non increasing function on  $[1, \infty)$  such that  $\lim_{s \rightarrow \infty} \ell^*(s) = 0$ , and let  $\ell$  be a function on  $[1, \infty)$  such that  $|\ell(s)| \leq \ell^*(s)$  for all  $s \in [1, \infty)$ . Define*

$$L(x) = c \exp\left\{\int_1^x \frac{\ell(s)}{s} ds\right\} \quad \text{and} \quad \omega_j = \frac{2^{-\rho j} L(2^j)}{\sum_{j'=J_0+1}^{J_1} 2^{-\rho j'} L(2^{j'})}.$$

Then, as  $J_0 \rightarrow \infty$  and for any  $\epsilon > 0$

$$(5.49) \quad \sum_{j=J_0+1}^{J_1} \omega_j j = J_0 + 1 + \rho'(1 + O(\ell^*(2^{J_0}))) + O(J_1(2 - \epsilon)^{J_0 - J_1})$$

$$(5.50) \quad \sum_{j=J_0+1}^{J_1} \omega_j j^2 = J_0^2 + 2J_0(1 + \rho') + 2\rho'^2 + 3\rho' + 1 + \rho' O(\ell^*(2^{J_0})) + O(J_1^2(2 - \epsilon)^{J_0 - J_1})$$

*Proof.* Denote  $S_k = \sum_{j=k+1}^{\infty} 2^{-\rho j} L(2^j)$ . Then,

$$(5.51) \quad S_k^{-1} \sum_{j=k+1}^{\infty} j L(2^j) 2^{-\rho j} = k + S_k^{-1} \sum_{j=k+1}^{\infty} (j-k) L(2^j) 2^{-\rho j} = k+1 + S_k^{-1} \sum_{j=k+1}^{\infty} S_j .$$

Note now that  $2^{-\rho j} = \rho'(2^{-\rho(j-1)} - 2^{-\rho j})$ . Hence

$$S_j = \rho' \sum_{j'=j+1}^{\infty} (2^{-\rho(j'-1)} - 2^{-\rho j'}) L(2^{j'}) = \rho' 2^{-\rho j} L(2^{j+1}) + R_j ,$$

with

$$R_j = \rho' \sum_{j'=j+1}^{\infty} 2^{-\rho j'} \{L(2^{j'+1}) - L(2^{j'})\} = \rho' \sum_{j'=j+1}^{\infty} 2^{-\rho j'} L(2^{j'}) \left\{ \exp \int_{2^{j'}}^{2^{j'+1}} \frac{\ell(s)}{s} ds - 1 \right\} .$$

Thus  $S_j = 2^{-\rho j} L(2^{j+1}) \{1 - R_j/S_j\}^{-1}$ . Since  $|\ell| \leq \ell^*$  and  $\ell^*$  is non increasing, we obtain:

$$\left| \frac{L(2^{j+1})}{L(2^j)} - 1 \right| = \left| \exp \int_{2^j}^{2^{j+1}} \frac{\ell(s)}{s} ds - 1 \right| \leq C \ell^*(2^j) .$$

Hence  $|R_j| \leq C \rho' S_j \ell^*(2^j)$  and for any  $j \geq k$ ,

$$(5.52) \quad S_j = \rho' 2^{-\rho j} L(2^{j+1}) \{1 + O(\ell^*(2^j))\} = \rho' 2^{-\rho j} L(2^j) \{1 + O(\ell^*(2^k))\} .$$

Using this expression in (5.51) yields

$$(5.53) \quad S_k^{-1} \sum_{j=k+1}^{\infty} j L(2^j) 2^{-\rho j} = k+1 + \rho'(1 + O(\ell^*(2^k))) .$$

The left-hand side of (5.49) writes

$$S_{J_0}^{-1} \sum_{j>J_0} j 2^{-\rho j} L(2^j) + \frac{U}{1-U} \{S_{J_0}^{-1} \sum_{j>J_0} j 2^{-\rho j} L(2^j) - S_{J_1}^{-1} \sum_{j>J_1} j 2^{-\rho j} L(2^j)\} .$$

with  $U := S_{J_1}/S_{J_0}$ . By (5.52),  $U = 2^{J_0-J_1} \frac{L(2^{J_1})}{L(2^{J_0})} \{1 + O(\ell^*(2^{J_0}))\}$ . Noting that  $|\frac{L(2^{J_1})}{L(2^{J_0})}| \leq \exp\{\ell^*(2^{J_0})(J_1 - J_0)\}$ , we have  $U \leq (2e^{\ell^*(2^{J_0})})^{J_0-J_1} \{1 + O(\ell^*(2^{J_0}))\}$ . Eq. (5.49) follows. Similarly to (5.53), we have

$$S_k^{-1} \sum_{j=k+1}^{\infty} j^2 L(2^j) 2^{-\rho j} = k^2 + 2k(1 + \rho') + 2(1 + \rho')^2 - (1 + \rho') + \rho' O(\ell^*(2^k)) .$$

and (5.50) is derived along the same lines. □

## REFERENCES

- C. Barakat, P. Thiran, G. Iannaccone, C. Diot, and P. Owezarski. A flow-based model for internet backbone traffic. In *Proceedings of the 2nd ACM SIGCOMM Workshop on Internet measurement*, pages 35 – 47. ACM Press, 2002.
- A. Cohen. *Numerical analysis of wavelet methods*, volume 32 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, 2003.
- N. G. Duffield, C. Lund, and M. Thorup. Properties and prediction of flow statistics from sampled packet streams. In *Proceedings of the 2nd ACM SIGCOMM Workshop on Internet measurement*, pages 159 – 171. ACM Press, 2002.
- G. Faÿ, F. Roueff, and P. Soulier. Estimation of the memory parameter of the infinite source poisson process. Full version. [arXiv:math.ST/0509371](https://arxiv.org/abs/math/0509371).
- P. Hall and A. H. Welsh. Best attainable rates of convergence for estimates of regular variation. *Ann. Statist.*, 3:1079–1084, 1984.
- H. R. Künsch. Statistical aspects of self-similar processes. In Yu.A. Prohorov and V.V. Sazonov (eds), *Proceedings of the first World Congress of the Bernoulli Society*, volume 1, pages 67–74. Utrecht, VNU Science Press, 1987.
- K. Maulik, S. Resnick, and H. Rootzén. Asymptotic independence and a network traffic model. *J. Appl. Probab.*, 39(4):671–699, 2002.
- Y. Meyer. *Wavelets and operators*, volume 37 of *Cambridge Studies in Advanced Mathematics*. Cambridge, Cambridge University Press, 1992.
- T. Mikosch, S. Resnick, H. Rootzén, and A. Stegeman. Is network traffic approximated by stable Levy motion or fractional Brownian motion? *Ann. Appl. Probab.*, 12:23–68, 2002.
- M. Parulekar and A. M. Makowski. M/G/Infinity input processes: A versatile class of models for network traffi. In *Proceedings of the INFOCOM '97*, 1997.
- S. Resnick. *Extreme values, regular variation and point processes*. Applied Probability, Vol. 4,. New York, Springer-Verlag, 1987.
- S. Resnick and H. Rootzén. Self-similar communication models and very heavy tails. *Ann. Appl. Probab.*, 10(3):753–778, 2000.
- G. A. Wornell and A. V. Oppenheim. Wavelet-based representations for a class of self-similar signals with application to fractal modulation. *IEEE Trans. Inform. Theory*, 38(2):785–800, 1992.

LABORATOIRE PAUL-PAINLEVÉ, UNIVERSITÉ LILLE 1, 59655 VILLENEUVE D’ASCQ CEDEX, FRANCE

*E-mail address:* gilles.fay@univ-lille1.fr

MODAL’X, UNIVERSITÉ PARIS X, 200 AVENUE DE LA RÉPUBLIQUE, 92000 NANTERRE, FRANCE

*E-mail address:* philippe.soulier@u-paris10.fr

ECOLE NATIONALE SUPÉRIEURE DES TÉLÉCOMMUNICATIONS, 46, RUE BARRAULT, 75634 PARIS  
CEDEX 13, FRANCE.

*E-mail address:* roueff@tsi.enst.fr