



HAL
open science

Continuum limit of the Volterra model, separation of variables and non standard realizations of the Virasoro Poisson bracket.

Olivier Babelon

► **To cite this version:**

Olivier Babelon. Continuum limit of the Volterra model, separation of variables and non standard realizations of the Virasoro Poisson bracket.. 2005. hal-00012185v1

HAL Id: hal-00012185

<https://hal.science/hal-00012185v1>

Preprint submitted on 17 Oct 2005 (v1), last revised 21 Mar 2006 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Continuum limit of the Volterra model, separation of variables and non standard realizations of the Virasoro Poisson bracket.

O. Babelon ¹

Laboratoire de Physique Théorique et Hautes Energies. ²

Abstract

The classical Volterra model, equipped with the Faddeev-Takhtadjan Poisson bracket provides a lattice version of the Virasoro algebra. The Volterra model being integrable, we can express the dynamical variables in terms of the so called separated variables. Taking the continuum limit of these formulae, we obtain the Virasoro generators written as determinants of infinite matrices, the elements of which are constructed with a set of points lying on an infinite genus Riemann surface. The coordinates of these points are separated variables for an infinite set of Poisson commuting quantities including L_0 . The scaling limit of the eigenvector can also be calculated explicitly, so that the associated Schroedinger equation is in fact exactly solvable.

¹Member of CNRS.

²L.P.T.H.E. Universités Paris VI–Paris VII (UMR 7589), Boîte 126, Tour 24-25, 5^{eme} étage, 4 place Jussieu, F-75252 PARIS CEDEX 05

1 Introduction.

The relation between integrable systems and conformal field theory has long been recognized [1, 2]. Although the emphasis has been put rightfully on Baxter Q operator and therefore on Sklyanin's separated variables [3], to the best of our knowledge there is no explicit expressions of the Virasoro generators in terms of these variables.

We make here a first step in this direction by considering the classical version of this problem. Our strategy will be to start with the Volterra model on the lattice [4, 6] equipped with the Faddeev-Takhtadjan [7, 8] Poisson bracket. Since the Volterra model is integrable, we can rewrite everything in terms of separated variables. Now, the Faddeev-Takhtadjan bracket goes directly to the Virasoro Poisson bracket in the continuum limit, and therefore by taking this limit in the separated variables formulae we will obtain the Virasoro generators expressed in terms of separated variables.

This leads to the following rather new type of formula for the Virasoro generators:

$$u(x) = \sum L_n e^{2in\pi x} = p_0^2 + 2\partial_x^2 \log \det \Theta(x) + (L_0 - p_0^2)\delta(x)$$

Here p_0 is the zero mode and Poisson commutes with everything, the term $(L_0 - p_0^2)\delta(x)$ will be explained later and the formula for L_0 is given in eq.(59). The infinite matrix $\Theta(x)$ reads ($k, m \in \{1, \dots, \infty\}$):

$$\Theta_{km}(x) = \frac{W_k(x)\partial_x E_m(x) - \partial_x W_k(x)E_m(x)}{Z_k^2 - m^2\pi^2}, \quad 0 \leq x \leq 1 \quad (1)$$

with

$$W_k(x) = \frac{\sin Z_k x}{Z_k} + \mu_k \frac{\sin Z_k(1-x)}{Z_k}, \quad E_m(x) = 2m\pi \sin m\pi x \quad (2)$$

The above formula for $u(x)$ is valid on the interval $0 \leq x \leq 1$, and should be extended outside this interval by periodicity (in particular the $\delta(x)$ term in a Dirac comb).

The result of this paper is that if the variables Z_k, μ_k have Poisson bracket ³

$$\{Z_k, Z_{k'}\} = 0, \quad \{Z_k, \mu_{k'}\} = 2(Z_k - p_0^2 Z_k^{-1})\mu_k \delta_{kk'}, \quad \{\mu_k, \mu_{k'}\} = 0 \quad (3)$$

then $u(x)$ does satisfies the Virasoro Poisson bracket.

$$\{u(x), u(y)\} = 4(u(x) + u(y)) \delta'(x-y) + 2\delta'''(x-y) \quad (4)$$

Moreover, the variables Z_k, μ_k are separated variables for an infinite set of higher commuting quantities, including L_0 .

Since the separated variables are also the ones which solve the classical inverse problem, the Schroedinger equation with the potential $u(x)$

$$(-\partial_x^2 - u(x))\psi(x, \Lambda) = \Lambda^2 \psi(x, \Lambda)$$

³Notice that if we redefine $\Lambda_k = \sqrt{Z_k^2 - p_0^2}$, the Poisson bracket becomes a standard quadratic bracket $\{\Lambda_k, \mu_k\} = 2\Lambda_k \mu_k$. However p_0 will then enter the formula for $\Theta(x)$ and, in this work, we prefer to keep that formula simple at the expense of a slightly more complicated Poisson bracket.

is exactly solvable, that is we have explicit formulae for both the potential $u(x)$ and a basis of solutions $\psi(x, \Lambda)$. Constructing the linear combination which is quasi periodic (the so called Bloch waves) introduces an infinite genus Riemann surface. The coefficients in the expression of this curve define a complete set of Poisson commuting Hamiltonians including L_0 . The separated variables are points on this curve.

The paper is organised as follows. In the first three sections we recall some known facts about the Volterra model on the lattice. In particular we recall the formulae expressing the dynamical degrees of freedom in terms of the separated variables.

In section 5 we compute the continuum (scaling) limit of the spectral curve. The result is eq.(41). We then show that the Hamiltonians H_m in this formula are in involution. Moreover we show that the scaling limit of the dynamical divisor still belongs to that curve, and hence define separated variables for these Hamiltonians.

In sections 6, 7 and 8, we compute the scaling limit of the eigenvector of the Lax matrix at each point of the spectral curve. The result is rather simple and is given in eq.(47). We then show that the obtained expression does satisfy a second order Schroedinger equation and we compute its potential $u(x)$. Finally, we construct the two quasi periodic solutions of that equation, the Bloch waves, and recover in this way exactly the same spectral curve as the one obtained previously.

In sections 9 and 10 we make a pause devoted to performing a few checks in a certain perturbative scheme, and to preparing the ground for the serious calculations coming next.

In sections 11 and 12 we prove that the potential $u(x)$ does satisfy the Virasoro Poisson bracket. An essential use is made of Hirota–Sato bilinear identities indicating once more, if needed, that all this construction is deeply rooted to integrability.

2 The Volterra model.

In this and the following two sections we recall some well known facts about the Volterra model. The Volterra model, as an integrable system, was introduced in [4]. It is a restricted version of the Toda lattice. We consider a periodic lattice with $N + 1$ sites. On each lattice site, we attach a dynamical variable a_i on which we impose the Faddeev-Takhtadjan Poisson bracket [7]:

$$\{a_i, a_j\} = a_i a_j \left[(4 - a_i - a_j)(\delta_{i,j+1} - \delta_{j,i+1}) - a_{j+1} \delta_{i,j+2} + a_{i+1} \delta_{j,i+2} \right] \quad (5)$$

This bracket⁴ is interesting because taking the continuum limit as

$$a_i \simeq 1 + \Delta^2 u(x), \quad \Delta = \frac{1}{(N+1)}$$

it becomes the Virasoro Poisson bracket eq.(4). For precisely this reason, and in this perspective, the lattice model has been extensively studied both at the classical level [7, 8], and at the quantum level [9, 10, 11, 13, 12]. The present paper is one more contribution this series of works.

⁴In terms of Toda Hamiltonian structures, it is a linear combination of restrictions of the second and fourth Poisson brackets.

The Lax matrix for the Volterra model is defined by:

$$L(\mu) = \begin{pmatrix} 0 & \sqrt{a_1} & 0 & \cdots & \mu^{-1}\sqrt{a_{N+1}} \\ \sqrt{a_1} & 0 & \sqrt{a_2} & \cdots & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & \sqrt{a_{i-1}} & 0 & \sqrt{a_i} & 0 \\ \vdots & & & \ddots & \vdots \\ 0 & \cdots & \sqrt{a_{N-1}} & 0 & \sqrt{a_N} \\ \mu\sqrt{a_{N+1}} & \cdots & 0 & \sqrt{a_N} & 0 \end{pmatrix} \quad (6)$$

It is well known that $\text{Tr}L^n(\mu)$ are in involution with respect to the Poisson bracket eq.(5). Hence we have an integrable system on the lattice whose continuum limit is directly related to conformal field theory.

The spectral curve Γ is defined as usual:

$$\Gamma \quad : \quad \det(L(\mu) - \lambda) = 0 \quad (7)$$

Expanding the determinant we see that it is of the form:

$$\mu + \mu^{-1} - t(\lambda) = 0 \quad (8)$$

where $t(\lambda)$ is polynomial of degree $N + 1$.

$$t(\lambda) = \mathcal{A}^{-1}\lambda^{N+1} - \mathcal{A}^{-1}\left(\sum_i a_i\right)\lambda^{N-1} + \cdots \quad (9)$$

where

$$\mathcal{A} = \sqrt{a_1 a_2 \cdots a_{N+1}}$$

Assuming $N = 2n$ even, $t(\lambda)$ is an *odd* polynomial $t(-\lambda) = -t(\lambda)$ and has exactly $n + 1$ independent coefficients. However, in that case, there is one Casimir function $K = t(2)$:

$$\{t(2), a_i\} = 0, \quad \forall i$$

Hence, the dimension of phase space is $N = 2n$ and we have n commuting quantities. The genus of the curve Γ is $g = N$.

At each point (λ, μ) of the spectral curve, we can attach an eigenvector $\Psi(\lambda, \mu) = (\psi_i(x))$, $i = 1, \dots, N + 1$, corresponding to the eigenvalue λ of $L(\mu)$. Explicitly, the equation $(L(\mu) - \lambda)\Psi = 0$ reads

$$\begin{aligned} \sqrt{a_1}\psi_2 + \mu^{-1}\sqrt{a_{N+1}}\psi_{N+1} &= \lambda\psi_1 \\ \sqrt{a_{i-1}}\psi_{i-1} + \sqrt{a_i}\psi_{i+1} &= \lambda\psi_i \\ \mu\sqrt{a_{N+1}}\psi_1 + \sqrt{a_N}\psi_N &= \lambda\psi_{N+1} \end{aligned} \quad (10)$$

We extend the definition of the coefficients a_i by periodicity $a_{i+N+1} = a_i$, and introduce a second order difference operator \mathcal{D} :

$$\left(\mathcal{D}\Psi\right)_i \equiv \sqrt{a_{i-1}}\psi_{i-1} + \sqrt{a_i}\psi_{i+1}$$

This operator is a discrete version of a Schroedinger operator with periodic potential. Eqs.(10) are then equivalent to:

$$\left(\mathcal{D}\Psi\right)_i = \lambda\psi_i, \quad \text{with} \quad \psi_{i+N+1} = \mu\psi_i \quad (11)$$

Thus, Ψ is a Bloch wave for the difference operator \mathcal{D} with a Bloch momentum μ .

In the continuum limit, eq.(11) becomes the Schroedinger equation.

$$(-\partial_x^2 - u(x))\psi(x) = \Lambda^2\psi(x), \quad \lambda \simeq 2 - \Delta^2\Lambda^2$$

3 The free case.

Since in the continuum limit $a_i \rightarrow 1$, it is useful to first recall some formulae in the trivial case $a_i = 1$. They will be generalized to the full case in the next section. To introduce the zero mode from the start, we consider the slightly more general case $a_i = a$:

$$\begin{aligned} \sqrt{a}(\psi_2 + \mu^{-1}\psi_{N+1}) &= \lambda\psi_1 \\ \sqrt{a}(\psi_{i-1} + \psi_{i+1}) &= \lambda\psi_i \\ \sqrt{a}(\mu\psi_1 + \psi_N) &= \lambda\psi_{N+1} \end{aligned}$$

The solution of the bulk equations is $\psi_i = \alpha x_+^i + \beta x_-^i$ where x_{\pm} are solutions of the equation

$$x^2 - zx + 1 = 0, \quad x_{\pm}(\lambda) = \frac{1}{2}(z \pm \sqrt{z^2 - 4}), \quad z = \frac{\lambda}{\sqrt{a}}$$

Imposing the two boundary equations, we get

$$\begin{aligned} (x_+^{N+1} - \mu)\alpha + (x_-^{N+1} - \mu)\beta &= 0 \\ (x_+^{N+1} - \mu)x_+\alpha + (x_-^{N+1} - \mu)x_-\beta &= 0 \end{aligned}$$

The compatibility of this system yields the spectral curve

$$\mu + \mu^{-1} = x_+^{N+1} + x_-^{N+1} \equiv t(\lambda) \quad (12)$$

We now impose that the curve passes through the point $(\lambda = 2, \mu = \mu_0)$ where μ_0 is related to the value of the Casimir function

$$K = \mu_0 + \mu_0^{-1} \quad (13)$$

Setting

$$x_{\pm}(2) = \frac{1}{\sqrt{a}} \pm \sqrt{\frac{1}{a} - 1} = e^{\pm\alpha}, \quad \mu_0 = e^{ip_0}$$

eq.(12) gives $\alpha = i\frac{p_0}{N+1}$. Hence the constant a is related to the value of the zero mode p_0 by:

$$\sqrt{a} = \frac{1}{\cos \frac{p_0}{N+1}}$$

The components of the eigenvector, properly normalized, are meromorphic functions on the spectral curve eq.(12). Choosing the normalisation $\psi_{N+1} = \mu$, they read

$$\psi_i(\lambda, \mu) = \frac{P_{N+1-i}(z) + \mu P_i(z)}{P_{N+1}(z)}, \quad z = \frac{\lambda}{\sqrt{a}} \quad (14)$$

We have introduced the polynomials of degree $j - 1$:

$$P_j(z) = \frac{x_+^j - x_-^j}{x_+ - x_-} \quad (15)$$

One has $P_j(z) = z^{j-1} + O(z^{j-3})$. The first few polynomials are

$$P_0 = 0, \quad P_1 = 1, \quad P_2 = z, \quad P_3 = z^2 - 1, \quad P_4 = z^3 - 2z$$

They are essentially the Tchebitchev polynomials of the second kind.

As we will see, eq.(14) is the general form of the meromorphic function $\psi_i(\lambda, \mu)$ even when $a_i \neq a$. In particular, in order to take the continuum limit, the poles of the eigenvector⁵ will have to be close to the roots of the equation $P_{N+1}(\frac{\lambda}{\sqrt{a}}) = 0$, that is

$$\lambda_k^{(0)} = 2\sqrt{a} \cos \frac{Z_k^{(0)}}{N+1}, \quad Z_k^{(0)} = k\pi \quad k = 1, \dots, N \quad (16)$$

For these special values $\lambda_k^{(0)}$, we have $x_{\pm} = e^{\pm i \frac{k\pi}{N+1}}$ and eq.(12) gives $\mu_k^{(0)} = (-1)^k$. The set of points $(\lambda_k^{(0)}, \mu_k^{(0)})$, $k = 1, \dots, N$ will be called the free configuration and will play an important role below.

It is simple to take the continuum limit in this free case. We set

$$\lambda \simeq 2 - \Delta^2 \Lambda^2, \quad z \simeq 2 - \Delta^2 Z^2, \quad \psi_{i\pm 1} = \psi(x \pm \Delta), \quad \Delta = \frac{1}{N+1}$$

where we have introduced the variable

$$Z = \sqrt{\Lambda^2 + p_0^2} \quad (17)$$

The eigenvector equation becomes the Schroedinger equation

$$-\psi''(x) - p_0^2 \psi(x) = \Lambda^2 \psi(x), \quad x = \Delta j \quad (18)$$

In this limit, we also have $x_{\pm} = 1 \pm i\Delta Z$ and the equation of the spectral curve reads:

$$\mu + \mu^{-1} = (1 + i\Delta Z)^{1/\Delta} + (1 - i\Delta Z)^{-1/\Delta}$$

In the limit $\Delta \rightarrow 0$, it becomes

$$\mu + \mu^{-1} = 2 \cos Z, \quad \mu_{\pm} = e^{\pm iZ} \quad (19)$$

⁵In fact in this simple case the eigenvector has no poles at finite distance because they are compensated by zeroes in the numerator. This degeneracy is lifted as soon as the a_i are not all equal.

Similarly, the eigenvector becomes a Baker-Akhiezer function

$$\psi(x) = \frac{\sin Z(1-x) + \mu \sin Zx}{\sin Z} \quad (20)$$

When $\mu = e^{\pm iZ}$ we get $\psi(x) = e^{\pm iZx}$ as it should be. Notice that when μ is kept as a free parameter, the above formula gives two independent solutions of eq.(18), but when μ belongs to the spectral curve eq.(19) one has

$$\psi(x+1) = \mu\psi(x)$$

Eq.(20) presents the two Bloch waves as a single function on the hyperelliptic spectral curve.

Another example important to us will be the Dirac comb.

$$[-\partial^2 - H_0\delta(x)]\psi(x) = \Lambda^2\psi(x), \quad \delta(x+1) = \delta(x)$$

On each interval $x_j = j < x < x_{j+1} = j+1$, one has

$$\psi(x) = \alpha_j e^{i\Lambda x} + \beta_j e^{-i\Lambda x}, \quad x_j < x < x_{j+1}$$

The Bloch condition

$$\psi(x+x_j) = \mu^j \psi(x), \quad 0 < x < 1$$

gives $\alpha_j = (\mu e^{-i\Lambda})^j \alpha_0$, $\beta_j = (\mu e^{i\Lambda})^j \beta_0$. The continuity of $\psi(x)$ gives $\alpha_0 = -\frac{(\mu - e^{-i\Lambda})}{(\mu - e^{i\Lambda})} \beta_0$, while the gap equation on the first derivative, $-\psi'(x_j+0) + \psi'(x_j-0) - H_0\psi(x_j) = 0$, gives the spectral curve

$$\mu + \mu^{-1} = 2 \cos \Lambda - H_0 \frac{\sin \Lambda}{\Lambda} \quad (21)$$

The Bloch wave itself is $\psi(x) = \mu^j \psi_{vac}(x-x_j)$, $x_j < x < x_{j+1}$, where $\psi_{vac}(x)$ is given by eq.(20) (with $p_0 = 0$) but Λ, μ now belonging to eq.(21).

4 Separated variables.

In this section, we generalise the previous analysis when $a_i \neq a$ and express the dynamical variables of the Volterra model in terms of the separated variables. Equivalent formulae were already obtained a long time ago in [4, 5]. A quantum version for the Toda chain can be found in [19].

We have to reconstruct the eigenvectors Ψ of $L(\mu)$. Let us set $\Psi = (\psi_i)$, $i = 1, \dots, N+1$. We normalize the last component $\psi_{N+1} = \mu$. Notice that due to eq.(8), μ does not vanish for finite λ . The components ψ_i are meromorphic functions on the spectral curve and are uniquely characterised by their poles and behavior at infinity which we now describe.

We will call $P^+(\lambda = \infty, \mu = \infty)$ and $P^-(\lambda = \infty, \mu = 0)$ the two points above $\lambda = \infty$. In the neighbourhood of P^\pm the local parameter is λ^{-1} and we have by direct expansion of eq.(8):

$$P^+ : \mu = \mathcal{A}^{-1} \lambda^{N+1} \left(1 + O(\lambda^{-2}) \right) \quad (22)$$

$$P^- : \mu = \mathcal{A} \lambda^{-N-1} \left(1 + O(\lambda^{-2}) \right) \quad (23)$$

At the points P^+ and P^- , the eigenvector $\Psi(P)$ behaves as:

$$\psi_i(P) = \frac{1}{\sqrt{a_{N+1}a_1a_2\cdots a_{i-1}}} \lambda^i \left(1 + O(\lambda^{-2})\right), \quad P \sim P^+ \quad (24)$$

$$\psi_i(P) = \sqrt{a_{N+1}a_1a_2\cdots a_{i-1}} \lambda^{-i} \left(1 + O(\lambda^{-2})\right), \quad P \sim P^- \quad (25)$$

This is easily deduced by inspection of eqs.(10).

From the general results of the classical inverse scattering theory, we expect $g + (N + 1) - 1 = 2N$ poles for the eigenvector (see e.g. [6, 15]). From eq.(24), we see that we have a fixed pole of order N at P^+ (on the component ψ_N), and there remains $g = N$ poles at finite distance, the so called *dynamical poles*. But we notice the symmetry property

$$\psi_i(-\lambda, -\mu) = (-1)^i \psi_i(\lambda, \mu)$$

so that the dynamical poles come in pairs

$$\lambda_{N+1-k} = -\lambda_k, \quad \mu_{N+1-k} = -\mu_k$$

and only $(\lambda_k, \mu_k), k = 1 \cdots n$ are independent parameters.

Everything can be expressed in terms of these $2n = N$ quantities $(\lambda_k, \mu_k), k = 1 \cdots n$. In fact, they can be viewed as coordinates on (an open set of) phase space.

First, the commuting Hamiltonians are easy to reconstruct. Indeed the spectral curve is determined by requiring that it passes through the points $(\lambda_k, \mu_k), k = 1 \cdots n$ and the point $(2, \mu_0)$, where μ_0 is related to the Casimir function as in eq.(13).

The equation of the curve itself can be written as a determinant

$$\det \begin{pmatrix} \lambda & \lambda^3 & \cdots & \lambda^{N+1} & \mu + \mu^{-1} \\ 2 & 2^3 & \cdots & 2^{N+1} & \mu_0 + \mu_0^{-1} \\ \lambda_1 & \lambda_1^3 & \cdots & \lambda_1^{N+1} & \mu_1 + \mu_1^{-1} \\ \vdots & & & & \vdots \\ \lambda_n & \lambda_n^3 & \cdots & \lambda_n^{N+1} & \mu_n + \mu_n^{-1} \end{pmatrix} = 0 \quad (26)$$

Expanding over the first row, we obtain a curve of the form eq.(8), and we can read directly the Hamiltonians as the coefficients of $t(\lambda)$. They appear as functions of the (λ_k, μ_k) and can be shown to Poisson commute (see [16, 17, 18] for a proof and for the quantum generalization of this fact).

Eqs.(24,25) and the data of the N dynamical poles also determine the functions ψ_i uniquely. Being meromorphic functions on a hyperelliptic curve, we can write quite generally

$$\psi_i = \frac{Q^{(i)}(\lambda) + \mu R^{(i)}(\lambda)}{\prod_{k=1}^n (\lambda^2 - \lambda_k^2)} \quad (27)$$

where $Q^{(i)}$ and $R^{(i)}$ are polynomials such that

$$Q^{(i)}(-\lambda) = (-1)^i Q^{(i)}(\lambda), \quad R^{(i)}(-\lambda) = (-1)^{i+1} R^{(i)}(\lambda)$$

Above λ_k , we have two points on the curve: (λ_k, μ_k) and (λ_k, μ_k^{-1}) . We want the poles to be at (λ_k, μ_k) only so that the numerator in eq.(27) should vanish at the points (λ_k, μ_k^{-1}) . This gives n conditions

$$Q^{(i)}(\lambda_k) + \mu_k^{-1} R^{(i)}(\lambda_k) = 0, \quad k = 1 \cdots n \quad (28)$$

To have a pole of order i at P^+ and a zero of order i at P^- we must choose

$$\text{degree } Q^{(i)} = N - i, \quad \text{degree } R^{(i)} = i - 1$$

Hence these two polynomials depend altogether on $n + 1$ coefficients which are determined by imposing the n conditions eq.(28) and requiring that the normalizations coefficients are inverse to each other at P^\pm as in eqs.(24,25).

It is convenient to use the basis of polynomials $P_j(\lambda)$ eq.(15). We write the formulae for ψ_i in the case i *odd*, the case i *even* is similar.

The polynomial $Q^{(i)}(\lambda)$ can be expanded over

$$Q^{(i)}(\lambda) : P_2(\lambda), P_4(\lambda), \cdots P_{N+1-i}$$

and the polynomial $R^{(i)}(\lambda)$ can be expanded over

$$R^{(i)}(\lambda) : P_1(\lambda), P_3(\lambda), \cdots P_i(\lambda)$$

Solving the linear system eq.(28), the eigenvector can be written as

$$\psi_i = \frac{K_i}{\prod(\lambda^2 - \lambda_k^2)} \det \begin{pmatrix} \mu P_1(\lambda) & \cdots & \mu P_i(\lambda) & -P_{N+1-i}(\lambda) & \cdots & -P_2(\lambda) \\ P_1(\lambda_1) & \cdots & P_i(\lambda_1) & -\mu_1 P_{N+1-i}(\lambda_1) & \cdots & -\mu_1 P_2(\lambda_1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_1(\lambda_k) & \cdots & P_i(\lambda_k) & -\mu_k P_{N+1-i}(\lambda_k) & \cdots & -\mu_k P_2(\lambda_k) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_1(\lambda_n) & \cdots & P_i(\lambda_n) & -\mu_n P_{N+1-i}(\lambda_n) & \cdots & -\mu_n P_2(\lambda_n) \end{pmatrix} \quad (29)$$

where K_i are constants independent of λ, μ . Defining

$$\Theta_i = \begin{pmatrix} P_1(\lambda_1) & \cdots & P_{i-2}(\lambda_1) & -\mu_1 P_{N+1-i}(\lambda_1) & \cdots & -\mu_1 P_2(\lambda_1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_1(\lambda_k) & \cdots & P_{i-2}(\lambda_k) & -\mu_k P_{N+1-i}(\lambda_k) & \cdots & -\mu_k P_2(\lambda_k) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_1(\lambda_n) & \cdots & P_{i-2}(\lambda_n) & -\mu_n P_{N+1-i}(\lambda_n) & \cdots & -\mu_n P_2(\lambda_n) \end{pmatrix} \quad (30)$$

We can compute the leading terms in eq.(29) when $\lambda \rightarrow \infty$. At $P^{(+)}$ the leading term comes from $\mu P_i(\lambda)$, while at $P^{(-)}$ it comes from $P_{N+1-i}(\lambda)$.

$$\begin{aligned} \psi_i &\simeq (-1)^{\frac{i-1}{2}} \mathcal{A}^{-1} K_i \det \Theta_i \lambda^i, & P^+ \\ \psi_i &\simeq (-1)^{\frac{i-1}{2}} K_i \det \Theta_{i+2} \lambda^{-i}, & P^- \end{aligned}$$

hence, imposing that the two coefficients of λ^i and λ^{-i} are inverse to each other, we get

$$K_i^2 = \frac{\mathcal{A}}{\det \Theta_i \det \Theta_{i+2}}$$

Comparing with eqs.(24, 25), we finally obtain

$$a_i = \frac{\det \Theta_i \det \Theta_{i+3}}{\det \Theta_{i+1} \det \Theta_{i+2}}, \quad a_N = \frac{\det \Theta_N}{\det \Theta_{N+2}} \mathcal{A}, \quad a_{N+1} = \frac{\det \Theta_3}{\det \Theta_1} \mathcal{A} \quad (31)$$

Here \mathcal{A}^{-1} is the coefficient of λ^{N+1} in $t(\lambda)$ eq.(9), computed from eq.(26).

We impose the Poisson bracket on the variables λ_k, μ_k

$$\{\lambda_k, \lambda_{k'}\} = 0, \quad \{\lambda_k, \mu_{k'}\} = -\frac{1}{2} \delta_{kk'} (4\lambda_k - \lambda_k^3) \mu_k, \quad \{\mu_k, \mu_{k'}\} = 0 \quad (32)$$

One can then check that the Hamiltonians defined by eq.(26) are all in involution (this is a general result), and that the a_i defined above do satisfy the Faddeev-Takhtadjan Poisson bracket. The fact that the expressions for a_N, a_{N+1} are different from the ones in the bulk is due to the choice of normalisation of the eigenvector. However, the Poisson bracket of the a_i is periodic. All this can be proved using techniques similar to the ones in [19].

5 Continuum limit of the spectral curve.

We now take the continuum limit of the spectral curve eq.(26). The result is eq.(41). We set

$$\lambda = \sqrt{a}z, \quad \lambda_k = \sqrt{a}z_k, \quad \frac{2}{\sqrt{a}} = 2 \cos \frac{p_0}{N+1} = z_0$$

From these, the scaled variables are defined like this:

$$Z = \sqrt{\Lambda^2 + p_0^2}, \quad z = 2 \cos \frac{Z}{N+1}, \quad z_k = 2 \cos \frac{Z_k}{N+1} \quad (33)$$

In the following, we will refer to the terminology "perturbation theory" when the points (Z_k, μ_k) are small deviations from the free configuration. The formulae we will write will make sense in this perturbative setting. This however does not exclude the possibility to have a finite number of points which are large deviation. We will also be interested in the deviation from the zero mode configuration. That is we make the substitution $\sqrt{a_i} \rightarrow \sqrt{a} \sqrt{\tilde{a}_i}$ everywhere on the lattice. Alternatively this amounts to using the variable $z = \lambda/\sqrt{a}$.

Using the basis of polynomials $P_j(z)$ defined in eq.(15) instead of the z^j , we can write the spectral curve as (it has the right form and passes through the right points)

$$\det \begin{pmatrix} \mu + \mu^{-1} & P_{N+2}(z) & \cdots & P_4(z) & P_2(z) \\ \mu_0 + \mu_0^{-1} & P_{N+2}(z_0) & \cdots & P_4(z_0) & P_2(z_0) \\ \mu_1 + \mu_1^{-1} & P_{N+2}(z_1) & \cdots & P_4(z_1) & P_2(z_1) \\ \vdots & & & & \vdots \\ \mu_n + \mu_n^{-1} & P_{N+2}(z_n) & \cdots & P_4(z_n) & P_2(z_n) \end{pmatrix} = 0$$

Without changing the determinant, we can subtract to the first column the linear combination of the next two columns:

$$P_{N+2}(z_k) - P_N(z_k) = 2 \cos Z_k$$

The first column becomes

$$\begin{pmatrix} \mu + \mu^{-1} - 2 \cos Z \\ \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix}$$

where we set

$$\gamma_k = \mu_k + \mu_k^{-1} - 2 \cos Z_k \quad (34)$$

Notice that $\gamma_0 = 0$. The reason for this subtraction is that for the free configuration we also have $\gamma_k^{(0)} = 0$, so that the spectral curve becomes simply $\mu + \mu^{-1} = 2 \cos Z$ as it should be. The subtraction gives sense to the spectral curve in perturbation theory. Expanding the determinant over the first row, we can write

$$\mu + \mu^{-1} - 2 \cos Z = \sum_{j=1}^{n+1} H_{2j} P_{2j}(z)$$

The H_{2j} are given by

$$H = N^{-1}V$$

where we defined

$$H = \begin{pmatrix} H_{N+2} \\ H_N \\ \vdots \\ H_2 \end{pmatrix}, \quad N = \begin{pmatrix} P_{N+2}(z_0) & \cdots & P_4(z_0) & P_2(z_0) \\ P_{N+2}(z_1) & \cdots & P_4(z_1) & P_2(z_1) \\ \vdots & & & \vdots \\ P_{N+2}(z_n) & \cdots & P_4(z_n) & P_2(z_n) \end{pmatrix}, \quad V = \begin{pmatrix} \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix}$$

We will need to treat separately the first row and column in the matrix N . Let us write it as

$$N = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

where

$$A = P_{N+2}(z_0), \quad B = (P_N(z_0) \quad \cdots \quad P_4(z_0) \quad P_2(z_0))$$

$$C = \begin{pmatrix} P_{N+2}(z_1) \\ \vdots \\ P_{N+2}(z_n) \end{pmatrix}, \quad D = \begin{pmatrix} P_N(z_1) & \cdots & P_4(z_1) & P_2(z_1) \\ \vdots & & & \vdots \\ P_N(z_n) & \cdots & P_4(z_n) & P_2(z_n) \end{pmatrix}$$

To zero-th order in perturbation theory, we denote $N = N^{(0)}$ and similarly $A^{(0)}, B^{(0)}, C^{(0)}, D^{(0)}$. To take the continuum limit we have to consider the matrix $NN^{(0)-1}$.

Lemma 1 *In the continuum limit, we have*

$$NN^{(0)-1} = \left(\frac{1}{\frac{\sin Z_k}{Z_k} \frac{p_0}{\sin p_0}} \quad \frac{\sin Z_k}{Z_k} \left\{ \frac{1}{Z_k^2 - m^2 \pi^2} - \frac{0}{p_0^2 - m^2 \pi^2} \right\} 2(-1)^m m^2 \pi^2 \right), \quad k, m = 1, \dots, \infty \quad (35)$$

Proof. Since $P_{N+2}(z) = zP_{N+1}(z) - P_N(z)$ and $P_{N+1}(z_k^{(0)}) = 0$, we have

$$C_k^{(0)} = -D_{k1}^{(0)} \implies (D^{(0)-1}C^{(0)})_k = -\delta_{k1}$$

so that

$$N^{(0)-1} = \frac{1}{A+B_1} \begin{pmatrix} 1 & -BD^{(0)-1} \\ F & (A+B_1)D^{(0)-1} - F \otimes BD^{(0)-1} \end{pmatrix}$$

where F is the column vector

$$F = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

Let us compute

$$NN^{(0)-1} = \begin{pmatrix} 1 & 0 \\ \frac{1}{A+B_1}(C+DF) & DD^{(0)-1} - \frac{1}{A+B_1}(C+DF) \otimes BD^{(0)-1} \end{pmatrix}$$

Now

$$\begin{aligned} A+B_1 &= P_{N+2}(z_0) + P_N(z_0) \\ (C+DF)_k &= P_{N+2}(z_k) + P_N(z_k) \end{aligned}$$

so that in the continuum limit we get

$$\frac{1}{A+B_1}(C+DF)_k \rightarrow \frac{\sin Z_k}{Z_k} \frac{p_0}{\sin p_0}$$

The main trick to proceed is an explicit formula for the inverse of the matrix $D^{(0)}$. It is not difficult to check that

$$(D^{(0)-1})_{jk} = \frac{4}{N+1} \sin^2 \frac{k\pi}{N+1} P_{2j}(z_k^{(0)})$$

With this, we find

$$\begin{aligned} (BD^{(0)-1})_m &= \frac{4}{N+1} \sin^2 \frac{m\pi}{N+1} \sum_{j=1}^n P_{2j}(z_0) P_{2j}(z_k^{(0)}) \\ &= \frac{\sin^2 \frac{m\pi}{N+1}}{\sin \frac{p_0}{N+1} \sin \frac{m\pi}{N+1}} \frac{4}{N+1} \sum_{j=1}^n \sin \frac{2jp_0}{N+1} \sin \frac{2jm\pi}{N+1} \rightarrow 2 \frac{m\pi}{p_0} \int_0^1 dx \sin p_0 x \sin m\pi x \end{aligned}$$

and the last integral is easily evaluated with the result

$$(BD^{(0)-1})_m \rightarrow (-1)^m \frac{2m^2\pi^2 \sin p_0}{p_0(p_0^2 - m^2\pi^2)}$$

Similarly we compute

$$(DD^{(0)-1})_{km} = \frac{\sin Z_k}{Z_k} \frac{(-1)^m 2m^2\pi^2}{Z_k^2 - m^2\pi^2}$$

From all this we get eq.(35). ■

We introduce the important infinite matrix

$$M_{km} = \frac{1}{Z_k^2 - m^2\pi^2}, \quad k, m = 1, \dots, \infty \quad (36)$$

and the important vector $|\eta\rangle$

$$|\eta\rangle = M^{-1} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \quad (37)$$

With these notations we can write the inverse of the matrix $NN^{(0)-1}$

Lemma 2

$$(NN^{(0)-1})^{-1} = \begin{pmatrix} \frac{p_0}{\sin p_0} \left(\frac{1}{1 - \langle \chi(p_0) | \eta \rangle} \right) \frac{(-1)^{m+1}}{2m^2\pi^2} \eta_m & \frac{(-1)^m}{2m^2\pi^2} \left\{ \left(1 + \frac{|\eta\rangle\langle\chi(p_0)|}{1 - \langle\chi(p_0)|\eta\rangle} \right) M^{-1} \right\}_{mk} \frac{Z_k}{\sin Z_k} \end{pmatrix}$$

where we have defined the vector $\langle\chi(p_0)|_m = \frac{1}{p_0^2 - \pi^2 m^2}$.

Proof. With the above notations we can write

$$NN^{(0)-1} = \begin{pmatrix} \frac{1}{\frac{\sin Z_k}{Z_k} \frac{p_0}{\sin p_0}} & 0 \\ \frac{\sin Z_k}{Z_k} M_{kl} (\delta_{lm} - \eta_l \chi_m(p_0)) & 2(-1)^m m^2 \pi^2 \end{pmatrix}$$

Letting

$$NN^{(0)-1} = \begin{pmatrix} 1 & 0 \\ Y & X \end{pmatrix} \implies (NN^{(0)-1})^{-1} = \begin{pmatrix} 1 & 0 \\ -X^{-1}Y & X^{-1} \end{pmatrix}$$

we find

$$(X^{-1})_{mk} = \frac{(-1)^m}{2m^2\pi^2} \left\{ \left(1 + \frac{|\eta\rangle\langle\chi(p_0)|}{1 - \langle\chi(p_0)|\eta\rangle} \right) M^{-1} \right\}_{mk} \frac{Z_k}{\sin Z_k} \quad (38)$$

and

$$(-X^{-1}Y)_m = \frac{p_0}{\sin p_0} \left(\frac{1}{1 - \langle\chi(p_0)|\eta\rangle} \right) \frac{(-1)^{m+1}}{2m^2\pi^2} \eta_m$$

■

Let us return to the formula for the spectral curve. Denote

$$\eta(Z) = 1 - \sum_{m=1}^{\infty} \frac{\eta_m}{Z^2 - m^2\pi^2} \quad (39)$$

and

$$|\Gamma\rangle_m = \sum_k M_{mk}^{-1} \frac{Z_k}{\sin Z_k} \gamma_k \quad (40)$$

One has

Proposition 1 *In the continuum limit, the equation of the spectral curve becomes:*

$$\mu + \mu^{-1} = 2 \cos Z + \frac{\sin Z}{Z} \left(-H_0 + \sum_m \frac{H_m}{Z^2 - m^2 \pi^2} \right) \quad (41)$$

where the conserved quantities H_m can be taken as

$$H_m = \Gamma_m + \frac{1}{\eta(p_0)} \langle \chi(p_0) | \Gamma \rangle \eta_m, \quad H_0 = \sum_m \frac{H_m}{p_0^2 - \pi^2 m^2} = \frac{1}{\eta(p_0)} \sum_m \frac{\Gamma_m}{p_0^2 - \pi^2 m^2} \quad (42)$$

Proof. We have

$$\mu + \mu^{-1} - 2 \cos Z = \sum_{i=0}^n P_{N+2-2i}(z) H_{N+2-2i} = \sum_{i=0}^n P_{N+2-2i}(z) (N^{-1} V)_i$$

we rewrite by inserting $1 = N^{(0)-1} N^{(0)}$

$$\mu + \mu^{-1} - 2 \cos Z = \sum_{i,j,k} P_{N+2-2i}(z) (N^{(0)-1})_{ik} (N^{(0)})_{kj} (N^{-1} V)_j \quad (43)$$

Hence, we need to compute

$$\begin{aligned} \sum_i P_{N+2-2i}(z) (N^{(0)-1})_{im} &= \\ & \left(\frac{P_{N+2}(z) + P_N(z)}{P_{N+2}(z_0) + P_N(z_0)}, -\frac{P_{N+2}(z) + P_N(z)}{P_{N+2}(z_0) + P_N(z_0)} B D^{(0)-1} + \sum_i P_{N+2-2i}(z) (D^{(0)-1})_{im} \right) \end{aligned}$$

The limit $N \rightarrow \infty$ is easy to take

$$\sum_i P_{N+2-2i}(z) (N^{(0)-1})_{im} \rightarrow \frac{\sin Z}{Z} \left(\frac{p_0}{\sin p_0}, 2(-1)^m m^2 \pi^2 \left\{ \frac{1}{Z^2 - m^2 \pi^2} - \frac{1}{p_0^2 - m^2 \pi^2} \right\} \right)$$

We are now ready to take the limit $N \rightarrow \infty$ in eq.(43)

$$\mu + \mu^{-1} - 2 \cos Z = \frac{\sin Z}{Z} \left(\frac{p_0}{\sin p_0} \tilde{H}_0 + \sum_{m=1}^{\infty} 2(-1)^m m^2 \pi^2 \left\{ \frac{1}{Z^2 - m^2 \pi^2} - \frac{1}{p_0^2 - m^2 \pi^2} \right\} \tilde{H}_m \right)$$

where

$$\tilde{H}_m = (N N^{(0)-1})^{-1} \begin{pmatrix} \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix} = \begin{pmatrix} 0 \\ X^{-1} \begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix} \end{pmatrix}$$

where X^{-1} is given in eq.(38) and we remembered that $\gamma_0 = 0$. Since $1 - \langle \chi(p_0) | \eta \rangle = \eta(p_0)$ the equation of the spectral curve becomes

$$\mu + \mu^{-1} - 2 \cos Z = \frac{\sin Z}{Z} \left(\sum_m \left(\frac{1}{Z^2 - m^2 \pi^2} - \frac{1}{p_0^2 - m^2 \pi^2} \right) \left(\Gamma_m + \frac{1}{\eta(p_0)} \eta_m \langle \chi(p_0) | \Gamma \rangle \right) \right)$$

■

Another useful expression of this result is:

$$\mu + \mu^{-1} = 2 \cos Z + \frac{\sin Z}{Z} \left(\sum_m \frac{\Gamma_m}{Z^2 - m^2 \pi^2} - \frac{\eta(Z)}{\eta(p_0)} \frac{\Gamma_m}{p_0^2 - m^2 \pi^2} \right) \quad (44)$$

The next proposition performs a few consistency checks.

Proposition 2 *The points $Z = p_0, \mu_0^{\pm 1}$, and $Z = Z_k, \mu_k^{\pm 1}$, all belong to the curve eq.(44).*

Proof. When $Z = p_0$, we find $\mu + \mu^{-1} = 2 \cos p_0$, hence the curve passes through the point $\Lambda = 0, \mu_0^{\pm 1}$, as it should be. When $Z = Z_k$, and recalling that $\eta(Z_k) = 0$, we find

$$\begin{aligned} \mu + \mu^{-1} &= 2 \cos Z_k + \frac{\sin Z_k}{Z_k} \sum_m \frac{1}{Z_k^2 - m^2 \pi^2} M_{ml}^{-1} \frac{Z_l}{\sin Z_l} (\mu_l + \mu_l^{-1} - 2 \cos Z_l) \\ &= 2 \cos Z_k + \frac{\sin Z_k}{Z_k} \sum_m M_{km} M_{ml}^{-1} \frac{Z_l}{\sin Z_l} (\mu_l + \mu_l^{-1} - 2 \cos Z_l) = \mu_k + \mu_k^{-1} \end{aligned}$$

Hence the curve passes through the points $Z_k, \mu_k^{\pm 1}$. ■

We now show that the H_m Poisson commute. We need the following result:

Lemma 3 *One has*

$$\begin{aligned} \{\Gamma_n, \Gamma_m\} &= 0 \\ \{\Gamma_n, \eta_m\} &= \{\Gamma_m, \eta_n\} \\ \{\eta_n, \eta_m\} &= 0 \end{aligned}$$

Proof. The last relation is obvious because the η_m depend only on the Z_k . Consider the second relation. We have

$$\eta_m = M_{mk}^{-1} |1\rangle_k, \quad \Gamma_m = M_{mk}^{-1} |\tilde{\gamma}\rangle_k$$

Hence

$$\begin{aligned} \{\eta_m, \Gamma_n\} &= \{M_{nk}^{-1} 1_k, M_{ml}^{-1} \tilde{\gamma}_l\} = M_{m,l}^{-1} \{M_{nk}^{-1}, \tilde{\gamma}_l\} 1_k \\ &= -M_{m,l}^{-1} M_{n,k'}^{-1} \{M_{k'p}, \tilde{\gamma}_l\} M_{pk}^{-1} |1\rangle_k = -M_{m,l}^{-1} M_{n,l}^{-1} \{M_{lp}, \tilde{\gamma}_l\} \eta_p \end{aligned}$$

where in the last step we used that $\{M_{k'p}, \tilde{\gamma}_l\} = 0$ if $k' \neq l$. The result is obviously symmetric in m and n . Finally the first statement is simple. Recalling the definition of Γ_m in eq.(40), we denote $\xi_k = \frac{Z_k}{\sin Z_k} \gamma_k$. One has

$$\{\Gamma_m, \Gamma_n\} = \{M_{mk}^{-1} \xi_k, M_{nl}^{-1} \xi_l\} = -M_{mr}^{-1} M_{nl}^{-1} [\{M_{rs}, \xi_l\} - \{M_{ls}, \xi_r\}] M_{sk}^{-1} \xi_k$$

but because of the structure of M , we have $\{M_{rs}, \xi_l\} = 0$ if $r \neq l$ and for $r = l$ the term in the square bracket obviously vanishes. This is a special case of a general theorem [17, 18]. \blacksquare

We are now ready to prove

Proposition 3 *The quantities H_0, H_m , Poisson commute*

$$\{H_0, H_n\} = 0, \quad \{H_n, H_m\} = 0$$

Proof. Using eq.(42), one has

$$\{H_n, H_m\} = \eta_n(\{C, \Gamma_m\} + C\{C, \eta_m\}) - \eta_m(\{C, \Gamma_n\} + C\{C, \eta_n\})$$

where we denoted

$$C = \frac{1}{\eta(p_0)} \langle \chi(p_0) | \Gamma \rangle = \frac{1}{\eta(p_0)} \sum_l \frac{\Gamma_l}{p_0^2 - \pi^2 l^2}$$

One has

$$\{\Gamma_m, C\} = \frac{1}{\eta(p_0)} C \sum_l \frac{\{\Gamma_m, \eta_l\}}{p_0^2 - \pi^2 l^2}, \quad \{\eta_m, C\} = \frac{1}{\eta(p_0)} \sum_l \frac{\{\eta_m, \Gamma_l\}}{p_0^2 - \pi^2 l^2}$$

hence

$$\{C, \Gamma_m\} + C\{C, \eta_m\} = -\frac{1}{\eta(p_0)} C \sum_l \frac{\{\Gamma_m, \eta_l\} + \{\eta_m, \Gamma_l\}}{p_0^2 - \pi^2 l^2} = 0$$

\blacksquare

All this means that (Z_k, μ_k) are separated coordinates for the Hamiltonians H_m .

6 Continuum limit of the eigenvector.

Having found the continuum limit of the spectral curve, we now consider the limit of the eigenvector. Again, the continuum limit can be computed, the result being eq.(47).

As seen from eq.(29), the eigenvector can be written as (for i odd)

$$\psi_i = \frac{\sqrt{\mathcal{A}}}{\prod (z^2 - z_j^2)} \frac{\det N_i}{\sqrt{\det \Theta_i \det \Theta_{i+2}}}$$

where

$$N_i = \begin{pmatrix} \mu P_i(z) + P_{N+1-i}(z) & \mu P_1(z) & \cdots & -P_{N+1-i}(z) & \cdots & -P_2(z) \\ P_i(z_1) + \mu_1 P_{N+1-i}(z_1) & P_1(z_1) & \cdots & -\mu_1 P_{N+1-i}(z_1) & \cdots & -\mu_1 P_2(z_1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_i(z_k) + \mu_j P_{N+1-i}(z_k) & P_1(z_k) & \cdots & -\mu_k P_{N+1-i}(z_k) & \cdots & -\mu_k P_2(z_k) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ P_i(z_n) + \mu_n P_{N+1-i}(z_n) & P_1(z_n) & \cdots & -\mu_n P_{N+1-i}(z_n) & \cdots & -\mu_n P_2(z_n) \end{pmatrix} \quad (45)$$

Compared to eq.(29) we have subtracted the i -th column to the first one for the same reason than in the previous section. Also we have used the variable z , z_k instead of λ , λ_k . Let us decompose the matrix N_i in blocs

$$N_i = \begin{pmatrix} U_i & V_i \\ W_i & \Theta_i \end{pmatrix}$$

where

$$\begin{aligned} U_i &\equiv \mu P_i(z) + P_{N+1-i}(z) \\ (W_i)_k &\equiv \mu_k P_i(z_k) + P_{N+1-i}(z_k) \\ (V_i)_j &= \mu P_j(z)\theta(i-j) - P_{N+1-j}(z)\theta(j-i), \quad i, j \text{ odd} \end{aligned}$$

To order zero, we have $-(-1)^k P_j(z_k^{(0)}) = P_{N+1-j}(z_k^{(0)})$ so that

$$N_i^{(0)} = \begin{pmatrix} U_i & V_i \\ 0 & \Theta_i^{(0)} \end{pmatrix}$$

where $\Theta_i^{(0)}$ is the matrix eq.(30) evaluated on the free configuration. It is in fact independent of i and we will denote it by $\Theta^{(0)}$. The appearance of zero in the lower left corner was the reason for the subtraction in eq.(45) and makes things better behaved in perturbation.

The matrix $N_i^{(0)}$ being bloc triangular we can compute its inverse

$$N_i^{(0)-1} = \begin{pmatrix} U_i^{-1} & -U_i^{-1}V_i\Theta^{(0)-1} \\ 0 & \Theta^{(0)-1} \end{pmatrix}$$

so that

$$N_i N_i^{(0)-1} = \begin{pmatrix} 1 & 0 \\ U_i^{-1}W_i & \Theta_i\Theta^{(0)-1} - U_i^{-1}W_i \otimes V_i\Theta^{(0)-1} \end{pmatrix}$$

Returning to the formula for ψ_i , we multiply all the matrices by $\Theta^{(0)-1}$. The factors $\det \Theta^{(0)-1}$ cancel between the numerator and denominator. We arrive at

$$\psi_i = \frac{\sqrt{\mathcal{A}} U_i}{\prod_k z^2 - z_k^2} \frac{\det \left(\Theta_i\Theta^{(0)-1} - U_i^{-1}W_i \otimes V_i\Theta^{(0)-1} \right)}{\sqrt{\det \left(\Theta_i\Theta^{(0)-1} \right) \det \left(\Theta_{i+2}\Theta^{(0)-1} \right)}}$$

We want to take the scaling limit of this expression. Again, the main trick is an explicit formula for the inverse of $\Theta^{(0)}$. It is not difficult to check that

$$(\Theta^{(0)-1})_{jk} = \frac{4}{N+1} \sin^2 \frac{k\pi}{N+1} P_j(z_k^{(0)})$$

Let us compute $\Theta_i\Theta^{(0)-1}$. Using the parametrisation eq.(33) we find (recall that i is assumed to be odd)

$$(\Theta_i\Theta^{(0)-1})_{km} = \frac{4}{N+1} \frac{\sin \frac{m\pi}{N+1}}{\sin \frac{Z_k}{N+1}} \left\{ \sum_{j=1, \text{odd}}^{i-2} \sin \frac{jZ_k}{N+1} \sin \frac{j m \pi}{N+1} \right.$$

$$-\mu_k \sum_{j=i,\text{odd}}^{N-1} \sin \frac{(N+1-j)Z_k}{N+1} \sin \frac{j m \pi}{N+1} \left. \vphantom{\sum} \right\}$$

Defining $\Theta(x)$ as the scaling limit of $\Theta_i \Theta^{(0)-1}$, we find (there is a factor 1/2 because the sum is over j odd only)

$$(\Theta(x))_{km} = 2 \frac{m\pi}{Z_k} \left\{ \int_0^x dy \sin Z_k y \sin m\pi y - \mu_k \int_x^1 dy \sin Z_k(1-y) \sin m\pi y \right\}$$

Similarly we define $U(x)$ and $W_k(x)$ by

$$U_i = (N+1)U(x), \quad (W_i)_k = (N+1)W_k(x)$$

We find

$$U(x) = \mu \frac{\sin x Z}{Z} + \frac{\sin(1-x)Z}{Z}$$

and

$$W_k(x) = \frac{\sin Z_k x}{Z_k} + \mu_k \frac{\sin Z_k(1-x)}{Z_k}$$

Finally, we have (again there is a factor 1/2 because the sum is over j odd only)

$$(V_i \Theta^{(0)-1})_m = V_m(x)$$

where

$$V_m(x) = 2m\pi \left\{ \int_0^x dy \mu \frac{\sin y Z}{Z} \sin m\pi y - \int_x^1 \frac{\sin(1-y)Z}{Z} \sin m\pi y \right\} \quad (46)$$

Putting everything together, we arrive at (up to a factor independent of x ⁶)

Proposition 4

$$\psi(x) = U(x) - \langle V(x) | \Theta^{-1}(x) | W(x) \rangle \quad (47)$$

where we denoted by $\langle V(x) |$ the row vector with components $V_m(x)$ and by $|W(x)\rangle$ the column vector with components $W_k(x)$.

Eq.(47) is the generalization of eq.(20). Here, the Z and μ dependence is hidden in the function $U(x)$ and the vector $V(x)$. For the moment they are free complex parameters.

We now want to specialize to $Z = p_0, \mu_0 = e^{\pm i p_0}$. We have

$$U(x)|_{p_0, e^{\pm i p_0}} = \frac{\sin p_0}{p_0} e^{\pm i p_0 x}, \quad V_m(x)|_{p_0, e^{\pm i p_0}} = U(x)|_{p_0, e^{\pm i p_0}} \tilde{V}_m^{(\pm)}(x)$$

⁶In this factor we will include in particular $\frac{1}{\prod_k (1 - Z_k^2/Z^2)}$ which produces the poles at $Z^2 = Z_k^2$. This is important for the analyticity properties of $\psi(x)$ but plays little role for the considerations of this paper.

where

$$\tilde{V}_m^{(\pm)}(x) = -m\pi \left[\frac{e^{im\pi x}}{m\pi \pm p_0} + \frac{e^{-im\pi x}}{m\pi \mp p_0} \right]$$

Hence, up to a constant

$$\psi^{(\pm)}(x) = e^{\pm ip_0 x} \left[1 - \langle \tilde{V}^{(\pm)}(x) | \Theta^{-1}(x) | W(x) \rangle \right]$$

These are the primary fields of CFT. Their logarithmic derivatives are the free fields of the Coulomb gaz representation. Notice that we have two such fields playing a completely symmetrical role: we go from one to the other by changing $p_0 \rightarrow -p_0$. This circumstance was recognized and used with great profit in [14]. The separated variables make this symmetry explicit and built in.

7 Schroedinger equation.

Having found a formula for the wave function $\psi(x)$, the next question is to find the potential in the Schroedinger equation that $\psi(x)$ is expected to satisfy. At this point it is simpler to forget the lattice model and work directly with eq.(47).

Let us denote by $|E(x)\rangle$ the vector with components

$$E_m(x) = 2m\pi \sin m\pi x$$

Calculating explicitly the integrals in eq.(46), we find (' denotes the derivative with respect to x)

$$V_m(x) = \frac{U(x)E'_m(x) - U'(x)E_m(x)}{Z^2 - m^2\pi^2} \quad (48)$$

and similarly for $\Theta(x)$, we find

$$\Theta_{km}(x) = \frac{W_k(x)E'_m(x) - W'_k(x)E_m(x)}{Z_k^2 - m^2\pi^2} \quad (49)$$

Notice the important formulae

$$\langle V'(x) | = U(x)\langle E(x) |, \quad \Theta'(x) = |W(x)\rangle\langle E(x) |$$

The derivative of the matrix $\Theta(x)$ is a rank one projector. The matrix $\Theta(x)$ has a form familiar in the theory of integrable systems, and we know that it leads to non linear differential equations. Indeed, let us define the vector $|\Phi(x)\rangle$ by

$$|\Phi(x)\rangle = \Theta^{-1}(x)|W(x)\rangle \quad (50)$$

and let K be the diagonal matrix

$$K_{mm'} = m\pi \delta_{mm'}$$

Proposition 5 *The vector $|\Phi(x)\rangle$ satisfies the set of coupled non linear second order differential equations*

$$|\Phi''\rangle + 2(\langle\Phi|E\rangle)'|\Phi\rangle + K^2|\Phi\rangle = 0 \quad (51)$$

where

$$\langle\Phi|E\rangle(x) = \sum_m \Phi_m(x)E_m(x)$$

Proof. By derivation, and using the formula for Θ' , we get

$$\begin{aligned} |W\rangle &= \Theta|\Phi\rangle \\ |W'\rangle &= \Theta'|\Phi\rangle + \Theta|\Phi'\rangle = \langle\Phi|E\rangle|W\rangle + \Theta|\Phi'\rangle \\ |W''\rangle &= \langle\Phi|E\rangle|W'\rangle + \Theta|\Phi''\rangle + (\langle\Phi|E\rangle' + \langle\Phi'|E\rangle)|W\rangle \end{aligned}$$

But we also have

$$\mathcal{Z}^2\Theta - \Theta K^2 = |W\rangle\langle E'| - |W'\rangle\langle E| \quad (52)$$

where we defined the diagonal matrix \mathcal{L}

$$\mathcal{Z}_{kk'} = Z_k\delta_{kk'}$$

Applying this identity to $|\Phi\rangle$, we get

$$\mathcal{Z}^2\Theta|\Phi\rangle - \Theta K^2|\Phi\rangle = \langle E'|\Phi\rangle|W\rangle - \langle E|\Phi\rangle|W'\rangle$$

But

$$\mathcal{Z}^2\Theta|\Phi\rangle = \mathcal{L}^2|W\rangle = -|W''\rangle$$

and therefore

$$|W''\rangle = \langle E|\Phi\rangle|W'\rangle - \langle E'|\Phi\rangle|W\rangle - \Theta K^2|\Phi\rangle$$

Comparing these two expressions of $|W''\rangle$, we find

$$\Theta|\Phi''\rangle + (\langle\Phi|E\rangle' + \langle\Phi'|E\rangle)|W\rangle = -\langle E'|\Phi\rangle|W\rangle - \Theta K^2|\Phi\rangle$$

Multiplying by Θ^{-1} yields eq.(51). ■

We are now ready to find the Schroedinger equation satisfied by ψ .

Proposition 6 *The function $\psi(x)$ defined by eq.(47) satisfies the linear second order differential equation*

$$-\psi''(x) - [p_0^2 + 2(\langle E|\Phi\rangle)']\psi(x) = \Lambda^2\psi(x) \quad (53)$$

Proof. We have

$$\psi(x) = U(x) - \langle V(x)|\Theta^{-1}|W(x)\rangle = U(x) - \langle V(x)|\Phi(x)\rangle$$

Using the formula for $\langle V(x)|$, we get

$$\psi(x) = U \left(1 - \langle E'| \frac{1}{Z^2 - K^2} |\Phi\rangle \right) + U' \langle E| \frac{1}{Z^2 - K^2} |\Phi\rangle$$

Next, remembering that $U''(x) = -Z^2U(x)$, $|E''(x)\rangle = -K^2|E(x)\rangle$, we obtain

$$\psi'(x) = -U \left(\langle E|\Phi\rangle + \langle E'|\frac{1}{Z^2 - K^2}|\Phi'\rangle \right) + U' \left(1 + \langle E|\frac{1}{Z^2 - K^2}|\Phi'\rangle \right)$$

and

$$\begin{aligned} \psi''(x) = & -U \left(Z^2 + \langle E|\Phi'\rangle + (\langle E|\Phi\rangle)' + \langle E'|\frac{1}{Z^2 - K^2}|\Phi''\rangle \right) \\ & + U' \left(-\langle E|\Phi\rangle + \langle E|\frac{1}{Z^2 - K^2}|\Phi''\rangle \right) \end{aligned}$$

Using now the equation for $|\Phi''\rangle$, we get eq.(53). ■

The potential $T(x) = 2(\langle E|\Phi\rangle)'$ can also be written directly in terms of Θ . In fact, we have

$$\partial_x^2 \log \det \Theta = \partial_x \operatorname{Tr} \Theta^{-1} \Theta' = \partial_x \langle E|\Theta^{-1}W\rangle = \partial_x \langle E|\Phi\rangle$$

hence

$$T(x) = 2\partial_x \langle E|\Phi\rangle = 2\partial_x^2 \log \det \Theta(x)$$

The Schroedinger equation therefore also reads

$$\psi''(x) + [p_0^2 + 2 \partial_x^2 \log \det \Theta] \psi(x) = -\Lambda^2 \psi(x) \quad (54)$$

In this formula both the potential and the function $\psi(x)$ are known. The potential therefore belongs to the class of exactly solvable potentials. It is strongly reminiscent of the formula for finite zones potentials [20, 21, 22, 23]. It can probably also be obtained by an infinite sequence of Darboux transformations [24].

The parameter μ which enters the function $U(x)$ and the vector $\langle V(x)|$ was, up to now, a free parameter. Eq.(47) therefore provides two linearly independent solutions of eq.(53). We now introduce the spectral curve by imposing the quasiperiodicity of $\psi(x)$.

8 Bloch Waves and Spectral Curve.

So far, $\psi(x)$ was defined on the interval $[0, 1]$. We extend its definition by imposing

$$\psi(x+1) = \mu\psi(x)$$

This extension is continuous as we now show.

Proposition 7

$$\psi(1) - \mu\psi(0) = 0$$

Proof. This follows immediately from

$$W_k(1) = \mu_k^{-1}W_k(0), \quad \Theta_{km}(1) = \mu_k^{-1}\Theta_{km}(0)(-1)^m \quad U(1) = \mu U(0)$$

■

It is worth computing explicitly $\psi(0)$. In terms of the matrix M introduced in eq.(36), we have

$$\Theta_{km}(0) = W_k(0)M_{km}E'_m(0), \quad U(0) = \frac{\sin Z}{Z}, \quad V_m(0) = U(0)\frac{1}{Z^2 - m^2\pi^2}E'_m(0)$$

It follows that

$$\psi(0) = \frac{\sin Z}{Z} \left(1 - \sum_m \frac{\eta_m}{Z^2 - m^2\pi^2} \right) = \frac{\sin Z}{Z} \eta(Z) \quad (55)$$

where η_m and $\eta(Z)$ are defined in eqs.(37, 39). Notice that when $Z^2 = Z_k^2$, we have $\psi(0) = 0$.

We now turn to the derivative of $\psi(x)$.

Lemma 4 *One has*

$$\begin{aligned} \psi'(1) - \mu\psi'(0) &= \mu \tilde{\Gamma}(\Lambda, \mu) \\ \tilde{\Gamma}(\Lambda, \mu) &= \mu + \mu^{-1} - 2 \cos Z - \frac{\sin Z}{Z} \sum_{m=1}^{\infty} \frac{E'_m(0)\Phi'_m(0) - E'_m(1)\Phi'_m(1)}{Z^2 - m^2\pi^2} \end{aligned} \quad (56)$$

Proof. We have

$$U(1) = \mu \frac{\sin Z}{Z}, \quad U(0) = \frac{\sin Z}{Z}$$

and

$$U'(1) = \mu \cos Z - 1, \quad U'(0) = \mu - \cos Z$$

Using $E_k(1) = E_k(0) = 0$, we get

$$\psi'(1) = -U(1)\langle E'(1) | \frac{1}{Z^2 - K^2} | \Phi'(1) \rangle + U'(1)$$

$$\psi'(0) = -U(0)\langle E'(0) | \frac{1}{Z^2 - K^2} | \Phi'(0) \rangle + U'(0)$$

From this the result follows. ■

At this point it is tempting to choose for the spectral curve $\tilde{\Gamma}(\Lambda, \mu) = 0$. However, this cannot be correct because the point $Z = p_0, \mu = \mu_0$ does not belong to it.

We have to change the Schroedinger equation. The only possible modification is at the edges. We consider the equation

$$\psi''(x) + \left[p_0^2 + 2\langle E\Phi \rangle' + H_0\delta(x) \right] \psi(x) = -\Lambda^2\psi(x) \quad (57)$$

The bulk formula for $\psi(x)$ does not change. The continuity of $\psi(x)$ at $x = 1$ is still valid, but the derivative now has a discontinuity

$$\int_{1-}^{1+} \psi'' + H_0\psi(1) = 0$$

Using

$$\psi(1) = \mu \eta(Z) \frac{\sin Z}{Z}$$

the Bloch condition becomes

$$\psi'(1) - \mu\psi'(0) - H_0\mu\eta(Z) = 0$$

that is

$$\mu + \mu^{-1} = 2 \cos Z + \frac{\sin Z}{Z} \left(\sum_m \frac{\Gamma_m}{Z^2 - m^2\pi^2} - H_0\eta(Z) \right)$$

where we have set

$$\Gamma_m = E'_m(0)\Phi'_m(0) - E'_m(1)\Phi'_m(1) \quad (58)$$

We now determine the coefficient H_0 by requiring that the curve passes through the points $p_0, \mu_0^{\pm 1}$. We find

$$H_0 = \frac{1}{\eta(p_0)} \sum_m \frac{\Gamma_m}{p_0^2 - m^2\pi^2}$$

Hence the curve takes the form

$$\mu + \mu^{-1} = 2 \cos Z + \frac{\sin Z}{Z} \left(\sum_m \frac{\Gamma_m}{Z^2 - m^2\pi^2} - \frac{\eta(Z)}{\eta(Z_0)} \frac{\Gamma_m}{p_0^2 - m^2\pi^2} \right)$$

In order to compare with eq.(44), we must compute Γ_m . We have

$$W_k(0) = \mu_k \frac{\sin Z_k}{Z_k}, \quad W_k(1) = \frac{\sin Z_k}{Z_k}, \quad W'_k(0) = 1 - \mu_k \cos Z_k, \quad W'_k(1) = \cos Z_k - \mu_k$$

and

$$\Theta_{km}(0) = \frac{W_k(0)E'_m(0)}{Z_k^2 - \pi^2 m^2} = W_k(0)M_{km}E'_m(0), \quad \Theta_{km}(1) = \frac{W_k(1)E'_m(1)}{Z_k^2 - \pi^2 m^2} = W_k(1)M_{km}E'_m(1)$$

where M is the matrix introduced in eq.(36). Since $|\Phi'(0)\rangle = \Theta^{-1}(0)|W'(0)\rangle$ and $|\Phi'(1)\rangle = \Theta^{-1}(1)|W'(1)\rangle$, one has

$$E'_m(0)\Phi'_m(0) = \sum_k M_{mk}^{-1} (\mu_k^{-1} - \cos Z_k) \frac{Z_k}{\sin Z_k}, \quad E'_m(1)\Phi'_m(1) = \sum_k M_{mk}^{-1} (\cos Z_k - \mu_k) \frac{Z_k}{\sin Z_k}$$

hence

$$\Gamma_m = E'_m(0)\Phi'_m(0) - E'_m(1)\Phi'_m(1) = \sum_k M_{mk}^{-1} \frac{Z_k}{\sin Z_k} \gamma_k$$

where we recall that $\gamma_k = \mu_k + \mu_k^{-1} - 2 \cos Z_k$. This is exactly eq.(40) and shows that we have recovered the spectral curve eq.(44).

Proposition 8 *The function Ψ has no pole at the point $Z = Z_k, \mu_k^{-1}$*

Proof. Here we restaure the factor $\frac{1}{\prod_k(1-Z_k^2/Z^2)}$.

$$U(x)|_{Z_k, \mu_k^{-1}} = \mu_k^{-1} W_k(x), \implies V_{k'}(x)|_{Z_k, \mu_k^{-1}} = \mu_k^{-1} \Theta_{kk'}(x)$$

$$\psi(x)|_{Z_k, \mu_k^{-1}} \simeq \frac{1}{\prod_k(Z^2 - Z_k^2)} \mu_k^{-1} [W_k(x) - \Theta_{kk'}(x) \Theta_{k'k''}^{-1}(x) W_{k''}(x)] = \text{regular}$$

■

9 Perturbation theory.

Now that we have found an expression for the potential $u(x)$ in terms of a countable set of variables Z_k, μ_k , we would like to check the Virasoro Poisson bracket directly. Recall that

$$u(x) = p_0^2 + T(x) + H_0 \delta(x), \quad T(x) = 2\partial_x \langle E | \Phi \rangle$$

Notice first that $T(x)$ has no Fourier component T_0 :

$$T_0 = \int_0^1 dx T(x) = 2 \int_0^1 dx \langle \langle E | \Phi \rangle \rangle' = 2(\langle E(1) | \Phi(1) \rangle - \langle E(0) | \Phi(0) \rangle) = 0$$

The Fourier expansion of the potential $u(x)$ reads

$$u(x) = \sum_n L_n e^{2i\pi n x} = p_0^2 + \sum_n (T_n + H_0) e^{2i\pi n x}$$

we must therefore identify

$$L_0 = p_0^2 + H_0 = p_0^2 + \frac{1}{\eta(Z_0)} \sum_m \frac{\Gamma_m}{p_0^2 - m^2 \pi^2} \quad (59)$$

and

$$L_n = T_n + H_0, \implies T_n = L_n - L_0 + p_0^2, \quad n \neq 0$$

If $u(x)$ has Poisson bracket eq.(4), the algebra of the L_n reads

$$\{L_n, L_m\} = 8i\pi(n-m)L_{n+m} - 16i\pi^3 n^3 \delta_{n+m,0}$$

The Poisson algebra for the T_n is then closed:

$$\{T_n, T_m\} = 8i\pi(n-m)T_{n+m} - 8i\pi n T_n + 8i\pi m T_m - 16i(\pi^3 n^3 - \pi p_0^2 n) \delta_{n+m,0}$$

or, in a form that will be useful later

$$\{T(x), T(y)\} = 2\delta'''(x-y) + 4(2p_0^2 + T(x) + T(y))\delta'(x-y) - 4T'(x)\delta(y) + 4T'(y)\delta(x) \quad (60)$$

In this section, we consider the situation where the variables (Z_k, μ_k) are close to the free configuration eq.(16). We set

$$Z_k = k\pi(1 - \delta\lambda_k), \quad \mu_k = (-1)^k e^{\delta\mu_k} \quad (61)$$

and perform a perturbation theory in $\delta\lambda_k, \delta\mu_k$. If $Z_k = k\pi, \mu_k = (-1)^k$, we have $\Theta^{(0)}(x) = \text{Id}$ by construction. So, we can write the expansion

$$\begin{aligned} \Theta(x) &= \text{Id} + \Theta^{(1)}(x) + \Theta^{(2)}(x) + \dots \\ |W(x)\rangle &= |W^{(1)}(x)\rangle + |W^{(2)}(x)\rangle + \dots \end{aligned}$$

where we have taken into account that $|W^{(0)}(x)\rangle = 0$. It follows that

$$|\Phi(x)\rangle = |\Phi^{(1)}(x)\rangle + |\Phi^{(2)}(x)\rangle + \dots$$

where

$$|\Phi^{(1)}\rangle = |W^{(1)}\rangle, \quad |\Phi^{(2)}\rangle = |W^{(2)}\rangle - \Theta^{(1)}|W^{(1)}\rangle, \dots$$

To lowest order, we find easily

$$T^{(1)}(x) = 2\langle E|\Phi^{(1)}\rangle' = -4 \sum_k \delta\lambda_k k^2 \pi^2 \cos 2k\pi x + \delta\mu_k k\pi \sin 2k\pi x$$

This shows in particular that $\delta\lambda_k$ and $\delta\mu_k$ are just the Fourier components of the potential in this first approximation. The Poisson bracket eq.(3) becomes to leading order

$$\{\delta\lambda_k, \delta\mu_k\} = -2 \left(1 - \frac{p_0^2}{k^2 \pi^2} \right)$$

To define modes independent of the zero mode p_0 we introduce

$$a_k = \alpha_k(\delta\mu_k + ik\pi\delta\lambda_k), \quad a_k^\dagger = \bar{\alpha}_k(\delta\mu_k - ik\pi\delta\lambda_k)$$

where the coefficients $\alpha_k, \bar{\alpha}_k$ satisfy⁷

$$\alpha_k \bar{\alpha}_k = \frac{1}{4\pi} \frac{k^2 \pi^2}{p_0^2 - k^2 \pi^2}$$

With this choice, one has

$$\{a_k, a_k^\dagger\} = ik$$

we can then rewrite

$$T^{(1)}(x) = 2i \sum_k k\pi \left(\frac{a_k}{\alpha_k} e^{2ik\pi x} - \frac{a_k^\dagger}{\bar{\alpha}_k} e^{-2ik\pi x} \right)$$

It is now straightforward to compute the Poisson bracket

$$\{T^{(1)}(x), T^{(1)}(y)\} = 2\delta'''(x-y) + 8p_0^2 \delta'(x-y)$$

This is the correct result for the Virasoro Poisson bracket in this approximation. Notice that the term $\delta'''(x-y)$ is exact already at this level.

⁷It is known [14] that the poles at $p_0^2 = k^2 \pi^2$ are classical remnants of the zeroes of the Kac determinant.

Next, we look at the conserved quantities. The leading terms in the expansions of η_m and H_m are easy to find:

$$\eta_m \simeq -2m^2\pi^2\delta\lambda_m, \quad \Gamma_m \simeq 2m^2\pi^2(\delta\mu_m^2 + m^2\pi^2\delta\lambda_m^2)$$

To see it, consider the defining relations of η_m

$$\sum_m \frac{\eta_m}{Z_k^2 - \pi^2 m^2} = 1, \forall k$$

When Z_k is given by eq.(61), the dominant term in the above sum is $m = k$. The equation becomes $\eta_k/(-2k^2\pi^2\delta\lambda_k) = 1$. The same argument starting with the equation of the spectral curve, eq.(41), taken at the point Z_k, μ_k which belongs to it, yields the formula for H_m . Written with the oscillators a_m, a_m^\dagger , we find

$$H_m = 8\pi(p_0^2 - \pi^2 m^2)a_m^\dagger a_m, \quad H_0 = 8\pi \sum_m a_m^\dagger a_m$$

It is clear that the H_m are in involution at this order. As we see, in first approximation, the dynamical system reduces to a set of decoupled harmonic oscillators. The generator L_0 is given by

$$L_0 = p_0^2 + 8\pi \sum_m a_m^\dagger a_m$$

It is easy to verify that

$$\{L_0, T^{(1)}(x)\}_0 = -4\partial_x T^{(1)}(x)$$

These perturbative arguments are a good indication that $u(x)$ indeed satisfies the Virasoro Poisson bracket. Clearly we will not go very far in perturbation and we now look for a formal proof of this fact. For that purpose, we need some preparation.

10 Some identities.

Before computing Poisson brackets to check the Virasoro algebra, we collect a number of useful identities. We start with a formula for the inverse matrix $\Theta^{-1}(x)$. It has the same form as $\Theta(x)$.

Proposition 9 *Let us define*

$$\langle F| = \langle E|\Theta^{-1} \tag{62}$$

Then, we can write

$$\Theta_{mk}^{-1} = \frac{\Phi_m F'_k - \Phi'_m F_k}{Z_k^2 - \pi^2 m^2} \tag{63}$$

Proof. Multiplying eq.(52) on both sides by Θ^{-1} , we get

$$\Theta^{-1} \mathcal{Z}^2 - K^2 \Theta^{-1} = \Theta^{-1} |W\rangle \langle E'| \Theta^{-1} - \Theta^{-1} |W'\rangle \langle E| \Theta^{-1}$$

and so

$$\Theta_{mk}^{-1} = \frac{(\Theta^{-1} |W'\rangle)_m (\langle E| \Theta^{-1})_k - (\Theta^{-1} |W\rangle)_m (\langle E'| \Theta^{-1})_k}{\pi^2 m^2 - Z_k^2}$$

But

$$\Theta^{-1}|W'\rangle = |\Phi'\rangle + \langle E|\Phi\rangle|\Phi\rangle, \quad \langle E'|\Theta^{-1} = \langle F'| + \langle E|\Phi\rangle\langle F|$$

Plugging into the above formula, we obtain eq.(63). ■

Proposition 10 *The vector $\langle F|$ satisfies a set of differential equations.*

$$\langle F''| + 2(\langle E|\Phi\rangle)'\langle F| + \langle F|Z^2 = 0 \quad (64)$$

Proof. The proof is the same as for $|\Phi\rangle$.

From this we easily deduce

$$(\Theta^{-1})' = -|\Phi\rangle\langle F|$$

which can also be proved using the similar property of Θ . Let us define

$$A_k(x) = \sum_m \frac{E_m(x)\Phi_m(x)}{Z_k^2 - m^2\pi^2}, \quad B_k(x) = \sum_m \frac{E'_m(x)\Phi_m(x)}{Z_k^2 - m^2\pi^2}$$

$$C_k(x) = \sum_m \frac{E_m(x)\Phi_m(x)}{(Z_k^2 - m^2\pi^2)^2}, \quad D_k(x) = \sum_m \frac{E'_m(x)\Phi_m(x)}{(Z_k^2 - m^2\pi^2)^2}$$

Proposition 11 *We have the identity*

$$(1 - B_k)W_k + A_kW'_k = 0 \quad (65)$$

Proof. This is just a rewriting of $|W\rangle = \Theta|\Phi\rangle$ using eq.(49) for $\Theta(x)$. ■
Similarly, one has

Proposition 12 *The following two identities hold*

$$(1 - B_k + A'_k)F_k - A_kF'_k = 0 \quad (66)$$

$$(B'_k + Z_k^2 A_k)F_k + (1 - B_k)F'_k = 0 \quad (67)$$

Proof. The first identity is a rewriting of $\langle F| = \langle E|\Theta^{-1}$ using eq.(63) for $\Theta^{-1}(x)$. The second identity is just a rewriting of $\langle F'| = \langle E'|\Theta^{-1} - \langle E|\Phi\rangle\langle F|$. ■

The above two identities form a linear system for F_k and F'_k . Its compatibility implies the following:

Proposition 13

$$(1 - B_k)^2 + A_kB'_k - A'_k B_k + A'_k + Z_k^2 A_k^2 = 0 \quad (68)$$

Let us define

$$\tilde{F}_k = W'_k(1 - B_k) + W''_k A_k \quad (69)$$

An important consequence of eqs.(66,67) is

Proposition 14 *The functions $F_k(x)$ and \tilde{F}_k are proportional.*

$$F_k(x) = -\frac{\zeta_k}{\mu_k \gamma_k} \tilde{F}_k(x) \quad (70)$$

The proportionality coefficient is written in this specific way for later convenience. The quantity γ_k is the one defined in eq.(34).

Proof. Let us compute the Wronskian

$$\begin{aligned} Wr(F_k, \tilde{F}_k) &= F_k \left(W''_k(1 - B_k) - W'_k B'_k + W'''_k A_k + W''_k A'_k \right) - F'_k \left(W'_k(1 - B_k) + W''_k A_k \right) \\ &= -\Lambda_k^2 W_k \left((1 - B_k + A'_k) F_k - A_k F'_k \right) - W'_k \left((B'_k + Z_k^2 A_k) F_k + (1 - B_k) F'_k \right) = 0 \end{aligned}$$

■

Proposition 15 *The following quantities are constants independent of x*

$$\eta_m = \Phi_m E'_m - \Phi'_m E_m - \langle E\Phi \rangle \Phi_m E_m - \sum_{p \neq m} \frac{1}{\pi^2(p^2 - m^2)} (\Phi_m \Phi'_p - \Phi'_m \Phi_p) (E'_m E_p - E_m E'_p) \quad (71)$$

The quantities η_m defined in eq.(71) are in fact the same as the ones introduced in eq.(37).

Proof. To prove the first statement, just take the derivative with respect to x and use eq.(51). To prove the second statement, use the η_m defined in eq.(71) to rewrite eq.(68) as

$$\sum_m \frac{\eta_m}{(Z_k^2 - m^2 \pi^2)} = 1, \quad \forall k \quad (72)$$

which is exactly the same as eq.(37). ■

A straightforward consequence is the "trace" formula that will be useful later:

$$\langle E\Phi' \rangle - \langle E'\Phi \rangle + \langle E\Phi \rangle^2 = - \sum_m \eta_m \quad (73)$$

We now compute the coefficients ζ_k appearing in eq.(70)

Proposition 16 *The coefficients ζ_k in eq.(70) are determined by the set of equations*

$$\sum_k \frac{\zeta_k}{Z_k^2 - m^2 \pi^2} = 1, \quad \forall m \quad (74)$$

These equations are dual to eq.(72).

Proof. Start with

$$\langle F|W \rangle = \langle E|\Theta^{-1}|W \rangle = \langle E|\Phi \rangle$$

Using eq.(65, 69,70), we have

$$F_k W_k = -\frac{\zeta_k}{\mu_k \gamma_k} (-W_k'^2 + W_k'' W_k) A_k = \zeta_k A_k \quad (75)$$

hence

$$\langle F|W \rangle = \sum_k \zeta_k A_k = \sum_m \left(\sum_k \frac{\zeta_k}{Z_k^2 - m^2 \pi^2} \right) E_m \Phi_m = \langle E|\Phi \rangle = \sum_m E_m \Phi_m$$

Since this has to hold for all x , the only possibility is eq.(74). ■

We collect below a few more identities of the type of eq.(75) that will be important later

Proposition 17

$$F_k W_k = \zeta_k A_k \quad (76)$$

$$F_k W_k' = -\zeta_k (1 - B_k) \quad (77)$$

$$F_k' W_k = \zeta_k (1 - B_k + A_k') \quad (78)$$

$$F_k' W_k' = \zeta_k (B_k' + Z_k^2 A_k) \quad (79)$$

Next, we relate the η_m and ζ_k

Proposition 18 *The following relation holds:*

$$\sum_m \frac{\eta_m}{(Z_k^2 - m^2 \pi^2)^2} = \frac{1}{\zeta_k}$$

Proof. We start from

$$\sum_m \Theta_{km} \Theta_{mk'}^{-1} = \delta_{kk'}$$

When $k = k'$ this gives

$$W_k F_k' D_k - W_k F_k (D_k' - A_k + Z_k^2 C_k) - W_k' F_k' C_k + W_k' F_k (C_k' - D_k) = 1$$

or using eqs.(76-79)

$$\zeta_k \left\{ 2D_k(1 - B_k) + D_k A_k' - D_k' A_k + A_k^2 - 2Z_k^2 C_k A_k - C_k B_k' - C_k'(1 - B_k) \right\} = 1$$

Expanding this formula using ($p \neq m$)

$$\frac{1}{(Z_k^2 - \pi^2 m^2)^2 (Z_k^2 - \pi^2 p^2)} = -\frac{1}{\pi^2 (p^2 - m^2)} \frac{1}{(Z_k^2 - \pi^2 m^2)^2} - \frac{1}{\pi^4 (p^2 - m^2)^2} \left\{ \frac{1}{(Z_k^2 - \pi^2 m^2)} - \frac{1}{(Z_k^2 - \pi^2 p^2)} \right\}$$

we get the result with η_m represented by eq.(71). ■

An immediate consequence is an expansion of the function $\eta(Z)$ near $Z^2 = Z_k^2$

$$\eta(Z) = (Z^2 - Z_k^2) \frac{1}{\zeta_k} + \dots \quad (80)$$

Returning to the formula for $\psi(x)$, we can write it as

$$\psi(x) = \frac{\mu - e^{-iZ}}{2iZ} w(x, Z) - \frac{\mu - e^{iZ}}{2iZ} w^*(x, Z)$$

where we define

$$w(x, Z) = e^{iZx} \left(1 - \langle E' | \frac{1}{Z^2 - K^2} | \Phi \rangle + \langle E | \frac{iZ}{Z^2 - K^2} | \Phi \rangle \right) \quad (81)$$

$$w^*(x, Z) = e^{-iZx} \left(1 - \langle E' | \frac{1}{Z^2 - K^2} | \Phi \rangle - \langle E | \frac{iZ}{Z^2 - K^2} | \Phi \rangle \right) \quad (82)$$

The functions w and w^* are defined on the Riemann sphere with a puncture at ∞ . One can compute the Wronskian

$$w'w^* - w^{*'}w = 2iZ\eta(Z)$$

This Wronskian vanishes precisely when $Z^2 = Z_k^2$. Hence, when $Z = Z_k$, $w(x, Z_k)$ becomes proportional to $w^*(x, Z_k)$. Indeed

$$\begin{aligned} w(x, Z_k) &= e^{iZ_k x} (1 - B_k + iZ_k A_k) \\ w^*(x, Z_k) &= e^{-iZ_k x} (1 - B_k - iZ_k A_k) \end{aligned}$$

Then eq.(65) becomes

$$w(x, Z_k) = \frac{\alpha_k^{(+)}}{\alpha_k^{(-)}} w^*(x, Z_k) \quad (83)$$

where

$$\alpha_k^{(\pm)} = 1 - \mu_k e^{\pm iZ_k}, \quad \alpha_k^{(+)} \alpha_k^{(-)} = \mu_k \gamma_k \quad (84)$$

The function \tilde{F}_k is solution of eq.(64). It is not difficult to see that the other solution of this equation is

$$\tilde{G}_k = A_k W_k' + 2Z_k (W_k D_k - W_k' C_k) + x \tilde{F}_k$$

In terms of w, w^* defined in eqs.(81,82), we have

$$\tilde{F}_k = \frac{1}{2} (\alpha_k^{(-)} w|_{Z_k} + \alpha_k^{(+)} w^*|_{Z_k}) = \alpha_k^{(-)} w|_{Z_k} = \alpha_k^{(+)} w^*|_{Z_k} \quad (85)$$

$$\tilde{G}_k = -\frac{i}{2} (\alpha_k^{(-)} \partial_Z w|_{Z_k} - \alpha_k^{(+)} \partial_Z w^*|_{Z_k}) \quad (86)$$

This will play an important role below.

11 Virasoro algebra.

We are now ready to compute the Poisson bracket $\{T(x), T(y)\}$. The result is precisely the algebra of the $T_n = L_n - L_0$, eq.(60).

Proposition 19 *Let $T(x) = 2\partial_x \langle E(x) | \Phi(x) \rangle$, then we have*

$$\left\{ \int_0^1 dx X(x) T(x), \int_0^1 dy Y(y) T(y) \right\} = - \int_0^1 dx (X''' Y - X Y''') - 4 \int_0^1 dx (X' Y - X Y') (p_0^2 + T) + 4 \int_0^1 dx X(x) \delta(x) \int_0^1 dy Y(y) T'(y) - 4 \int_0^1 dx X(x) T'(x) \int_0^1 dy Y(y) \delta(y) \quad (87)$$

The proof is rather long and we will split it in several lemmas. Since $|\Phi\rangle = \Theta^{-1}|W\rangle$, we have

$$\{|\Phi\rangle_1, |\Phi\rangle_2\} = \Theta_1^{-1} \Theta_2^{-1} \left[\{\Theta_1, \Theta_2\} |\Phi_1\rangle |\Phi_2\rangle - \{\Theta_1, W_2\} |\Phi_1\rangle - \{W_1, \Theta_2\} |\Phi_2\rangle + \{W_1, W_2\} \right]$$

In this expression, all Poisson brackets can be computed explicitly. Using only eq.(65), we arrive at

$$\{\Phi_m(x), \Phi_n(y)\} = -2 \sum_k \Theta_{mk}^{-1}(x) \Theta_{nk}^{-1}(y) \frac{\sin Z_k}{Z_k \gamma_k} \left(1 - \frac{p_0^2}{Z_k^2} \right) \left[\tilde{F}_k(x) \tilde{G}_k(y) - \tilde{F}_k(y) \tilde{G}_k(x) \right] \quad (88)$$

where γ_k are defined in eq.(34). Multiplying by $E_m(x) E_n(y)$ and remembering that $\langle E | \Theta^{-1} = \langle F |$ we get

$$\{\langle E(x) | \Phi(x) \rangle, \langle E(y) | \Phi(y) \rangle\} = 2 \sum_k \frac{\zeta_k^2 \sin Z_k}{Z_k \gamma_k^3 \mu_k^2} \left(1 - \frac{p_0^2}{Z_k^2} \right) \times \left(\mathcal{A}_k(y) \mathcal{B}_k(x) - \mathcal{A}_k(x) \mathcal{B}_k(y) \right) \quad (89)$$

where

$$\mathcal{A}_k(x) = \tilde{F}_k^2(x), \quad \mathcal{B}_k(x) = \tilde{F}_k(x) \tilde{G}_k(x)$$

Using eqs.(81,82), we can write

$$\begin{aligned} \mathcal{A}_k(x) &= \mu_k \gamma_k w(x, Z) w^*(x, Z) |_{Z=Z_k} \\ \mathcal{B}_k(x) &= \frac{\mu_k \gamma_k}{2i} (w^*(x, Z) \partial_Z w(x, Z) - w(x, Z) \partial_Z w^*(x, Z)) |_{Z=Z_k} \end{aligned}$$

The strategy to evaluate the right hand side of eq.(89) is to rewrite it as a sum over the residues of certain poles of a function on the Riemann Z -sphere. This sum can then be transformed as a sum over the residues of the other poles (there will be none in our case) plus a integral over a small circle at infinity surrounding an essential singularity. This last integral can then be evaluated using the known asymptotics of the function. Let us define

$$a(Z) = \frac{Z}{2i \sin Z} e^{-iZ}, \quad b(Z) = -\frac{Z}{2i \sin Z} \cos Z, \quad c(Z) = \frac{Z}{2i \sin Z} e^{iZ} \quad (90)$$

and introduce the functions

$$\begin{aligned}\Omega_1 &= w(x, Z)w^*(y, Z) - w^*(x, Z)w(y, Z) \\ \Omega_2 &= a(Z)w(x, Z)w(y, Z) \\ &\quad + b(Z)(w(x, Z)w^*(y, Z) + w^*(x, Z)w(y, Z)) \\ &\quad + c(Z)w^*(x, Z)w^*(y, Z)\end{aligned}$$

These functions are defined on the Riemann Z -sphere have poles at the points $\pm m\pi$ and have an essential singularity at infinity. Let

$$\Omega = \Omega_1\Omega_2$$

and we recall the definition of $\eta(Z)$ eq.(39).

Lemma 5

$$\sum_{\pm Z_k} \text{Res} \frac{\Omega}{\eta^2(Z)} \left(1 - \frac{p_0^2}{Z^2}\right) = -2 \sum_k \frac{\zeta_k^2 \sin Z_k}{Z_k \gamma_k^3 \mu_k^2} \left(1 - \frac{p_0^2}{Z_k^2}\right) \times \left(\mathcal{A}_k(y)\mathcal{B}_k(x) - \mathcal{A}_k(x)\mathcal{B}_k(y)\right) \quad (91)$$

Proof. The factor $\eta^2(Z)$ introduces double poles at $\pm Z_k$ because $\eta(Z_k) = 0$. However using eq.(83), we see immediately that $\Omega_1|_{\pm Z_k} = 0$, so that the poles are in fact simple. Remembering eq.(80), we have

$$\sum_{\pm Z_k} \text{Res} \frac{\Omega}{\eta^2(Z)} \left(1 - \frac{p_0^2}{Z^2}\right) = \sum_k \frac{\zeta_k^2}{4Z_k^2} \left(1 - \frac{p_0^2}{Z_k^2}\right) \left(\partial_Z \Omega|_{Z_k} + \partial_Z \Omega|_{-Z_k}\right)$$

We need to compute $\partial_Z \Omega|_{\pm Z_k} = \partial_Z \Omega_1|_{\pm Z_k} \Omega_2|_{\pm Z_k}$. Evaluating at Z_k gives

$$\partial_Z \Omega|_{Z_k} = \frac{2i}{\mu_k^2 \gamma_k^2} \left(a(Z_k) \frac{\alpha_k^+}{\alpha_k^-} + 2b(Z_k) + c(Z_k) \frac{\alpha_k^-}{\alpha_k^+} \right) \left(\mathcal{A}_k(y)\mathcal{B}_k(x) - \mathcal{A}_k(x)\mathcal{B}_k(y) \right)$$

while using $w(-Z_k) = w^*(Z_k)$, $\partial_Z w|_{-Z_k} = -\partial_Z w^*|_{Z_k}$, we also have

$$\partial_Z \Omega|_{-Z_k} = \frac{2i}{\mu_k^2 \gamma_k^2} \left(c(-Z_k) \frac{\alpha_k^+}{\alpha_k^-} + 2b(-Z_k) + a(-Z_k) \frac{\alpha_k^-}{\alpha_k^+} \right) \left(\mathcal{A}_k(y)\mathcal{B}_k(x) - \mathcal{A}_k(x)\mathcal{B}_k(y) \right)$$

The result follows from the identities

$$\begin{aligned}a(Z_k) \frac{\alpha_k^+}{\alpha_k^-} + 2b(Z_k) + c(Z_k) \frac{\alpha_k^-}{\alpha_k^+} &= 2i \frac{Z_k \sin Z_k}{\gamma_k} \\ c(-Z_k) \frac{\alpha_k^+}{\alpha_k^-} + 2b(-Z_k) + a(-Z_k) \frac{\alpha_k^-}{\alpha_k^+} &= 2i \frac{Z_k \sin Z_k}{\gamma_k}\end{aligned}$$

■

Next we have to examine the poles at $\pm m\pi$ in the expression

$$\frac{\Omega}{\eta^2(Z)} \left(1 - \frac{p_0^2}{Z^2}\right)$$

We rewrite Ω_1 and Ω_2 as

$$\begin{aligned}\Omega_1 &= (w(x) - w^*(x))w^*(y) - (w(y) - w^*(y))w^*(x) \\ \Omega_2 &= \frac{Z}{4i \sin Z} \times \left\{ (w^*(x) - w(x))(e^{iZ}w^*(y) - e^{-iZ}w(y)) + \right. \\ &\quad \left. + (w^*(y) - w(y))(e^{iZ}w^*(x) - e^{-iZ}w(x)) \right\}\end{aligned}$$

Recalling the formula eq.(81, 82) for $w(x, Z)$ and $w^*(x, Z)$, we see that when $Z = 0$, we have $w = w^*$ so that $\Omega_1 = 0(Z)$ and $\Omega_2 = 0(Z^2)$. Hence we have no pole at $Z = 0$. When $Z = \pm\pi m + \epsilon$

$$w = \frac{1}{\epsilon}w_{\pm m}^{(-1)} + w_{\pm m}^{(0)} + \dots, \quad w^* = \frac{1}{\epsilon}w_{\pm m}^{*(-1)} + w_{\pm m}^{*(0)} + \dots$$

with

$$w_{\pm m}^{(-1)}(x) = w_{\pm m}^{*(-1)}(x) = \mp\pi m \Phi_m(x)$$

Because the two leading terms are the same, both $w^*(x, Z) - w(x, Z)$ and $e^{iZ}w^*(x, Z) - e^{-iZ}w(x, Z)$ are regular. So Ω_1 and Ω_2 both behaves like $1/\epsilon$. Since $1/\eta^2(Z)$ behaves like ϵ^2 , the whole thing is in fact regular.

We come to the conclusion that everything happens at infinity. We want to compute

$$\left\{ \int dx X(x)T(x), \int dy Y(y)T(y) \right\} = 4 \int_0^1 dx X(x) \int_0^1 dy Y(y) \int_{C_\infty} dZ \left(1 - \frac{p_0^2}{Z^2}\right) \frac{1}{\eta^2(Z)} \partial_x \partial_y \Omega \quad (92)$$

Let

$$\begin{aligned}\tilde{w}(x, Z) &= \eta^{-1/2}(Z) w(x, Z) \\ \tilde{w}^*(x, Z) &= \eta^{-1/2}(Z) w^*(x, Z)\end{aligned}$$

The wronskian of \tilde{w} and \tilde{w}^* is $2iZ$ and therefore these functions coincide with the Baker–Akhiezer functions which are usually introduced in the pseudo-differential approach to the KdV hierarchy (see e.g. [15]). At infinity, we have

$$\begin{aligned}\tilde{w}(x) &= e^{iZx} \left(1 - \frac{\omega(x)}{iZ} + \frac{\omega'(x) + \omega^2(x)}{2(iZ)^2} + \dots\right) \\ \tilde{w}^*(x) &= e^{-iZx} \left(1 + \frac{\omega(x)}{iZ} + \frac{\omega'(x) + \omega^2(x)}{2(iZ)^2} + \dots\right)\end{aligned}$$

where we set

$$\omega(x) = \langle E\Phi \rangle(x)$$

Eq.(73) is needed to verify this formula. Using theses asymptotic forms, we find

$$\begin{aligned}\partial_x (\tilde{w}^2(x)) &= \left(\sum_{n=-\infty}^1 A_n(x)(iZ)^n \right) e^{2iZx} \\ \partial_x (\tilde{w}^{*2}(x)) &= \left(\sum_{n=-\infty}^1 (-1)^n A_n(x)(iZ)^n \right) e^{-2iZx} \\ \partial_x (\tilde{w}(x)\tilde{w}^*(x)) &= \sum_{n=-\infty}^{-1} C_{2n}(x)(iZ)^{2n}\end{aligned}$$

where

$$A_1 = 2, \quad A_0 = -4\omega(x), \quad A_{-1} = 4\omega^2(x), \quad A_{-2} = -\frac{8}{3}\omega^3 - 2 \int^x \omega'^2, \quad A_{-3} = \frac{4}{3}\omega^4 + 4\omega \int^x \omega'^2$$

$$C_{-2}(x) = \omega''(x)$$

Consider the term proportional to $b(Z)$ in eq.(92):

$$4 \int_{C_\infty} \frac{dZ}{2i\pi} b(Z) \left(1 - \frac{p_0^2}{Z^2}\right) \int dx dy (X(x)Y(y) - X(y)Y(x)) \partial_x \tilde{w}^2(x) \partial_y \tilde{w}^{*2}(y)$$

Lemma 6 *Let us define*

$$I_p(z) = \int_{C_\infty} \frac{dZ}{2i\pi} b(Z) (iZ)^p e^{2iZz}$$

We have

$$I_{-1}(z) = -\frac{1}{2}\delta(z), \quad I_0(z) = -\frac{1}{4}\delta'(z), \quad I_1(z) = -\frac{1}{8}\delta''(z), \quad I_2(z) = -\frac{1}{16}\delta'''(z)$$

$$I_{-n}(z) = -\frac{2^{n-3}}{(n-2)!} \epsilon(z) z^{n-2}, \quad n \geq 2$$

Proof. It is clear that

$$\partial_z I_p(z) = 2I_{p+1}(z)$$

hence we can determine all the $I_p(z)$ recursively. For $p \geq 0$ this is done by successively differentiating $I_{-1}(z)$ which is easy to calculate

$$I_{-1}(z) = \int_{C_\infty} \frac{dZ}{2i\pi} b(Z) \frac{1}{iZ} e^{2iZz} = \frac{1}{2} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{\cos Z}{\sin Z} e^{2iZz} = -\frac{1}{2} \sum_{n \in \mathbb{Z}} e^{2in\pi z} = -\frac{1}{2} \delta(z)$$

For $p \leq -2$ we have to successively integrate $I_{-1}(z)$. For this we need boundary conditions which are provided by

$$\int_{C_\infty} \frac{dZ}{2i\pi} b(Z) Z^{-p} = 0, \quad p \geq 2 \quad (93)$$

This is because

$$\begin{aligned} \int_{C_\infty} \frac{dZ}{2i\pi} b(Z) \sum_{p=2}^{\infty} \zeta^p Z^{-p} &= -\frac{\zeta}{2i} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{\cos Z}{\sin Z} \sum_{p=2}^{\infty} \zeta^{p-1} Z^{-p+1} \\ &= -\frac{\zeta^2}{2i} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{\cos Z}{\sin Z} \frac{1}{Z - \zeta} = \frac{\zeta^2}{2i} \left(\cot \zeta - \frac{1}{\zeta} - \sum_{n=0}^{\infty} \frac{2\zeta}{\zeta^2 - n^2\pi^2} \right) = 0 \end{aligned}$$

■

Denoting

$$F_p(x, y) = \sum_{m+n=p} (-1)^m A_n(x) A_m(y)$$

we get

$$4 \int_0^1 dx \int_0^1 dy (X(x)Y(y) - X(y)Y(x)) \sum_{-\infty}^{p=2} F_p(x, y) (I_p(x-y) + p_0^2 I_{p-2}(x-y))$$

In this expression, we separate the terms with a $\delta(x-y)$ function or its derivative which will lead to local terms (L_b) and the non local terms (NL_b) with are proportional to $\epsilon(x-y)$.

$$\begin{aligned} L_b &= 4 \int dx dy (X(x)Y(y) - X(y)Y(x)) (F_2 I_2 + F_1 I_1 + (F_0 + p_0^2 F_2) I_0 + (F_{-1} + p_0^2 F_1) I_{-1}) \\ NL_b &= 4 \int dx dy (X(x)Y(y) - X(y)Y(x)) \sum_{n=0}^{\infty} (F_{-2-n} + p_0^2 F_{-n}) I_{-n-2} \end{aligned} \quad (94)$$

We have

$$\begin{aligned} F_2(x, y) &= -4, \quad F_1(x, y) = 8(\omega(x) - \omega(y)) \\ F_0(x, y) &= -8(\omega(x) - \omega(y))^2, \quad F_{-1}(x, y) = \frac{16}{3}(\omega(x) - \omega(y))^3 + 4 \int_y^x \omega'^2 \end{aligned}$$

The local terms are

$$\begin{aligned} L_b &= 4 \int_0^1 dx \int_0^1 dy (X(x)Y(y) - X(y)Y(x)) \left\{ \frac{1}{4} \delta'''(x-y) + p_0^2 \delta'(x-y) + \right. \\ &\quad \left. - (\omega(x) - \omega(y)) \delta''(x-y) - 2(\omega(x) - \omega(y))^2 \delta'(x-y) - \frac{1}{2} F_{-1}(x, y) \delta(x-y) \right\} \end{aligned}$$

The last two terms obviously vanish and what remains is

$$L_b = \int_0^1 dx \{ -(X'''Y - XY''') - 4((X'Y - XY')(p_0^2 + 2\omega') \}$$

Lemma 7 *The non local term eq.(94) is identically zero.*

Proof. The non local term reads

$$\begin{aligned} NL_b &= -2 \int_0^1 dx \int_0^1 dy (X(x)Y(y) - X(y)Y(x)) \epsilon(x-y) \\ &\quad \left[\sum_{n=0}^{\infty} (F_{-2-n}(x, y) + p_0^2 F_{-n}(x, y)) \frac{2^n}{(n)!} (x-y)^n \right] \end{aligned}$$

The first sum is just the coefficient of $(iZ)^{-2}$ in the formal expansion of $\partial \tilde{w}^2(x) \partial \tilde{w}^{*2}(y)$ while the second sum is the coefficient of $(iZ)^0$. Hence we have

$$NL_b = 2 \int_0^1 dx \int_0^1 dy (X(x)Y(y) - X(y)Y(x)) \epsilon(x-y) \int_{C_\infty} \frac{dZ}{2i\pi} (Z - p_0^2 Z^{-1}) \partial \tilde{w}^2(x) \partial \tilde{w}^{*2}(y)$$

The above expression is zero in the following sense. Let us write

$$\int_{C_\infty} \frac{dZ}{2i\pi} (Z - p_0^2 Z^{-1}) \partial \tilde{w}^2(x) \partial \tilde{w}^{*2}(y) = \sum_{i=1}^{\infty} \frac{(y-x)^i}{i!} \int_{C_\infty} \frac{dZ}{2i\pi} (Z - p_0^2 Z^{-1}) \partial \tilde{w}^2(x) \partial^{i+1} \tilde{w}^{*2}(x)$$

We will show that all the integrals around C_∞ in the right hand side are identically zero.

Since the function \tilde{w} satisfies the Schroedinger equation, its square \tilde{w}^2 satisfies a third order differential equation

$$\mathcal{D}\tilde{w}^2 = -4Z^2\partial\tilde{w}^2, \quad \mathcal{D} = \partial^3 + 8\omega'\partial + 4\omega''$$

Let us introduce a pseudo differential operator Φ such that

$$\tilde{w}^2 = \Phi e^{2iZx}$$

Then

$$\mathcal{D} = \partial\Phi\partial^2\Phi^{-1}$$

Since \mathcal{D} is anti self-adjoint, we also have

$$\mathcal{D} = -\mathcal{D}^* = \Phi^{*-1}\partial^2\Phi^*\partial$$

it follows that \tilde{w}^{*2} which is solution of

$$(-\mathcal{D}^*)\tilde{w}^{*2} = -4Z^2\partial\tilde{w}^{*2}$$

can be written as

$$\tilde{w}^{*2} = \partial^{-1}\Phi^{*-1}\partial e^{-2iZx}$$

Hence

$$\partial\tilde{w}^2 = \partial\Phi e^{2iZx}, \quad \partial^{i+1}\tilde{w}^{*2} = \partial^i\Phi^{*-1}\partial e^{-2iZx}$$

Finally, we have to compute

$$\begin{aligned} \int_{C_\infty} \frac{dZ}{2i\pi} (Z - p_0^2 Z^{-1}) (\partial\tilde{w}^2)\partial^{i+1}\tilde{w}^{*2} &= \int_{C_\infty} \frac{dZ}{2i\pi} (Z - p_0^2 Z^{-1}) (\partial\Phi e^{2iZx})\partial^i\Phi^{*-1}\partial e^{-2iZx} \\ &= \frac{-1}{2i} \int_{C_\infty} \frac{dZ}{2i\pi} (\partial\Phi e^{2iZx})\partial^i\Phi^{*-1}(\partial^2 + 4p_0^2)e^{-2iZx} \end{aligned}$$

We recall the formula (see e.g.[15])

$$\int_{C_\infty} \frac{dZ}{2i\pi} (De^{iZx})(Fe^{-iZx}) = \text{Res}_\partial(DF^*)$$

where Res_∂ is Adler's residue [25]. So our expression is equal to

$$\text{Res}_\partial(\partial\Phi(\partial^2 + 4p_0^2)\Phi^{-1}\partial^i) = \text{Res}_\partial((\mathcal{D} + 4p_0^2\partial)\partial^i) = 0$$

because $(\mathcal{D} + 4p_0^2\partial)\partial^i$ is a differential operator. ■

Consider next the term proportional to $a(Z)$ in eq.(92).

$$4 \int_0^1 dx \int_0^1 dy \left(X(x)Y(y) - X(y)Y(x) \right) \int_{C_\infty} \frac{dZ}{2i\pi} a(Z) \left(1 - \frac{p_0^2}{Z^2} \right) \partial_x \tilde{w}^2(x) \partial_y \left(\tilde{w}(y) \tilde{w}^*(y) \right)$$

Lemma 8 *Let us define*

$$J_p(x) = \int_{C_\infty} \frac{dZ}{2i\pi} a(Z) (iZ)^p e^{2iZx}, \quad p = -1, -2, \dots$$

One has

$$J_{-1}(x) = \frac{1}{2}\delta(x), \quad J_{-p}(x) = 2^{p-2} \frac{x^{p-2}}{(p-2)!} (\epsilon(x) - 1), \quad p \geq 2$$

Proof. One has

$$\partial_x J_p(x) = 2J_{p+1}(x)$$

The calculation of $J_{-1}(x)$ is easy. Next, we need boundary conditions to determine the other J_p by integration

$$\begin{aligned} \sum_{n=2}^{\infty} (i\zeta)^n J_{-n}(0) &= \frac{\zeta^2}{2i} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{e^{-iZ}}{\sin Z} \frac{1}{Z - \zeta} \\ &= \frac{\zeta^2}{2i} \left[-\frac{e^{-i\zeta}}{\sin \zeta} + \sum_{n \in \mathbb{Z}} \frac{1}{\zeta - n\pi} \right] = \frac{\zeta^2}{2i} \left[-\frac{e^{-i\zeta}}{\sin \zeta} + \cot \zeta \right] = \frac{\zeta^2}{2} \end{aligned}$$

It follows that all the non local terms containing $J_p(x)$ for $p \leq -2$ vanish when $0 < x < 1$. The $a(Z)$ term is ■

$$L_a = 4 \int_0^1 dx \int_0^1 dy \left(X(x)Y(y) - X(y)Y(x) \right) A_1(x) C_{-2}(y) J_{-1}(x)$$

or

$$L_a = 4 \int_0^1 dx X(x) \delta(x) \int_0^1 dy Y(y) \omega''(y) - 4 \int_0^1 dx Y(x) \delta(x) \int_0^1 dy X(y) \omega''(y)$$

Finally, the term in $c(Z)$ is just equal to the $a(Z)$ one and double it.

Putting everything together, we arrive at eq.(87).

12 Poisson bracket $\{L_0, u(y)\}$.

In the previous section, we have obtained the correct Poisson bracket for the generators $T_n = L_n - L_0$. We now have to reintroduce L_0 and check that it has the correct Poisson brackets. The candidate for L_0 was given in eq.(59). Let us recall it:

$$L_0 = p_0^2 + \frac{1}{\eta(p_0)} \sum_k \frac{\Gamma_k}{p_0^2 - k^2 \pi^2}$$

where

$$\Gamma_k = E'_k(0) \Phi'_k(0) - E'_k(1) \Phi'_k(1)$$

and

$$\eta(Z) = 1 - \sum_m \frac{\eta_m}{Z^2 - m^2 \pi^2} = 1 - \sum_m \frac{E'_m(0) \Phi_m(0)}{Z^2 - m^2 \pi^2}$$

where we have used eq.(71), evaluated at $x = 0$, to express η_m . We will show that

Proposition 20

$$\left\{ L_0, \int_0^1 dy Y(y) u(y) \right\} = -4 \int_0^1 dy Y(y) \left[(L_0 - p_0^2) \delta'(y) + T'(y) \right] \quad (95)$$

This shows that $\{L_0, \cdot\}$ acts on $u(y)$ as ∂_y as it should be.

Again, the proof is long and we will split it in several lemmas. We need to compute $\{\Phi_m(x), T(y)\}$ and $\{\Phi'_m(x), T(y)\}$ for $x = 0$ and $x = 1$.

Multiplying eq.(88) by $E_n(y)$ and remembering that $\langle E | \Theta^{-1} = \langle F |$ and $F_k(x) = -\frac{\zeta_k}{\gamma_k \mu_k} \tilde{F}_k(x)$, we get

$$\begin{aligned} \{\Phi_m(x), \langle E(y) \Phi(y) \rangle\} &= \quad (96) \\ &= -2\Phi_m(x) \sum_k \frac{\sin Z_k}{Z_k} \frac{\zeta_k^2 (1 - p_0^2 Z_k^{-2})}{\mu_k^2 \gamma_k^3 (Z_k^2 - \pi^2 m^2)} \left[\tilde{F}'_k(x) \tilde{F}_k(x) \tilde{F}_k(y) \tilde{G}_k(y) - \tilde{F}_k^2(y) \tilde{F}'_k(x) \tilde{G}_k(x) \right] \\ &\quad + 2\Phi'_m(x) \sum_k \frac{\sin Z_k}{Z_k} \frac{\zeta_k^2 (1 - p_0^2 Z_k^{-2})}{\mu_k^2 \gamma_k^3 (Z_k^2 - \pi^2 m^2)} \left[\tilde{F}_k^2(x) \tilde{F}_k(y) \tilde{G}_k(y) - \tilde{F}_k^2(y) \tilde{F}_k(x) \tilde{G}_k(x) \right] \end{aligned}$$

As before, this can be expressed in terms of Ω_1 and Ω_2 . We find

$$\{\Phi_m(x), \langle E(y) \Phi(y) \rangle\} = - \sum_k \text{Res}_{\pm Z_k} \frac{(1 - p_0^2 Z^{-2})}{\eta^2(Z) (Z^2 - \pi^2 m^2)} \Omega_1 (\Phi_m(x) \partial_x \Omega_2 - \Phi'_m(x) \Omega_2)$$

We start by computing $\{\eta(p_0), \langle E(y) \Phi(y) \rangle\}$. Setting $x = 0$ in eq.(96), the term in $\Phi'_m(x)$ vanishes because $F_k(0) = 0$. We are left with ($x = 0$, but we keep it for a while)

$$\{\Phi_m(x), \langle E(y) \Phi(y) \rangle\} = -\Phi_m(x) \sum_k \text{Res}_{\pm Z_k} \frac{(1 - p_0^2 Z^{-2})}{\eta^2(Z) (Z^2 - \pi^2 m^2)} \Omega_1 \partial_x \Omega_2 \quad (97)$$

Multiplying by $\frac{E'_m(0)}{(p_0^2 - m^2 \pi^2)}$ and summing over m , we get in the right hand side the sum

$$\begin{aligned} - \sum_m \frac{E'_m(0) \Phi_m(0)}{(p_0^2 - m^2 \pi^2) (Z_k^2 - \pi^2 m^2)} &= - \sum_m \frac{\eta_m}{(p_0^2 - \pi^2 m^2) (Z_k^2 - \pi^2 m^2)} \\ &= \frac{-1}{Z_k^2 - p_0^2} \sum_m \left(\frac{\eta_m}{p_0^2 - m^2 \pi^2} - \frac{\eta_m}{Z_k^2 - \pi^2 m^2} \right) \\ &= \frac{1}{Z_k^2 - p_0^2} \eta(p_0) \end{aligned}$$

where we used that $\sum_m \frac{\eta_m}{Z_k^2 - \pi^2 m^2} = 1$, $\forall k$. The factor $1/(Z_k^2 - p_0^2)$ cancel with the factor $(1 - p_0^2 Z_k^{-2})$ in eq.(97) and we are left with

$$\{\eta(p_0), \langle E \Phi \rangle(y)\} = -\eta(p_0) \sum_k \text{Res}_{\pm Z_k} \frac{1}{\eta^2(Z) Z^2} \Omega_1 \partial_x \Omega_2$$

Finally

$$\left\{ \eta(p_0), \int_0^1 dy Y(y) T(y) \right\} = 2\eta(p_0) \int_0^1 dy Y(y) \int_{C_\infty} \frac{dZ}{2i\pi} \frac{1}{Z^2} \partial_y (\tilde{\Omega}_1 \partial_x \tilde{\Omega}_2) \quad (98)$$

where we used the asymptotics expressions for \tilde{w} and \tilde{w}^* inside Ω_1 and Ω_2 , hence removing the $\eta^2(Z)$ factor.

Lemma 9

$$\left\{ \eta(p_0), \int_0^1 dy Y(y) T(y) \right\} = 4\eta(p_0) \int_0^1 dy Y(y) \delta'(y) \quad (99)$$

Proof. Let us consider the integral over C_∞ in eq.(98). The term containing $b(Z)$ can be written as

$$L_b = \int_{C_\infty} \frac{dZ}{2i\pi} \frac{b(Z)}{Z^2} \times \left\{ (\tilde{w}(x)\tilde{w}^{*'}(x) - \tilde{w}'(x)\tilde{w}^*(x))\partial_y(\tilde{w}(y)\tilde{w}^*(y)) + \right. \\ \left. \tilde{w}(x)\tilde{w}'(x)\partial_y\tilde{w}^{*2}(y) - \tilde{w}^*(x)\tilde{w}^{*'}(x)\partial_y\tilde{w}^2(y) \right\}$$

On the first line, we recognize the wronskian of \tilde{w} and \tilde{w}^* , which is just equal to $-2iZ$. Since $\partial_y(\tilde{w}(y)\tilde{w}^*(y)) = Z^{-2}S_2'(y) + \dots$ this term vanish by eq.(93). Hence

$$L_b = \frac{1}{2} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{b(Z)}{Z^2} \left\{ \partial_x \tilde{w}^2(x) \partial_y \tilde{w}^{*2}(y) - \partial_x \tilde{w}^{*2}(x) \partial_y \tilde{w}^2(y) \right\} \quad (100)$$

$$= - \sum_{p=-2}^{\infty} F_{-p}(x, y) I_{-p-2}(x - y) \\ = \delta'(y - x) + \frac{1}{2} \epsilon(x - y) \int_{C_\infty} \frac{dZ}{2i\pi} \frac{1}{Z} \partial_x \tilde{w}^2(x) \partial_y \tilde{w}^{*2}(y) \quad (101)$$

The $\epsilon(x - y)$ term is zero because

$$\int_{C_\infty} \frac{dZ}{2i\pi} \frac{1}{Z} \partial^{i+1} \tilde{w}^2(x) \partial \tilde{w}^{*2}(x) = \int_{C_\infty} \frac{dZ}{2i\pi} \frac{1}{Z} (\partial^{i+1} \Phi e^{2iZx}) (\Phi^{*-1} \partial e^{-2iZx}) \\ = \int_{C_\infty} \frac{dZ}{2i\pi} (\partial^{i+1} \Phi e^{2iZx}) (\Phi^{*-1} e^{-2iZx}) \\ = \text{Res}_\partial (\partial^{i+1} \Phi \cdot \Phi^{-1}) = 0$$

The $a(Z)$ term reads

$$L_a = \partial_y \int_{C_\infty} \frac{dZ}{2i\pi} \frac{a(Z)}{Z^2} (\tilde{w}(x)\tilde{w}^*(y) - \tilde{w}^*(x)\tilde{w}(y)) \tilde{w}'(x)\tilde{w}(y) \\ = \int_{C_\infty} \frac{dZ}{2i\pi} \frac{a(Z)}{Z^2} \left(\frac{1}{2} \partial_x \tilde{w}^2(x) \partial_y (\tilde{w}(y)\tilde{w}^*(y)) \right. \\ \left. - \frac{1}{2} (\tilde{w}^*(x)\tilde{w}'(x) + \tilde{w}^{*'}(x)\tilde{w}(x)) \partial_y \tilde{w}^2(y) - \frac{1}{2} (\tilde{w}^*(x)\tilde{w}'(x) - \tilde{w}^{*'}(x)\tilde{w}(x)) \partial_y \tilde{w}^2(y) \right)$$

Again, the last term is the wronskian and so

$$L_a = \int_{C_\infty} \frac{dZ}{2i\pi} \frac{a(Z)}{Z^2} \left(-iZ \partial_y (\tilde{w}^2(y)) + \frac{1}{2} \partial_x \tilde{w}^2(x) \partial_y (\tilde{w}(y) \tilde{w}^*(y)) - \frac{1}{2} \partial_x (\tilde{w}^*(x) \tilde{w}(x)) \partial_y (\tilde{w}^2(y)) \right)$$

The first term is

$$\begin{aligned} \int_{C_\infty} \frac{dZ}{2i\pi} a(Z) (iZ)^{-p-1} A_{-p}(y) e^{2iZy} &= \sum_{p=-1}^{\infty} A_{-p}(y) J_{-p-1}(y) \\ &= \frac{1}{2} \delta'(y) - \omega(y) \delta(y) + (\epsilon(y) - 1) \sum_{p=1}^{\infty} A_{-p}(y) \frac{(2y)^{p-1}}{(p-1)!} \end{aligned}$$

The last sum is zero because $0 < y < 1$. The $\delta(y)$ term vanishes because $\omega(0) = 0$. The second term is

$$\frac{1}{2} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{a(Z)}{Z^2} A_{-p}(x) C_{-q}(y) (iZ)^{-p-q} e^{2iZx} = -\frac{1}{2} (\epsilon(x) - 1) \left(A_1(x) C_{-2}(y) (2x) + \dots \right) \quad (102)$$

This vanishes when $x = 0$. The third term is

$$-\frac{1}{2} \int_{C_\infty} \frac{dZ}{2i\pi} \frac{a(Z)}{Z^2} A_{-p}(y) C_{-q}(x) (iZ)^{-p-q} e^{2iZy} = \frac{1}{2} (\epsilon(y) - 1) \left(A_1(y) C_{-2}(x) (2y) + \dots \right)$$

and this vanishes when $0 < y < 1$.

Finally, it is easy to see that the $c(Z)$ term is equal to the $a(Z)$ one. Putting everything together, we get eq.(99). ■

Lemma 10

$$\left\{ \sum_m \frac{E'_m(0) \Phi'_m(0) - E'_m(1) \Phi'_m(1)}{p_0^2 - \pi^2 m^2}, \int_0^1 dy Y(y) T(y) \right\} = -4\eta(p_0) \int_0^1 dy Y(y) T'(y) \quad (103)$$

Proof. Taking the derivative with respect to x of eq.(96) and remembering that $F_k(0) = F_k(1) = 0$, the remaining terms are (there is a cancellation in the $\Phi'_m(x)$ term)

$$\begin{aligned} \{ \Phi'_m(x), \langle E(y) \Phi(y) \rangle \} &= -2\Phi_m(x) \sum_k \frac{\sin Z_k}{Z_k} \frac{\zeta_k^2 (1 - p_0^2 Z_k^{-2})}{\mu_k^2 \gamma_k^3 (Z_k^2 - \pi^2 m^2)} \times \\ &\quad \partial_x \left[\tilde{F}'_k(x) \tilde{F}_k(x) \tilde{F}_k(y) \tilde{G}_k(y) - \tilde{F}_k^2(y) \tilde{F}'_k(x) \tilde{G}_k(x) \right] \end{aligned}$$

where it is understood that $x = 0$ or $x = 1$. By exactly the same argument as before

$$\left\{ \sum_m \frac{E'_m(x) \Phi'_m(x)}{p_0^2 - \pi^2 m^2}, \int_0^1 dy Y(y) T(y) \right\} = 2\eta(p_0) \int_0^1 dy Y(y) \partial_y \partial_x \int_{C_\infty} \frac{dZ}{2i\pi} \frac{1}{Z^2 \eta(Z)} \Omega_1 \partial_x \Omega_2$$

Hence, we just have to take the derivative with respect to x of the previous result, before setting $x = 0$ or $x = 1$. At $x = 0$ we get

$$2\eta(p_0) \int_0^1 dy Y(y) \left[-\delta''(y) + 4\omega''(y) \right]$$

The $\delta''(y)$ term comes from eq.(101) while the second term comes from eq.(102) doubled by the $c(Z)$ contribution. At $x = 1$ only the periodic $\delta''(y)$ remains. Taking the difference we obtain eq.(103). ■

Putting everything together, we arrive at eq.(95).

13 Conclusion.

We have succeeded to take the continuum limit in the formulae expressing the dynamical variables of the Volterra model in terms of the separated variables. This yields exactly solvable potentials and formulae for the Virasoro generators of a rather unusual type. Still, we were able to check that they have the correct Poisson brackets. Of course the most interesting thing now is to try to quantize this approach. As a first step, a semiclassical analysis along the lines of [26] should be very enlightening. The full quantum theory however may reserve some surprise. The bracket eq.(3) being in fact an ordinary quadratic bracket, it is natural to quantize it with Weyl type commutation relations. This opens up the possibility of phenomena as those advocated in [9, 27].

Acknowledgements. This work was supported in part by the European Network ENIGMA, Contract number : MRTN-CT-2004-5652.

References

- [1] J.L.Gervais, *Transport matrices associated with the Virasoro algebra*. Phys.Lett. B160 (1985) 279.
- [2] V. Bazhanov, S. Lukyanov, A. Zamolodchikov *Integrable Structure of Conformal Field Theory, Quantum KdV Theory and Thermodynamic Bethe Ansatz*. hep-th/9412229, Commun.Math.Phys. 177 (1996) 381-398. *Integrable Structure of Conformal Field Theory II. Q-operator and DDV equation*. hep-th/9604044, Commun.Math.Phys. 190 (1997) 247-278. *Integrable Structure of Conformal Field Theory III. The Yang-Baxter Relation*. hep-th/9805008 , Commun.Math.Phys. 200 (1999) 297-324.
- [3] E.K. Sklyanin, *The quantum Toda chain*. Lect. Notes in Phys., (1985) 226, 196-233. *Separation of variables*. Prog. Theor. Phys. (suppl), 185 (1995) p.35.
- [4] M. Kac, P. van Moerbeke, *Some probabilistic aspect of scattering theory*. Proceedings of the Conference on functional integration and its applications, Cumberland lodge (London) 1974. *On some periodic Toda lattices*, Proc. Nat. Acad. Sci., USA, **72**, (4), 1627-1629 (1975). *A complete solution of the periodic Toda problem*, Proc. Nat. Acad. Sci., USA, **72**, (8), 2879-2880 (1975).

- [5] P. van Moerbeke, *The spectrum of Jacobi matrices*, Invent. Math. **37**, 45-81 (1976).
- [6] B.A. Dubrovin, I.M. Krichever, S.P. Novikov, *Integrable Systems I*. Encyclopedia of Mathematical Sciences, Dynamical systems IV. Springer (1990) p.173–281.
- [7] L.D.Faddeev, L.Takhtajan, *Liouville model on the lattice*. Springer Lectures notes in Physics, 246 (1986) 66.
- [8] A.Volkov, *A Hamiltonian interpretation of the Volterra model*. Zapiski.Nauch.Semin. LOMI 150 (1986) 17. *Liouville theory and sh-Gordon model on the lattice*. Zapiski.Nauch.Semin. LOMI 151 (1987) 24. *Miura transformation on the lattice*. Theor.Math.Phys. 74 (1988) 96.
- [9] O.Babelon, *Exchange formula and lattice deformation of the Virasoro algebra*. Physics letters 238B (1990) 234.
- [10] Volkov, A. Yu. *Quantum Volterra Model*. Phys. Lett. A167, 345 (1992), *Noncommutative Hypergeometry*. math.QA/0312084
- [11] Faddeev, L. D., Volkov, A. Yu. *Abelian current algebra and the Virasoro algebra on the lattice*. Phys. Lett. B315, 311318 (1993)
- [12] L. Faddeev, A. Yu. Volkov *Shift Operator for Nonabelian Lattice Current Algebra*. hep-th/9606088, Publ.Res.Inst.Math.Sci.Kyoto 40 (2004) 1113-1125.
- [13] L.D. Faddeev, R.M. Kashaev, A.Yu. Volkov, *Strongly coupled quantum discrete Liouville theory. I: Algebraic approach and duality*. hep-th/0006156, Commun.Math.Phys. 219 (2001) 199-219
- [14] J.L.Gervais, A.Neveu, *Novel triangle relation and absence of tachyon in Liouville string field theory*. Nucl.Phys. B238 (1984) 125. *Oscillator representations of the two-dimensional conformal algebra*. Commun.Math.Phys.100:15,1985
- [15] O.Babelon, D. Bernard, M. Talon, **Introduction to Classical Integrable Systems**. Cambridge University Press (2003).
- [16] F. V. Atkinson, *Multiparameter spectral theory*. Bull. Amer. Math. Soc. 74:1-27 (1968). **Multiparameter Eigenvalue Problems**. Academic, 1972.
- [17] B. Enriquez, V. Rubtsov, *Commuting families in skew fields and quantization of Beauville's fibration*. math.AG/0112276.
- [18] O. Babelon, M. Talon, *Riemann surfaces, separation of variables and classical and quantum integrability*. hep-th/0209071. Phys. Lett. A. 312 (2003) 71-77.
- [19] O. Babelon, *On the Quantum Inverse Problem for the Closed Toda Chain*. hep-th/0304052. J. Phys A. Math. Gen. 37 (2004) pp.303-316.
- [20] S.P. Novikov, *The periodic problem for the Korteweg-de Vries equation*. Funkt. Anal. i Ego Pril. **8** (1974) 54-66. Translation in Funct. Anal. Jan.1975, 236-246.
- [21] B. Dubrovin, S.P. Novikov, *Periodic and conditionally periodic analogues of multisoliton solutions of the Korteweg-de Vries equation*. Dokl. Akad. Nauk. USSR **6** (1974) 2131-2144.

- [22] A. Its, V. Matveev, *On Hill operators with finitely many lacunae*. Funkt. Anal. i ego Pril. **9** (1975).
- [23] H.P. McKean, P. van Moerbeke, *The Spectrum of Hill's Equation*. Inventiones Math. **30** (1975), 217-274.
- [24] V.B. Matveev and M.A. Salle, **Darboux Transformations and Solitons**, Springer-Verlag, (1990).
- [25] M. Adler, *On a trace functional for formal pseudo-differential operators and the symplectic structure of the Korteweg-de-Vries equations*. Inventiones Mathematicae 50 (1979) 219.
- [26] Feodor A. Smirnov *Quasi-classical Study of Form Factors in Finite Volume*. hep-th/9802132. *Dual Baxter equations and quantization of Affine Jacobian*. math-ph/0001032
- [27] L. D. Faddeev, *Discrete Heisenberg-Weyl Group and Modular Group*. hep-th/9504111, Lett.Math.Phys. 34 (1995) 249-254, *Modular Double of Quantum Group*. math.QA/9912078