



Stable directions for small nonlinear Dirac standing waves

Nabile Boussaid

► To cite this version:

| Nabile Boussaid. Stable directions for small nonlinear Dirac standing waves. 2005. hal-00016041v1

HAL Id: hal-00016041

<https://hal.science/hal-00016041v1>

Preprint submitted on 16 Dec 2005 (v1), last revised 27 Apr 2006 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stable directions for small nonlinear Dirac standing waves

Nabile Boussaid

Ceremade, Université Paris Dauphine, Place du Maréchal De Lattre De Tassigny F-75775 Paris Cédex 16 - France
E-mail: boussaid@ceremade.dauphine.fr

December 16, 2005

Abstract: We prove that for a Dirac operator with no resonance at thresholds nor eigenvalue at thresholds the propagator satisfies propagation and dispersive estimates.

When this linear operator has only two simple eigenvalues close enough, we study an associated class of nonlinear Dirac equations which have stationary solutions. As an application of our decay estimates, we show that these solutions have stable directions which are tangent to the subspaces associated with the continuous spectrum of the Dirac operator. This result is the analogue, in the Dirac case, of a theorem by Tsai and Yau about the Schrödinger equation. To our knowledge, the present work is the first mathematical study of the stability problem for a nonlinear Dirac equation.

Introduction

We study the stability of stationary solutions of a time-dependent nonlinear Dirac equation.

Usually, a localized stationary solution of a given time-dependent equation represents the bound state of a particle. Like Ranada [Ran], we call it a *particle like solutions* (PLS). In the literature, the term *soliton* is also found instead of PLS, but this additionally means that the particle keeps its form after a collision. Many works have been devoted to the proof of the existence of such solutions for a large variety of equations. Although their stability is a crucial problem (in particular in numerical computation or experiment), a smaller attention has been deserved to this issue.

There are different definitions of stability. The first one is commonly called *orbital stability*. It means that the orbit of the perturbation of a PLS stays close to the PLS or a manifold of PLS but does not necessarily converge. A stronger notion is *asymptotic stability*, which means that the perturbation of the PLS relaxes asymptotically towards the initial PLS. In practice, it is often replaced by the asymptotic stability of the manifold of PLS.

In fact in many conservative problems asymptotic stability does not hold. But one has asymptotic stability for a restricted class of perturbations, forming the so-called *stable manifold*.

In this paper, we deal with the problem of stability of small PLS of the following nonlinear Dirac equation:

$$i\partial_t\psi = (D_m + V)\psi + \nabla F(\psi). \quad (\text{NLDE})$$

Here, D_m is the usual Dirac operator [Tha92] acting on $L^2(\mathbb{R}^3, \mathbb{C}^4)$

$$D_m = \alpha \cdot (-i\nabla) + m\beta = -i \sum_{k=1}^3 \alpha_k \partial_k + m\beta$$

where $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ and β are \mathbb{C}^4 hermitian matrices satisfying the following properties:

$$\begin{cases} \alpha_i \alpha_k + \alpha_k \alpha_i = 2\delta_{ik} \mathbf{1}_{\mathbb{C}^4}, & i, k \in \{1, 2, 3\} \\ \alpha_i \beta + \beta \alpha_i = \mathbf{0}_{\mathbb{C}^4}, & i \in \{1, 2, 3\} \\ \beta^2 = \mathbf{1}_{\mathbb{C}^4}. \end{cases}$$

In (NLDE), V is the external potential field and F is a nonlinearity such that

$$\forall(\theta, z) \in \mathbb{R} \times \mathbb{C}^4, \quad F(e^{i\theta} z) = F(z).$$

Some additional assumptions on F will be made in the sequel. Stationary solutions (PLS) of (NLDE) take the form $\psi(t, x) = e^{-iEt} \phi(x)$ where ϕ satisfies

$$E\phi = (D_m + V)\phi + \nabla F(\phi). \quad (\text{PLSE})$$

We prove the existence of a manifold of small solutions to (PLSE), interpreted as particle like solutions to (NLDE). Then, we construct a stable manifold around this manifold. At the origin, it is tangent

to the sum of the eigenspace associated with the first eigenvalue and the continuous spectral subspace of $D_m + V$. This is the analogue in the Dirac case of [TY02d, Theorem 1.1, non-resonant case]. The interpretation is that radiations (described by the continuous spectrum) do not destabilize too much the PLS manifold. To prove stabilization towards the PLS manifold, we shall need linear decay estimates associated with the continuous spectral subspace of $D_m + V$.

To our knowledge, this is the first stability result on a nonlinear Dirac equation.

The problem of stability has been extensively studied for Schrödinger and Klein-Gordon equations. The methods used to treat these cases cannot be easily adapted to our problem, due to the fact that the Dirac operator D_m is not bounded-below, contrarily to $-\Delta$. The non-negativity of the latter permits to use minimization and concentration-compactness methods to prove the existence of orbitally stable standing waves, see e.g. Cazenave and Lions [CL82] or more recently Cid and Felmer [CF01].

In his review on nonlinear Dirac models, Ranada [Ran] writes that physicists first claimed that PLS (particle like solutions) of the nonlinear Dirac equation couldn't be stable since the second derivative of the energy functional is not positive-definite. Actually, in a very general setting (not related to the Dirac case), Shatah and Straus [SS85] and Grillakis, Shatah and Straus [GSS87] proved a general orbital stability condition even if the hessian of the energy functional is not positive-definite. Their conditions allow only one simple negative eigenvalue (and a kernel of dimension one also) for the second variation. It therefore cannot be directly applied to the Dirac case. However, it gave rise to an interesting discussion about the application of this method to the Dirac equation in some physical papers [SV86, AS86, BSV87]. Ranada also refers to numerical experiments which seem to confirm that some PLS are asymptotically stable in the Dirac case.

In the Schrödinger case, the asymptotic stability has been extensively studied during the last decade. A fundamental work is the one of Soffer and Weinstein [SW90, SW92], which is devoted to the study of a small nonlinear perturbation of a Schrödinger operator having one simple eigenvalue. They proved that the perturbed small PLS relaxes to a PLS. Later, Pillet and Wayne [PW97] proposed a different proof in the spirit of the central manifold theorem. In all these works, asymptotic stability is a direct consequence of propagation or dispersive estimates on the Schrödinger operator. Hence the considered perturbation is necessarily localized at initial time. To avoid such an assumption, Gustafson, Nakanishi and Tsai [GNT04] proposed to use the Strichartz estimates.

Generalizations have been considered for instance by Tsai and Yau [TY02a, TY02c, TY02d, TY02b, Tsa03], who treated the case of a Schrödinger operator having two simple eigenvalues. An interesting phenomenon appeared: if the two eigenvalues are sufficiently distant, then after linearization around the excited state, one obtains a resonance. Tsai and Yau showed that if there is no resonance, the manifold of ground state has stable directions. In the resonant case, the manifold of ground states is asymptotically stable, whereas the manifold of excited states has stable and unstable directions (in case of instability, under some conditions, one has relaxation to the ground state). Notice that earlier Soffer and Weinstein [SW99] studied a similar resonance phenomenon in the case of the Klein-Gordon equation with a simple eigenvalue; they showed that it induced 'metastability'. Another problem has been studied by Cuccagna [Cuc01, Cuc03, Cuc05]. He considered the case of big PLS, when the linearized operator has only one eigenvalue and obtained the asymptotical stability of the manifold of ground states. Tsai, Yau and Cuccagna also need propagation or dispersive estimates. The latter is proved by generalizing the work of Yajima [Yaj95] on wave operator.

Interesting development are also given by Rodnianski, Schlag and Soffer [RSS05a] who proved asymptotic stability of an arbitrary number of weakly interacting big PLS. Schlag [Sch04] and Krieger and Schlag [KS05] proved the existence of stable direction for unstable big PLS. We point out that some of the works of Schlag [ES04, GS04, RSS05b] or Soffer [HSS99, JSS91, RSS05b] are dedicated to prove dispersive estimate.

We also would like to mention the works of Buslaev and Perelmann [BP95, BP92b, BP92c, BP92a], Buslaev and Sulem [BS03, BS02] or Weder [Wed00], in the one dimensional Schrödinger case.

The paper is organized as follows.

In the next section, we define the important objects and state our main results. We start with the propagation and dispersive linear estimates which will be crucial tools for this study. Then, we consider the nonlinear equation (NLDE) and state the existence of the PLS manifold. Eventually, we present our main theorem in which the stable manifold is constructed.

The third section is devoted to the proof of the propagation estimate, which uses spectral techniques. This is a time decay estimate in weighted L^2 spaces, expressing the fact that states associated with the continuous spectrum are not stationary. We use Mourre estimate similarly to Hunziker, Sigal and Soffer [HSS99] (for a generalization of the method, see e.g. [BdMGS96]). This method cannot be used in the

neighborhood of the thresholds which need a specific treatment. In particular, problems can occur in the presence of eigenvalues at thresholds or resonance, and we shall assume in the whole paper that we are not in this situation. For the Schrödinger case, a similar problem has been studied by Jensen and Kato [JK79], Jensen and Nenciu [JN01, JN04]. Our arguments near the thresholds are inspired of these works. For a related study, see the article of Fournais and Skibsted [FS04] dealing with long range perturbations of Schrödinger operators.

In Section 4, we then prove the dispersive estimate, using the propagation estimate established in Section 3. For an interesting survey on dispersive estimates for Schrödinger operator, see Schlag [Sch05]. It seems that the Dirac equation with a potential behaves like a Klein-Gordon equation with a magnetic potential and we have not been able to generalize the methods used in the Schrödinger case. Instead, we used techniques inspired of works by Cuccagna and Shirmer [CS01].

Finally, the last sections are devoted to the proof of our main result concerning the stability of the stationary solutions of (NLDE). We assume that the Dirac operator $D_m + V$ only has two simple eigenvalues and that it has no eigenvalue at thresholds nor resonance at thresholds. Note that our assumptions exclude electric potentials, for which the eigenvalues are always degenerate, see [Par90, BH92]. In Section 4, this permits us to construct a manifold of PLS and then to study the spectrum of the linearized operator. This in turn, in Section 5, will allow us to decompose a solution of (NLDE) in three parts: the PLS part, the dispersive part associated with the continuous spectrum and a part corresponding to “excited states”. This last part needs a particular treatment since it is not dispersive and hence disturbs the relaxation towards the PLS manifold.

Acknowledgements. I would like to thank Éric Séré for fruitful discussions and advices during the preparation of this work.

1. Main results

This section is devoted to the presentation of the model and the statement of our main results.

1.1. Decay estimates for a Dirac operator with potential. Let us first state our results concerning the time decay of $e^{-it(D_m+V)}$ in weighted L^2 spaces and Besov spaces. This kind of estimates are called respectively propagation and dispersive estimates. As mentioned in the introduction, these results will be very important tools for the study of our nonlinear time-dependent Dirac equation.

The following spaces will be needed to state the main result of this subsection.

Definition 1.1 (Weighted Sobolev space). *The weighted Sobolev spaces are defined by*

$$\mathcal{H}_\sigma^t(\mathbb{R}^3, \mathbb{C}^4) = \{f \in \mathcal{S}'(\mathbb{R}^3), \|\langle Q \rangle^\sigma \langle P \rangle^t f\|_2 < \infty\}$$

for $\sigma, t \in \mathbb{R}$. We endow it with the norm

$$\|f\|_{\mathcal{H}_\sigma^t} = \|\langle Q \rangle^\sigma \langle P \rangle^t f\|_2.$$

If $t = 0$, we write L_σ^2 instead of \mathcal{H}_σ^0 .

We have used the usual notations $\langle u \rangle = \sqrt{1+u^2}$, $P = -i\nabla$, and Q is the operator of multiplication by x in \mathbb{R}^3 . For the sake of clarity, let us also recall the

Definition 1.2 (Besov space). *For $s \in \mathbb{R}$ and $1 \leq p, q \leq \infty$, the Besov space $B_{p,q}^s(\mathbb{R}^3, \mathbb{C}^4)$ is the space of all $f \in \mathcal{S}'(\mathbb{R}^3, \mathbb{C}^4)$ (dual of the Schwartz space) such that*

$$\|f\|_{B_{p,q}^s} = \left(\sum_{j \in \mathbb{N}} 2^{jsq} \|\varphi_j * f\|_p^q \right)^{\frac{1}{q}} < +\infty$$

with $\widehat{\varphi} \in \mathcal{D}(\mathbb{R}^n \setminus \{0\})$ such that $\sum_{j \in \mathbb{Z}} \widehat{\varphi}(2^{-j}\xi) = 1$ for all $\xi \in \mathbb{R}^3 \setminus \{0\}$, $\widehat{\varphi}_j(\xi) = \widehat{\varphi}(2^{-j}\xi)$ for all $j \in \mathbb{N}^*$ and for all $\xi \in \mathbb{R}^3$, and $\widehat{\varphi}_0 = 1 - \sum_{j \in \mathbb{N}^*} \widehat{\varphi}_j$. We endow it with the norm $f \in B_{p,q}^s(\mathbb{R}^3, \mathbb{C}^4) \mapsto \|f\|_{B_{p,q}^s}$.

In the whole paper, we shall work within the following

Assumption 1.1. *The potential $V : \mathbb{R}^3 \mapsto S_4(\mathbb{C})$ (self-adjoint 4×4 matrices) is a C^∞ function such that there exists $\rho > 5$ with*

$$\forall \alpha \in \mathbb{N}^3, \exists C > 0, \forall x \in \mathbb{R}^3, |\partial^\alpha V|(x) \leq \frac{C}{\langle x \rangle^{\rho+|\alpha|}}.$$

Notice that by [Tha92, Theorem 4.2], the operator

$$H := D_m + V$$

is essentially self-adjoint on $C_0^\infty(\mathbb{R}^3 \setminus \{0\}, \mathbb{C}^4)$ and self-adjoint on $H^1(\mathbb{R}^3, \mathbb{C}^4)$. We also work with the

Assumption 1.2. *The operator H presents no resonance at thresholds and no eigenvalue at thresholds.*

A resonance is an eigenvector in $L_{-\sigma}^2(\mathbb{R}^3, \mathbb{C}^4) \setminus L^2(\mathbb{R}^3, \mathbb{C}^4)$ for some $\sigma > 0$. Let

$$\mathbf{P}_c(H) = \mathbf{1}_{(-\infty, -m] \cup [m, +\infty)}(H) \quad (1.1)$$

be the projector associated with the continuous spectrum of H and

$$\mathcal{H}_c = \mathbf{P}_c(H)L^2(\mathbb{R}^3, \mathbb{C}^4).$$

We are now able to state our

Theorem 1.1 (Propagation for perturbed Dirac dynamics). *Assume that Assumption 1.1 and 1.2 hold and let be $\sigma > 5/2$. Then one has*

$$\|e^{-itH}\mathbf{P}_c(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)} \leq C \langle t \rangle^{-3/2}.$$

The proof of this result will be given in Section 2. We notice that it is still true if we assume $\rho > 3$ in Assumption 1.1.

Our next result is the following theorem, proved in Section 3.

Theorem 1.2 (Dispersion for perturbed Dirac dynamics). *Assume that Assumptions 1.1 and 1.2 hold. Then for $\theta \in [0, 1]$, $p \in [1, 2]$ and $s - s' \geq (2 + \theta)(\frac{2}{p} - 1)$ there exists a constant $C > 0$ such that*

$$\|e^{-itH}\mathbf{P}_c(H)\|_{B_{p,q}^s, B_{p',q}^{s'}} \leq CK(t)^{(\frac{2}{p}-1)}$$

with $\frac{1}{p} + \frac{1}{p'} = 1$ and

$$K(t) = \begin{cases} |t|^{-1+\theta/2} & \text{if } |t| \in (0, 1], \\ |t|^{-1-\theta/2} & \text{if } |t| \in [1, \infty). \end{cases}$$

1.2. The stable manifold around the PLS for the nonlinear Dirac equation. We now want to study the following nonlinear Dirac equation

$$\begin{cases} i\partial_t \psi = H\psi + \nabla F(\psi) \\ \psi(0, \cdot) = \psi_0. \end{cases} \quad (1.2)$$

with $\psi \in C^1(I, H^1(\mathbb{R}^3, \mathbb{C}^4))$ for some open interval I which contains 0 and where we recall that $H = D_m + V$. The nonlinearity $F : \mathbb{C}^4 \mapsto \mathbb{R}$ is a differentiable map for the real structure of \mathbb{C}^4 and hence the ∇ symbol has to be understood for the real structure of \mathbb{C}^4 . For the usual hermitian product of \mathbb{C}^4 , one has

$$DF(v)h = \Re \langle \nabla F(v), h \rangle.$$

We work within the following

Assumption 1.3. *The operator H has only two simple eigenvalues $\lambda_0 < \lambda_1$, with ϕ_0 and ϕ_1 as associated normalized eigenvectors. Moreover, the non resonant condition*

$$|\lambda_1 - \lambda_0| < \min\{|\lambda_0 + m|, |\lambda_1 - m|\}$$

holds.

Assumption 1.4. *The function $F : \mathbb{C}^4 \mapsto \mathbb{R}$ is in $\mathcal{C}^\infty(\mathbb{R}^8, \mathbb{R})$, satisfies either $D^\alpha F(z) = 0$ for $|\alpha| = 5$ or $F(z) = O(|z|^5)$ as $z \rightarrow 0$ and has the gauge invariance property:*

$$F(e^{i\theta}z) = F(z), \forall z \in \mathbb{C}^4, \forall \theta \in \mathbb{R}.$$

We will prove in Theorem 1.3 that some solutions of the equation (1.2) are global and can be decomposed as the sum of a PLS plus a remainder part which is vanishing. Since the PLS part may change during the evolution, we need to track it. So we prove that around the origin, PLS form a manifold. We have the

Proposition 1.1 (PLS manifold). *Suppose that Assumptions 1.1–1.4 hold. Then for any $\sigma \in \mathbb{R}^+$, there exists Ω a neighborhood of $0 \in \mathbb{C}$, a \mathcal{C}^∞ map*

$$h : \Omega \mapsto \{\phi_0\}^\perp \cap H^2(\mathbb{R}^3, \mathbb{C}^4) \cap L_\sigma^2(\mathbb{R}^3, \mathbb{C}^4)$$

and a \mathcal{C}^∞ map $E : \Omega \mapsto \mathbb{R}$ such that $S(u) = u\phi_0 + h(u)$ satisfy for all $u \in \Omega$,

$$HS(u) + \nabla F(S(u)) = E(u)S(u), \quad (1.3)$$

with the following properties

$$\begin{cases} h(e^{i\theta}u) = e^{i\theta}h(u), & \forall \theta \in \mathbb{R}, \\ h(u) = O(|u|^2), \\ E(u) = E(|u|), \\ E(u) = \lambda_0 + O(|u|^2). \end{cases}$$

Proof. This kind of results is now classical and left to the reader. For more details, see Subsection 4.1.

We are now able to write the main theorem of this paper. Its proof is given in Section 6.

Theorem 1.3 (Stable manifold). *Suppose that Assumptions 1.1–1.4 hold. Let $s, s', \beta \in \mathbb{R}_+^*$ be such that $s' \geq s + 3 \geq \beta + 6$. There exists $\varepsilon_0 > 0$, $R > 0$, $K > 0$ and a Lipschitz map*

$$\Psi : B_{\mathbb{C}}(0, \varepsilon) \times \left(\mathcal{H}_c \cap B_{H_\sigma^{s'}}(0, R) \right) \mapsto \mathbb{C}$$

with $\Psi(v, 0) = 0$,

$$|\Psi(v, \xi)| \leq K \left(|v| + \|\xi\|_{H_\sigma^{s'}} \right)^2,$$

and such that the following hold. For any initial condition of the form

$$\psi_0 = S(v_0) + \xi_0 + \Psi(v_0, \xi_0)\phi_1$$

with $v_0 \in B_{\mathbb{C}}(0, \varepsilon)$ and $\xi_0 \in \mathcal{H}_c \cap B_{H_\sigma^{s'}}(0, R)$, one has

(i) there exists a unique global solution ψ of (1.2) in

$$\mathcal{C}^k \left(\mathbb{R}, H^{s'}(\mathbb{R}^3, \mathbb{C}^4) \cap H_{-\sigma}^s(\mathbb{R}^3, \mathbb{C}^4) \cap B_{\infty, 2}^\beta(\mathbb{R}^3, \mathbb{C}^4) \right),$$

for any $k \in \mathbb{N}$ with $k \leq s'$;

(ii) there exists $(v_\infty; \xi_\infty; E_\infty) \in B_{\mathbb{C}}(0, \varepsilon) \times B_{H_\sigma^{s'}}(0, R) \times B_{\mathbb{R}}(0, \varepsilon)$ with

$$|v_\infty - v_0| \leq K \|\xi_0\|_{H_\sigma^{s'}}^2, \quad |E_\infty| \leq K \|\xi_0\|_{H_\sigma^{s'}}^2, \quad \|\xi_\infty - \xi_0\|_{H_\sigma^{s'}} \leq K \|\xi_0\|_{H_\sigma^{s'}}^2$$

such that

$$\psi(t) = e^{-i(tE(v_\infty) + E_\infty)} S(u_\infty) + e^{-itD^m} \xi_\infty + \varepsilon(t),$$

where

$$\begin{cases} \|\varepsilon(t)\|_{H^{s'}} \leq K \|\xi_0\|_{H_\sigma^{s'}}^2 \\ \|\varepsilon(t)\|_{H_{-\sigma}^s} \leq \frac{K}{\langle t \rangle^2} \|\xi_0\|_{H_\sigma^{s'}}^2 \\ \|\varepsilon(t)\|_{B_{\infty, 2}^\beta} \leq \frac{K}{\langle t \rangle^2} \|\xi_0\|_{H_\sigma^{s'}}^2. \end{cases}$$

as $t \rightarrow +\infty$.

We notice that the stabilization is faster than the propagation and the dispersion: it is of order $\langle t \rangle^{-2}$ whereas $e^{-itD_m}\xi_\infty$ is of order $\langle t \rangle^{-3/2}$ by Theorems 1.1 and 1.2. This permits to give a physical meaning to the theorem. Indeed, we state the existence of a family of initial states which form a manifold tangent at the origin to the sum of the eigenspace of H associated to λ_0 and the subspace associated with the continuous spectrum of H . This family of initial states gives rise to solution of (1.2) which asymptotically split in two parts. The first one is a PLS: $e^{-i(tE(u_\infty)+E_\infty)}S(u_\infty)$ the other is a free radiation: $e^{-itD_m}\xi_\infty$. Hence if one perturbs a PLS with a radiation then this PLS is moved, proportionally to the energy of the radiation, to another PLS and relaxes by emitting a free radiation.

This phenomenon is due to the propagation and the dispersion property of the subspace associated with the continuous spectrum of H . We don't think that such a phenomenon could take place for perturbation in the direction of the excited states ϕ_1 . Indeed, on this subspace, the dynamic seems to be conservative. The fact that we used propagation and dispersive estimates restricted the family of perturbations to regular and localized ones.

We now turn to the proof of our results.

2. Proof of Theorem 1.1: propagation estimates

Here we prove the propagation estimates of Theorem 1.1. The method used by Jensen and Kato [JK79] to prove this kind of estimates for Schrödinger operator works only for initial state which are spectrally localized near the thresholds $\pm m$. They used the spectral density as the Fourier transform of the propagator. But the Dirac resolvent

$$R_V(\lambda \pm i\varepsilon) = (H - \lambda \mp i\varepsilon)^{-1}$$

does not decay in $B(L_\sigma^2, L_{-\sigma}^2)$ as $|\lambda| \rightarrow +\infty$ for any $\sigma > 0$, see [Yam93]. So we cannot use its Fourier transform. To our knowledge, this method is the only one that permits to treat the problem of propagation for energies near thresholds. Hence with this method, we only prove (in the section 2.1) the

Proposition 2.1 (Propagation near thresholds). *Suppose that Assumptions 1.1 and 1.2 hold and let $\chi \in C_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$ be such that its support is in a sufficiently small neighborhood of $[-m; m]$. Then one has for $\sigma > 5/2$*

$$\|e^{-itH}\mathbf{P}_c(H)\chi(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)} \leq C \langle t \rangle^{-3/2}.$$

We recall that $\mathbf{P}_c(H)$ is defined by (1.1).

We also need to treat the propagation estimates for initial state which spectrum does not contain any threshold. We cannot use the spectral density. So we work directly with the propagator. This is exactly the method used by Hunziker, Sigal and Soffer in [HSS99]. But in our case, their result needs some adaptation. Hence we need to generalize [HSS99, Theorem 1.1] to the case of unbounded energy. In Section 2.2, we prove the

Proposition 2.2 (Propagation far from thresholds). *Suppose that Assumption 1.1 holds. Then for any $\chi \in C^\infty(\mathbb{R}^3, \mathbb{C}^4)$ bounded with support in $(-\infty; -m) \cup (m; +\infty)$ and for any $\sigma \geq 0$, there is $C > 0$ such that*

$$\|e^{-itH}\chi(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)} \leq C \langle t \rangle^{-\sigma}.$$

The proof of Theorem 1.1 is then a consequence of the above propositions

Proof (Proof of Theorem 1.1).

We choose $\chi_0 \in C^\infty(\mathbb{R}^3, \mathbb{C}^4)$ satisfying the assumptions of Proposition 2.1, $\chi_\infty \in C^\infty(\mathbb{R}^3, \mathbb{C}^4)$ satisfying assumptions of Proposition 2.2. Hence the continuous spectrum of H is divided in two parts. We obtain the inequality

$$\|e^{-itH}\mathbf{P}_c(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)} \leq \|e^{-itH}\chi_0(H)\mathbf{P}_c(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)} + \|e^{-itH}\chi_\infty(H)\|_{B(L_\sigma^2, L_{-\sigma}^2)}.$$

Hence from Proposition 2.1, and 2.2, we deduce Theorem 1.1.

It therefore remains to prove Propositions 2.1, and 2.2.

2.1. Step 1 : Propagation near thresholds.

2.1.1. Proof of Proposition 2.1. We now prove Proposition 2.1. Let be $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{C}^4)$, then the operator $e^{-itH} \mathbf{P}_c(H) \chi(H)$ as a function of t is the Fourier transform with respect to λ of

$$\lambda \mapsto \Im R_V^+(\lambda) \mathbf{1}_{(-\infty, -m] \cup [m, \infty)}(\lambda) \chi(\lambda),$$

where

$$R_V^\pm(\lambda) = \lim_{\varepsilon \rightarrow 0^+} R_V(\lambda \pm i\varepsilon), \quad (2.1)$$

we will prove in Section 2.2 that the limit exists in $\mathcal{B}(L_\sigma^2, L_{-\sigma}^2)$. So Proposition 2.1 is a consequence of the

Proposition 2.3. *Suppose that Assumptions 1.1 and 1.2 hold. Then for $\lambda > m$ close enough to m , one has*

$$R_V^\pm(\lambda) = \lim_{\varepsilon \rightarrow 0^+} R_V(\lambda \pm i\varepsilon)$$

exists in $\mathcal{B}(\mathcal{H}_\sigma^{-1/2}, \mathcal{H}_{-\sigma}^{1/2})$ for $\sigma > 3/2$. It is \mathcal{C}^l if $\sigma > 1/2 + l$ and $0 < l \leq 2$ with

$$\frac{d^l}{d\lambda^l} \Im R_V^\pm(\lambda) = O(\sqrt{\lambda - m}^{1/2-l}), \quad (2.2)$$

as $\lambda \rightarrow m^+$.

The same holds for $\lambda < -m$ if m is replaced by $-m$.

We prove it in Section 2.1.2. The idea is then to apply to

$$\lambda \mapsto \Im R_V^+(\lambda) \mathbf{1}_{(-\infty, -m] \cup [m, \infty)}(\lambda) \chi(\lambda). \quad (2.3)$$

with $k = 1$ and $\theta = 1/2$, the following

Lemma 2.1 (Lemma 10.2 of [JK79]). *Suppose $F(\lambda) = 0$ for $\lambda > a > 0$, $F^{(k+1)}(\lambda) \in L^1([\delta, +\infty[)$ for any $\delta > 0$ and an integer $k \geq 0$ and that $F^{k+1} = O(\lambda^{\theta-2})$ near 0 for some $\theta \in]0, 1[$. Assume further that $F^{(j)}(0) = 0$ for $j \leq k - 1$, then*

$$\widehat{F}(t) = O(t^{-k-\theta}).$$

The symbol O may be replaced by o throughout.

We refer to [JK79] for the proof of Lemma 2.1. In fact to apply this lemma to (2.3), one should split this function in two parts, one supported in \mathbb{R}^+ and the other in \mathbb{R}^- . Then one translates the first one by $-m$ and applies the lemma. To deal with the other part, one works exactly in the same way after a symmetry with respect to the origin. To end the proof of Proposition 2.1, it remains to prove Proposition 2.3. This the goal of the next section.

2.1.2. Behavior near thresholds of the Dirac resolvent: proof of Proposition 2.3. In this section, our aim is to prove Proposition 2.3. First of all, we notice that if the limits (2.1) exist then we have

$$R_V^-(\lambda)^* = R_V^+(\lambda),$$

and since

$$\alpha_5(D_m + V - z)^{-1} \alpha_5 = -(D_m + \alpha_5 V \alpha_5 + z)^{-1},$$

for $\alpha_5 = \prod_{i=1}^3 \alpha_i \beta$, one obtains

$$\alpha_5 R_V^\pm(\lambda)^{-1} \alpha_5 = -R_{\alpha_5 V \alpha_5}^\mp(-\lambda).$$

So we only need to study the behavior of $R^+(\lambda)$ near $+m$. Let us introduce

$$\mathbb{C}_{++} = \{z \in \mathbb{C}, \Im z > 0, \Re z > 0\}$$

then the behavior for the free case ($V = 0$) is given by the

Proposition 2.4 (Dirac's resolvent expansion). *Let be $s, s' > 1/2$ with $s + s' > 2$ and $t \in \mathbb{R}$. Then $R_0(z) \in \mathcal{B}(\mathcal{H}_{s,2}^{t-1}, \mathcal{H}_{-s',2}^t)$ is uniformly continuous in \mathbb{C}_{++} and so it can be continuously extended to $\overline{\mathbb{C}_{++}}$.*

Moreover, the formal series $z \in \mathbb{C}_{++}$,

$$R_0(z) = \sum_{j=0}^{\infty} (i\sqrt{z^2 - m^2})^j D_m G_j + \sum_{j=0}^{\infty} z(i\sqrt{z^2 - m^2})^j G_j \text{ with } \Im(\sqrt{z^2 - m^2}) > 0,$$

is an asymptotic expansion for $z \rightarrow m$ in the following sense:

Let $k \in \mathbb{N}$, if $R_0(z)$ is approximated by the correspondent finite series up to $j = k$, the remainder is $o(|z - m|^{k/2})$ in the norm of $\mathcal{B}(\mathcal{H}_{s,2}^{t-1}, \mathcal{H}_{-s',2}^t)$ with $s, s' > k + 1/2$ (and $s + s' > 2$ if $k = 0$) and $t \in \mathbb{R}$. In the same sense, this identity can be differentiated in z any number of times. More precisely, for $l \in \mathbb{N}^$ the l^{th} derivative in z of the said finite series is equal to $\frac{d^l}{dz^l} R(z)$ up to an error $o(|z - m|^{k/2-l})$ in the norm of $\mathcal{B}(\mathcal{H}_{s,2}^{t-1}, \mathcal{H}_{-s',2}^t)$ with $s, s' > k + l + 1/2$ and $t \in \mathbb{R}$.*

Proof. It is an adaptation of lemmas of [JK79]. We rewrite [JK79, Lemma 2.1], [JK79, Lemma 2.2] and [JK79, Lemma 2.3] in the Dirac case with help of the identity

$$(D_m - z)^{-1}(D_m + z)^{-1} = (-\Delta + m^2 - z^2)^{-1},$$

or in $(\mathbb{C}^2)^2$

$$(D_m - z)^{-1} = \begin{pmatrix} \frac{z + m}{-\Delta - z^2 + m^2} & \frac{\sigma \cdot \nabla}{-\Delta - z^2 + m^2} \\ \frac{\sigma \cdot \nabla}{-\Delta - z^2 + m^2} & \frac{z - m}{-\Delta - z^2 + m^2} \end{pmatrix}$$

where σ are the two dimensional Pauli matrices.

To obtain the behavior of the Dirac resolvent in the general case, we would like to use the formula

$$R_V(z) = M(z)^{-1} R_0(z) \quad (2.4)$$

with

$$M(z) = (1 + R_0(z)V).$$

To give a meaning to Identity (2.4), we have to prove that $M(z)$ is invertible in $B(\mathcal{H}_{-\sigma}^{1/2})$ for $\sigma > 1/2$ with $\sigma + 1/2 < \rho$. We also give the asymptotic behavior of $R_V^+(z)$ and some of its derivatives as $\lambda \rightarrow m^+$. By means of Proposition 2.4, one has

$$z \in \overline{\mathbb{C}_{++}} \mapsto M(z) \in B(\mathcal{H}_{-\sigma}^{1/2})$$

is uniformly continuous for $1/2 < \sigma$ and $2 < \sigma + \sigma' \leq \rho$ and some $\sigma' > 1/2$. Hence we have

$$M(z) = M(m) + A(z),$$

with $A(z)$ uniformly continuous in $B(\mathcal{H}_{-\sigma'}^{1/2}, \mathcal{H}_{-\sigma}^{1/2})$ near m in \mathbb{C}_{++} for $1/2 < \sigma$ and $2 < \sigma + \sigma' \leq \rho$ and some $\sigma' > 1/2$ and tending to 0 as $\lambda \rightarrow m$. We now prove the

Lemma 2.2 (Threshold's eigenvector and resonance). *Suppose that Assumption 1.1 holds. Let be $\mathcal{M}(s)$ the kernel of $M(m)$ in $\mathcal{H}_{-s}^{1/2}$ and $\mathcal{K}(s)$ the kernel of $(H - m)$ in $\mathcal{H}_{-s}^{1/2}$. Then $\mathcal{M}(s)$ and $\mathcal{K}(s)$ are finite dimensional and do not depend on $s \in (1/2, \rho - 1/2)$. So we write \mathcal{M} and \mathcal{K} and we have*

$$\mathcal{M} = \mathcal{K}$$

Proof. Let be $u \in \mathcal{K}(s)$, then $(D_m + V - m)u = 0$ and $u \in \mathcal{H}_{-s}^{1/2}$, so $Vu \in \mathcal{H}_{\rho-s}^{-1/2}$ and since $\rho - s > 1/2$, $s > 1/2$, and $s + \rho - s > 2$, we obtain, by Proposition 2.4, $(D_m - m)^{-1}Vu \in \mathcal{H}_{-s}^{1/2}$. For any $\phi \in \mathcal{C}_0^\infty$,

$$\langle \phi, (D_m - m)(D_m - m)^{-1}Vu \rangle = \langle (D_m - m)(D_m - m)^{-1}\phi, Vu \rangle = \langle \phi, Vu \rangle,$$

we obtain $(D_m - m)(D_m - m)^{-1}Vu = Vu$ and $(D_m - m)(u + (D_m - m)^{-1}Vu) = 0$. Since $D_m - m$ has no kernel in $\mathcal{H}_{-s}^{1/2}$, because there's no harmonic function in L^2_{-s} , we obtain $u + (D_m - m)^{-1}Vu = 0$. Hence, we have $\mathcal{K}(s) \subset \mathcal{M}(s)$.

Conversely $I + (D_m - m)^{-1} V$ defines a Fredholm operator of $\mathcal{B}(\mathcal{H}_{-s}^{1/2})$. If $u \in \mathcal{M}(s)$ then $u \in \mathcal{H}_{-s}^{1/2}$ and $(D_m - m)^{-1} V u \in \mathcal{H}_{-s}^{1/2}$. So we write $0 = (D_m - m)(u + (D_m - m)^{-1} V u) = (D_m - m + V)u$ and we obtain $\mathcal{K}(s) = \mathcal{M}(s)$.

Now we introduce $I + V(D_m - m)^{-1} \in \mathcal{B}(\mathcal{H}_s^{-1/2})$, and its kernel $\mathcal{N}(s)$ which is finite dimensional is a Fredholm operator. We have that $\mathcal{N}(s)$ is decreasing with s and $\mathcal{M}(s)$ is increasing. Since, by duality, $\dim \mathcal{M}(s) = \dim \mathcal{N}(s)$, we deduce that $\mathcal{N}(s)$ and $\mathcal{K}(s) = \mathcal{M}(s)$ do not depend on s .

We are now able to conclude the proof of Proposition 2.3.

Proof (Proof of Proposition 2.3). Assumption 1.2 gives $\mathcal{K} = 0$ and so with Lemma 2.2, one obtains $\mathcal{M} = 0$. Hence $M(m)$ is invertible since it is a Fredholm operator. We use Von Neumann series to obtain that $M(z)$ is invertible in $B(\mathcal{H}_{-\sigma}^{1/2})$ for $\sigma > 1/2$, $2 < \sigma + \sigma' \leq \rho$ and some $\sigma' > 1/2$ and

$$M(z)^{-1} = M(m)^{-1} \sum_{j \geq 0} (A(z)M(m)^{-1})^j.$$

So for $\lambda \geq m$ close enough to m ,

$$M^+(\lambda)^{-1} = \lim_{\varepsilon \rightarrow 0^+} M(\lambda + i\varepsilon)^{-1}$$

exists in $B(\mathcal{H}_{-\sigma}^{1/2})$ for $\sigma > 1/2$ with $2 < \sigma + \sigma' \leq \rho$ and some $\sigma' > 1/2$. We obtain that $\lim_{\varepsilon \rightarrow 0^+} R_V(\lambda + i\varepsilon)$ exists in $B(\mathcal{H}_{\sigma''}^{-1/2}, \mathcal{H}_{-\sigma}^{1/2})$ for $\sigma > 1/2$ and $\sigma \geq \sigma'' > 1/2$ with $\sigma + \sigma'' > 2$, $2 < \sigma + \sigma' \leq \rho$ and some $\sigma' > 1/2$.

Using Proposition 2.4, we prove that if $1/2 + k < \sigma$, and $\sigma' + 1/2 + k < \rho$ then

$$\frac{d^k}{d\lambda^k} M^+(\lambda) = O(\sqrt{\lambda - m}^{1/2-k})$$

in $B(\mathcal{H}_{-\sigma'}^{-1/2}, \mathcal{H}_{-\sigma}^{1/2})$ for $k \in \mathbb{N}^*$ as $\lambda \rightarrow m^+$. Since we have

$$\frac{d}{d\lambda} F(\lambda)^{-1} = -F(\lambda)^{-1} \left(\frac{d}{d\lambda} F(\lambda) \right) F(\lambda)^{-1},$$

for matrix valued differentiable function F with invertible values, we obtain for $k \in \mathbb{N}^*$ the estimate

$$\frac{d^k}{d\lambda^k} M^+(\lambda)^{-1} = O(\sqrt{\lambda - m}^{1/2-k}),$$

in $B(\mathcal{H}_{-\sigma}^{1/2})$ with $1/2 + k < \sigma$ and $\sigma + 1/2 + k < \rho$ as $\lambda \rightarrow m^+$. So by Leibniz formula, we also have for $k \in \mathbb{N}^*$

$$\frac{d^k}{d\lambda^k} R_V^+(\lambda) = O(\sqrt{\lambda - m}^{1/2-k}),$$

in $B(\mathcal{H}_{\sigma'}^{-1/2}, \mathcal{H}_{-\sigma}^{1/2})$ with $1/2 + j < \sigma$, $1/2 + k - j < \sigma'$ and $1/2 + k - j + \sigma < \rho$ for all $j \in \{0, \dots, k\}$ as $\lambda \rightarrow m^+$. For the case $k = 0$, we have the formula

$$R_V(z) = R_0(z) (1 + V R_0(z))^{-1}.$$

Since $R^+(m) = R^-(m)$, this leads to

$$\Im R_V^+(m) = 0,$$

and so

$$\Im R_V^\pm(\lambda) = O(\sqrt{\lambda - m}^{1/2}),$$

as $\lambda \rightarrow m^+$ in $B(\mathcal{H}_{\sigma}^{-1/2}, \mathcal{H}_{-\sigma}^{1/2})$ with $3/2 < \sigma$ and $\sigma + 3/2 < \rho$. Hence (2.2) is proved.

2.2. Step 2: Propagation Far from thresholds. In this section, we give the proof of Proposition 2.2. We prove the propositions for $t \geq 0$. Then using $(e^{-itH})^* = e^{itH}$, the result easily follows for $t \leq 0$.

2.2.1. *Proof of Proposition 2.2.* Let us introduce

$$A = \frac{1}{2} \{D_m^{-1}P \cdot Q + Q \cdot PD_m^{-1}\}.$$

[IM99, Lemma 3.1] gives that A is an essentially self-adjoint operator and the domain of its closure contains the domain of $\langle Q \rangle$. Proposition 2.2 is then a consequence of the

Theorem 2.1 (Minimal escape velocity). *Suppose that Assumption 1.1 holds. Then for any $\chi \in C_0^\infty$ bounded with support in $(-\infty, -m) \cup (m, +\infty)$, there exists $\theta > 0$ such that for any $l \in \mathbb{R}$, for any $v \in (0, \theta)$, and any $a \in \mathbb{R}$ one has*

$$\forall t > 0, \left\| \mathbb{1}_{A-a-vt \leq 0} e^{-itH} \chi(H) \mathbb{1}_{A-a \geq 0} \right\| \leq Ct^{-l}.$$

The proof will be given in Section 2.2.2. Let us now show that Theorem 2.1 implies Proposition 2.2.

Proof (Proof of Proposition 2.2). We notice that

$$\langle A \rangle^{-\alpha} = \langle A \rangle^{-\alpha} \mathbb{1}_{\pm A \geq ct} + O(t^{-\alpha}),$$

this leads to

$$\langle A \rangle^{-\alpha} e^{-itH} \chi(H) \langle A \rangle^{-\alpha} = \langle A \rangle^{-\alpha} \mathbb{1}_{A \leq \frac{(\theta-\varepsilon)t}{2}} e^{-itH} \chi(H) \mathbb{1}_{A \geq \frac{\theta t}{2}} \langle A \rangle^{-\alpha} + O(t^{-\alpha}),$$

So if we choose $a = -\frac{\theta t}{2}$ and $v = \theta - \frac{\varepsilon}{2}$ in Theorem 2.1, we obtain

$$\|\langle A \rangle^{-\alpha} e^{-itH} \chi(H) \langle A \rangle^{-\alpha}\| \leq Ct^{-\min(\alpha, l)}.$$

Then we prove that $\langle A \rangle^\alpha \langle Q \rangle^{-\alpha}$ is bounded for any positive α . It is quite immediate for integer α using multi-commutator expansion [HS00, Identity (B.24)]. To prove it for any positive real, we use [SS98, Identity (1.2)]. This identity states that for a self adjoint with $B \geq 1$ and a positive real β , we have on domain of $B^{[\alpha]+1}$

$$B^\beta = \frac{\sin(\pi\{\beta\})}{\pi} \int_0^{+\infty} \frac{w^{\{\beta\}-1}}{1+wB^{-1}} dw B^{[\beta]},$$

where $\{\beta\} = \beta - [\beta]$ and $[\beta]$ is the integer part. With this formula for $B = \langle A \rangle^{2k}$ for any $k \in \mathbb{N}$, we prove for any $\beta \in]0, 1[$ that

$$\langle A \rangle^{2k\beta} \leq C \langle Q \rangle^{2k\beta}$$

which ends the proof of Proposition 2.2.

2.2.2. *Proof of Theorem 2.1.* Our proof of Theorem 2.1 is an adaptation of the one of [HSS99], we make some modification.

For any self-adjoint operator B with domain $D(B)$, we write $Ad_A(B)$ for the operator $[A, B]$ with domain $D(A) \cap D(B)$ dense in $D(B)$, defined by

$$\forall u, v \in D(A) \cap D(B), \langle i[A, B]u, v \rangle = i(\langle Bu, Av \rangle - \langle Au, Bv \rangle).$$

First of all, we have

Lemma 2.3. *Suppose that Assumption 1.1 holds. Then $Ad_A^k(H)$ is bounded and can be written as a finite sum of terms of the form*

$$f(P)g(Q)h(P)$$

where f and h are rational fractions with coefficients in $\mathcal{M}_4(\mathbb{C})$ of degree at most 0 with no poles, and g is a function that satisfies Assumption 1.1.

Proof. The proof is a simple calculation based on the fact that $Ad_{P_j}(f(Q)) = -i(\nabla_j f)(Q)$.

Let us introduce the wave operators associated with the couple (D_m, H) :

$$W^\pm = \lim_{t \rightarrow \pm\infty} e^{itH} e^{-itD_m}.$$

By [GM01, Theorem 1.5], these operators exist and their range is given by $\mathbf{P}_c(H)$. Moreover, they are unitary and

$$(W^\pm)^* = \lim_{t \rightarrow \pm\infty} e^{itD_m} e^{-itH} \mathbf{P}_c(H),$$

which also exists, by [RS79, Proposition XI.3]. Since for $W_t = e^{itH} e^{-itD_m}$, $\partial_t W_t f$ also converges for $f \in H^1(\mathbb{R}^3, \mathbb{C}^4)$, we deduce

$$HW^\pm = W^\pm D_m$$

and hence the intertwining property:

$$f(H)W^\pm = W^\pm f(D_m) \quad (2.5)$$

for any bounded borelian function f . Then we introduce

$$A_V = W^+ A (W^+)^*$$

and we now obtain the

Lemma 2.4 (Mourre estimate). *If V tends to 0 at infinity then for any $\nu \geq 0$, one has*

$$\mathbb{1}_{|H| \geq m+\nu} i[H, A_V] \mathbb{1}_{|H| \geq m+\nu} \geq \frac{\nu^2}{m^2 + \nu^2} \mathbb{1}_{|H| \geq m+\nu}.$$

Proof. This is a consequence of

$$i[D_m, A] = \frac{-\Delta}{-\Delta + m^2}.$$

We now adapt [HSS99, Theorem 1.1] to the case of unbounded energy since here the multi-commutators $Ad_A^k(H)$ are bounded operators. We introduce the

Definition 2.1. *We call generalized indicator function of \mathbb{R}^- a function of the form*

$$x \mapsto e^{\frac{u(x)}{x}} \mathbb{1}_{\mathbb{R}^-}(x)$$

with $u \in C_0^\infty(\mathbb{R})$ supported in $[-\eta, \eta]$ (for some $\eta > 0$), nonnegative, and such that $u(0) = 1$.

Note that our generalized indicator function of \mathbb{R}^- are of infinite order in the sense of [HSS99, Section 2]. Using the commutators expansion presented in [HS00, Section B] and the Mourre estimate of Lemma 2.4, we have the

Lemma 2.5. *Assume V satisfies Assumption 1.1 and that f is a generalized indicator function of \mathbb{R}^- . Let be $A_s = s^{-1}\{A - a\}$ and $0 < \varepsilon \leq 1$. Then for any $n \in \mathbb{N}$ and $\delta > 0$ there exists a $C > 0$ independent of $a \in \mathbb{R}$ such that for $s \geq 1$*

$$g(H) i[H, f(A_s)] g(H) \leq s^{-1} \theta g(H) f'(A_s) g(H) + C s^{-1-\varepsilon} g(H) f^{1-\delta}(A_s) g(H) + C s^{-(2n-1-\varepsilon)} g^2(H).$$

Proof. See [HSS99, Lemma 2.1], in our case we don't need to replace H by $b(H)H$ with $b \in C_0^\infty$. Indeed, our commutators $Ad_A^k(H)$ are bounded by means of Lemma 2.3. Then we replace the notion of function of order p by the one of generalized indicator function. Finally, we use the fact that a generalized indicator function f satisfies

$$\forall k \in \mathbb{N}, \forall \delta \in (0, 1), \exists C > 0, \quad |f^{(k)}| \leq C |f|^{1-\delta}.$$

We are now able to give the

Proof (Proof of Theorem 2.1). We notice that if $0 < v < \theta - \eta$ and if F is a positive non increasing \mathcal{C}^∞ -function which equals 0 on \mathbb{R}^+ and 1 on $(-\infty, -\eta)$, we have

$$\mathbb{1}_{(A-a-vs)<0} \leq F\left(\frac{A-a}{s} - \theta\right).$$

Now suppose F is a generalized indicator function of R^- . We consider

$$F(s^{-1}\{A-a-\theta t\})$$

and study the time evolution of the observable $g(H)f(A_{ts})g(H)$, where $f = F^2$, with respect to $e^{-itH}\mathbb{1}_{A-a>0}$. That is to say we study

$$\langle e^{-itH}\mathbb{1}_{A-a>0}\psi, g(H)f(A_{ts})g(H)e^{-itH}\mathbb{1}_{A-a>0}\psi \rangle.$$

We work exactly as in the proof of [HSS99, Theorem 1.1]. Hence using Lemma 2.5 we obtain for $0 \leq t \leq s$ and $s > 1$

$$\begin{aligned} \langle e^{-itH}\mathbb{1}_{A-a>0}\psi, g(H)f(A_{ts})g(H)e^{-itH}\mathbb{1}_{A-a>0}\psi \rangle &\leq Cs^{-(2n-2-\varepsilon)}\|\psi\|^2 \\ &\quad + Cs^{-1-\varepsilon} \int_0^t \langle e^{-itH}\mathbb{1}_{A-a>0}\psi, g(H)f(A_{ts})g(H)e^{-itH}\mathbb{1}_{A-a>0}\psi \rangle^{1-\delta} \|\psi\|^{2\delta}. \end{aligned}$$

Then using the Gronwall's lemma (see [ABdMG96, Lemma 7.A.1]), we obtain

$$\langle e^{-itH}\mathbb{1}_{A-a>0}\psi, g(H)f(A_{ts})g(H)e^{-itH}\mathbb{1}_{A-a>0}\psi \rangle \leq \left\{ Cs^{-\delta(2n-2-\varepsilon)}\|\psi\|^{2\delta} + \delta Cs^{-\varepsilon}\|\psi\|^{2\delta} \right\}^{1/\delta},$$

so if we choose a small δ and a big n , the proof is done if we choose $s = \max\{1, t\}$.

3. Proof of Theorem 1.2: dispersive estimates

Dispersive estimates for Schrödinger operators with electric potentials take place in Lebesgue spaces. This fact permits to use simple perturbation methods (like Duhamel's formula) to prove the decay estimates for perturbed Schrödinger equations. Unfortunately, we have only been able to prove dispersive estimates for Dirac operators in Besov spaces, so it was not possible for us to use Duhamel's formula or other perturbation method used for Schrödinger operators.

We notice that in the case of a Dirac operator with scalar potential (matrix valued function colinear with β), the square of the Dirac equation gives four coupled Klein-Gordon equations with an electrostatic potential. This permits to use results on the Klein-Gordon equation. For example, Yajima [Yaj95] proved dispersive estimates for the Klein-Gordon equation by using wave operators associated with Schrödinger operators including an electrostatic potential. But in the general case, by taking the square of a Dirac operator with a potential, we obtain also a magnetic potential. Hence the method used by Yajima does not work in our case.

We have not been able to find any reference mentioning dispersive estimates for the free Dirac equation. As we said, it could be deduced from dispersive estimates of Klein-Gordon equation. Here, to give a sketch of the proof for the general case, we first prove the free case estimates (see Section 3.2), using estimates on oscillatory integrals of Section 3.1. In Section 3.3, following Cuccagna and Schirmer [CS01], we introduce the distorted plane waves. This permits us to tackle the proof of the general case in Section 3.3.2.

3.1. Estimates on some oscillatory integrals. Here, we state some stationary phase type results which will be useful for the rest of the proof. We denote by S^2 the unit sphere of \mathbb{R}^3 .

Lemma 3.1. *Let be $l \in \mathbb{N}$ and $f \in \mathcal{C}^1(S^2)$ and for any $v \in S^2$ and any $k \in \mathbb{R}$ define*

$$J_v(k) = \int_{S^2} e^{ik\{1-v \cdot \omega\}} f(\omega) d\omega.$$

Then we have

$$|J_v(k)| \leq \frac{C}{\langle k \rangle} \left\{ \sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^\alpha f(\omega)|}{|\omega - v|^{|\alpha|}} d\omega + \sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^\alpha f(\omega)|}{|\omega + v|^{|\alpha|}} d\omega \right\}, \quad (3.1)$$

where C does not depend on f , k or v .

If f is in $\mathcal{C}^2(S^2)$ with $f(v) = f(-v) = 0$, we have

$$|J_v(k)| \leq \frac{C}{\langle k \rangle^{3/2}} \left\{ \sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^{2\alpha} f(\omega)|}{|\omega - v|^{|\alpha|}} d\omega + \sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^{2\alpha} f(\omega)|}{|\omega + v|^{|\alpha|}} d\omega \right\}, \quad (3.2)$$

where C does not depend on f , k or v .

If f is in $\mathcal{C}^2(S^2)$ and vanishes in a neighborhood of v and $-v$, we have

$$|J_v(k)| \leq \frac{C}{\langle k \rangle^2} \left\{ \frac{\sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^{2\alpha} f(\omega)|}{|\omega - v|^{|\alpha|}} d\omega}{\text{dist}(\text{supp}(f), v)} + \frac{\sum_{|\alpha| \leq 1} \int_{S^2} \frac{|\nabla^{2\alpha} f(\omega)|}{|\omega + v|^{|\alpha|}} d\omega}{\text{dist}(\text{supp}(f), -v)} \right\}, \quad (3.3)$$

where C does not depend on f , k or v .

Proof. We can suppose $v = (0, 0, 1)$ since estimate (3.1), (3.2) and (3.3) are invariant under the action of rotations. We have

$$J_v(k) = \int_0^{2\pi} \int_0^\pi e^{ik\{1-\cos(\phi)\}} f(\theta, \phi) \sin(\phi) d\phi d\theta,$$

then we make an integration by parts in ϕ

$$J_v(k) = -\frac{i}{k} \int_0^{2\pi} \left[e^{ik\{1-\cos(\phi)\}} f(\theta, \phi) \right]_0^\pi d\theta + \frac{i}{k} \int_0^{2\pi} \int_0^\pi e^{ik\{1-\cos(\phi)\}} \partial_\phi f(\theta, \phi) d\phi d\theta,$$

If we suppose that f vanishes in a neighborhood of v or $-v$, then we use

$$|f(\theta, \phi)| \leq \int_0^\pi |\partial_\phi f(\theta, \psi)| d\psi$$

to obtain (3.1) in this case. Otherwise with help of a smooth cut-off, we split the integral in two parts, each one has a support far from v or $-v$. Repeating the previous proof for each part, we prove the estimate (3.1) in the general case.

If moreover we have $f(v) = f(-v) = 0$ then we have for any $\alpha > 0$

$$\begin{aligned} J_v(k) &= \frac{i}{k} \int_0^{2\pi} \int_0^\alpha e^{ik\{1-\cos(\phi)\}} \partial_\phi f(\theta, \phi) d\phi d\theta \\ &\quad + \frac{i}{k} \int_0^{2\pi} \int_{\pi-\alpha}^\pi e^{ik\{1-\cos(\phi)\}} \partial_\phi f(\theta, \phi) d\phi d\theta \\ &\quad + \frac{i}{k} \int_0^{2\pi} \int_\alpha^{\pi-\alpha} e^{ik\{1-\cos(\phi)\}} \partial_\phi f(\theta, \phi) d\phi d\theta. \end{aligned}$$

We use an integration by parts to obtain for the second term of the right hand side

$$\begin{aligned} \int_\alpha^{\pi-\alpha} e^{ik\{1-\cos(\phi)\}} \partial_\phi f(\theta, \phi) d\phi &= \frac{i}{k} \left[e^{ik\{1-\cos(\phi)\}} \frac{\partial_\phi f(\theta, \phi)}{\sin(\phi)} \right]_\alpha^{\pi-\alpha} \\ &\quad - \frac{i}{k} \int_\alpha^{\pi-\alpha} e^{ik\{1-\cos(\phi)\}} \left\{ \frac{\partial_\phi^2 f(\theta, \phi)}{\sin(\phi)} - \frac{\cos(\phi) \partial_\phi f(\theta, \phi)}{\sin(\phi)^2} \right\} d\phi, \end{aligned}$$

for the other terms of the right hand side direct estimations give us

$$|J_v(k)| \leq \frac{C\alpha}{|k|} \int_0^{2\pi} \sup_\phi |\partial_\phi f(\theta, \phi)| d\theta + \frac{C}{\alpha|k|^2} \int_0^{2\pi} \left\{ \sup_\phi |\partial_\phi f(\theta, \phi)| d\theta + \int_0^{2\pi} \int_0^\pi |\partial_\phi^2 f(\theta, \phi)| d\phi d\theta \right\},$$

choosing $\alpha = \sqrt{|k|}^{-1}$ and working like in the proof of the estimate (3.1), we obtain estimate (3.2). The reader recognized the proof of the well known Van der Corput Lemma with modification in order to give precise estimates.

For the estimate (3.3), we first split the integral $J_v(k)$ in two hemispheres with respect to the pole v and we choose $\alpha = \text{dist}(\text{supp}(f), v)$ or $\alpha = \text{dist}(\text{supp}(f), -v)$.

We obtain first the

Proposition 3.1. *Let $h \in \mathcal{C}(\mathbb{R})$ and $g \in \mathcal{C}^2(\mathbb{R}^3)$ be such that the integrals appearing in the following estimate are finite. Then defining*

$$I(k, u) = \int_{\mathbb{R}^3} e^{ik\{h(|\xi|) - \xi \cdot u\}} g(\xi) d\xi,$$

for any $u \in \mathbb{R}^3$ and any $k \in \mathbb{R}$, we have

$$|I(k, u)| \leq \frac{C}{|ku|} \max_{|\alpha| \leq 1} \left\{ \int_{\mathbb{R}^3} |\xi|^{|\alpha|-1} \frac{|\nabla^\alpha g(\xi)|}{\left| \frac{u}{|u|} - \frac{\xi}{|\xi|} \right|^{|\alpha|}} d\xi, \int_{\mathbb{R}^3} |\xi|^{|\alpha|-1} \frac{|\nabla^\alpha g(\xi)|}{\left| \frac{u}{|u|} + \frac{\xi}{|\xi|} \right|^{|\alpha|}} d\xi \right\} \quad (3.4)$$

where C does not depend on h , g , k or u .

If moreover g vanishes in a cone of axis $D = \text{Span}(u)$, we have

$$|I(k, u)| \leq \frac{C}{|ku|^2 \text{dist}(\text{supp}(g) \cap S^2, D \cap S^2)} \times \\ \times \max_{|\alpha| \leq 1} \left\{ \int_{\mathbb{R}^3} |\xi|^{2|\alpha|-2} \frac{|\nabla^{2\alpha} g(\xi)|}{\left| \frac{u}{|u|} - \frac{\xi}{|\xi|} \right|^{|\alpha|}} d\xi, \int_{\mathbb{R}^3} |\xi|^{2|\alpha|-2} \frac{|\nabla^{2\alpha} g(\xi)|}{\left| \frac{u}{|u|} + \frac{\xi}{|\xi|} \right|^{|\alpha|}} d\xi \right\}$$

where C does not depend on h , g , k or u .

Proof. We write

$$I(k, u) = \int_{\mathbb{R}^3} e^{ik\{h(|\xi|) - \xi \cdot u\}} g(\xi) d\xi = \int_{\mathbb{R}^+} e^{ik\{h(\rho) - \rho|u|\}} J_{\frac{u}{|u|}, \rho}(\rho k|u|) \rho^2 d\rho,$$

where $J_{v, \rho}(k) = \int_{S^2} e^{ik\{1-v \cdot \omega\}} g(\rho\omega) d\omega$ and we apply Lemma 3.1.

We introduce a first useful variant with the

Proposition 3.2. *Let $g \in \mathcal{C}^{1+k}(\mathbb{R}^3)$ be such that the integrals appearing in the following estimate are finite. We introduce*

$$F(x) = \int_{\mathbb{R}^3} e^{i\{|\xi||x| - \xi \cdot x\}} g(\xi) d\xi$$

for any $x \in \mathbb{R}^3$. Then for all $\alpha \in \mathbb{N}^3$ such that $|\alpha| \leq k$ we have

$$|\nabla^\alpha F(x)| \leq \frac{C}{|x|^{|\alpha|+1}} \max_{|\beta| \leq 1+|\alpha|} \left\{ \int_{\mathbb{R}^3} |\xi|^{|\beta|-1} \frac{|\nabla^\beta g(\xi)|}{\left| \frac{x}{|x|} - \frac{\xi}{|\xi|} \right|^{|\beta|}} d\xi \right\}. \quad (3.5)$$

If moreover g vanishes in a half cone of axis $D^+ = \left\{ \rho \frac{x}{|x|}, \rho \in \mathbb{R}^+ \right\}$. Then for all $\alpha \in \mathbb{N}^3$ such that $|\alpha| \leq k$ we have

$$|\nabla^\alpha F(x)| \leq \frac{C}{|x|^{|\alpha|+2} \text{dist}(\text{supp}(g) \cap S^2, D^+ \cap S^2)} \max_{|\beta| \leq 2+|\alpha|} \left\{ \int_{\mathbb{R}^3} |\xi|^{|\beta|-2} \frac{|\nabla^\beta g(\xi)|}{\left| \frac{x}{|x|} - \frac{\xi}{|\xi|} \right|^{|\beta|}} d\xi \right\}. \quad (3.6)$$

Proof. The critical points correspond to the the semi axis spanned by x . We treat the part of the integrals which is far from critical points by using an integration by parts with help of the operator

$L = \frac{\frac{\xi}{|\xi|} - \frac{x}{|x|}}{|x| \left| \frac{\xi}{|\xi|} - \frac{x}{|x|} \right|^2} \cdot \nabla_\xi$. Let be $\phi(x, \xi) = \{|\xi||x| - \xi \cdot x\}$, we have

$$F(x) = \langle L e^{i\phi(x, \cdot)}, g \rangle = \langle e^{i\phi(x, \cdot)}, L^* g \rangle,$$

with

$$L^* = -L - \frac{2}{|x||\xi| \left| \frac{\xi}{|\xi|} - \frac{x}{|x|} \right|^2}.$$

This gives the bound

$$\frac{C}{|x|} \max_{|\beta| \leq 1} \left\{ \int_{\mathbb{R}^3} |\xi|^{|\beta|-1} |\nabla^\beta g(\xi)| d\xi \right\},$$

we obtain Estimate (3.5) for $\alpha = 0$. The method to treat the other part of the integral is exactly the one we used in the proof of Proposition 3.1.

For higher order derivatives, we have

$$\nabla_x e^{ik\phi(x,\xi)} = \frac{|\xi|}{|x|} \nabla_\xi e^{ik\phi(x,\xi)}$$

and so

$$\langle \nabla_x e^{i\phi(x,\cdot)}, g \rangle = -\frac{1}{|x|} \langle e^{i\phi(x,\cdot)}, \nabla |Q|g \rangle.$$

the result is then obtained by applying this trick $|\alpha|$ times and then repeating our proof for the case $\alpha = 0$, we obtain Estimates (3.5) and (3.6) for $\nabla^\alpha F(x)$.

And finally, we need the

Proposition 3.3. *Let be $g \in C^2(\mathbb{R}^3)$ with compact support. Then for any $u \in \mathbb{R}^3$, $k \in \mathbb{R}$ and*

$$I(k, u) = \int_{\mathbb{R}^3} e^{ik\{\sqrt{\xi^2+m^2}-\xi \cdot u\}} g(\xi) d\xi,$$

we have

$$|I(k, u)| \leq \frac{C}{|k|^{3/2}} \max_{|\alpha| \leq 2} \left\{ \int_{\mathbb{R}^3} |\langle \xi \rangle^{|\alpha|-1} \nabla^\alpha g(\xi)| d\xi \right\};$$

$$\frac{1}{|u| \sqrt{\inf_{x \in \text{supp}(g)} \left\{ \frac{m^2}{\sqrt{x^2 + m^2}^3} \right\}}} \max_{l \leq 1} \left\{ \int_{\mathbb{R}^3} |\xi|^{l-2} \frac{|\partial_{|\xi|}^l \partial_\omega g(\xi)|}{\left| \frac{u}{|u|} - \frac{\xi}{|\xi|} \right|} d\xi \right\}. \quad (3.7)$$

Proof. We can suppose $u = (0, 0, |u|)$ since estimate (3.7) is invariant under the action of rotations.

The oscillatory integral $I(k, u)$ is bounded and critical points of the phase of $I(k, u)$ are supported by the semi axis spanned by u . With help of a smooth cut-off function χ , we split the integral in two parts $I(k, u) = I_1(k, u) + I_2(k, u)$, where $I_1(k, u)$ is supported in a half cone around u . We then use multiple integrations by parts with help of the operator

$$L = \frac{\frac{\xi}{\sqrt{\xi^2+m^2}} - u}{|k| \left| \frac{\xi}{\sqrt{\xi^2+m^2}} - u \right|^2} \cdot \nabla_\xi.$$

Since $(1 - \chi)g \in C^2(\mathbb{R}^3)$ has support far from critical points and since for $\lambda(\xi) = \sqrt{\xi^2 + m^2}$ we have $\left\| \nabla_\xi^\alpha \lambda(\xi) \right\| \leq C_\alpha \lambda(\xi)^{1-|\alpha|}$, we obtain

$$|I_1(k, u)| \leq \frac{C}{|k|} \sum_{|\alpha| \leq 1} \|\lambda(Q)^{|\alpha|-1} \nabla^\alpha g\|_{L^1}$$

and

$$|I_1(k, u)| \leq \frac{C}{k^2} \sum_{|\alpha| \leq 2} \|\lambda(Q)^{|\alpha|-1} \nabla^\alpha g\|_{L^1}.$$

Otherwise $I_2(k, u)$ has support in a small cone around u , and we have

$$\begin{aligned} I_2(k, u) &= \int_{\mathbb{R}^3} e^{ik\{\sqrt{\xi^2+m^2}-\xi \cdot u\}} \tilde{g}(\xi) d\xi d\omega \rho^2 d\rho \\ &= \int_{\mathbb{R}^+} e^{ik\{\sqrt{\xi^2+m^2}-|\xi||u|\}} J_{\frac{u}{|u|}, \rho}(\rho k|u|) \rho^2 d\rho, \end{aligned}$$

with

$$J_{v,\rho}(k) = \int_{S^2} e^{ik\{1-v\cdot\omega\}} \tilde{g}(\rho\omega) d\omega,$$

where $\tilde{g} = \chi g$. We obtain after an integration by parts

$$J_{v,\rho}(k) = -\frac{i}{k} \int_0^{2\pi} \left[e^{ik\{1-\cos(\phi)\}} \tilde{g}(\rho\omega(\theta, \phi)) \right]_0^\pi d\theta + \frac{i}{k} \int_0^{2\pi} \int_0^\pi e^{ik\{1-\cos(\phi)\}} \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) d\phi d\theta,$$

Since we assumed \tilde{g} is supported in half cone around u , we have $\tilde{g}(\rho\omega(\theta, \pi)) = 0$. Hence we obtain

$$J_{v,\rho}(k) = \frac{i}{k} \int_0^{2\pi} \int_0^\pi \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) d\theta + \frac{i}{k} \int_0^{2\pi} \int_0^\pi e^{ik\{1-\cos(\phi)\}} \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) d\phi d\theta,$$

and so

$$\begin{aligned} I_2(k, u) &= \frac{i}{|k||u|} \int_{\mathbb{R}^+} \int_0^{2\pi} \int_{-\pi}^\pi e^{ik\{\sqrt{\rho^2+m^2}-\rho|u|\}} \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) d\phi d\theta \rho d\rho \\ &\quad + \frac{i}{|k||u|} \int_{\mathbb{R}^+} \int_0^{2\pi} \int_{-\pi}^\pi e^{i\{\sqrt{\rho^2+m^2}-\rho|u|\cos(\phi)\}} \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) d\phi d\theta \rho d\rho. \end{aligned} \quad (3.8)$$

Let us now study the decay resulting from the dispersive behavior of the radial part. To this end, we follow the proof of the well-known Van Der Corput lemma. We study

$$L(k, u, \phi, \phi', \theta) = \int_{\mathbb{R}^+} e^{i|k|\{\sqrt{\rho^2+m^2}-\rho|u|\cos(\phi')\}} \partial_\phi \tilde{g}(\rho\omega(\theta, \phi)) \rho d\rho.$$

Notice that, in view of (3.8), we are only interested by $L(k, u, \phi, \phi', \theta)$ and $L(k, u, \phi, 0, \theta)$. First, for any differentiable function on \mathbb{R} such that $f' \geq 1$, we have for any $\alpha \in \mathbb{R}^+$

$$|\{t \in \mathbb{R}; |f(t)| \leq \alpha\}| \leq \alpha. \quad (3.9)$$

We introduce

$$h(\rho) = \sqrt{\rho^2 + m^2} - \rho|u|\cos(\phi'),$$

and we apply (3.9) to

$$f(\rho) = \frac{\partial_\rho h(\rho)}{\inf_{x \in \text{supp}(g)} \left\{ |\partial_{|x|}^2 h(|x|)| \right\}}.$$

We notice that $\partial_\rho^2 h(\rho)$ does not depend on u or ϕ' . With help of a smooth cut-off function, we split the integral L in two parts, one has support

$$\{\rho \in \mathbb{R}^+; |f(\rho)| < \alpha\}$$

and the other is its complementary. In fact, we obtain exactly three interval corresponding to

$$\{\rho \in \mathbb{R}^+; f(\rho) < -\alpha\}, \{\rho \in \mathbb{R}^+; -\alpha \leq f(\rho) \leq \alpha\}, \{\rho \in \mathbb{R}^+; \alpha < f(\rho)\}.$$

In the first and third interval, we make an integration by parts and in the second interval, we use Estimate (3.9). Hence we obtain the bound

$$\begin{aligned} |L(k, u, \phi, \phi', \theta)| &\leq \max \left\{ \alpha, \frac{1}{\alpha|k| \inf_{x \in \text{supp}(\tilde{g})} \left\{ |\partial_{|x|}^2 h(|x|)| \right\}} \right\} \max_{\rho \in \mathbb{R}^+} \{\rho |\partial_\phi \tilde{g}(\rho\omega(\theta, \phi))|\} \\ &\quad + \frac{1}{\alpha|k| \inf_{x \in \text{supp}(\tilde{g})} \left\{ |\partial_{|x|}^2 h(|x|)| \right\}} \max \left\{ \int_{\mathbb{R}^+} |\partial_\phi \tilde{g}(\rho\omega(\theta, \phi))| d\rho, \int_{\mathbb{R}^+} \rho |\partial_\rho \partial_\phi \tilde{g}(\rho\omega(\theta, \phi))| d\rho \right\}. \end{aligned}$$

We use

$$\rho |\partial_\phi \tilde{g}(\rho\omega(\theta, \phi))| \leq 2 \max \left\{ \int_{\mathbb{R}^+} |\partial_\phi \tilde{g}(r\omega(\theta, \phi))| dr, \int_{\mathbb{R}^+} r |\partial_\rho \partial_\phi \tilde{g}(r\omega(\theta, \phi))| dr \right\}$$

and then choose

$$\alpha = \frac{1}{\sqrt{|k| \inf_{x \in \text{supp}(\tilde{g})} \left\{ |\partial_{|x|}^2 h(|x|)| \right\}}},$$

and plugging the resulting estimates for $\phi' = 0$ and $\phi' = \phi$ in (3.8), we obtain estimate (3.7).

3.2. Dispersive estimates for the free case equation. Thanks to the tools introduced in Section 3.1, we are able to state the

Theorem 3.1 (Dispersive estimates for free Dirac operator). *For any $p \in [1, 2]$, for all $\theta \in [0, 1]$, for all $s, s' \in \mathbb{R}$, such that $s - s' \geq (\frac{2}{p} - 1)(2 + \theta)$ and $q \in [1, \infty]$, we have*

$$\|e^{-itD_m}\|_{B_{p,q}^s, B_{p',q}^{s'}} \leq K(t)^{\frac{2}{p}-1},$$

with

$$K(t) = \begin{cases} |t|^{-1+\theta/2} & \text{if } |t| \in [0, 1], \\ |t|^{-1-\theta/2} & \text{if } |t| \in [1, \infty), \end{cases}$$

and $p' = \frac{p}{p-1}$.

Proof. We only need to prove the case $p = 1$, since the general case follows by interpolation of the case $p = 1$ and the charge conservation which corresponds to the case $p = 2$. Then using $D_m = \sqrt{-\Delta + m^2}(\pi_+ - \pi_-)$ with $\pi_{\pm} = \mathbb{1}_{\mathbb{R}^{\pm}}(D_m) = \frac{1}{2} \{1 \pm |D_m|^{-1} D_m\}$, we obtain the estimates from those relative to the relativistic Schrödinger operator $\sqrt{-\Delta + m^2}$:

$$\|e^{-it\sqrt{-\Delta+m^2}}\|_{B_{1,q}^s, B_{\infty,q}^{s'}} \leq K(t)$$

which in turn follows from

Proposition 3.4. *For any $\chi \in \mathcal{D}(\mathbb{R}^3, \mathbb{C}^4)$, we define $\chi_j(x) = \chi(2^{-j}|x|)$. Then for $\theta' \in [0, 1]$, we have:*

1. if $0 \notin \text{supp}(\chi)$,

$$\|e^{-it\sqrt{-\Delta+m^2}}\chi_j(\sqrt{-\Delta+m^2})\|_{L^1, L^\infty} \leq C2^{2j}|t|^{-(1+\theta'/2)} \quad (3.10)$$

where C is independent of t and j .

2. if $0 \in \text{supp}(\chi)$,

$$\|e^{-it\sqrt{-\Delta+m^2}}\chi(-\Delta+m^2)\|_{L^1, L^\infty} \leq C\langle t \rangle^{-3/2} \quad (3.11)$$

where C is independent t .

We postpone the proof of Proposition 3.4 until the end of the proof of Theorem 3.1.

Indeed one has

$$2^{js'} \left\| e^{-it\sqrt{-\Delta+m^2}} \chi_j(\sqrt{-\Delta+m^2}) f \right\|_{\infty} \leq C2^{3j} \|f\|_1$$

interpolating with Estimate (3.10) of Proposition 3.4 for $\theta' = 0$ when $t \leq 1$ and using Estimate (3.10) for $t \geq 1$, one obtains

$$2^{js'} \left\| e^{-it\sqrt{-\Delta+m^2}} \chi_j(\sqrt{-\Delta+m^2}) f \right\|_{\infty} \leq C\kappa_j(t) 2^{js} \|f\|_1$$

with

$$\kappa_j(t) = 2^{j(2+\theta+s'-s)} \begin{cases} |t|^{-1+\theta/2} & \text{if } |t| \leq 1, \\ |t|^{-1-\theta/2} & \text{if } |t| \geq 1. \end{cases}$$

We use

$$\sup_{j \in \mathbb{N}} \kappa_j \leq CK(t)$$

if $2 + \theta \leq s - s'$ and Estimate (3.11) to prove Theorem 3.1. Hence to conclude the proof, we need to give the

Proof (Proof of Proposition 3.4). The kernel of $e^{-it\sqrt{-\Delta+m^2}}\chi_j(\sqrt{-\Delta+m^2})$ is given by $(2\pi)^{-3/2}K_j(x-y)$ where

$$K_j(x, t) = \int_{\mathbb{R}^3} e^{-it\sqrt{\xi^2+m^2}+x \cdot \xi} \chi_j(\sqrt{\xi^2+m^2}) d\xi$$

Hence, we estimate the L^∞ norm of K_j .

If $|x|/|t| \ll 2^{j-1}/\sqrt{2^{2j-2}+m^2}$ or $|x|/|t| \gg 1$, we use non stationary phase lemma in \mathbb{R}^3 with help of the operator $L = \frac{\frac{\xi}{\sqrt{\xi^2+m^2}}-x}{t \left| \frac{\xi}{\sqrt{\xi^2+m^2}}-x \right|^2} \cdot \nabla$. Hence, in this case, we obtain the estimate

$$\left| \int_{\mathbb{R}^3} e^{-it\sqrt{\xi^2+m^2}+x\cdot\xi} \chi_j(\sqrt{\xi^2+m^2}) d\xi \right| \leq C_n 2^{-(n-3)j} |t|^{-n},$$

for any $n \in \mathbb{N}$. Otherwise, we apply Proposition 3.1 with $h(r) = \sqrt{r^2+m^2}$, $k = t$, $u = x/t$ and $g(x) = \chi_j(|x|)$. So if $0 \notin \text{supp}(\chi)$, to obtain

$$\left| \int_{\mathbb{R}^3} e^{-it\sqrt{\xi^2+m^2}+x\cdot\xi} \chi_j(\sqrt{\xi^2+m^2}) d\xi \right| \leq \frac{C}{|t|} \max_{|\beta| \leq 1} \int_{\mathbb{R}^3} |\xi|^{|\beta|-1} \frac{|\nabla^\beta \chi_j(\sqrt{\xi^2+m^2})|}{\left| \frac{x}{|x|} \pm \frac{\xi}{|\xi|} \right|} d\xi \leq \frac{C 2^{2j}}{|t|}.$$

Notice that in our case, $|x|/|t| \geq c' > 0$. If instead of Proposition 3.1, we use Proposition 3.3 with $g = \chi_j$, $k = t$ and $u = x/t$, we prove the estimate

$$|K_j(x, t)| \leq \frac{C 2^{2j}}{|t|^{3/2}}.$$

The estimate (3.10) is then obtained by interpolation. For (3.11), we use the classical stationary (Morse lemma...) and non-stationary phase methods (integration by parts...) in \mathbb{R}^3 . For more details about the method one can look at the end of the proof of Proposition 3.9. This ends the proof of 3.4.

This ends the proof of Theorem 3.1.

3.3. Distorted Plane Waves. Our aim is now to generalize the previous method to the perturbed case. Let us introduce the wave operators

$$W^\pm = \lim_{t \rightarrow \pm\infty} e^{it(D_m+V)} e^{-itD_m}$$

(for the existence and the completeness : $\text{Ran}(W^\pm) = \text{Ran}(\mathbf{P}_c(H))$ of these operator see [GM01, Theorem 1.5]). With the intertwining property (see Identity (2.5)), and Fourier transform \mathcal{F} , we shall obtain for $h(\xi) = \alpha \cdot \xi + m\beta$

$$e^{-itH} \mathbf{P}_c(H) = W^\pm e^{-itD_m} W^{\pm*} = W^\pm \mathcal{F} e^{-ith(Q)} (W^\pm \mathcal{F})^*.$$

So we can adapt the previous method if we are able to prove some estimates about the kernel ψ_V of $W^\pm \mathcal{F}$. The kernel ψ_V is called *distorted plane wave*. We just notice that ψ_V is a 4×4 matrix valued function.

We will show that the previous method work with $\psi_V \psi_V^* \chi_j$ in place of χ_j with small modifications. So we need estimates on ψ_V . Generally, distorted plane waves are studied like perturbations of free plane waves. So we will prove estimates on the perturbative part, written w in the sequel.

3.3.1. Definition and properties. We need to introduce the free plane wave, let be

$$h(k) = \alpha \cdot k + m\beta$$

for any $k \in \mathbb{R}^3$, we notice $D_m = h(P)$. This hermitian matrix has for eigenvectors the

$$\psi_0^j(k, x) = e^{ik \cdot x} u(k) e_j$$

where

$$u(k) = \frac{(m + \lambda(k)) Id - \beta \alpha \cdot k}{\sqrt{2\lambda(k)(m + \lambda(k))}} \quad (3.12)$$

with $\lambda(k) = \sqrt{k^2 + m^2}$ and $(e_j)_j$ are vectors of the canonical basis of \mathbb{C}^4 . For more details see [Tha92, Section 1.4, Section 1.F].

By definition, a distorted plane wave is a solution of the PDO equation

$$(D_m + V) \psi = \pm \sqrt{k^2 + m^2} \psi \quad (3.13)$$

with for some j and any $k \in \mathbb{R}^3$, $\psi(k, x) - \psi_0^j(k, x)$ tending to zero as x goes to infinity (in some sense), see [Agm75, section 5].

A solution of (3.13) is a function $\psi(k, x)$ of two variables here k is a 3-dimensional vector which is called the wave vector. A free plane wave ψ_0^j satisfies the PDO equation (3.13) in the case $V = 0$. Following [Agm75], we introduce two families of function

$$\psi_V^j(k, x) = \psi_0^j(k, x) - \left\{ R_V^+(\lambda(k))V(\cdot)\psi_0^j(k, \cdot) \right\}(x)$$

for $j \in \{1, 2\}$ and

$$\psi_V^j(k, x) = \psi_0^j(k, x) - \left\{ R_V^+(-\lambda(k))V(\cdot)\psi_0^j(k, \cdot) \right\}(x)$$

for $j \in \{3, 4\}$. The rest of the proof works also for R_V^- instead of R_V^+ (the trace of the resolvent R_V^\pm was introduced in (2.1)).

In case there is no resonance at thresholds and no eigenvalue at thresholds, Theorem 1.1 gives us that $R_V^+(\lambda(p))$ is in $B(L_\sigma^2, L_{-\sigma}^2)$ for any $\sigma > 5/2$, this also work if $\sigma \geq 1$ see Proposition 3.10 below. So the previous definition make sense if Assumption 1.1 holds and we have the

Proposition 3.5. *Suppose that Assumptions 1.1 and 1.2 hold. Then for any $k \in \mathbb{R}^3 \setminus \{0\}$, $\psi_V^j(k, x)$ satisfies equation (3.13).*

Distorted plane waves define a generalized Fourier transform. We introduce $\psi_V(k, x) \in \mathcal{M}_4(\mathbb{C})$ the matrix with vector column $\psi_V^j(k, x)$ and we define

$$(\mathcal{F}_V f)(k) = \int_{\mathbb{R}^3} \psi_V(k, x) f(x) dx,$$

which is *a priori* defined on the Schwartz space $\mathcal{S}(\mathbb{R}^3, \mathbb{C}^4)$ but will be extended to L^2 .

Distorted plane waves are also called generalized eigenfunctions, since they correspond to “eigenvalues” associated with the continuous spectrum. Indeed, we can prove the

Theorem 3.2 (Eigenfunction Expansion). *The operator \mathcal{F}_V defines a bounded linear map from L^2 into itself. Its kernel is given by the the sum of the eigenspaces of H . Moreover it is a unitary map from $\mathbf{P}_c(H)L^2$ onto $L^2(\mathbb{R}^3)$ with*

$$(\mathcal{F}_V^* f)(x) = \lim_{n \rightarrow \infty} \int_{K_n} \psi_V(k, x)^* f(k) dk,$$

for any $(K_n)_{n \in \mathbb{N}}$ a family of compact sets with $K_n \subset K_{n+1}$ and $\cup_{n \in \mathbb{N}} K_n = \mathbb{R}^3$. Finally, for any interval $I \subset \mathbb{R}$, one has

$$\|\mathbf{1}_I(H)f\|^2 = \int_{\sigma(h(k)) \cap I \neq \emptyset} |\mathcal{F}_V f|^2 dk \quad (3.14)$$

where $\sigma(h(k))$ is the spectrum of $h(k)$.

Proof. The proof is an easy adaptation of the proof of [Agm75, Theorem 6.2] (see also [RS79, Theorem XI.41]), the main difference is that here we insert the unitary matrix u defined in (3.12). Formula (3.14) is nothing more than an adaptation of [Agm75, Formula (6.6)] or [RS79, Formula 82e’].

We also have the

Lemma 3.2 (Intertwining Property). *Let g be a bounded borelian function with support in $\mathbb{R} \setminus (-m, m)$, we have*

$$\mathcal{F}_V g(H) = (g \circ h) \mathcal{F}_V. \quad (3.15)$$

Proof. Using (3.14), we obtain that (3.15) is true for $g = \mathbf{1}_I$ with I an interval of $\mathbb{R} \setminus (-m, m)$. We then obtain it for bounded borelian function with support in $\mathbb{R} \setminus (-m, m)$, usual density arguments and properties of functional calculus. More precisely, we use the fact that a bounded sequence of borelian functions which converges everywhere gives a sequence of bounded operators which converge strongly.

Hence we deduce that, for any $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$, the kernel of $e^{-itH}\chi(H)$ is given by

$$[e^{-itH}\chi(H)\mathbf{P}_c(H)](x, y) = \int_{\mathbb{R}^3} \psi_V(k, x)^* e^{-it h(k)} \chi(h(k)) \psi_V(k, y) dk.$$

which exactly means

$$e^{-itH}\chi(H)\mathbf{P}_c(H) = (\mathcal{F}_V)^* e^{-it h} \chi(h) \mathcal{F}_V \quad (3.16)$$

We recall that we want to prove the decay of $e^{-itH}\chi(H)$ as $t \rightarrow +\infty$ in some Besov spaces. We observe that

$$e^{-it h(k)} \chi(h(k)) = e^{-it \lambda(k)} \chi(\lambda(k)) P_+(k) + e^{it \lambda(k)} \chi(-\lambda(k)) P_-(k),$$

where $P_+(k)$ (resp. $P_-(k)$) is the projector associated with the positive (resp. negative) part of the spectrum of $h(k)$, *i.e.*

$$P_\pm(k) = \frac{1}{2} \left(1 \pm \frac{h(k)}{\lambda(k)} \right).$$

Hence, in the following we study the functions

$$(x, y) \in \mathbb{R}^3 \times \mathbb{R}^3 \mapsto \int_{\mathbb{R}^3} e^{\mp i t \lambda(k)} (P_\pm \psi_V(k, x))^* (P_\pm \psi_V(k, y)) \chi(h(k)) dk.$$

3.3.2. End of the proof of Theorem 1.2. We now prove Theorem 1.2 with help of three propositions which will be proven in Section 3.3.3. These propositions give some estimates on the perturbed part of the distorted plane wave. Following Cuccagna and Schirmer in [CS01], we write $\psi_V(k, x) = e^{ik \cdot x} (u(k) + w(k, x))$ where w is the perturbation part which satisfies

$$w(k, x)_i = \begin{cases} e^{-ik \cdot Q} R_V^+(+\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{1, 2\}, \\ e^{-ik \cdot Q} R_V^+(-\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{3, 4\}, \end{cases} \quad (3.17)$$

and we now state our propositions.

Proposition 3.6. *Suppose that Assumptions 1.1 and 1.2 hold. Then there exists $C > 0$ such that for any $k, x \in \mathbb{R}^3 \setminus \{0\}$, and any $\beta \in \mathbb{N}^3$ with $|\beta| \leq 1$, one has*

$$|\nabla_k^\beta w(k, x)| \leq \frac{C}{\langle k \rangle^{|\beta|}} \frac{\langle x \rangle^{|\beta|}}{\left\langle |x| \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle}. \quad (3.18)$$

Moreover one has

$$|\{\nabla_k w(k, x) \langle Q \rangle^{-\sigma} q\}(x)| \leq C \frac{\langle \min\{|x|, |k|\} \rangle}{\langle k \rangle \left\langle \min\{|x|, |k|\} \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle^2}. \quad (3.19)$$

We use this to prove the time decay in $|t|^{-1}$. Unfortunately this doesn't work for the $|t|^{-3/2}$ decay, hence we then study

$$v(k, x)_i = \begin{cases} e^{i|k||Q|} R_V^+(+\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{1, 2\}, \\ e^{-i|k||Q|} R_V^+(-\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{3, 4\}, \end{cases} \quad (3.20)$$

and

$$\tilde{v}(k, x)_i = \begin{cases} e^{i|k||Q|} \nabla_k R_V^+(+\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{1, 2\}, \\ e^{-i|k||Q|} \nabla_k R_V^+(-\lambda(k)) V e^{ik \cdot Q} u(k)_i, & \text{if } i \in \{3, 4\}. \end{cases} \quad (3.21)$$

One has the

Proposition 3.7. *Suppose that Assumptions 1.1 and 1.2 hold. Then if $\rho > 3 + |\beta|$ for some $\beta \in \mathbb{N}^3$, there exists $C > 0$ such that for any $k, x \in \mathbb{R}^3 \setminus \{0\}$, one has*

$$|\nabla_k^\beta v(k, x)| \leq \frac{C}{\left\langle |x| \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle}.$$

Proposition 3.8. *Suppose that Assumptions 1.1 and 1.2 hold. Then if $\rho > 3 + |\beta|$ for some $\beta \in \mathbb{N}^3$, there exists $C > 0$ such that for any $k, x \in \mathbb{R}^3 \setminus \{0\}$, one has*

$$|\nabla_k^\beta \tilde{v}(k, x)| \leq \frac{\langle \min\{|x|, |k|\} \rangle}{\langle k \rangle \left\langle \min\{|x|, |k|\} \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle^2}.$$

Using Proposition 3.6, 3.7 and 3.8 (which are proved in Section 3.3.3 below), let us prove the following

Proposition 3.9. *Suppose that Assumptions 1.1 and 1.2 hold. Then we have for $\chi \in C_0^\infty(\mathbb{R})$ with support in $\mathbb{R} \setminus [-m, m]$ for any $\theta \in [0, 1]$ and $j \in \mathbb{N}$,*

$$\|e^{-itH} \chi(2^{-j}H)\|_{L^1 \rightarrow L^\infty} \leq \frac{C 2^{(2+\theta)j}}{|t|^{1+\theta/2}}, \quad (3.22)$$

with C independent of t and j .

We also have for $\chi \in C_0^\infty(\mathbb{R})$, for any $\theta \in [0, 1]$

$$\|e^{-itH} \chi(H) \mathbf{P}_c(H)\|_{L^1 \rightarrow L^\infty} \leq \frac{C}{|t|^{1+\theta/2}}, \quad (3.23)$$

with C independent of t .

Proof. The proof works like the one of Proposition 3.4 with some modification due to the fact that high derivatives in k of $w(k, x)$ grow with respect to x .

We need the L^∞ norm of the kernel of $e^{-itH} \chi(2^{-j}H)$ which thanks to (3.16) is given by

$$\begin{aligned} I_j(t, x, y) &= \int_{\mathbb{R}^3} e^{-it\sqrt{\xi^2+m^2}} e^{-i\xi \cdot (x-y)} \{P_+(\xi)(u^*(\xi) + w^*(x, \xi))\} \{P_+(\xi)(u(\xi) + w(y, \xi))\} \chi(2^{-j}\lambda(\xi)) d\xi \\ &+ \int_{\mathbb{R}^3} e^{+it\sqrt{\xi^2+m^2}} e^{-i\xi \cdot (x-y)} \{P_-(\xi)(u^*(\xi) + w^*(x, \xi))\} \{P_-(\xi)(u(\xi) + w(y, \xi))\} \chi(-2^{-j}\lambda(\xi)) d\xi. \end{aligned}$$

We notice that if we expand each integrand in terms of u and w , we obtain the sum of the integrals

$$I_j^\pm[z, z'](t, x, y) = \int_{\mathbb{R}^3} e^{\mp it\sqrt{\xi^2+m^2}} e^{-i\xi \cdot (x-y)} \{P_\pm(\xi)z^*(x, \xi)P_\pm(\xi)z'(y, \xi)\} \chi(\pm 2^{-j}\lambda(\xi)) d\xi.$$

with $z, z' \in \{u, w\}$. We notice that $I_j^+[u, u](t, x, y) + I_j^-[u, u](t, x, y)$ is the kernel of $e^{-itD_m} \chi_j(D_m)$, hence we only treat the other integrals.

For the $|t|^{-1}$ decay, if $|x - y|/|t| \ll 2^{j-1}/\sqrt{2^{2j-2} + m^2}$ or $|x - y|/|t| \gg 1$, the phase has no critical point. We use an integration by parts in \mathbb{R}^3 with help of the operator

$$L = \frac{\left(\frac{\xi}{\sqrt{\xi^2+m^2}} - x \right)}{t \left| \frac{\xi}{\sqrt{\xi^2+m^2}} - x \right|^2} \cdot \nabla_\xi.$$

So with the estimate (3.18) of Proposition 3.6 and with

$$\left| \partial_i \frac{\xi}{\sqrt{\xi^2+m^2}} \right| \leq \frac{C}{|\xi|},$$

we obtain the estimate

$$|I_j^\pm[z, z'](t, x, y)| \leq C 2^{2j} |t|^{-1},$$

with C independent of j and t .

Otherwise if $|x - y|/|t| \geq c > 0$, using first (3.4) of Proposition 3.1 and then (3.19) of Proposition 3.6, we infer

$$|I_j^\pm[z, z'](t, x, y)| \leq C 2^{2j} |t|^{-1},$$

with C independent of j and t .

For the $|t|^{-3/2}$ decay, first if $|x - y|/|t| \geq c > 0$, we write

$$I_j^\pm[z, z'](t, x, y) = \int_{\mathbb{R}^+} e^{\mp it\sqrt{\rho^2+m^2}-i\rho|x-y|} J_{\frac{x-y}{|x-y|}}(\rho|x-y|) \rho^2 d\rho.$$

where

$$J_v(k) = \int_{S^2} e^{ik(1-\omega \cdot v)} \{P_\pm(\rho\omega)z^*(x, \rho\omega)P_\pm(\rho\omega)z'(y, \rho\omega)\} \chi(\pm 2^{-j}\lambda(\rho\omega)) d\omega.$$

We can suppose $v = (0; 0; 1)$ and so

$$J_v(k) = \int_0^{2\pi} \int_0^\pi e^{ik(1-\cos(\phi))} \left\{ P_\pm(\rho\omega(\theta, \phi))z^*(x, \rho\omega(\theta, \phi)) \times \right. \\ \left. \times P_\pm(\rho\omega(\theta, \phi))z'(y, \rho\omega(\theta, \phi)) \right\} \chi(\pm 2^{-j}\lambda(\rho\omega(\theta, \phi))) \sin(\phi) d\phi d\theta.$$

An integration by parts in ϕ gives

$$J_v(k) = \frac{1}{ik} \int_0^{2\pi} \int_0^\pi \left[e^{ik(1-\cos(\phi))} \left\{ P_\pm(\rho\omega(\theta, \phi))z^*(x, \rho\omega(\theta, \phi)) \times \right. \right. \\ \left. \left. \times P_\pm(\rho\omega(\theta, \phi))z'(y, \rho\omega(\theta, \phi)) \right\} \chi(\pm 2^{-j}\lambda(\rho\omega(\theta, \phi))) \right]_0^\pi d\theta \\ - \frac{1}{ik} \int_0^{2\pi} \int_0^\pi e^{ik(1-\cos(\phi))} \partial_\phi \left\{ P_\pm(\rho\omega(\theta, \cdot))z^*(x, \rho\omega(\theta, \cdot)) \times \right. \\ \left. \times P_\pm(\rho\omega(\theta, \cdot))z'(y, \rho\omega(\theta, \cdot)) \chi(\pm 2^{-j}\lambda(\rho\omega(\theta, \cdot))) \right\}(\phi) d\phi d\theta.$$

The integrand of the first term can be rewritten in order to obtain a sum of two integral in ϕ over the interval $[0, \pi]$. To this end, we introduce a smooth cut-off function which splits $[0, \pi]$ in two parts one is a neighborhood of 0 and the other a neighborhood of π . then most of the terms obtained after derivation could be treated by the method used for the $|t|^{-1}$ decay. Only the two terms where derivatives of z appear need a particular treatment. Now we have to distinguish the case $z = z' = w$ from the two others where $z = u$ or $z' = u$. If $z = z' = w$, the terms which need a particular treatment are bounded by $C|t|^{-1}$ times the supremum in ϕ' of the $L_{\phi, \theta}^1([0, \pi] \times [0, 2\pi])$ of

$$L_{j,n,m}^\pm(t, x, y, \phi, \phi') = \int_{\mathbb{R}^+} e^{\mp it\sqrt{\rho^2+m^2}-i\rho|x-y|\cos(\phi')} \left\{ \left\{ P_\pm(\rho\omega) \partial_\phi^n z^*(x, \rho\omega) \right\} \times \right. \\ \left. \times \left\{ P_\pm(\rho\omega(\theta, \phi)) \partial_\phi^m z'(y, \rho\omega(\theta, \phi)) \right\} \chi(\pm 2^{-j}\lambda(\rho\omega(\theta, \phi))) \right\} \rho d\rho,$$

with $n, m \in \mathbb{N}$ such that $n + m = 1$. It is a sum of terms of the form

$$\int_{\mathbb{R}^+} e^{\mp it\left\{ \mp \sqrt{\rho^2+m^2} - \rho \frac{|x-y|}{t} (\cos(\phi') - \cos(\phi)) - \varepsilon_i \rho \frac{|x|}{t} + \varepsilon_{i'} \rho \frac{|y|}{t} \right\}} \left\{ \left\{ P_\pm(\rho\omega)_{k,i} \left(e^{-i\psi(k,x)} \partial_\phi^n z^*(x, \rho\omega) \right)_{i,l} \right\} \times \right. \\ \left. \times \left\{ P_\pm(\rho\omega(\theta, \phi))_{l,k'} \left(\partial_\phi^m z'(y, \rho\omega(\theta, \phi)) e^{i\psi(k,y)} \right)_{k',i'} \right\} \chi(\pm 2^{-j}\lambda(\rho\omega(\theta, \phi))) \right\} \rho d\rho.$$

where $\psi(x, k) \in \mathcal{M}_4(\mathbb{C})$ is given by

$$\begin{pmatrix} (|x||k| + x \cdot k) I_{\mathbb{C}^2} & 0_{\mathbb{C}^2} \\ 0_{\mathbb{C}^2} & (-|x||k| + x \cdot k) I_{\mathbb{C}^2} \end{pmatrix}.$$

For the sake of simplicity, we work with $(i; i') = (1; 1)$ the other cases can be treated in the same way. We introduce

$$K(\rho) = \left\{ P_\pm(\rho\omega)_{k,i} \left(e^{-i\psi(k,x)} \partial_\phi^n z^*(x, \rho\omega) \right)_{i,l} \right\} \left\{ P_\pm(\rho\omega(\theta, \phi))_{l,k'} \left(\partial_\phi^m z'(y, \rho\omega(\theta, \phi)) e^{i\psi(k,y)} \right)_{k',i'} \right\}, \\ \phi(\rho) = \mp \sqrt{\rho^2 + m^2} - \rho \frac{|x-y|}{t} (\cos(\phi') - \cos(\phi)) - \varepsilon_i \rho \frac{|x|}{t} + \varepsilon_{i'} \rho \frac{|y|}{t}$$

and

$$f(\rho) = \frac{\partial_\rho \phi(\rho)}{\inf_{x \in \text{supp}(\chi_j)} \{|\partial_\rho^2 \phi(|x|, \lambda)|\}}.$$

With help of a smooth cut-off function, we split the integral in two parts, one has support $\{t \in \mathbb{R}; |f(t)| \leq \alpha\}$ on which we use the estimate

$$|\{t \in \mathbb{R}; |f(t)| \leq \alpha\}| \leq \alpha,$$

the other is its complementary, in which we make an integration by parts. We obtain the estimate

$$\begin{aligned} |J_j^+[r, r'](t, x, y)| &\leq C\alpha \max_{\rho \in A_j} \{\rho |K(\rho)|\} \\ &+ \frac{1}{\alpha t \inf_{\rho \in A_j} \{|\partial_\rho^2 \phi(\rho)|\}} \max \left\{ \max_{\rho \in A_j} \{\rho |K(\rho)|\}; \int_{A_j} \{\rho |(\partial_\rho K)(\rho)|\}; \right. \\ &\quad \left. \int_{A_j} |K(\rho)| d\rho; 2^{-j} \int_{A_j} \{\rho |K(\rho)|\} \right\}, \end{aligned}$$

where $A_j = g^{-1}\{\text{supp}(\chi_j)\}$ with $g(\rho) = \sqrt{\rho^2 + m^2}$. Hence with (3.19) of Proposition 3.6, Proposition 3.8 and decay of derivatives of P_\pm , we can choose $\alpha = 2^{2j} \sqrt{t}^{-1}$ and we obtain the bound of (3.22) in this case.

For the case $(z, z') = (u, w)$ (the case $(z, z') = (w, u)$ is similar), we study by the same way the integral

$$\begin{aligned} \int_{\mathbb{R}^+} e^{\mp i t \left\{ \mp \sqrt{\rho^2 + m^2} - \rho \frac{|x-y|}{t} (\cos(\phi') - \cos(\phi)) - \varepsilon_i \rho \frac{\omega \cdot x}{t} + \varepsilon_{i'} \rho \frac{|y|}{t} \right\}} &\left\{ \left\{ P_\pm(\rho\omega)_{k,i} (\partial_\phi^n z^*(x, \rho\omega))_{i,l} \right\} \times \right. \\ &\times \left. \left\{ P_\pm(\rho\omega(\theta, \phi))_{l,k'} (\partial_\phi^m r'(y, \rho\omega(\theta, \phi)) e^{i\psi(k,y)})_{k',i'} \right\} \chi(\pm 2^{-j} \lambda(\rho\omega(\theta, \phi))) \right\} \rho d\rho. \end{aligned}$$

If $|x - y|/|t| \ll 1$, we can suppose $|x - y|/|t| < |\xi|/(2\sqrt{\xi^2 + m^2})$ for any $\xi \in \text{supp}(\chi_j)$ and instead of applying the trick of the proof of Lemma 3.1 (integration by parts with respect to an angular variables) to the integral $I_j^\pm[z, z'](t, x, y)$, we make an integration by parts with help of

$$\frac{1}{\frac{|\xi|}{\sqrt{\xi^2 + m^2}} \pm \frac{\xi}{|\xi|} \cdot \frac{x-y}{t}} \partial_{|\xi|}.$$

The rest of the proof is the same.

We now turn to the proof of estimate (3.23), the kernel of the operator is given by a sum of term of the form

$$I_j^\pm[z, z'](t, x, y) = \int_{\mathbb{R}^3} e^{\mp i t \sqrt{\xi^2 + m^2}} e^{-i \xi \cdot (x-y)} \{P_\pm(\xi) z^*(x, \xi) P_\pm(\xi) z'(y, \xi)\} \chi(\pm \lambda(\xi)) d\xi.$$

We first notice that Proposition 3.6 implies that this integral is bounded. Then we split the integral in two parts. One is supported in a small neighborhood of the critical point of the phase, the other is its complementary. To treat this last integral we work exactly like the case “ $\frac{|x-y|}{t} \ll 1$ ”, just mentioned above. For the other one applies the Morse lemma to reduced the study to

$$\begin{aligned} \int_{\mathbb{R}^3} e^{\mp i t \xi^2} \{P_\pm(f(\xi)) z^*(x, f(\xi)) P_\pm(f(\xi)) z'(y, f(\xi))\} \tilde{\chi}((f(\xi))) d\xi = \\ \int_{S^2} \int_{\mathbb{R}^+} \rho e^{\mp i t \rho^2} \{P_\pm(f(\rho\omega)) z^*(x, f(\rho\omega)) P_\pm(f(\rho\omega)) z'(y, f(\rho\omega))\} \tilde{\chi}((f(\rho\omega))) d\rho d\omega, \end{aligned}$$

where $\tilde{\chi}$ is the product of an indicator of a small neighborhood of the critical point with $\chi(\pm \lambda(\cdot))$. Then an integration by parts in ρ and the Van Der Corput lemma give (3.23) when $\theta = 1$. Since we have that the integral is bounded the general case easily follows.

We are now able to write the proof of Theorem 1.2, using Proposition 3.9.

Proof (Proof of Theorem 1.2). We notice that

$$\begin{aligned}\phi_k(D_m)\phi_j(H) &= D_m^{-1}\phi_k(D_m)H\phi_j(H) - D_m^{-1}\phi_k(D_m)V\phi_j(H) \\ &= 2^{-k}\tilde{\phi}_k(D_m)2^j\tilde{\phi}_j(H) - 2^{-k}\tilde{\phi}_k(D_m)V\phi_j(H)\end{aligned}$$

We can also use H^{-1} since the support of ϕ_j is far from 0

$$\phi_k(D_m)\phi_j(H) = D_m\phi_k(D_m)H^{-1}\phi_j(H) - \phi_k(H)VH^{-1}\phi_j(H)$$

or higher power in D_m^{-1} or H^{-1} to obtain with (3.23)

$$\|\phi_i(D_m)e^{-it(H)}\phi_j(H)\phi_k(D_m)\|_{L^1, L^\infty} \leq C2^{-r'|j-i|}\frac{C2^{(2+\theta)j}}{t^{1+\theta}}2^{-r|j-k|}$$

for any reals r, r' with C independent of i, j . Hence if $r, r' > 0$, we work like in the proof of Theorem 3.1 to conclude the proof.

It now remains to prove Proposition 3.6, 3.7 and 3.8.

3.3.3. Some estimates.

Estimates for w . We remind us of the definition of w in (3.17) and we introduce

$$\tilde{R}_V^\pm(k) = e^{-ik \cdot Q} R_V^\pm(\pm\lambda(k)) e^{ik \cdot Q}. \quad (3.24)$$

We have

Lemma 3.3. *For any $\alpha \in \mathbb{N}^3$, let be $\sigma > 4 + |\alpha|$. Then there exists $C > 0$ such that for any $k, x \in \mathbb{R}^3 \setminus \{0\}$*

$$\left| \left\{ \nabla_k^\alpha \tilde{R}_0^\pm(k) \langle Q \rangle^{-\sigma} q \right\} (x) \right| \leq \frac{C}{\langle k \rangle^{|\alpha|}} \frac{\langle x \rangle^{|\alpha|}}{\left\langle |x| \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle} \|q\|_{W^{2+|\alpha|}, \infty}. \quad (3.25)$$

We also have that there exists $C > 0$ such that for any $k, x \in \mathbb{R}^3 \setminus \{0\}$

$$\left| \left\{ \nabla_k^\alpha \tilde{R}_0^\pm(k) \langle Q \rangle^{-\sigma} q \right\} (x) \right| \leq C \frac{\langle x \rangle^{\alpha-1}}{\langle k \rangle^{\alpha-1}} \frac{\langle \min\{|x|, |k|\} \rangle}{\langle k \rangle \left\langle \min\{|x|, |k|\} \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle^2} \|q\|_{W^{2+|\alpha|}, \infty}. \quad (3.26)$$

Proof. We write

$$\begin{aligned}\left(\tilde{R}_0^\pm(k) \langle Q \rangle^{-\sigma} q \right) (x) &= \int_{\mathbb{R}^3} \frac{e^{i\{\pm|k||y|+k \cdot y\}}}{4\pi|y|} \left\{ \frac{\alpha \cdot (x-y)q(x-y)}{\langle x-y \rangle^{\sigma+2}} + \frac{\alpha \cdot \nabla q(x-y)}{\langle x-y \rangle^\sigma} \right\} dy \\ &\quad + (\alpha \cdot k + m\beta \pm \lambda(k)) \int_{\mathbb{R}^3} \frac{e^{i\{\pm|k||y|+k \cdot y\}}}{4\pi|y|} \frac{q(x-y)}{\langle x-y \rangle^\sigma} dy.\end{aligned}$$

We restrict our study to $\tilde{R}_0^+(k)$ since the two case are similar. Hence we only need to estimate an integral of the form

$$R(k)(x) = \int_{\mathbb{R}^3} e^{i\{|k||y|+k \cdot y\}} \frac{u(x-y)}{|y|} dy$$

with $u \in W_\sigma^{1+|\alpha|, \infty}(\mathbb{R}^3, \mathbb{C})$.

In a first step, a straightforward calculation shows that

$$|\nabla_k^\alpha R(k)(x)| \leq C \langle x \rangle^{|\alpha|-1} \|u\|_{L^\infty} \quad (3.27)$$

if $\sigma > 3 + \max\{|\alpha| - 1; 0\}$. Then using the trick we used in the proof of Proposition 3.2, we obtain

$$\nabla_k^\alpha R(k)(x) = \frac{i^{|\alpha|}}{|k|^{|\alpha|}} \int_{\mathbb{R}^3} e^{i\{|k||y|+k \cdot y\}} \left\{ (\nabla|Q|)^\alpha \frac{u(x-\cdot)}{|Q|} \right\} (y) dy,$$

and so with (3.27), we infer

$$|\nabla_k^\alpha R(k)(x)| \leq \frac{C \langle x \rangle^{|\alpha|-1}}{\langle k \rangle^{|\alpha|}} \|u\|_{W_\sigma^{|\alpha|, \infty}}, \quad (3.28)$$

since $\sigma > 3 + \max\{|\alpha| - 1, 0\}$.

In a second step, we apply Estimate (3.5) of Proposition 3.2 to $R(k)(x)$, this gives

$$|\nabla_k^\alpha R(k)(x)| \leq \frac{C}{|k|^{|\alpha|+1}} \max_{|\beta| \leq 1+|\alpha|} \left\{ \int_{\mathbb{R}^3} |y|^{|\beta|-1} \frac{1}{\left| \frac{k}{|k|} - \frac{y}{|y|} \right|} \left| \nabla^\beta \left\{ \frac{u(x-\cdot)}{|Q|} \right\} (y) \right| dy \right\}.$$

Hence we need to estimate on integral of the form

$$G(x, \omega) = \int_{\mathbb{R}^3} \frac{1}{|y|^n} \frac{1}{\langle x-y \rangle^s} \frac{1}{\left| \frac{y}{|y|} - \omega \right|} dy$$

with $\omega \in S^2$, $-|\alpha| + 1 \leq n \leq 2$ and $s > \sigma$. To obtain appropriate estimates, we use

$$|x-y| \geq \frac{1}{4} \max\{|y|, |x|\} \left| \frac{x}{|x|} - \frac{y}{|y|} \right| + \frac{1}{2} ||x| - |y||$$

to write for $\theta, \theta' \geq 0$ such that $\theta + \theta' = 1$,

$$\begin{aligned} G(x, \omega) &\leq \int_{\mathbb{R}^3} \frac{1}{|y|^n} \frac{1}{\langle |x| - |y| \rangle^{\theta s}} \frac{1}{\left\langle |x| \left| \frac{x}{|x|} - \frac{y}{|y|} \right| \right\rangle^{\theta' s}} \frac{1}{\left| \frac{y}{|y|} - \omega \right|} dy \\ &\leq \frac{C}{|x| \left\langle |x| \left| \omega - \frac{x}{|x|} \right| \right\rangle} \int_{\mathbb{R}^+} \frac{1}{r^{n-2}} \frac{1}{\langle |x| - r \rangle^{\theta s}} dr \\ &\leq \frac{C \langle x \rangle^{|\alpha|+1}}{|x| \left\langle |x| \left| \omega - \frac{x}{|x|} \right| \right\rangle}, \end{aligned}$$

if $\theta' s > 2$ and $\theta s > 1 + \max\{2 - n; 0\}$. Since $G(0, \omega)$ is bounded, we obtain

$$G(x, \omega) \leq \frac{C \langle x \rangle^{|\alpha|}}{\left\langle |x| \left| \omega - \frac{x}{|x|} \right| \right\rangle},$$

Hence, we obtain with estimate (3.28)

$$|\nabla_k^\alpha R(k)(x)| \leq \frac{C}{\langle k \rangle^{|\alpha|+1}} \frac{\langle x \rangle^{|\alpha|}}{\left\langle |x| \left| \frac{k}{|k|} - \frac{x}{|x|} \right| \right\rangle} \|u\|_{W_\sigma^{|\alpha|+1, \infty}}, \quad (3.29)$$

which gives estimate (3.25).

In a third step, if $k/|k| \neq x/|x|$, we split the integral for $\nabla_k^\alpha R(k)(x)$ in two parts with help of a smooth cut-off function defined in S^2 the support of which is a half cone determined by the bisector plane of the couple $(k/|k|; x/|x|)$. So we obtain $\nabla_k^\alpha R(k)(x) = R_1(k)(x) + R_2(k)(x)$ with $R_1(k)(x)$ having a support containing $x/|x|$ and $R_2(k)(x)$ having a support containing $k/|k|$. We then apply the estimate (3.6) of Proposition 3.2 to $R_1(k)(x)$ to obtain

$$|\nabla_k^\alpha R_1(k)(x)| \leq \frac{C}{|k|^{|\alpha|+2} \left| \frac{k}{|k|} - \frac{x}{|x|} \right|} \max_{|\beta| \leq 2+|\alpha|} \left\{ \int_{\mathbb{R}^3} |y|^{|\beta|-2} \frac{1}{\left| \frac{k}{|k|} - \frac{y}{|y|} \right|} \left| \nabla^\beta \left\{ \frac{u(x-\cdot)}{|Q|} \right\} (y) \right| dy \right\}.$$

This gives the estimate

$$|\nabla_k^\alpha R_1(k)(x)| \leq \frac{C}{|k|^{|\alpha|+2}} \frac{\langle x \rangle^{|\alpha|-1}}{\left| \frac{k}{|k|} - \frac{x}{|x|} \right|^2} \|u\|_{W_\sigma^{|\alpha|+2, \infty}},$$

since $\sigma > 2 + |\alpha|$. Using (3.29), we infer

$$|\nabla_k^\alpha R_1(k)(x)| \leq \frac{C}{\langle k \rangle^{|\alpha|+1}} \frac{\langle x \rangle^{|\alpha|}}{\langle \sqrt{|k||x|} \left| \frac{k}{|k|} - \frac{x}{|x|} \right| \rangle^2} \|u\|_{W_\sigma^{|\alpha|+2, \infty}},$$

or, using (3.28)

$$|\nabla_k^\alpha R_1(k)(x)| \leq \frac{C}{\langle k \rangle^{|\alpha|}} \frac{\langle x \rangle^{|\alpha|-1}}{\langle |k| \left| \frac{k}{|k|} - \frac{x}{|x|} \right| \rangle^2} \|u\|_{W_\sigma^{|\alpha|+2,\infty}}.$$

For $R_2(x)(k)$, we use the inequality $|x - y| \geq \frac{|x| \left| \frac{x}{|x|} - \frac{y}{|y|} \right|}{2}$ to obtain

$$R_2(k)(x) \leq \frac{C \langle x \rangle^{|\alpha|-1}}{\langle k \rangle^{|\alpha|} \langle |x| \left| \frac{k}{|k|} - \frac{x}{|x|} \right| \rangle^s} \|u\|_{W_\sigma^{|\alpha|,\infty}},$$

since $\sigma > 3 + s + \max\{|\alpha| - 1, 0\}$. So now we easily deduce estimate (3.26).

For the sequel, we need the following

Lemma 3.4. *Let be $s \in \mathbb{R}$ and ϕ a \mathcal{C}^∞ function such that there is $\sigma > 0$ with*

$$\forall \alpha \in \mathbb{N}^3, |\nabla^\alpha \phi(x)| \leq \frac{C_\alpha}{\langle x \rangle^\sigma}.$$

We have that $[\langle P \rangle^s, \phi(Q)]$ is bounded from H_q^t into $H_{q'}^{t'}$ with $q' + \sigma \geq q$ and $t' + 1 \geq t + s$.

Proof. We want to prove that

$$\langle Q \rangle^q \langle P \rangle^t [\langle P \rangle^s, \phi(Q)] \langle P \rangle^{-t'} \langle Q \rangle^{-q'} \quad (3.30)$$

is bounded in $\mathcal{B}(L^2)$. Using the identity

$$[\langle P \rangle^s, \phi(Q)] = [\langle P \rangle^{s/2}, \phi(Q)] \langle P \rangle^{s/2} + \langle P \rangle^{s/2} [\langle P \rangle^{s/2}, \phi(Q)]$$

we reduce the proof to the case $|s| < 1$. And with the identity

$$[\langle P \rangle^s, \phi(Q)] = -\langle P \rangle^s [\langle P \rangle^{-s}, \phi(Q)] \langle P \rangle^s$$

we only need to study the case $-1 < s < 0$. The proof in this case is based on the following identity for $-1 < s < 0$

$$\langle P \rangle^s = (-\Delta + 1)^{s/2} = \frac{-\sin(\pi \{\frac{s}{2}\})}{\pi} \int_0^{+\infty} \frac{w^{\{\frac{s}{2}\}}}{-\Delta + 1 + w} dw.$$

So we have

$$[\langle P \rangle^s, \phi(Q)] = \sum_{k=1}^m \frac{\Gamma(s/2 + 1)}{\Gamma(s/2 + 1 - k)} (-\Delta + 1)^{s/2-k} Ad_{-\Delta+1}^k(\phi(Q)) + R_m$$

with

$$R_m = \frac{(-1)^m \sin(\pi \{\frac{s}{2}\})}{\pi} \int_0^{+\infty} \frac{w^{\{\frac{s}{2}\}}}{(-\Delta + 1 + w)^{m+1}} Ad_{-\Delta+1}^{m+1}(\phi(Q)) \frac{dw}{-\Delta + 1 + w}.$$

Then we use $\frac{-\Delta+1}{-\Delta+1+w} = 1 - \frac{w}{-\Delta+1+w}$, and we commute powers of $\langle P \rangle$ with operators of the form $\nabla^\alpha \phi(Q)$. Hence we can repeat the previous computation until we obtain only non positive powers of $\langle P \rangle$ in (3.30). So we only need to prove that operators of the form

$$[\langle Q \rangle^q, \phi(P)] \langle Q \rangle^{-q'}$$

with $q \leq q' + 1$ and ϕ satisfying the assumption of the lemma are bounded in $\mathcal{B}(L^2)$, we just repeat the previous calculation but we switch the role of P and Q . This ends the proof.

We now state a particular version of the Limiting Absorption Principle for H .

Proposition 3.10. *We assume that Assumptions 1.1 and 1.2 hold. Then for any $\sigma \geq 1$ there exists $C > 0$ such that for any $k \in \mathbb{R}^3$*

$$\|\tilde{R}_V^\pm(k)\|_{\mathcal{B}(L_\sigma^2, L_{-\sigma}^2)} \leq C.$$

Proof. In fact, we just need to prove that for any $\sigma \geq 1$ there exists $C > 0$ such that for any $\lambda \in \mathbb{R} \setminus (-m, m)$

$$\|R_V^+(\lambda)\|_{\mathcal{B}(L_\sigma^2, L_{-\sigma}^2)} \leq C.$$

Using Theorem 1.1, we have that it is true if $\sigma > 5/2$. Then we use Born expansion

$$R_V^+(\lambda) = R_0^+(\lambda) - R_0^+(\lambda)V R_0^+(\lambda) + R_0^+(\lambda)V R_V^+(\lambda)V R_0^+(\lambda)$$

and [IM99, Theorem 2.1(i)] to end the proof.

We are now able to give the

Proof (Proof of Proposition 3.6). We only give a the general idea of the proof and we leave the details to the reader. We notice that with \tilde{R}_V^\pm defined by (3.24), we obtain

$$w = \tilde{R}_V V u$$

with an abuse of notation since we avoid to distinguish the case where we have \tilde{R}_V^+ or \tilde{R}_V^- . We recall the identities

$$\tilde{R}_V V = \tilde{R}_0 V - \tilde{R}_0 V \tilde{R}_V V = \tilde{R}_0 V - \tilde{R}_0 V \tilde{R}_0 + \tilde{R}_0 V \tilde{R}_V V \tilde{R}_0 V. \quad (3.31)$$

Since, we have

$$\tilde{R}_V = (1 + \tilde{R}_0 V)^{-1} \tilde{R}_0, \quad (1 + \tilde{R}_0 V)^{-1} = 1 - \tilde{R}_V V,$$

for $|\alpha| = 1$, we obtain

$$\nabla^\alpha \tilde{R}_V = (1 - \tilde{R}_V V) \nabla^\alpha \tilde{R}_0 (1 - V \tilde{R}_V). \quad (3.32)$$

Using (3.32), we obtain a formula where only derivatives of \tilde{R}_0 appear (if there is derivatives). Then between a derivative of \tilde{R}_0 and \tilde{R}_V , we insert a \tilde{R}_0 with the identity (3.31):

$$\tilde{R}_V V \nabla^\alpha \tilde{R}_0 V = \tilde{R}_0 V \nabla^\alpha \tilde{R}_0 V - \tilde{R}_0 V \tilde{R}_0 V \nabla^\alpha \tilde{R}_0 V + \tilde{R}_0 V \tilde{R}_V V \tilde{R}_0 V \nabla^\alpha \tilde{R}_0 V.$$

This ensures that if $\rho > 5$, V or its derivatives decays enough to use Estimate (3.25) and Proposition 3.10. Since these estimates need derivatives and Sobolev's injections, we apply Lemma 3.4 to conclude the proof.

Estimates for v . We remind us of the definition of v in (3.20) and we introduce

$$S_V^{\varepsilon_1, \varepsilon_2}(k) = e^{-\varepsilon_1 \varepsilon_2 i |k| |Q|} R_V^{\varepsilon_1}(\varepsilon_2 \lambda(k)) e^{ik \cdot Q},$$

where $\varepsilon_i \in \{-1, 1\}$. With an abuse of notation, we will write $v = S_V V u$. We have

Lemma 3.5. *There exists $C > 0$, such that for any $k \in \mathbb{R}^3 \setminus \{0\}$ and $\beta \in \mathbb{N}^3$*

$$\left| \left(\nabla_k^\beta S_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q \right) (x) \right| \leq \frac{C}{\left\langle |x| \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle} \|q\|_{W^{2+|\beta|}, \infty}$$

for any $\sigma > 3 + |\beta|$.

Proof.

$$\begin{aligned} (S_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q) (x) &= \int_{\mathbb{R}^3} \frac{e^{i\varepsilon_1 \varepsilon_2 \{|k||x-y| - |k||x| + \varepsilon_1 \varepsilon_2 k \cdot y\}}}{4\pi|x-y|} \left\{ \frac{\alpha \cdot y q(y)}{\langle y \rangle^{\sigma+2}} + \frac{\alpha \cdot \nabla q(y)}{\langle y \rangle^\sigma} \right\} dy \\ &\quad + (\alpha \cdot k + m\beta \pm \lambda(k)) \int_{\mathbb{R}^3} \frac{e^{i\varepsilon_1 \varepsilon_2 \{|k||x-y| - |k||x| + \varepsilon_1 \varepsilon_2 k \cdot y\}}}{4\pi|x-y|} \frac{q(y)}{\langle y \rangle^\sigma} dy. \end{aligned}$$

For the sake of simplicity, we only write the proof when $\beta = 0$. The proof for derivatives works in the same way using $||x-y| - |x|| \leq |y|$ and $\sigma > 3 + |\beta|$. But the proof for the case $\beta = 0$, has been already done since

$$|(S_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q) (x)| = |(R_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q) (x)|.$$

Hence using Proposition 3.10, we able to write the

Proof (Proof of Proposition 3.7). We write with an abuse of notation

$$v = S_V V u,$$

and we use the Born formula

$$S_V V = S_0 V - S_0 V \tilde{R}_V V,$$

together with Lemma 3.5, Propositions 3.4 and 3.10. The proof works like the one for w .

Estimates for \tilde{v} : We remind us of the definition of \tilde{v} in (3.21) and we introduce

$$T_V^{\varepsilon_1, \varepsilon_2}(k) = e^{-\varepsilon_1 \varepsilon_2 i |k| |Q| + i k \cdot Q} \nabla_k \tilde{R}_V^{\varepsilon_1}(\varepsilon_2 \lambda(k)) e^{i k \cdot Q},$$

where $\varepsilon_i \in \{-1, 1\}$. With another abuse of notation, here we will write $\tilde{v} = T_V V u$. We have

Lemma 3.6. *There exists $C > 0$, such that for any $k \in \mathbb{R}^3 \setminus \{0\}$ and $\beta \in \mathbb{N}^3$*

$$\left| \left(\nabla_k^\beta T_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q \right) (x) \right| \leq C \frac{\langle \min\{|x|, |k|\} \rangle}{\langle k \rangle \left\langle \min\{|x|, |k|\} \left| \frac{x}{|x|} - \frac{k}{|k|} \right| \right\rangle^2} \|q\|_{W^{2+|\beta|, \infty}},$$

for any $\sigma > 4 + |\beta|$.

Proof. This is an obvious adaptation of the proof of Lemma 3.5, we just notice that one has

$$\left| \left(T_0^{\varepsilon_1, \varepsilon_2}(k) \langle Q \rangle^{-\sigma} q \right) (x) \right| = \left| \left(\nabla_k \tilde{R}_V^{\varepsilon_1}(\varepsilon_2 \lambda(k)) \langle Q \rangle^{-\sigma} q \right) (x) \right|.$$

Hence, we have

Proof (Proof of Proposition 3.8). One more time, we write with an abuse of notation

$$v = T_V V u + S_V V \nabla_k u,$$

The second term of the right hand side could be studied exactly as we done in proof of Proposition 3.7 and for the first one we use the formula

$$T_V V = T_0 V - T_0 V \tilde{R}_V V + S_0 V \nabla_k \tilde{R}_V V,$$

together with Lemma 3.6, Propositions 3.4 and 3.10. The proof works like the one for w .

4. The linearized operator

In this section, we study the spectral properties of the linearized operator, associated with Equation (1.2), around a stationary state. This will be useful since we compare the dynamics associated with Equation (1.2) to the dynamic of the linear Dirac equation associated with H . This comparison is possible only because when the PLS is small, the linearized operator is a small perturbation of H .

4.1. The manifold of the particle like solutions. First we notice that Proposition 1.1, which gives the existence of stationary states, is a consequence of

Proposition 4.1. *Let H be a self adjoint operator on $L^2(\mathbb{R}^3, \mathbb{C}^4)$ and with a simple eigenvalue λ_0 associated with a normalized eigenvector ϕ_0 . Assume that there is a neighborhood $\mathcal{O} \subset \mathbb{R}$ of λ_0 such that for all $\lambda \in \mathcal{O}$ the operator $(H - \lambda)^{-1} P_0$ is in $\mathcal{B}(L_\sigma^2(\mathbb{R}^3, \mathbb{C}^4))$, for any $\sigma \in \mathbb{R}^+$, and $\mathcal{B}(H^l(\mathbb{R}^3, \mathbb{C}^4), H^{l+1}(\mathbb{R}^3, \mathbb{C}^4))$, for any $l \in \mathbb{N}$, and where P_0 is the projector into the orthogonal space of ϕ_0 . Let be $F \in \mathcal{C}^{k+1}(\mathbb{C}^4, \mathbb{C}^4)$ such that $F(z) = O(|z|^3)$.*

Then for any $\sigma \in \mathbb{R}^+$, there exists Ω a neighborhood of $0 \in \mathbb{C}$, a \mathcal{C}^k map

$$h : \Omega \mapsto \{\phi_0\}^\perp \cap H^2(\mathbb{R}^3, \mathbb{C}^4) \cap L_\sigma^2(\mathbb{R}^3, \mathbb{C}^4)$$

and a \mathcal{C}^k map $E : \Omega \mapsto \mathbb{R}$ such that $S(u) = u\phi_0 + h(u)$ satisfy for all $u \in \Omega$,

$$HS(u) + \nabla F(S(u)) = E(u)S(u),$$

with the following properties

$$\begin{cases} h(e^{i\theta} u) = e^{i\theta} h(u), & \forall \theta \in \mathbb{R}, \\ h(u) = O(|u|^2), \\ E(u) = E(|u|), \\ E(u) = \lambda_0 + O(|u|^2). \end{cases}$$

The proof of this proposition is an obvious adaptation of the one of [PW97, Proposition 2.2], and we don't repeat it here. One can also obtain it by means of the Crandall-Rabinowitz theorem but it doesn't give immediately the decomposition associated to the spectrum of $H = D_m + V$.

To show that $(H - \lambda)^{-1}P_0$ is in $\mathcal{B}(L^2_\sigma(\mathbb{R}^3, \mathbb{C}^4))$ for any $\sigma > 0$, we just need to prove that $\alpha \mapsto e^{\alpha\langle Q \rangle}(H - \lambda)^{-1}P_0 e^{-\alpha\langle Q \rangle}$ is of class \mathcal{C}^k near 0 in $\mathcal{B}(L^2(\mathbb{R}^3, \mathbb{C}^4))$ for any $k \in \mathbb{N}$, this can be proved with help of [His00, Lemma 5.1]. To prove that $(H - \lambda)^{-1}P_0$ for any $l \in \mathbb{N}$ is in $\mathcal{B}(H^l(\mathbb{R}^3, \mathbb{C}^4), H^{l+1}(\mathbb{R}^3, \mathbb{C}^4))$ for any $l \in \mathbb{N}$, we first notice that $(D_m - \lambda)^{-1}$ is in $\mathcal{B}(H^l(\mathbb{R}^3, \mathbb{C}^4), H^{l+1}(\mathbb{R}^3, \mathbb{C}^4))$ then we use wave operator (see [GM01, Theorem 1.5]) and the intertwining property (see (2.5)) to conclude.

We shall need some properties of stationary solutions of (1.2). We have the

Lemma 4.1 (exponential decay). *Let $l \in \mathbb{N}$, $s \in \mathbb{R}^+$ and $p, q \in [1, \infty]$ there is $\gamma > 0$, $\varepsilon > 0$ and $C > 0$ such that for all $u \in B_{\mathbb{C}}(0, \varepsilon)$ one has*

$$\|e^{\gamma\langle Q \rangle} d^l S(u)\|_{B_{p,q}^s} \leq C \|S(u)\|_2.$$

Proof. In fact we prove that for any k in \mathbb{N} there is $\gamma > 0$ and $\varepsilon > 0$ and $C > 0$ such that for all $u \in B_{\mathbb{C}}(0, \varepsilon)$ one has

$$\|e^{\gamma\langle Q \rangle} d^l S(u)\|_{H^k} \leq C \|S(u)\|_2.$$

then interpolation and the following property of Besov spaces over \mathbb{R}^3 permit to conclude: $B_{2,2}^s = H^s$, $B_{p,r}^s \subset B_{p,q}^{s'}$ if $s' < s$, $B_{r,q}^u \subset B_{p,q}^s$ if $1 \leq r \leq p \leq \infty$ and $u - n/r = s - n/p$ and $\|uv\|_{B_{p,q}^s} \leq C \|u\|_{B_{q,t}^s} \|v\|_{B_{r,t}^s}$ if $\frac{1}{p} + \frac{s}{3} > \frac{1}{q} + \frac{1}{r}$.

We only make the proof for $l = 0$, the other cases are similar. We have

$$D_m S(u) + V S(u) + \nabla F(S(u)) = E(u) S(u).$$

let us write

$$W S(u) = \nabla F(S(u)) + V S(u).$$

We have that W decays and is in L^1 ; we can write $W = W_c + W_\delta$ where W_c is compactly supported and $\|W_\delta\|_{L^1 \cap L^\infty} \leq \delta$.

We have that $D_m + W_\delta - E(u)$ is invertible for δ sufficiently small and

$$e^{\gamma\langle Q \rangle} S(u) = e^{\gamma\langle Q \rangle} (D_m + W_\delta - E(u))^{-1} e^{-\gamma\langle Q \rangle} \{e^{\gamma\langle Q \rangle} W_c S(u)\}.$$

For γ small, $\left(D_m + \gamma \frac{\alpha \cdot Q}{\langle Q \rangle} + W_\delta - E(u)\right)$ is invertible in L^2 and

$$e^{\gamma\langle Q \rangle} S(u) = \left(D_m + \gamma \frac{\alpha \cdot Q}{\langle Q \rangle} + W_\delta - E(u)\right)^{-1} e^{\gamma\langle Q \rangle} W_c S(u).$$

This proves the lemma for $k = 0$ since $e^{\gamma\langle Q \rangle} W_c$ is bounded. Now we notice that

$$\begin{aligned} |P| \left(D_m + \gamma \frac{\alpha \cdot Q}{\langle Q \rangle} + W_\delta - E(u)\right)^{-1} \\ = \frac{|P|}{D_m} - \frac{|P|}{D_m} \left(2\gamma \frac{\alpha \cdot Q}{\langle Q \rangle} + W_\delta - E(u)\right) \left(D_m + \gamma \frac{\alpha \cdot Q}{\langle Q \rangle} + W_\delta - E(u)\right)^{-1}. \end{aligned}$$

Hence we obtain

$$\|e^{\gamma\langle Q \rangle} S(u)\|_{H^k} \leq C \|S(u)\|_{H^{k-1}}.$$

This identity proves the lemma by induction.

4.2. *The spectrum of the linearized operator.* Here we study the spectrum of the linearized operator associated with Equation (1.2) around a stationary state $S(u)$. Let us introduce

$$H(u) = H + d^2F(S(u)) - E(u)$$

where d^2F is the differential of ∇F . The operator $H(u)$ is \mathbb{R} -linear but not \mathbb{C} -linear. Replacing $L^2(\mathbb{R}^3, \mathbb{C}^4)$ by $L^2(\mathbb{R}^3, \mathbb{R}^4 \times \mathbb{R}^4)$ with the inner product obtained by taking the real part of the inner product of $L^2(\mathbb{R}^3, \mathbb{C}^4)$, we obtain a symmetric operator. We then complexify this real Hilbert space and obtain $L^2(\mathbb{R}^3, \mathbb{C}^4 \times \mathbb{C}^4)$ with its canonical hermitian product. This process transforms the operator $-i$ into

$$J = \begin{pmatrix} 0 & -Id_{\mathbb{C}^4} \\ Id_{\mathbb{C}^4} & 0 \end{pmatrix}.$$

For $\phi \in L^2(\mathbb{R}^3, \mathbb{R}^4 \times \mathbb{R}^4) \subset L^2(\mathbb{R}^3, \mathbb{C}^4 \times \mathbb{C}^4)$, we still write ϕ instead of

$$\begin{pmatrix} \Re \phi \\ \Im \phi \end{pmatrix}.$$

The extension of $H(u)$ over $L^2(\mathbb{R}^3, \mathbb{C}^4 \times \mathbb{C}^4)$ is also written $H(u)$ and is now a real operator.

The linearized operator associated with Equation (1.2) around the stationary state $S(u)$ is given by $JH(u)$. We shall now study its spectrum. Differentiating (1.3), we have that

$$\mathcal{H}_0 = \text{Span} \left\{ \frac{\partial}{\partial \Re u} S(u), \frac{\partial}{\partial \Im u} S(u) \right\}$$

is stable under the action of $JH(u)$. We notice (see [GNT04]) that

$$\mathcal{H}_0(u) = \text{Span} \{ JS(u), \partial_{|u|} S(u) \}. \quad (4.1)$$

Using gauge invariance and differentiating, we obtain

$$JH(u)JS(u) = 0 \quad \text{and} \quad JH(u)\partial_{|u|} S(u) = \partial_{|u|} E(u)JS(u).$$

Hence $\mathcal{H}_0(u)$ is contained in the geometric null space of $JH(u)$, in fact it is exactly the geometric null space as proved in the sequel of this subsection. First, we show that $JH(u)$ has two other simple eigenvalues, as stated in the following

Lemma 4.2. *Let be*

$$S_1^+(0) = \begin{pmatrix} \phi_1 \\ -i\phi_1 \end{pmatrix} \quad \text{and} \quad S_1^-(0) = \begin{pmatrix} \overline{\phi_1} \\ i\phi_1 \end{pmatrix}.$$

Suppose that Assumptions 1.1–1.4 hold, then there are $\varepsilon > 0$ and four $\mathcal{C}^{[r]-1}$ maps $E_1^\pm : B_{\mathbb{C}}(0, \varepsilon) \mapsto \mathbb{C}$ and $k_1^\pm : B_{\mathbb{C}}(0, \varepsilon) \mapsto \{S_1^\pm(0)\}^\perp$ such that

$$JH(u)S_1^\pm(u) = E_1^\pm(u)S_1^\pm(u),$$

with $\|S_1^\pm(u)\| = 1$,

$$S_1^\pm(u) = S_1^\pm(0) + k_1^\pm(u),$$

with $E_1^\pm(u) = \pm i(\lambda_1 - \lambda_0) + O(|u|^2)$ and $k_1^\pm(0) = 0$.

Proof. This can be proved in the same fashion as [PW97, Proposition 2.2].

We also obtain

Lemma 4.3 (exponential decay in Besov spaces). *Suppose that Assumptions 1.1–1.4 hold, then for any $l \in \mathbb{N}$, $s \in \mathbb{R}$ and $p, q \in [1, \infty]$ there is $\gamma > 0$, $\varepsilon > 0$ and a positive constant C such that for all $u \in B_{\mathbb{C}}(0, \varepsilon)$,*

$$\|e^{\gamma \langle Q \rangle} d^l S_1^\pm(u)\|_{B_{p,q}^s} \leq C \|S_1^\pm(u)\|_2.$$

Proof. The proof is exactly the same as the one of Lemma 4.1.

Let $\mathcal{H}_{\pm 1}(u)$ be the space spanned by $S_1^\pm(u)$. Let us now prove that the orthogonal space with respect to the hermitian product associated to J

$$\mathcal{H}_c(u) = \{\mathcal{H}_0(u) \oplus \mathcal{H}_{+1}(u) \oplus \mathcal{H}_{-1}(u)\}^\perp$$

contains no eigenvector. We notice that $\mathcal{H}_c(u)$ is stable under the action of $JH(u)$. We have

Lemma 4.4 (Continuous subspace property). *Suppose that Assumptions 1.1–1.4 hold, let $\mathbf{P}_c(u)$ be the orthogonal projector onto $\mathcal{H}_c(u)$. Then there exists $\varepsilon > 0$ such that for $u', u \in B_{\mathbb{C}}(0, \varepsilon)$*

$$\mathbf{P}_c((u))|_{\mathcal{H}_c(u')} : \mathcal{H}_c(u') \mapsto \mathcal{H}_c(u)$$

is an isomorphism from $B_{p,q}^s(\mathbb{R}^3, \mathbb{C}^8) \cap \mathcal{H}_c(u')$ into $B_{p,q}^s(\mathbb{R}^3, \mathbb{C}^8) \cap \mathcal{H}_c(u)$, for any $s \in \mathbb{R}^+$ and any $p, q \in [1, \infty]$. The inverse $R(u', u)$ is continuous with respect to u and u' .

Proof. This proof is a straightforward adaptation of the one of [GNT04, Lemma 2.2].

So we have

Lemma 4.5. *Under the assumptions of Proposition 1.1, there exists $\varepsilon > 0$ such that for any $u \in B_{\mathbb{C}}(0, \varepsilon)$ and any $\psi \in L_{\sigma}^2$,*

$$\|\langle Q \rangle^{-\sigma} e^{sJH(u)} \mathbf{P}_c(u) \psi\| \leq C \langle s \rangle^{-\min\{\sigma, 3/2\}} \|\langle Q \rangle^{\sigma} \psi\|.$$

As a consequence $\mathcal{H}_c(u)$ does not contain any eigenvector.

Proof. For the sake of clarity, we introduce

$$\zeta(u) = \left(J \frac{\partial}{\partial \Re u} S(u), J \frac{\partial}{\partial \Im u} S(u), JS_1^+(u), JS_1^-(u) \right).$$

Writing Duhamel's formula for $H(u)$ with respect to $H - E(u)$, we obtain

$$e^{tJH(u)} \mathbf{P}_c(u) = e^{tJ(H-E(u))} \mathbf{P}_c(u) + \int_0^t e^{(t-s)J(D_m+V-E(u))} J d^2 F(S(u)) e^{sJH(u)} \mathbf{P}_c(u) ds.$$

We have

$$\mathbf{P}_c(0)|_{\mathcal{H}_c(u')}^{-1} = R(u', 0) = Id_{L^2} + \sum_i |\alpha_i(u', 0)\rangle \langle \zeta_i(0)|$$

where the coordinates of $\alpha_i(u', u)$ are linear combination of the coordinates of $\zeta(u)$, so it can be extended to $L_{-\sigma}^2$ and we have

$$\begin{aligned} & \|\langle Q \rangle^{-\sigma} e^{-tJH(u)} \mathbf{P}_c(u) \psi\| \\ & \leq \|\langle Q \rangle^{-\sigma} \mathbf{P}_c(0)|_{\mathcal{H}_c(u')}^{-1} \langle Q \rangle^{\sigma}\| \left\{ \|\langle Q \rangle^{-\sigma} \mathbf{P}_c(0) e^{-tJ(D_m+V-E(u))} \mathbf{P}_c(u) \psi\| \right. \\ & \quad \left. + \int_0^t \|\langle Q \rangle^{-\sigma} \mathbf{P}_c(0) e^{-J(t-s)(D_m+V-E(u))} J D \nabla F(S(u)) e^{-sJH(u)} \mathbf{P}_c(u) \psi\| ds \right\} \\ & \leq C \langle t \rangle^{-\min\{\sigma, 3/2\}} \|\langle Q \rangle^{\sigma} \psi\| \\ & \quad + C \int_0^t \langle t-s \rangle^{-\min\{\sigma, 3/2\}} \|\langle Q \rangle^{2\sigma} N(u)\| \times \|\langle Q \rangle^{-\sigma} e^{-isH(u)} \mathbf{P}_c(u) \psi\| ds \end{aligned}$$

We then introduce

$$M(t) = \sup_{s \in [0, t]} \{ \langle s \rangle^{-\min\{\sigma, 3/2\}} \|\langle Q \rangle^{-\sigma} e^{-sJH(u)} \mathbf{P}_c(u) \psi\| \}$$

and we obtain for $|z| \leq \varepsilon$

$$M(t) \leq C(\|\langle Q \rangle^{\sigma} \psi\| + \varepsilon M(t))$$

which gives for ε sufficiently small

$$M(t) \leq C \|\langle Q \rangle^{\sigma} \psi\|,$$

or

$$\|\langle Q \rangle^{-\sigma} e^{-sJH(u)} \mathbf{P}_c(u) \psi\| \leq C \langle s \rangle^{-\min\{\sigma, 3/2\}} \|\langle Q \rangle^{\sigma} \psi\|.$$

Since an eigenvector is necessarily in L_{σ}^2 , we obtain that there is no stationary state in the range of $\mathbf{P}_c(u)$ that is to say $\mathcal{H}_c(u)$.

This gives

Lemma 4.6. *We have for sufficiently small $u \in \mathbb{C}$ $E_1^{\pm}(u) \in i\mathbb{R}$ and $S_1^-(u) = \overline{S_1^+(u)}$ for the conjugation of \mathbb{C}^8 .*

Proof. We specify the essential spectrum of $JH(u)$. A straightforward study gives that the continuous spectrum of $JH(0)$ is given by

$$\{i\lambda; \lambda \in \mathbb{R}, |\lambda| \geq \min\{|m - \lambda_0|, |m + \lambda_0|\}\}.$$

Using Weyl's criterion (see [RS78, Theorem XIII.14, Corollary 1], the adaptation is quite easy in our case), we obtain that for sufficiently small u the essential spectrum is

$$\{i\lambda; \lambda \in \mathbb{R}, |\lambda| \geq \min\{|m - E|, |m + E|\}\}.$$

Hence $E_1^\pm(u)$ are necessarily purely imaginary. Indeed if $H(u) - E_1^\pm(u)J$ is not invertible then $H(u) + \overline{E_1^\pm(u)}J$ is not invertible too. Since $-\overline{E_1^\pm(u)}$ is not in the essential spectrum, it is necessarily an eigenvalue in the neighborhood of $\pm i(\lambda_1 - \lambda_0)$. Hence this gives $-\overline{E_1^\pm(u)} = E_1^\pm(u)$.

4.3. Decomposition of the system. We want to decompose a solution ϕ of the equation (1.2) with respect to the spectrum of $JH(u)$. And in fact, we only study the resulting equations for these different parts. First we isolate a part which corresponds to a PLS. For any solution of (1.2) over an interval of time I , we write for $t \in I$

$$\phi(t) = e^{-i \int_0^t E(u(s)) ds} (S(u(t)) + \eta(t)).$$

In order to give an equation for η , we introduce the following space

$$\mathcal{H}_0^\perp(u) = \left\{ \eta \in L^2(\mathbb{R}^3, \mathbb{C}^8), \left\langle J\eta, \frac{\partial}{\partial \Re u} S(u) \right\rangle = 0, \left\langle J\eta, \frac{\partial}{\partial \Im u} S(u) \right\rangle = 0 \right\}.$$

In fact it is the space

$$\mathcal{H}_{+1}(u) \oplus \mathcal{H}_{-1}(u) \oplus \mathcal{H}_c(u)$$

which is stable under the action of $JH(u)$ and we state the

Lemma 4.7 (decomposition lemma). *Let be $s \geq 0$ and $p \geq 1$ there exist $\delta > 0$ and a \mathcal{C}^∞ map $U : B_{W^{s,p}}(0, \delta) \mapsto B_{\mathbb{C}}(0, \varepsilon)$ which satisfies for $\psi \in B_{W^{s,p}}(0, \delta)$*

$$\psi = S(u) + \eta, \text{ with } \eta \in \mathcal{H}_0^\perp(u) \iff u = U(\psi)$$

Proof. It is [GNT04, Lemma 2.3].

This lemma ensures that we can impose the orthogonality condition

$$\eta(t) \in \mathcal{H}_0^\perp(u(t)). \quad (4.2)$$

So instead of solving the Equation (1.2) in ϕ , we want to solve the equation

$$\begin{aligned} i\partial_t \eta &= \{H - E(u)\} \eta + \{\nabla F(S(u) + \eta) - \nabla F(S(u))\} - i dS(u) \dot{u} \\ &= \{H + d^2 F(S(u)) - E(u)\} \eta + N(u, \eta) - i dS(u) \dot{u} \end{aligned} \quad (4.3)$$

for $\eta \in \mathcal{H}_0^\perp(u(t))$. Here $d^2 F$ is the differential of ∇F and dS the differential of S in \mathbb{R}^2 . To close the system, we need an equation for u . Let us now derive an equation for the path u , by means of (4.2):

$$\langle \eta(t), JdS(u(t)) \rangle = 0.$$

After a time derivation, we obtain

$$0 = \langle JH(u(t))\eta(t) + JN(S(u(t)), \eta(t)) + dS(u(t))\dot{u}(t), JdS(u(t)) \rangle + \langle \eta, Jd^2 S(u(t))\dot{u}(t) \rangle.$$

Since $S(u) \in J\mathcal{H}_0(u)$, we have

$$\langle H(u)\eta, dS(u) \rangle = \langle \eta, H(u)dS(u) \rangle = \langle \eta, dE(u)S(u) \rangle = 0,$$

or

$$[\langle JdS(u(t)), dS(u(t)) \rangle + \langle J\eta(t), d^2 S(u(t)) \rangle] \dot{u}(t) = -\langle N(u(t), \eta(t)), dS(u(t)) \rangle.$$

So we notice that

$$[\langle JdS(u(t)), dS(u(t)) \rangle + \langle J\eta(t), d^2 S(u(t)) \rangle] = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} + O(|u(t)| + \|\eta(t)\|_2),$$

which proves that $[\langle JdS(u(t)), dS(u(t)) \rangle + \langle J\eta(t), d^2S(u(t)) \rangle]$ is invertible for small $|u(t)|$ and $\|\eta(t)\|_2$, we therefore introduce its inverse

$$A(u, \eta) = [\langle JdS(u), dS(u) \rangle + \langle J\eta, d^2S(u) \rangle]^{-1}$$

and write

$$\partial_t u(t) = -A(u(t), \eta(t)) \langle N(u(t), \eta(t)), dS(u(t)) \rangle.$$

Plugging in Equation (4.3), and similarly to the linear case we decompose η with respect to the spectral decomposition of $H(u) = H + D\nabla F(S(u)) - E(u)$

$$\eta(t) = \alpha^+(t)S_1^+(u) + \alpha^-(t)S_1^-(u) + z(t)$$

with $z \in \mathcal{H}_c(u) \cap L^2(\mathbb{R}^3, \mathbb{R}^8)$ and $\alpha^- = \overline{\alpha^+}$. we obtain the system

$$\begin{cases} \dot{u} = -A(u, \eta) \langle N(u, \eta), dS(u) \rangle \\ \dot{\alpha}^\pm = E^\pm(u) \alpha^\pm + \langle JN(u, \eta), S_1^\pm(u) \rangle \\ \quad + \langle dS(u) A(u, \eta) \langle N(u, \eta), dS(u) \rangle S_1^\pm(u) \\ \quad - \langle (dS_1^\pm(u)) A(u, \eta) \langle N(u, \eta), dS(u) \rangle, S_1^\pm(u) \rangle \alpha^\pm, \\ \partial_t z = JH(u)z + \mathbf{P}_c(u) JN(u, \eta) \\ \quad + \mathbf{P}_c(u) dS(u) A(u, \eta) \langle N(u, \eta), dS(u) \rangle \\ \quad - (D\mathbf{P}_c(u)) A(u, \eta) \langle N(u, \eta), dS(u) \rangle z \end{cases},$$

which we will now study. We notice that this equation is defined only for z small with real values, $\alpha^- = \overline{\alpha^+}$ small and u small.

5. The stabilization towards the PLS manifold

We now build a solution which stabilizes toward the manifold of the stationary states. To this end, we will use Theorem 1.1 and Theorem 1.2 to prove that z tends to zero in L^∞ and L_{loc}^2 . But we think that it will be possible only if α^+ and α^- also tends to zero. It is not always true and here we build solutions for which we ensure this fact. We also notice that we look for a real solution $\phi = S(u) + \eta$, hence η should be real and therefore $\alpha^- = \overline{\alpha^+}$.

We impose the following condition

$$\begin{cases} \lim_{t \rightarrow +\infty} u(t) = u_\infty, \\ |u(t) - u_\infty| \leq \frac{C}{\langle t \rangle^2}, \\ |\alpha| \leq \frac{C}{\langle t \rangle^2}. \end{cases}$$

Under the assumptions of Theorem 1.3, let us define for any $\varepsilon, \delta > 0$

$$\mathcal{U}(\varepsilon, \delta) = \left\{ u \in \mathcal{C}^{k+1}(\mathbb{R}, B_{\mathbb{C}}(0, \varepsilon)), \lim_{t \rightarrow +\infty} u(t) = u_\infty \text{ exists, } |u(t) - u_\infty| \leq \frac{\delta}{\langle t \rangle^2}, \forall t \geq 0 \right\}$$

and for any $u \in \mathcal{U}(\varepsilon)$, let s, s', β be such that $s' \geq s + 3 \geq \beta + 6$ and $\sigma > 3$, we define

$$\mathcal{Z}(u, \delta) = \left\{ z \in \mathcal{C}^k(\mathbb{R}, L^2(\mathbb{R}^3, \mathbb{R}^8)), z(t) \in \mathcal{H}_c(u(t)), \right. \\ \left. \max \left[\sup_{v \in [0, t]} \{ \|z(v)\|_{H^{s'}} \}, \sup_{v \in [0, t]} \{ \langle v \rangle^{3/2} \|z(v)\|_{B_{\infty, 2}^\beta} \}, \sup_{v \in [0, t]} \{ \langle v \rangle^{3/2} \{ \|z(v)\|_{H_{-\sigma}^s} \} \} \right] < \delta \right\}.$$

Then we define the space

$$\Omega(\delta) = \left\{ \alpha = (\alpha^+, \alpha^-) \in \mathcal{C}^{k+1}(\mathbb{R}), \alpha^- = \overline{\alpha^+}, \sup_{t \in \mathbb{R}^+} \langle t \rangle^{3/2} |\alpha(t)| < \delta \right\}.$$

5.1. Step 1: Construction of α . For any $u \in \mathcal{U}(\varepsilon, \delta)$ and $z \in \mathcal{Z}(u, \delta)$, let us define a map $\mathcal{G}_{u,z}$ on $\Omega(\delta)$ by

$$\begin{aligned} \mathcal{G}_{u,z}(\alpha)^\pm(t) = & - \int_t^\infty e^{\int_s^t E_1^\pm(u(w)) dw} \{ \langle JN(u(v), \eta(v)), S_1^\pm(u(v)) \rangle \\ & + \langle dS(u(v))A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle S_1^\pm(u(v)) \rangle \\ & - \langle (dS_1^\pm(u(v)))A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle, S_1^\pm(u(v)) \rangle \alpha^\pm(v) \} dv. \end{aligned}$$

We want to show that $\mathcal{G}_{u,z}$ stabilizes $\Omega(\delta)$ and is a contraction for the L^∞ norm. We have the

Lemma 5.1. *Let be $\sigma \in \mathbb{R}$, $s > 1$ and $p, p_1, p_2, q \in [1, \infty]$ such that*

$$\frac{1}{p} + \frac{s}{3} > \frac{1}{p_1} + \frac{1}{p_2}.$$

Then there exists $\varepsilon_0 > 0$ and $C > 0$ such that for all $u \in B_{\mathbb{C}}(0, \varepsilon_0)$ and $\eta \in B_{p,q}^s(\mathbb{R}^3, \mathbb{R}^8) \cap L^\infty(\mathbb{R}^3, \mathbb{R}^8)$, such that

$$\|\langle Q \rangle^\sigma N(u, \eta)\|_{B_{p,q}^s} \leq C(s, F, |u| + \|\eta\|_{L^\infty}) \left(|u| + \|\eta\|_{B_{p_2,q}^s} \right) \|\eta\|_{L^\infty} \|\langle Q \rangle^\sigma \eta\|_{B_{p_1,q}^s}. \quad (5.1)$$

Proof. We recall the definition

$$N(u, \eta) = \nabla F(S(u) + \eta) - \nabla F(S(u)) - d^2 F(S(u))\eta.$$

We have

$$N(u, \eta) = \int_0^1 \int_0^1 d^3 F(S(u) + \theta' \theta \eta) \cdot \eta \cdot \theta \eta d\theta' d\theta.$$

Since for $s \in \mathbb{R}_+^*$, $p, p_1, p_2, q \in [1, \infty]$ such that $\frac{1}{p} + \frac{s}{3} > \frac{1}{p_1} + \frac{1}{p_2}$, we have $\|uv\|_{B_{p,q}^s} \leq C\|u\|_{B_{p_1,q'}^s} \|v\|_{B_{p_2,q'}^s}$. Then since $s > 1$, we use (see [EV97, Proposition 2.1])

$$\|d^3 F(\psi)\|_{B_{p',q}^s} \leq C(s, F, \|\psi\|_\infty) \|\psi\|_{B_{p',q}^s}.$$

Then using Lemma 4.1, we conclude the proof.

Hence we have the

Lemma 5.2. *There exists $\delta_0 > 0$ and $\varepsilon_0 > 0$ such that for any $\delta \in (0, \delta_0)$ and $\varepsilon \in (0, \varepsilon_0)$, for any $u \in \mathcal{U}(\varepsilon, \delta)$ and $z \in \mathcal{Z}(u, \delta)$, $\mathcal{G}_{u,z}(\alpha)$ maps $\Omega(\delta)$ into itself.*

Proof. We have by means of Estimate (5.1) with e.g. $\sigma < -3$, $s = 0$, $p = q = 2$ and $p_1 = p_2 = 4$ and if $u_0 \in \mathbb{C}$ and $z_0 \in \mathcal{H}_c(u_0) \cap H_c^{s'}(\mathbb{R}^3, \mathbb{R}^8)$ are small enough

$$\begin{aligned} |\mathcal{G}_{u,z}(\alpha)^\pm(t)| & \leq C \int_t^\infty \{ |\langle JN(u(s), \eta(s)), S_1^\pm(u(s)) \rangle| \\ & + |\langle dS(u(v))A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle S_1^\pm(u(v)) \rangle| \\ & + |\langle (dS_1^\pm(u(v)))A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle, S_1^\pm(u(v)) \rangle \alpha^\pm(v)| \} dv \\ & \leq C\delta^2 \langle t \rangle^{-2}. \end{aligned}$$

Hence for small δ and small ε , we have $\mathcal{G}_{u,z}(\Omega(\delta)) \subset \Omega(\delta)$.

To prove that $\mathcal{G}_{u,z}$ is a contraction for the L^∞ norm, we use the

Lemma 5.3. *Let be $\sigma \in \mathbb{R}$, $s > 0$ and $p, q \in [1, \infty]$ such that $sp > 3$. Then for any $\varepsilon > 0$ and $M > 0$ there exists $C > 0$ such that for all $u, u' \in B_{\mathbb{C}}(0, \varepsilon)$ and $\eta, \eta' \in B_{p,q}^s(\mathbb{R}^3, \mathbb{R}^8) \cap L^\infty(\mathbb{R}^3, B_{\mathbb{R}^8}(0, M))$, such that*

$$\begin{aligned} \|\langle Q \rangle^\sigma \{N(u, \eta) - N(u', \eta')\}\|_{B_{p,q}^s} & \leq C \left(s, F |u| + |u'| + \|\eta\|_{B_{p,q}^s} + \|\eta'\|_{B_{p,q}^s} \right) \times \\ & \times \left\{ \left(\|\langle Q \rangle^{\sigma_1} \eta\|_{B_{p,q}^s} + \|\langle Q \rangle^{\sigma_1} \eta'\|_{B_{p,q}^s} \right)^2 \left(|u - u'| + \|\langle Q \rangle^{\sigma_1} (\eta - \eta')\|_{B_{p,q}^s} \right) \right. \\ & + \left(|u| + |u'| + \|\langle Q \rangle^{\sigma'_1} \eta\|_{B_{p,q}^s} + \|\langle Q \rangle^{\sigma'_1} \eta'\|_{B_{p,q}^s} \right) \times \\ & \left. \times \left(\|\langle Q \rangle^{\sigma'_2} \eta\|_{B_{p,q}^s} + \|\langle Q \rangle^{\sigma'_2} \eta'\|_{B_{p,q}^s} \right) \|\langle Q \rangle^{\sigma'_3} (\eta - \eta')\|_{B_{p,q}^s} \right\}, \end{aligned}$$

with $2\sigma_1 + \sigma_2 = \sigma'_1 + \sigma'_2 + \sigma'_3 = \sigma$.

Proof. Since, we have

$$N(u, \eta) = \int_0^1 \int_0^1 d^3 F(S(u) + \theta' \theta \eta) \cdot \eta \cdot \theta \eta d\theta' d\theta.$$

we can also restrict the study to $d^3 F(\phi) - d^3 F(\phi')$. If $d^5 F \neq 0$, we have

$$\|\langle Q \rangle^\sigma (d^3 F(\phi) - d^3 F(\phi'))\|_{B_{p,q}^s} \leq \int_0^1 \|d^4 F(\phi + t(\phi - \phi'))\|_{B_{p,q}^s} \|\langle Q \rangle^\sigma (\phi - \phi')\|_{B_{p,q}^s} dt$$

then since $s > 0$, we use

$$\|d^4 F(\psi)\|_{B_{p,q}^s} \leq C(s, F, \|\psi\|_{B_{p,q}^s}).$$

Then using Lemma 4.1, we conclude the proof when $d^5 F \neq 0$. Otherwise the proof is easily adaptable since $d^4 F$ is a constant matrix of $\mathcal{M}_4(\mathbb{C})$.

We also need the

Lemma 5.4. *Let be $\sigma \in \mathbb{R}$, $s > 0$ and $p, q \in [1, \infty]$. For any $\varepsilon > 0$ and $M > 0$, there exists $C > 0$ such that for all $u, u' \in B_{\mathbb{C}}(0, \varepsilon)$ and $\eta, \eta' \in B_{p,q}^s(\mathbb{R}^3, \mathbb{R}^8) \cap L^\infty(\mathbb{R}^3, B_{\mathbb{R}^8}(0, M))$, one has*

$$|A(u, \eta) - A(u', \eta')| \leq C \left\{ |u - u'| + \|\langle Q \rangle^\sigma \{\eta - \eta'\}\|_{B_{p,q}^s} \right\} \quad (5.2)$$

Proof. We recall that

$$A(u, \eta) = [\langle JdS(u), dS(u) \rangle + \langle J\eta, d^2 S(u) \rangle]^{-1}.$$

We have

$$\begin{aligned} A(u, \eta) - A(u', \eta') &= -[\langle JdS(u), dS(u) \rangle + \langle J\eta, d^2 S(u) \rangle]^{-1} \times \\ &\quad \times \left\{ \langle JdS(u), dS(u) \rangle + \langle J\eta, d^2 S(u) \rangle - \langle JdS(u'), dS(u') \rangle + \langle J\eta', d^2 S(u') \rangle \right\} \times \\ &\quad \times [\langle JdS(u'), dS(u') \rangle + \langle J\eta', d^2 S(u') \rangle]^{-1}. \end{aligned}$$

The lemma then follows Lemma 4.1.

Hence we have the

Lemma 5.5. *There exists $\delta_0 > 0$ and ε_0 such that there exists $\kappa \in (0, 1)$ such that for any $\delta \in (0, \delta_0)$ and $\varepsilon \in (0, \varepsilon_0)$, for any $u, u' \in \mathcal{U}(\varepsilon)$ and $z \in \mathcal{Z}(u, \delta)$ and $z' \in \mathcal{Z}(u', \delta)$, for any $\alpha, \alpha' \in \Omega(\delta)$, one has*

$$\|\mathcal{G}_{u', z'}(\alpha') - \mathcal{G}_{u, z}(\alpha)\|_{L^\infty(\mathbb{R}^+)} \leq \kappa \left(|u' - u|_{L^\infty(\mathbb{R}^+)} + |\alpha' - \alpha|_{L^\infty(\mathbb{R}^+)} + \|z' - z\|_{L^\infty(\mathbb{R}^+, B_{\infty, 2}^\beta)} \right).$$

Proof. It is a straightforward computation based on Lemma 5.3 with e.g. $\sigma < -6$, $\sigma_3, \sigma_2' < -3$ and $s = 0$, $p = q = 2$, on Lemma 5.4, on Lemma 5.1 with e.g. $\sigma < -3$, $s = 0$, $p = q = 2$ and $p_1 = p_2 = 4$ and on Lemma 4.3.

We now state the

Lemma 5.6. *There exists $\delta_0 > 0$ and ε_0 such that for any $u \in \mathcal{U}(\varepsilon_0)$ and $z \in \mathcal{Z}(u, \delta_0)$, the equation*

$$\begin{cases} \alpha^\pm = E^\pm(u) \alpha^\pm + \langle JN(u, \eta), S_1^\pm(u) \rangle \\ \quad + \langle dS(u) A(u, \eta) \langle N(u, \eta), dS(u) \rangle S_1^\pm(u) \rangle \\ \quad - \langle (dS_1^\pm(u)) A(u, \eta) \langle N(u, \eta), dS(u) \rangle, S_1^\pm(u) \rangle \alpha^\pm, \\ \lim_{t \rightarrow +\infty} \alpha(t) = 0 \end{cases}$$

where $\eta(t) = \alpha^+(t) S_1^+(u) + \alpha^-(t) S_1^-(u) + z(t)$, has a unique solution in $\Omega(\delta_0)$.

Proof. The proof is now classical since we proved that the integral equation

$$\alpha(t) = \mathcal{G}_{u, z}(\alpha)(t)$$

can be solved by means of the fixed point theorem.

5.2. *Step 2: Construction of z .* Let be $u \in \mathcal{U}(\varepsilon, \delta)$ and $z_0 \in \mathcal{H}_c(u(0)) \cap H_\sigma^{s'}$. Let us write $u_\infty = \lim_{t \rightarrow +\infty} u(t)$, we define $\mathcal{T}_{u,z_0}(z)$ by

$$\begin{aligned} \mathcal{T}_{u,z_0}(z)(t) &= e^{JtH(u_\infty)} z_0 + \int_0^t e^{J(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) J \{d^2 F(S(u(v))) - d^2 F(S(u_\infty))\} z(v) dv \\ &\quad + \int_0^t e^{J(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) JN(u(v), \eta(v)) dv \\ &\quad + \int_0^t e^{J(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) dS(u(v)) A(u(v), \eta) \langle N(u(v), \eta(v)), dS(u(v)) \rangle dv \\ &\quad - \int_0^t e^{J(t-v)H(u_\infty)} (d\mathbf{P}_c(u(v))) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle z(v) dv. \end{aligned}$$

We have

Lemma 5.7. *There exists $\delta_0 > 0$ and $\varepsilon_0 > 0$ and $C > 0$ such that for any $\delta \in (0, \delta_0)$, for any $\varepsilon \in (0, \varepsilon_0)$ and for any $u \in \mathcal{U}(\varepsilon, \delta)$, the application \mathcal{T}_{u,z_0} maps $\mathcal{Z}(u, \delta)$ into itself if $\|z_0\|_{H_\sigma^{s'}} \leq C\delta$*

Proof. With Lemma A.1 and Lemma 4.4, we obtain

$$\begin{aligned} \|\mathcal{T}_{u,z_0}(t)\|_{H^{s'}} &\leq C \|\mathbf{P}_c(u_\infty) \mathcal{T}_{u,z_0}(t)\|_{H^{s'}} \\ &\leq C \|z_0\|_{H^{s'}} + C \int_0^t \|d^2 F(S(u(v))) - d^2 F(S(u_\infty)) z(v)\|_{H^{s'}} dv \\ &\quad + C \int_0^t \|N(u(v), \eta(v))\|_{H^{s'}} dv \\ &\quad + C \int_0^t \|dS(u(v)) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle\|_{H^{s'}} dv \\ &\quad + C \int_0^t \|(d\mathbf{P}_c(u(v))) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle z(v)\|_{H^{s'}} dv. \end{aligned}$$

Now, with Lemma 5.1, we obtain

$$\begin{aligned} \|\mathcal{T}_{u,z_0}(t)\|_{H^{s'}} &\leq C \|z_0\|_{H^{s'}} + C\varepsilon \int_0^t \|u(v) - u_\infty\| \|z\|_{H^{s'}} dv \\ &\quad + C \int_0^t (|u(v)| + \|\eta(v)\|_{H^{s'}}) \|\eta(v)\|_{L^\infty} \|\eta(v)\|_{H^{s'}} dv \\ &\quad + C \int_0^t (|u(v)| + \|\eta(v)\|_{H^{s'}}) \|\eta(v)\|_{L^\infty} \|\eta(v)\|_{H^{s'}} \|z\|_{H^{s'}} dv, \end{aligned}$$

and so

$$\|\mathcal{T}_{u,z_0}(t)\|_{H^{s'}} \leq C \|z_0\|_{H^{s'}} + C\varepsilon\delta^2 + C(\varepsilon + \delta)\delta^2 + C(\varepsilon + \delta)\delta^3.$$

Then, we also have

$$\begin{aligned} \mathcal{T}_{u,z_0}(t) &= e^{-itH} z_0 + \int_0^t e^{-i(t-v)H} \mathbf{P}_c(u(v)) d^2 F(S(u(v))) z(v) dv \\ &\quad + \int_0^t e^{-i(t-v)H} \mathbf{P}_c(u(v)) JN(u(v), \eta(v)) dv \\ &\quad + \int_0^t e^{-i(t-v)H} \mathbf{P}_c(u(v)) dS(u(v)) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle dv \\ &\quad - \int_0^t e^{-i(t-v)H} (d\mathbf{P}_c(u(v))) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle z(v) dv. \end{aligned}$$

Hence by Lemma 4.4 and Theorem 1.2, we have

$$\begin{aligned}
\|\mathcal{T}_{u,z_0}(t)\|_{B_{\infty,2}^\beta} &\leq C\langle t\rangle^{-3/2}\|z_0\|_{B_{1,2}^{\beta+3}} + C\int_0^t \langle t-v\rangle^{-3/2} \|d^2F(S(u(v)))z(v)\|_{B_{1,2}^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} \|N(u(v),\eta(v))\|_{B_{1,2}^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} \|dS(u(v))A(u(v),\eta(v))\langle N(u(v),\eta(v)),dS(u(v))\rangle\|_{B_{1,2}^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} \times \\
&\quad \times \|(d\mathbf{P}_c(u(v)))A(u(v),\eta(v))\langle N(u(v),\eta(v)),dS(u(v))\rangle z(v)\|_{B_{1,2}^{\beta+3}} dv.
\end{aligned}$$

With Lemma 5.1, we infer

$$\begin{aligned}
\|\mathcal{T}_{u,z_0}(t)\|_{B_{\infty,2}^\beta} &\leq \\
&C\langle t\rangle^{-3/2}\|z_0\|_{B_{1,2}^{\beta+3}} + C\int_0^t \langle t-v\rangle^{-3/2} |u(v)|^2 \|z(v)\|_{H_{-\sigma}^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} (|u(v)| + \|\eta(v)\|_{H^{\beta+3}}) \|\eta(v)\|_{L^\infty} \|\eta(v)\|_{H^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} (|u(v)| + \|\eta(v)\|_{H^{\beta+3}}) \|\eta(v)\|_{L^\infty} \|\eta(v)\|_{H^{\beta+3}} dv \\
&\quad + C\int_0^t \langle t-v\rangle^{-3/2} (|u(v)| + \|\eta(v)\|_{H^{\beta+3}}) \|\eta(v)\|_{L^\infty} \|\eta(v)\|_{H^{\beta+3}}^2 dv.
\end{aligned}$$

With the estimate

$$\int_0^t \langle t-v\rangle^{-3/2} \langle v\rangle^{-3/2} dv \leq C\langle t\rangle^{-3/2},$$

we infer

$$\langle t\rangle^{3/2}\|\mathcal{T}_{u,z_0}(t)\|_{B_{\infty,2}^\beta} \leq C\|z_0\|_{B_{1,2}^{\beta+3}} + C\varepsilon^2\delta + C(\varepsilon + \delta)\delta^2 + C(\varepsilon + \delta)\delta^3.$$

Then we also have

$$\begin{aligned}
\mathcal{T}_{u,z_0}(t) &= e^{-itH}z_0 + \int_0^t e^{-i(t-v)H}\mathbf{P}_c(u(v))\nabla F(\eta(v)) dv \\
&\quad + \int_0^t e^{-i(t-v)H}\mathbf{P}_c(u(v))J\{\nabla F(S(u(v)) + \eta(v)) - \nabla F(S(u(v)) - \nabla F(\eta(v))\} dv \\
&\quad + \int_0^t e^{-i(t-v)H}\mathbf{P}_c(u(v))dS(u(v))A(u(v),\eta(v))\langle N(u(v),\eta(v)),dS(u(v))\rangle dv \\
&\quad - \int_0^t e^{-i(t-v)H}(d\mathbf{P}_c(u(v)))A(u(v),\eta(v))\langle N(u(v),\eta(v)),dS(u(v))\rangle z(v) dv.
\end{aligned}$$

We now use Lemma 4.4 and Theorem 1.1, except for the third term of the right hand side for which we used Theorem 1.2 since $\sigma > 3$. We also use Lemma 5.1, except for the second term of the right hand side for which an obvious adaptation of the proof of Lemma 5.3 gives

$$\begin{aligned}
\|\nabla F(S(u(v)) + \eta(v)) - \nabla F(S(u(v)) - \nabla F(\eta(v))\|_{H_\sigma^s} \\
\leq C(|u(v)| + \|\eta(v)\|_{H^s})|u(v)|\|\eta(v)\|_{H_{-\sigma}^s}
\end{aligned}$$

and so we obtain

$$\langle t\rangle^{3/2}\|\mathcal{T}_{u,z_0}(t)\|_{H_{-\sigma}^s} \leq C\|z_0\|_{H_\sigma^s} + C\varepsilon^2\delta + C(\varepsilon + \delta)\delta^2 + C(\varepsilon + \delta)\delta^3.$$

Therefore we have that \mathcal{T}_{u,z_0} leaves $\mathcal{Z}(u, \delta)$ invariant if $\|z_0\|_{H_\sigma^s}$, δ and ε are small enough.

An important property of \mathcal{T} is given by the

Lemma 5.8. *There exists $\delta_0 > 0$ and $\varepsilon_0 > 0$ such that there exists $\kappa \in (0, 1)$ such that for any $\delta \in (0, \delta_0)$, for any $\varepsilon \in (0, \varepsilon_0)$, for any $u, u' \in \mathcal{U}(\varepsilon, \delta)$, for any $z_0 \in \mathcal{H}_c(u(0))$, for any $z'_0 \in \mathcal{H}_c(u'(0))$, for $z \in \mathcal{Z}(u, \delta)$ and for any $z' \in \mathcal{Z}(u', \delta)$, one has for all $t \in \mathbb{R}$*

$$\|\mathcal{T}_{u', z'_0}(z')(t) - \mathcal{T}_{u, z_0}(z)(t)\|_{H^{s'}} \leq \|z_0 - z'_0\|_{H^{s'}} + \kappa \{\|u - u'\|_{L^\infty} + \|\eta - \eta'\|_{H^{s'}}\}.$$

Proof. It is an easy consequences of straightforward estimates on the following identity

$$\begin{aligned} \mathcal{T}_{u, z_0}(t) &= e^{-itH(u_\infty)} z_0 \\ &+ \int_0^t e^{-i(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) (d^2 F(S(u(v))) - d^2 F(S(u_\infty))) z(v) dv \\ &+ \int_0^t e^{-i(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) JN(u(v), \eta(v)) dv \\ &+ \int_0^t e^{-i(t-v)H(u_\infty)} \mathbf{P}_c(u(v)) dS(u(v)) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle dv \\ &- \int_0^t e^{-i(t-v)H(u_\infty)} (d\mathbf{P}_c(u(v))) A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle z(v) dv, \end{aligned}$$

based only on Lemma 5.3, 5.4 and 5.5 and on the fact that

$$\begin{aligned} \mathbf{P}_c(u_\infty) \left(e^{-isH(u_\infty)} - e^{-isH(u'_\infty)} \right) \mathbf{P}_c(u(v)) &= \\ \mathbf{P}_c(u_\infty) \int_0^s \left(e^{-i(s-s')H(u_\infty)} (d^2 F(S(u'_\infty)) - d^2 F(S(u_\infty))) e^{-is'H(u'_\infty)} \right) ds' \mathbf{P}_c(u(v)) \end{aligned}$$

form a family of operator in $\mathcal{B}(H^{s'}(\mathbb{R}^3, \mathbb{C}^8), H^{s'}(\mathbb{R}^3, \mathbb{C}^8))$ (to this end we use Lemma A) such that

$$\left\| \mathbf{P}_c(u_\infty) \left(e^{-isH(u_\infty)} - e^{-isH(u'_\infty)} \right) \mathbf{P}_c(u(v)) \right\| \leq C\varepsilon |u'_\infty - u_\infty|,$$

with C independent of u, u' .

Lemma 5.9. *There exists $\delta > 0$ and $\varepsilon > 0$ such that for any $u \in \mathcal{U}(\delta, \varepsilon)$ there is a solution $z \in \mathcal{Z}(u, \delta)$ of the equation*

$$\begin{cases} \partial_t z = JH(u)z + \mathbf{P}_c(u) JN(u, \eta) \\ \quad + \mathbf{P}_c(u) dS(u) A(u, \eta) \langle N(u, \eta), dS(u) \rangle \\ \quad - (D\mathbf{P}_c(u)) A(u, \eta) \langle N(u, \eta), dS(u) \rangle z, \\ z(0) = z_0, \end{cases} \quad (5.3)$$

where $\eta(t) = \alpha^+(t)S_1^+(u) + \alpha^-(t)S_1^-(u) + z(t)$, whenever $z_0 \in H_\sigma^{s'}$ is small enough.

Proof. It is a consequence of the fix point theorem applied to \mathcal{T}_{u, z_0} .

Lemma 5.10. *Under the assumptions of Lemma 5.9, for any $u \in \mathcal{U}(\delta, \varepsilon)$ and solution z of (5.3), with $z_0 \in H_\sigma^{s'}$ small, the following limit*

$$z_\infty = \lim_{t \rightarrow \infty} e^{itD_m} z(t)$$

exists in $H^{s'} \cap B_{\infty, 2}^\beta \cap H_{-\sigma}^s$.

Proof. We first prove that $\lim_{t \rightarrow \infty} e^{itH} z(t)$ exists by using exactly the same method as the one of Lemma 5.7. Then we just use Lemma A.4 to prove the existence of z_∞ .

5.3. Step 3: Construction of u . Here we want to solve the equation for u . We notice that z and α have been built in the previous section and are functions of u and $z_0 \in \mathcal{H}_c(u(0))$. Let us introduce for any $\alpha \in \Omega(\delta)$ and $u_0 \in B_{\mathbb{C}}(0, \varepsilon)$ the function on $\mathcal{U}(\varepsilon, \delta)$:

$$f_{u_0}(u)(t) = u_0 - \int_0^t A(u(v), \eta(v)) \langle N(u(v), \eta(v)), dS(u(v)) \rangle dv,$$

where $\eta(t) = \alpha^+(t)S_1^+(u) + \alpha^-(t)S_1^-(u) + z(t)$. We have the

Lemma 5.11. *There exists $\delta_0 > 0$ and $\varepsilon_0 > 0$ such that for any $\delta \in (0, \delta_0)$, for any $\varepsilon \in (0, \varepsilon_0)$, the function f_{u_0} maps $\mathcal{U}(\varepsilon, \delta)$ into itself if u_0 and $z_0 \in H_{\sigma}^{s'}$ are small enough.*

Proof. By means of Lemma 5.1, we obtain

$$|f_{u_0}(u)(t)| \leq |u_0| + C \int_0^t \|N(u(v), \eta(v))\|_{H_{-\sigma}^s} \leq |u_0| + C(\varepsilon + \delta)\delta^2.$$

Hence for u_0 and δ small $f_{u_0}(u)(t) \in B_{\mathbb{C}}(0, \varepsilon)$. Estimate (5.1) also gives the existence of $(f_{u_0}(u))_{\infty} = \lim_{t \rightarrow +\infty} f_{u_0}(u)(t)$ and then

$$|(f_{u_0}(u))_{\infty} - f_{u_0}(u)(t)| \leq C \int_t^{+\infty} \|N(u(v), \eta(v))\|_{H_{-\sigma}^s} \leq \frac{C}{t^2} (\varepsilon + \delta) \delta^2.$$

The function f_{u_0} has also a Lipschitz property as proved by the

Lemma 5.12. *There exists $\delta_0 > 0$ and $\varepsilon_0 > 0$ such that there exists $\kappa \in (0, 1)$ such that for any $\delta \in (0, \delta_0)$, for any $\varepsilon \in (0, \varepsilon_0)$, for any $u, u' \in \mathcal{U}(\varepsilon, \delta)$, for any $z_0 \in \mathcal{H}_c(u(0)) \cap H_{\sigma}^{s'}$, for any $z'_0 \in \mathcal{H}_c(u'(0)) \cap H_{\sigma}^{s'}$ small enough, for u_0, u'_0 small enough, there exists $C > 0$ such that*

$$\forall t \in \mathbb{R}, |f_{u_0}(u) - f_{u'_0}(u')| \leq |u_0 - u'_0| + \kappa \left(\|u - u'\|_{L^{\infty}([-t, t])} \right)$$

Proof. This is a straightforward consequence of Lemma 5.3, 5.4, 5.5 and 5.8.

We are now able to prove the

Lemma 5.13. *There exists $\delta_0 > 0$ such that for any $\delta \in (0, \delta_0)$ such that for any $u_0 \in \mathbb{C}$ small and $z_0 \in \mathcal{H}_c(u_0) \cap H_{\sigma}^{s'}$ small, the equation*

$$\begin{cases} \dot{u} = -A(u, \eta) \langle N(u, \eta), dS(u) \rangle, \\ u(0) = u_0, \end{cases}$$

where $\eta(t) = \alpha^+(t)S_1^+(u) + \alpha^-(t)S_1^-(u) + z(t)$, has a unique solution in $\mathcal{U}(\delta, \varepsilon)$.

Proof. This is also a straightforward consequence of the fixed point theorem for f_{u_0} .

5.4. Step 4: End of the proof of Theorem 1.3. We now conclude our proof and we need the

Lemma 5.14 (Decomposition lemma 2). *Let be $s \geq 0$ and $p \geq 1$ there exist $\delta > 0$ and a \mathcal{C}^{∞} map $U_0 : B_{W^{s,p}}(0, \delta) \mapsto B_{\mathbb{C}}(0, \varepsilon)$ which satisfies for $\psi \in B_{W^{s,p}}(0, \delta)$*

$$\psi = S(u) + \eta, \text{ with } \eta \in \{\phi_0\}^{\perp} \iff u = U_0(\psi)$$

Proof. It is a consequence of the implicit function theorem like in proof of [GNT04, Lemma 2.3].

With respect to the notation of Theorem 1.3 for $\psi_0 = S(u_0) + \alpha^+(0)S_1^+(u_0) + \alpha^-(0)S_1^-(u_0) + \mathbf{P}_c(u_0)\tilde{z}_0$ where $\tilde{z}_0 \in \text{Ran}(\mathbf{P}_c)$, we introduce

$$\begin{cases} v_0 = U_0(\psi_0) \\ \xi_0 = \mathbf{P}_c(\psi_0 - S(v_0)) \end{cases}$$

and

$$\begin{cases} G(u_0, \tilde{z}_0)_1 = U_0(S(u_0) + \alpha^+(0)S_1^+(u_0) + \alpha^-(0)S_1^-(u_0) + \mathbf{P}_c(u_0)\tilde{z}_0) \\ G(u_0, \tilde{z}_0)_2 = \mathbf{P}_c\left(S(u_0) - S(G(u_0, \tilde{z}_0)_1) \right. \\ \quad \left. + \alpha^+(0)S_1^+(u_0) + \alpha^-(0)S_1^-(u_0) + \mathbf{P}_c(u_0)\tilde{z}_0\right) \end{cases}$$

Then using $U_0(S(u_0)) = u_0$, we write $G(u_0, \tilde{z}_0) = (u_0, \tilde{z}_0) + H(u_0, \tilde{z}_0)$, with

$$\|H(u_0, \tilde{z}_0) - H(u'_0, \tilde{z}'_0)\|_{H_\sigma^{s'}} \leq \kappa \left(|u_0 - u'_0| + \|\tilde{z}_0 - \tilde{z}'_0\|_{H_\sigma^{s'}} \right)$$

with $\kappa \leq 1/2$ if u_0, u'_0 and $\tilde{z}_0, \tilde{z}'_0$ small enough. Hence in this case G is invertible with a Lipshitz inverse F . Then we choose

$$\begin{aligned} \Psi(v_0, \xi_0) &= \langle S(F(v_0, \xi_0)_1) \\ &\quad + \alpha(F(v_0, \xi_0)_1)^+(0)S_1^+(F(v_0, \xi_0)_1) + \alpha(F(v_0, \xi_0)_1)^-(0)S_1^-(F(v_0, \xi_0)_1) \\ &\quad + \mathbf{P}_c(F(v_0, \xi_0)_1)F(v_0, \xi_0)_2 - S(v_0), \phi_1 \rangle \\ \xi_\infty &= (\mathbf{P}_c(F(v_0, \xi_0)_1)F(v_0, \xi_0)_2)_\infty \\ E_\infty &= \int_0^\infty \{E(F(v_0, \xi_0)_1(v)) - E((F(v_0, \xi_0)_1)_\infty)\} dv. \end{aligned}$$

In the proof of Lemma 5.7, we see that δ is of the same order as $\|\xi_0\|_{H_\sigma^{s'}}$. The rest of the Theorem easily follows. \square

APPENDICES

A. The wave operator and similarity for the linearized operator

Using an argument of similarity like [Kat66], we prove the

Lemma A.1. *For all $s \in \mathbb{R}^+$, there exists $C_s > 0$, such that*

$$\forall t \in \mathbb{R}, \|e^{tJH(z)}\|_{\mathcal{L}(H_s)} \leq C_s.$$

We will use the boundedness the wave operator

$$W_\pm = s - \lim_{t \rightarrow \pm\infty} e^{tJH(z)^*J} e^{-it(H-E(z))} \mathbf{P}_c(H).$$

To this end we prove the

Lemma A.2 (Smooth and small non-selfadjoint perturbations). *Let be $\psi \in L^2$ and $\sigma > 1$. Then there exists $\varepsilon > 0$ and $C > 0$ such that*

$$\forall z \in B_{\mathbb{C}}(0, \varepsilon), \int_0^\infty \| \langle Q \rangle^{-\sigma} e^{-tH(z)^*J} \mathbf{P}_c(z)\psi \|_2 \leq C \|\psi\|_2^2.$$

Proof. By Lemma (4.4), we prove $\mathbf{P}_c(z)R(z, 0)\mathbf{P}_c(0)\mathbf{P}_c(z) = \mathbf{P}_c(z)$. Taking the adjoint with respect to the real structure, we infer $\mathbf{P}_c(z)\mathbf{P}_c(0)R(z)^*\mathbf{P}_c(z) = \mathbf{P}_c(z)$. Then, we write

$$\begin{aligned} &\| \langle Q \rangle^{-\sigma} e^{-tH(z)^*J} \mathbf{P}_c(z) \| \\ &= C \| \langle Q \rangle^{-\sigma} \mathbf{P}_c(z) e^{-tH(z)^*J} \mathbf{P}_c(0)R(z)^*\mathbf{P}_c(z) \| \\ &\leq \| \langle Q \rangle^{-\sigma} \mathbf{P}_c(z) e^{-it(H-E(z))} \mathbf{P}_c(0)R(z)^*\mathbf{P}_c(z) \| \\ &\quad + \int_0^t \| \langle Q \rangle^{-\sigma} \mathbf{P}_c(z) e^{-(t-s)(H(z)^*)J} D\nabla F(S(z))^* e^{-is(H-E(z))} \mathbf{P}_c(0)R(z)^*\mathbf{P}_c(z) \| ds \\ &\leq \| \langle Q \rangle^{-\sigma} \mathbf{P}_c(z) e^{-it(H-E(z))} \mathbf{P}_c(0)R(z)^*\mathbf{P}_c(z) \| \\ &\quad + C|z|^{r-2} \int_0^t \| \langle Q \rangle^{-\sigma} \mathbf{P}_c(z) e^{-(t-s)(H(z)^*)J} \| \| \langle Q \rangle^{-\sigma} e^{-is(H-E(z))} \mathbf{P}_c(0) \| ds. \end{aligned}$$

Using Proposition 3.10, we obtain the claim (A.2) for z sufficiently small.

Lemma A.3. *Let be $W_t = e^{tH(z)J}e^{-it(H-E(z))}\mathbf{P}_c(H)$. Then the limits*

$$W^\pm = \lim_{t \rightarrow \pm\infty} W_t$$

exist in $B(H^s)$ and their range is $\text{Ran}(\mathbf{P}_c(z))$. The same is true for W_t^ and*

$$(W^\pm)^{-1} = (W^\pm)^* = \lim_{t \rightarrow \pm\infty} W_t^*.$$

Proof. Let us define $W_t = e^{-tH(z)^*J}e^{-it(H-E(z))}$. we have for $\phi \in H_c(z)$ and $\psi \in H_c(0)$

$$\langle \phi, W_t \psi \rangle = \langle \phi, \psi \rangle + \int_0^t \left\langle \phi, \frac{d}{ds} W_s \psi \right\rangle ds,$$

Since we have

$$\begin{aligned} \left\langle \phi, \frac{d}{ds} W_s \psi \right\rangle &= \left\langle e^{tJH(z)} \phi, D\nabla F(S(z)) e^{-it(H-E(z))} \psi \right\rangle \\ &\leq C|z|^{r-3} \|\langle Q \rangle^{-\sigma} e^{tJH(z)} \phi\| \|\langle Q \rangle^{-\sigma} e^{-it(H-E(z))} \psi\|. \end{aligned}$$

which gives $\langle \phi, \frac{d}{ds} W_s \psi \rangle \in L^1(\mathbb{R})$, so W_\pm exists and is bounded in $\mathcal{L}(H_c(0), H_c(z))$ by the previous lemma. Since for any vector ϕ in an eigenspace of $H(z)$, $W_t^* \phi$ tends weakly to zero, we obtain that the ran of W^\pm is a subspace of the range of $\mathbf{P}_c(z)$. Then statements about $(W^t)^*$ follows by the same way. The invertibility is immediate.

The proof of the H^k bounds follows from commutation argument, we apply the previous scheme to

$$\begin{aligned} \partial_i e^{tJH(z)} \mathbf{P}_c(z) &= [\partial_i, \mathbf{P}_c(z)] e^{tJH(z)} \mathbf{P}_c(z) + \mathbf{P}_c(z) e^{tJH(z)} [\partial_i, \mathbf{P}_c(z)] \\ &\quad + \mathbf{P}_c(z) [\partial_i, e^{tJH(z)}] \mathbf{P}_c(z) \\ &= [\partial_i, \mathbf{P}_c(z)] e^{tJH(z)} \partial_i + \mathbf{P}_c(z) e^{tJH(z)} [\partial_i, \mathbf{P}_c(z)] \\ &\quad + \int_0^t e^{-(t-s)JH(z)} \mathbf{P}_c(z) (\partial_i D\nabla F(S)) e^{sJH(z)} \mathbf{P}_c(z) dz. \end{aligned}$$

By the same way we prove the

Lemma A.4. *Let be $W_t = e^{tH(z)J}e^{-it(D_m-E(z))}$. Then the limits*

$$W^\pm = \lim_{t \rightarrow \pm\infty} W_t$$

exist in $B(H^s)$ and their range is $\text{Ran}(\mathbf{P}_c(z))$. The same is true for W_t^ and*

$$(W^\pm)^{-1} = (W^\pm)^* = \lim_{t \rightarrow \pm\infty} W_t^*.$$

Proof (Proof of Lemma A.1). The intertwining property

$$e^{tJH(z)} \mathbf{P}_c(z) W_\pm = W_\pm e^{-it(H-E(z))} \mathbf{P}_c(0)$$

gives

$$W_\pm^{-1} e^{tJH(z)} \mathbf{P}_c(z) W_\pm = e^{-it(H-E(z))} \mathbf{P}_c(0)$$

gives the lemma on $H_c(z)$. On $H_c(z)^\perp$, the lemma is obvious, since we have a sum of eigenspaces.

References

- ABdMG96. Werner O. Amrein, Anne Boutet de Monvel, and Vladimir Georgescu. *C_0 -groups, commutator methods and spectral theory of N -body Hamiltonians*, volume 135 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1996.
- Agm75. Shmuel Agmon. Spectral properties of Schrödinger operators and scattering theory. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 2(2):151–218, 1975.
- AS86. A. Alvarez and M. Soler. Stability of the minimum solitary wave of a nonlinear spinorial model. *Phys. Rev D*, 34:644–645, July 1986.
- BdMGS96. Anne Boutet de Monvel, Vladimir Georgescu, and Jaouad Sahbani. Boundary values of resolvent families and propagation properties. *C. R. Acad. Sci. Paris Sér. I Math.*, 322(3):289–294, 1996.
- BH92. E. Balslev and B. Helffer. Limiting absorption principle and resonances for the Dirac operator. *Adv. in Appl. Math.*, 13(2):186–215, 1992.
- BP92a. V. S. Buslaev and G. S. Perel'man. Nonlinear scattering: states that are close to a soliton. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 200(Kraev. Zadachi Mat. Fiz. Smezh. Voprosy Teor. Funktsii. 24):38–50, 70, 187, 1992.
- BP92b. V. S. Buslaev and G. S. Perel'man. On nonlinear scattering of states which are close to a soliton. *Astérisque*, (210):6, 49–63, 1992. Méthodes semi-classiques, Vol. 2 (Nantes, 1991).
- BP92c. V. S. Buslaev and G. S. Perel'man. Scattering for the nonlinear Schrödinger equation: states that are close to a soliton. *Algebra i Analiz*, 4(6):63–102, 1992.
- BP95. V. S. Buslaev and G. S. Perel'man. On the stability of solitary waves for nonlinear Schrödinger equations. In *Nonlinear evolution equations*, volume 164 of *Amer. Math. Soc. Transl. Ser. 2*, pages 75–98. Amer. Math. Soc., Providence, RI, 1995.
- BS02. Vladimir S. Buslaev and Catherine Sulem. Asymptotic stability of solitary waves for nonlinear Schrödinger equations. In *The legacy of the inverse scattering transform in applied mathematics (South Hadley, MA, 2001)*, volume 301 of *Contemp. Math.*, pages 163–181. Amer. Math. Soc., Providence, RI, 2002.
- BS03. Vladimir S. Buslaev and Catherine Sulem. On asymptotic stability of solitary waves for nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(3):419–475, 2003.
- BSV87. P. Blanchard, J. Stubbe, and L. Vázquez. Stability of nonlinear spinor fields with application to the Gross-Neveu model. *Phys. Rev. D*, 36:2422–2428, October 1987.
- CF01. Carlos Cid and Patricio Felmer. Orbital stability and standing waves for the nonlinear Schrödinger equation with potential. *Rev. Math. Phys.*, 13(12):1529–1546, 2001.
- CL82. T. Cazenave and P.-L. Lions. Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.*, 85(4):549–561, 1982.
- CS01. Scipio Cuccagna and Pedro Paulo Schirmer. On the wave equation with a magnetic potential. *Comm. Pure Appl. Math.*, 54(2):135–152, 2001.
- Cuc01. Scipio Cuccagna. Stabilization of solutions to nonlinear Schrödinger equations. *Comm. Pure Appl. Math.*, 54(9):1110–1145, 2001.
- Cuc03. Scipio Cuccagna. On asymptotic stability of ground states of NLS. *Rev. Math. Phys.*, 15(8):877–903, 2003.
- Cuc05. Scipio Cuccagna. Erratum: “Stabilization of solutions to nonlinear Schrödinger equations” [Comm. Pure Appl. Math. **54** (2001), no. 9, 1110–1145; mr1835384]. *Comm. Pure Appl. Math.*, 58(1):147, 2005.
- ES04. M. Burak Erdoğan and Wilhelm Schlag. Dispersive estimates for Schrödinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three. I. *Dyn. Partial Differ. Equ.*, 1(4):359–379, 2004.
- EV97. M. Escobedo and L. Vega. A semilinear Dirac equation in $H^s(\mathbf{R}^3)$ for $s > 1$. *SIAM J. Math. Anal.*, 28(2):338–362, 1997.
- FS04. S. Fournais and E. Skibsted. Zero energy asymptotics of the resolvent for a class of slowly decaying potentials. *Math. Z.*, 248(3):593–633, 2004.
- GM01. V. Georgescu and M. Măntoiu. On the spectral theory of singular Dirac type Hamiltonians. *J. Operator Theory*, 46(2):289–321, 2001.
- GNT04. Stephen Gustafson, Kenji Nakanishi, and Tai-Peng Tsai. Asymptotic stability and completeness in the energy space for nonlinear Schrödinger equations with small solitary waves. *Int. Math. Res. Not.*, (66):3559–3584, 2004.
- GS04. M. Goldberg and W. Schlag. A limiting absorption principle for the three-dimensional Schrödinger equation with L^p potentials. *Int. Math. Res. Not.*, (75):4049–4071, 2004.
- GSS87. Manoussos Grillakis, Jalal Shatah, and Walter Strauss. Stability theory of solitary waves in the presence of symmetry. I. *J. Funct. Anal.*, 74(1):160–197, 1987.
- His00. P. D. Hislop. Exponential decay of two-body eigenfunctions: a review. In *Proceedings of the Symposium on Mathematical Physics and Quantum Field Theory (Berkeley, CA, 1999)*, volume 4 of *Electron. J. Differ. Equ. Conf.*, pages 265–288 (electronic), San Marcos, TX, 2000. Southwest Texas State Univ.
- HS00. W. Hunziker and I. M. Sigal. Time-dependent scattering theory of N -body quantum systems. *Rev. Math. Phys.*, 12(8):1033–1084, 2000.
- HSS99. W. Hunziker, I. M. Sigal, and A. Soffer. Minimal escape velocities. *Comm. Partial Differential Equations*, 24(11-12):2279–2295, 1999.
- IM99. A. Iftimovici and M. Măntoiu. Limiting absorption principle at critical values for the Dirac operator. *Lett. Math. Phys.*, 49(3):235–243, 1999.
- JK79. Arne Jensen and Tosio Kato. Spectral properties of Schrödinger operators and time-decay of the wave functions. *Duke Math. J.*, 46(3):583–611, 1979.
- JN01. Arne Jensen and Gheorghe Nenciu. A unified approach to resolvent expansions at thresholds. *Rev. Math. Phys.*, 13(6):717–754, 2001.
- JN04. Arne Jensen and Gheorghe Nenciu. Erratum: “A unified approach to resolvent expansions at thresholds” [Rev. Math. Phys. **13** (2001), no. 6, 717–754; mr 1841744]. *Rev. Math. Phys.*, 16(5):675–677, 2004.
- JSS91. J.-L. Journé, A. Soffer, and C. D. Sogge. Decay estimates for Schrödinger operators. *Comm. Pure Appl. Math.*, 44(5):573–604, 1991.
- Kat66. Tosio Kato. Wave operators and similarity for some non-selfadjoint operators. *Math. Ann.*, 162:258–279, 1965/1966.

- KS05. J. Krieger and W. Schlag. Stable manifolds for all supercritical monic nls in one dimension. preprint, 2005.
- Par90. Bernard Parisse. Résonances paires pour l'opérateur de Dirac. *C. R. Acad. Sci. Paris Sér. I Math.*, 310(5):265–268, 1990.
- PW97. Claude-Alain Pillet and C. Eugene Wayne. Invariant manifolds for a class of dispersive, Hamiltonian, partial differential equations. *J. Differential Equations*, 141(2):310–326, 1997.
- Ran. A. F. Ranada. Classical nonlinear dirac field models of extended particles. In *Quantum theory, groups, fields and particles*, volume 198, pages 271–291. A. O. Barut, Amsterdam, Reidel.
- RS78. Michael Reed and Barry Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- RS79. Michael Reed and Barry Simon. *Methods of modern mathematical physics. III*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1979. Scattering theory.
- RSS05a. Igor Rodnianski, Wilhelm Schlag, and Avraham Soffer. Asymptotic stability of n-soliton states of nls. to appear in *Comm. Pure and Appl. Math.*, 2005.
- RSS05b. Igor Rodnianski, Wilhelm Schlag, and Avraham Soffer. Dispersive analysis of charge transfer models. *Comm. Pure Appl. Math.*, 58(2):149–216, 2005.
- Sch04. W. Schlag. Stable manifolds for an orbitally unstable nls. preprint, 2004.
- Sch05. W. Schlag. Dispersive estimates for schroedinger operators: a survey. preprint, 2005.
- SS85. Jalal Shatah and Walter Strauss. Instability of nonlinear bound states. *Comm. Math. Phys.*, 100(2):173–190, 1985.
- SS98. I.M. Sigal and A. Soffer. Local decay and velocity bounds for quantum propagation. preprint, 1998.
- SV86. W. A. Strauss and L. Vázquez. Stability under dilations of nonlinear spinor fields. *Phys. Rev. D*, 34:641–643, July 1986.
- SW90. A. Soffer and M. I. Weinstein. Multichannel nonlinear scattering for nonintegrable equations. *Comm. Math. Phys.*, 133(1):119–146, 1990.
- SW92. A. Soffer and M. I. Weinstein. Multichannel nonlinear scattering for nonintegrable equations. II. The case of anisotropic potentials and data. *J. Differential Equations*, 98(2):376–390, 1992.
- SW99. A. Soffer and M. I. Weinstein. Resonances, radiation damping and instability in Hamiltonian nonlinear wave equations. *Invent. Math.*, 136(1):9–74, 1999.
- Tha92. Bernd Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.
- Tsa03. Tai-Peng Tsai. Asymptotic dynamics of nonlinear Schrödinger equations with many bound states. *J. Differential Equations*, 192(1):225–282, 2003.
- TY02a. Tai-Peng Tsai and Horng-Tzer Yau. Asymptotic dynamics of nonlinear Schrödinger equations: resonance-dominated and dispersion-dominated solutions. *Comm. Pure Appl. Math.*, 55(2):153–216, 2002.
- TY02b. Tai-Peng Tsai and Horng-Tzer Yau. Classification of asymptotic profiles for nonlinear Schrödinger equations with small initial data. *Adv. Theor. Math. Phys.*, 6(1):107–139, 2002.
- TY02c. Tai-Peng Tsai and Horng-Tzer Yau. Relaxation of excited states in nonlinear Schrödinger equations. *Int. Math. Res. Not.*, (31):1629–1673, 2002.
- TY02d. Tai-Peng Tsai and Horng-Tzer Yau. Stable directions for excited states of nonlinear Schrödinger equations. *Comm. Partial Differential Equations*, 27(11-12):2363–2402, 2002.
- Wed00. Ricardo Weder. Center manifold for nonintegrable nonlinear Schrödinger equations on the line. *Comm. Math. Phys.*, 215(2):343–356, 2000.
- Yaj95. Kenji Yajima. The $W^{k,p}$ -continuity of wave operators for Schrödinger operators. *J. Math. Soc. Japan*, 47(3):551–581, 1995.
- Yam93. Osanobu Yamada. A remark on the limiting absorption method for Dirac operators. *Proc. Japan Acad. Ser. A Math. Sci.*, 69(7):243–246, 1993.