



HAL
open science

Squeezing and entangling nuclear spins in helium 3

Gael Reinaudi, Alice Sinatra, Aurelien Dantan, Michel Pinard

► **To cite this version:**

Gael Reinaudi, Alice Sinatra, Aurelien Dantan, Michel Pinard. Squeezing and entangling nuclear spins in helium 3. 2005. hal-00016227v1

HAL Id: hal-00016227

<https://hal.science/hal-00016227v1>

Preprint submitted on 21 Dec 2005 (v1), last revised 9 Jan 2006 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Squeezing and entangling nuclear spins in helium 3

G. Reinaudi, A. Sinatra

LKB, ENS, 24 Rue Lhomond, 75231 Paris Cedex 05, France

A. Dantan, M. Pinard

LKB, UPMC, case 74, place Jussieu, 75252 Paris Cedex 05, France

Abstract

We present a realistic model for transferring the squeezing or the entanglement of optical field modes to the collective ground state nuclear spin of ^3He using metastability exchange collisions. We discuss in detail the requirements for obtaining good quantum state transfer efficiency and study the possibility to readout the nuclear spin state optically.

PACS numbers: 03.67.-a, 03.67.Hk, 42.50.Dv, 67.65.+z

I. INTRODUCTION

Helium 3 atoms in their ground state possess a purely nuclear spin $I = 1/2$. Such spins are well-isolated from the environment and show extremely long coherence times. Longitudinal coherence times T_1 of several days are measured in room temperature samples [1]. The transverse coherence time T_2 , which would be as long as T_1 in zero magnetic field, is usually limited by magnetic field inhomogeneity if no special precaution is taken. Transverse coherence times of several hours are observed in a very low field [2]. These very long coherence times originate from the weakness of magnetic coupling on the one hand, and from the absence of electrical coupling on the other hand, as there is no electric quadrupole coupling within the ground state for spins $1/2$. It is tempting to exploit such long-lived coherence for quantum information purposes. In a previous letter [3] we studied the possibility to transfer the squeezing of a cavity mode to ^3He nuclear spins. We showed that the squeezing could be stocked and retrieved from the atoms at a later time realizing thus a quantum memory [4, 5, 6, 7, 8, 9]. For the sake of simplicity we presented in our letter a simplified model involving only two metastable levels and gave numerical results for the more complicated case of ^3He . In this paper we concentrate on ^3He and treat in detail this more realistic case.

In section II we derive metastability exchange collisions equations for populations and coherences of the metastable and ground states. In section III we describe the model for squeezing transfer from a squeezed vacuum mode of the electromagnetic field to the atoms. Numerical results are shown and discussed in section IV. In section V we obtain analytical results in the adiabatic elimination limit for the optical coherences and the cavity field. Section VI is devoted to the readout scheme of the quantum memory. In section VII, as a straightforward application of our scheme, we consider the possibility of creating long-lived quantum correlations between two macroscopic spins, in the move of the successful experiment in Copenhagen [10], in which two macroscopic spins were entangled for 0.5ms, but on a completely different timescale. Finally, in section VIII, we use a toy model to explore the consequences of an imperfect polarization of the atoms on our squeezing transfer scheme.

II. METASTABILITY EXCHANGE COLLISIONS IN HELIUM 3

Over forty years ago, Colegrove, Scheerer and Walters [11] demonstrated a technique to polarize helium ^3He relying on (i) an optical interaction on an infrared transition from the metastable 2^3S triplet state to the 2^3P triplet state, and (ii) *metastability exchange collisions* between atoms in the ground state and in the metastable state. During such a collision, two atoms exchange their electronic degrees of freedom so that the metastable atom, oriented by optical pumping and with a nuclear polarization due to hyperfine coupling in the metastable state, becomes a polarized ground state atom [12]. This technique called *metastability exchange optical pumping* is currently used to prepare polarized samples for nuclear physics experiments as well as in nuclear magnetic resonance imaging applications [13].

In what follows we suggest that metastability exchange collisions can also be used to transfer quantum correlations to the ground state nuclear spin of ^3He .

A. Equations for the one-body density matrix elements

Partridge and Series [12] describe a metastability exchange (ME) collision in terms of the one-body density matrices representing the internal states of two colliding atoms that we name ρ_g^{at} and ρ_m^{at} for the ground and metastable state, respectively. The density matrices after the collision, $\rho_g^{\text{at}'}$ and $\rho_m^{\text{at}'}$, are given by

$$\begin{cases} \rho_g^{\text{at}'} = \text{Tr}_e \rho_m^{\text{at}} \\ \rho_m^{\text{at}'} = \rho_g^{\text{at}} \otimes \text{Tr}_n \rho_m^{\text{at}} \end{cases} \quad (1)$$

where Tr_e and Tr_n are trace operators over the electronic and nuclear variables. ME collisions between these two atoms will occur at a rate γ_{exc} depending on the metastability exchange cross section, the relative velocity of the atoms and the volume explored by the atoms.

Let us consider n metastable atoms with density matrix ρ_m^{at} and N ground state atoms with density matrix ρ_g^{at} . The total number of atoms is $N + n$ and we neglect coherences between the ground and metastable states. The matrices ρ_g^{at} and ρ_m^{at} evolve according to

$$\begin{cases} \frac{d}{dt} \rho_g^{\text{at}} = -\gamma_f \rho_g^{\text{at}} + \gamma_f \text{Tr}_e \rho_m^{\text{at}} \\ \frac{d}{dt} \rho_m^{\text{at}} = -\gamma_m \rho_m^{\text{at}} + \gamma_m \rho_g^{\text{at}} \otimes \text{Tr}_n \rho_m^{\text{at}} \end{cases} \quad (2)$$

where γ_f and γ_m are the metastability exchange collision rates in the ground and metastable states, which satisfy

$$\frac{\gamma_f}{\gamma_m} = \frac{n}{N} \quad (3)$$

The calculation of $d\rho_m/dt$ is performed by expressing ρ in the decoupled spin basis of the nuclear spin $I = \frac{1}{2}$ and the total electronic spin $J = S = 1$ in the metastable state, followed by a projection onto the hyperfine states (eigenstates of the total momentum operator F and F_z) labelled from 1 to 6 as in figure 1. The explicit evolution equations for the density matrix elements are given in the Appendix. The fully polarized state in which all the atoms are in the sublevel with highest angular momentum projection along z is stationary for equations (2).




FIG. 1: Sublevels 1 to 6 are metastable; 9 and 0 are the ground state sublevels. The fully polarized stationary state is shown.

Starting from the one-atom classical equations (2) we proceed in two steps which will be detailed in the following:

1. We consider the equations for the collective density matrix elements (n metastable and N ground state atoms) and linearize these equations around the fully polarized steady state in which the only non-zero elements of ρ are $\rho_{44} = n$ and $\rho_{00} = N$.
2. From the linearized classical equations, interpreted as semiclassical equations for the mean values of the collective operators, we derive the corresponding Heisenberg-Langevin equations.

B. Linearized Heisenberg-Langevin equations

By linearization around the fully polarized solution we obtain equations for the “fluctuations” or deviations of ρ_{ij} from their steady-state values. Such linear equations coincide with the linearized semiclassical equations for the collective atomic operators operators mean values:

$$\rho_{kl} = \langle S_{lk} \rangle \quad k, l = 1, 2, 3 \quad (4)$$

$$\rho_{kl} = \langle I_{lk} \rangle \quad k, l = 9, 0 \quad (5)$$

where $S_{kl} = \sum_{i=1}^n |k\rangle_i \langle l|_i$ for $k, l = 1, 2, 3$ and $I_{kl} = \sum_{i=1}^N |k\rangle_i \langle l|_i$ for $k, l = 9, 0$ are the collective atomic operators in the metastable and ground state, respectively. The corresponding linearized Heisenberg-Langevin equation for the operators are obtained by adding zero-mean valued fluctuating terms which are the Langevin forces. Denoting by f_α the Langevin force for the operator α we get:

$$\dot{S}_{21} = -\gamma_m S_{21} + f_{21} \quad (6)$$

$$\dot{S}_{32} = \frac{2}{9}\gamma_m \left(-\frac{7}{2}S_{32} + \sqrt{3}S_{21} + S_{65} \right) + f_{32} \quad (7)$$

$$\dot{S}_{65} = -\frac{7}{9}\gamma_m \left(S_{65} - \frac{2}{7}\sqrt{3}S_{21} - \frac{2}{7}S_{32} \right) + f_{65} \quad (8)$$

$$\dot{S}_{43} = \gamma_m \left(-\frac{1}{3}S_{43} + \frac{2\sqrt{3}}{9}(S_{32} + S_{65}) \right) + \frac{\sqrt{3}}{3}\gamma_f I_{09} + f_{43} \quad (9)$$

$$\dot{I}_{09} = \frac{1}{3} \left[-3\gamma_f I_{09} + 2\gamma_m \left(S_{32} - \frac{1}{2}S_{65} + \frac{\sqrt{3}}{2}S_{43} + \frac{\sqrt{3}}{2}S_{21} \right) \right] + f_{09} \quad (10)$$

where $\gamma_m = \gamma_{exc}N$ and $\gamma_f = \gamma_{exc}n$ represent the exchange collisions rate for a metastable and a ground state atom, respectively. If α and β denote two system operators, $\langle f_\alpha(t)f_\beta(t') \rangle = D_{\alpha\beta}\delta(t-t')$ where $D_{\alpha,\beta}$ is the corresponding coefficient of the diffusion matrix which can be calculated using the generalized Einstein relations [2] for an ensemble of uncorrelated atoms. The non-zero coefficients are

$$D_{43,34} = \frac{2}{3}\gamma_m n, \quad D_{09,34} = D_{43,90} = -\frac{2\sqrt{3}}{3}\gamma_m n, \quad D_{09,90} = 2\gamma_m n. \quad (11)$$

Langevin forces are necessary to the consistency of the model. Otherwise, the non-Hamiltonian character of the exchange terms leads to a violation of the Heisenberg uncertainty relations. Physically, these forces originate from the fluctuating character of the ME collisions and their correlation time is the collision time, much shorter ($\sim 10^{-13}$ s) than all the times scales we are interested in.

C. Consequences of the Heisenberg-Langevin equations for ME collisions

We notice that Eqs. (6)-(8) for S_{21} , S_{32} , S_{65} form a closed subsystem of equations. This means that in the frequency domain each of these variables can be expressed as a linear combination of the Langevin forces f_{21} , f_{32} , f_{65} . However, in the fully polarized limit we consider here, these Langevin forces do not contribute to the diffusion matrix. It follows that these variables do not contribute to the spin noise and can be neglected. One is then left with only two equations

$$\dot{S}_{43} = -\frac{\gamma_m}{3}S_{43} + \frac{\sqrt{3}}{3}\gamma_f I_{09} + f_{43} \quad (12)$$

$$\dot{I}_{09} = -\gamma_f I_{09} + \gamma_m \frac{\sqrt{3}}{3}S_{43} + f_{09} \quad (13)$$

Let us introduce the transverse spin quadratures

$$S_x = (S_{34} + S_{43})/2, \quad S_y = i(S_{34} - S_{43})/2 \quad (14)$$

and similarly for the ground state, and let us assume that the ground state is initially squeezed, while the metastable atoms are in a coherent spin state. Integrating (12)-(13) with the initial conditions $\Delta \bar{I}_y^2(0) = \Delta I_y^2(0)/(N/4) = e^{-2r}$ and $\Delta \bar{S}_y^2(0) = \Delta S_y^2(0)/(n/4) = 1$ one finds the normalized steady state variances to be

$$\Delta \bar{S}_y^2 = 1 - [1 - e^{-2r}] \frac{3nN}{(3n + N)^2} \quad (15)$$

$$\Delta \bar{I}_y^2 = 1 - [1 - e^{-2r}] \frac{N^2}{(3n + N)^2} \quad (16)$$

Since $n \ll N$ (typically $n/N \sim 10^{-6}$), the ground state spin is still squeezed by approximately the same factor e^{-2r} , whereas the metastable atoms squeezing is negligible (in n/N). By introducing the two-spin correlation functions \mathcal{C}_S and \mathcal{C}_I for the metastable and ground state respectively:

$$\mathcal{C}_S = \frac{\Delta \bar{I}_y^2 - 1}{4n} \quad \text{and} \quad \mathcal{C}_I = \frac{\Delta \bar{I}_y^2 - 1}{4N} \quad (17)$$

this simple calculation shows that ME collisions tend to equalize the correlation function (up to some numerical constant): $\mathcal{C}_S = 3\mathcal{C}_I$. If the ground state spin is squeezed, \mathcal{C}_I has a negative value of order $1/N$, corresponding to maximal collective correlations for the N -particle ensemble. However, as $n \ll N$, this negative value of the correlation function in the metastable state is by far too small to induce squeezing into the n -particle metastable

state, which would require $\mathcal{C}_S \sim -1/n$. We notice that for $e^{-2r} = 1$ we recover the coherent spin state with no correlation between the ground state and the metastable spins.

Noise spectra can also be derived in a similar fashion. By defining the noise spectrum as

$$\mathcal{S}_{x_i x_j}(\omega) = \int d\tau e^{-i\omega\tau} \langle x_i(0)x_j(\tau) \rangle \quad (18)$$

where x_i, x_j are fluctuations of system operators and for the same initial conditions $\Delta \bar{I}_y^2(0) = \Delta I_y^2(0)/(N/4) = e^{-2r}$ and $\Delta \bar{S}_y^2(0) = \Delta S_y^2(0)/(n/4) = 1$ we get:

$$\mathcal{S}_{I_y, I_y}(\omega) = \frac{\pi(Ne^{-2r} + 3n)N^2\delta(\omega)}{2(N + 3n)^2} + \frac{9\gamma_{\text{exc}}nN}{18\omega^2 + 2(N + 3n)^2\gamma_{\text{exc}}^2} \quad (19)$$

$$\mathcal{S}_{S_y, S_y}(\omega) = \frac{3\pi(Ne^{-2r} + 3n)n^2\delta(\omega)}{2(N + 3n)^2} + \frac{3\gamma_{\text{exc}}nN}{18\omega^2 + 2(N + 3n)^2\gamma_{\text{exc}}^2} \quad (20)$$

The equal time correlations (15) and (16) can be recovered from these formulas by integration:

$$\langle x_i x_j \rangle = \frac{1}{2\pi} \int d\omega \mathcal{S}_{x_i x_j}(\omega). \quad (21)$$

For an initial coherent spin state ($e^{-2r} = 1$), the ME collisions process does not change the collective spin variances, but it affects their spectra. The δ -shaped atomic spectra of the two spins in absence of ME collisions acquire a width of order $\gamma_{\text{exc}}(N + 3n)$, that is, of order γ_m . The contribution to the total variance of the ‘‘broad’’ part of the spectrum which is not sensitive to initial squeezing in the system, is large for the metastable state and small for the ground state.

III. THE MODEL FOR SQUEEZING TRANSFER

In figure 2 are represented the ^3He energy levels which are relevant for our squeezing transfer scheme. The atoms interact with a coherent control field of Rabi frequency Ω and frequency ω_1 that we treat classically, and a cavity field described by operators A and A^\dagger . The field injected into the cavity, A_{in} with frequency ω_2 , is in an amplitude-squeezed vacuum state: $\langle A_{in} \rangle = 0$ and $\Delta X_{in}^2 = e^{-2r}$, $\Delta Y_{in}^2 = e^{2r}$, where we have introduced the field quadratures

$$X = A + A^\dagger \quad Y = i(A^\dagger - A). \quad (22)$$

The coherent field (π -polarized) and the squeezed vacuum (σ -polarized) are tuned to the blue side of the so-called C_9 transition ($\lambda = 1.08 \mu\text{m}$) from the $F = 3/2$ level of the 2^3S

metastable state to the 2^3P_0 state, the highest in energy of the 2^3P multiplicity. The atom-field Hamiltonian of the system is:

$$H = H_0 + \hbar \left\{ \Omega S_{73} e^{-i\omega_1 t} + \Omega S_{82} e^{-i\omega_1 t} + g_A A S_{74} + g_B A S_{83} + h.c. \right\} \quad (23)$$

where H_0 describes the atom-field free evolution, $g_{A,(B)} = d_{A,(B)} \sqrt{2\pi\omega_2/\hbar V}$ are the coupling constants between the atoms and the cavity field, V being the volume of the cavity mode and $d_{A,(B)}$ the atomic dipoles of the transitions $7 \leftrightarrow 4$, ($8 \leftrightarrow 3$). The system is initially prepared in the fully polarized state $\langle I_{00} \rangle = N$ and $\langle S_{44} \rangle = n$ by preliminary optical pumping.

Non-Hamiltonian contributions to the evolution of the system operators describe damping of the cavity mode, spontaneous emission from the excited state and the ME collisions described in detail in the previous section. Linearizing the equations in the rotating frame




FIG. 2: a) Metastable and excited sublevels of ^3He . Three coupling constants to the light are introduced. b) Squeezing transfer scheme using a control field Ω , a squeezed vacuum field A and metastability exchange collisions. 9 and 0 are the ground state sublevels.

around the fully polarized state solution we obtain the following closed set of equations:

$$\dot{S}_{21} = -(\gamma_m - i\delta_{12})S_{21} + i\Omega S_{81} + f_{21} \quad (24)$$

$$\dot{S}_{81} = -(\gamma - i(\Delta_{18} - 2\delta_{las}))S_{81} + i\Omega S_{21} + f_{81} \quad (25)$$

$$\dot{S}_{32} = \frac{2}{9}\gamma_m \left(-\frac{7}{2}S_{32} + \sqrt{3}S_{21} + S_{65} \right) + i\delta_{23}S_{32} - i\Omega(S_{38} - S_{72}) + f_{32} \quad (26)$$

$$\dot{S}_{72} = -(\gamma - i(\Delta_{27} - 2\delta_{las}))S_{72} - i\Omega(S_{78} - S_{32}) + f_{72} \quad (27)$$

$$\dot{S}_{43} = \gamma_m \left(-\frac{1}{3}S_{43} + \frac{2\sqrt{3}}{9}(S_{32} + S_{65}) \right) + \frac{\sqrt{3}}{3}\gamma_f I_{09} + i\delta_{34}S_{43} - i\Omega S_{47} + f_{43} \quad (28)$$

$$\dot{S}_{65} = -\frac{7}{9}\gamma_m \left(S_{65} - \frac{2}{7}\sqrt{3}S_{21} - \frac{2}{7}S_{32} \right) + i\delta_{56}S_{65} + f_{65} \quad (29)$$

$$\dot{S}_{47} = -(\gamma + i\Delta_{47})S_{47} - ig_A n A - i\Omega S_{43} + f_{47} \quad (30)$$

$$\dot{S}_{38} = -(\gamma + i\Delta_{38})S_{38} - i\Omega(S_{32} - S_{78}) + f_{38} \quad (31)$$

$$\dot{S}_{78} = -(2\gamma - i\delta_{87})S_{78} - i\Omega(S_{72} - S_{38}) + f_{78} \quad (32)$$

$$\dot{I}_{09} = \frac{1}{3} \left[-3\gamma_f I_{09} + 2\gamma_m \left(S_{32} - \frac{1}{2}S_{65} + \frac{\sqrt{3}}{2}S_{43} + \frac{\sqrt{3}}{2}S_{21} \right) \right] + i\delta_{90}I_{09} + f_{09} \quad (33)$$

$$\dot{A} = -(\kappa + i\Delta_C)A - ig_B S_{38} - ig_A S_{47} + \sqrt{2\kappa}A_{in} \quad (34)$$

where

$$\Delta_{ij} = (E_j - E_i) - \omega_2 \quad (35)$$

$$\delta_{ij} = (E_j - E_i) - \delta_{las} \quad (36)$$

$$\delta_{las} = \omega_1 - \omega_2 \quad (37)$$

and γ is the coherence decay rate due to spontaneous emission from the excited state and collisions. The non-zero diffusion coefficients are

$$D_{43,34} = \frac{2}{3}\gamma_m n, \quad D_{43,90} = -\frac{2\sqrt{3}}{3}\gamma_m n, \quad D_{47,74} = 2\gamma n, \quad D_{09,34} = -\frac{2\sqrt{3}}{3}\gamma_m n, \quad D_{09,09} = 2\gamma_m n \quad (38)$$

We notice that metastable variables S_{72} , S_{32} , S_{78} , S_{21} , S_{81} , S_{65} and S_{38} form a closed subsystem of equations. This means that in the frequency domain each of these variables can be expressed as a linear combination of the Langevin forces f_{72} , f_{32} , f_{78} , f_{21} , f_{81} , f_{65} and f_{38} . However, in the fully polarized limit we consider here, these Langevin forces do not give rise to non-zero diffusion coefficients. It follows that these variables do not contribute to the spin noise and can be neglected. One is then left with only four relevant equations

$$\dot{S}_{43} = -\frac{\gamma_m}{3}S_{43} + \frac{\sqrt{3}}{3}\gamma_f I_{09} + i\delta_{34}S_{43} - i\Omega S_{47} + f_{43} \quad (39)$$

$$\dot{S}_{47} = -(\gamma + i\Delta)S_{47} - ig_A n A - i\Omega S_{43} + f_{47} \quad (40)$$

$$\dot{I}_{09} = \frac{1}{3} \left(-3\gamma_f I_{09} + \gamma_m \sqrt{3} S_{43} \right) + i\delta_{90}I_{09} + f_{09} \quad (41)$$

$$\dot{A} = -(\kappa + i\Delta_C)A - ig_A S_{47} + \sqrt{2\kappa}A_{in} \quad (42)$$

with $\Delta = \Delta_{47}$.

IV. NUMERICAL RESULTS

Equations (39)-(42) can be used to find the variances of the metastable and ground state spin numerically. A typical result is displayed in figure 3, for which we assume that a squeezed vacuum field with $\Delta X_{in}^2 = 0.5$ is injected into the cavity with the coherent




FIG. 3: Symbols: numerical calculations for spin variances in ground state (squares) and metastable state (circles), as a function of the ratio Γ/γ_m (lower x -axis). The corresponding magnetic field needed to satisfy the resonance conditions (45) and (46) is shown in the upper x -axis. Numerical values of parameters are $e^{-2r} = 0.5$, $C = 500$, $\kappa = 100\gamma$, $\Delta = -2000\gamma$, $\gamma = 2 \times 10^7 \text{ s}^{-1}$, $\gamma_m = 5 \times 10^6 \text{ s}^{-1}$. The crossed squares correspond to calculations including an extra relaxation rate $\gamma_0 = 10^3 \text{ s}^{-1}$ for the metastable variables. The lines correspond to the analytical predictions (49) and (50).

control field in the squeezing-transfer configuration. In this figure $\Delta\bar{S}_y^2$ and $\Delta\bar{I}_y^2$ represent the variances of S_y and I_y , both normalized to their coherent spin state values. They are plotted as a function of the ratio Γ/γ_m , where Γ is the pumping parameter

$$\Gamma = \gamma 3\Omega^2(1 + C)/\Delta_{47}^2, \quad (43)$$

and $C = g^2n/(\kappa\gamma)$ the cooperativity. It is precisely this ratio Γ/γ_m which acts as a control parameter to decide how much of the available squeezing of the field is shared between the metastable and the ground state spin. If $\Gamma \gg \gamma_m$, correlations are established among the metastable-state spins, the leakage of correlation towards the ground state being negligible. The metastable collective spin is squeezed while the ground state spin remains unsqueezed. In the opposite limit $\Gamma \ll \gamma_m$, spin exchange is the dominant process for metastable atoms;

they transfer their correlations to the ground state which then becomes squeezed, while the metastable state remains unsqueezed.

In this plot we have chosen the best conditions for squeezing transfer:

1. The metastable coherence S_{43} is resonantly excited by the two fields in a Raman configuration. By introducing the effective two-photon detuning for this coherence

$$\tilde{\delta} = \delta_{34} + \Omega^2/\Delta \quad (44)$$

accounting for the light-shift of level 3, this condition is $\tilde{\delta} = 0$, or

$$(E_4 - E_3)/\hbar + \Omega^2/\Delta = \omega_1 - \omega_2. \quad (45)$$

2. The ground state coherence I_{09} should be resonantly excited by the metastable coherence, i.e.

$$(E_0 - E_9)/\hbar = \omega_1 - \omega_2. \quad (46)$$

In practice a magnetic field (shown as the upper x -axis) can be used to simultaneously fulfill (45) and (46). When the resonance conditions are fulfilled the difference in the Larmor frequencies in the metastable and in the ground state is exactly compensated by the light-shift induced by the coherent control field. Choosing $\Gamma = 0.1\gamma_m$ as a working point, the required field is about $B = 57$ mG, corresponding to $\omega_I = 184$ Hz.

The vapor parameters in the figure correspond to a 1 torr sample at 300 K, with $\gamma_m = 5 \times 10^6$ s⁻¹, and $\gamma = 2 \times 10^7$ s⁻¹, and a metastable atom density of 3.2×10^{10} atoms/cm³ which gives $n/N = 10^{-6}$. The symbols with a cross are a second calculation in which we added a finite relaxation rate in the metastable state γ_0 , to account for the fact that metastable atoms are destroyed as they reach the cell walls. We notice that only the ground state spin squeezing in the region $\Gamma \ll \gamma_m$ is affected.

V. ANALYTICAL RESULTS

In order to have a better physical insight it is possible to find simple analytical results within some reasonable approximation. By adiabatic elimination of the polarization S_{47} and the cavity field assuming $\gamma, \kappa \gg \gamma_m, \gamma_f$, one obtains

$$\dot{S}_{43} + \left(\frac{\gamma_m}{3} + \frac{\Gamma}{3} - i\tilde{\delta}\right)S_{43} = \frac{\gamma_f}{\sqrt{3}}I_{09} + f_{43} - \frac{\Omega}{\Delta}f_{47} + i\frac{\Omega gn}{\Delta}\sqrt{\frac{2}{\kappa}}A_{in} \quad (47)$$

$$\dot{I}_{09} + (\gamma_f - i\delta_{09})I_{09} = \frac{\gamma_m}{\sqrt{3}}S_{43} + f_{09} \quad (48)$$

In deriving (47) we assumed a Raman configuration $\Delta \gg \gamma$, $\frac{C\gamma}{\Delta} \ll 1$ and that the cavity detuning exactly compensates the cavity field dephasing due to the atoms: $\Delta_C = C\kappa\gamma/\Delta$. From equation (47) we see that $(\gamma_m + \Gamma)/3$ is the inverse of the characteristic time constant for the metastable coherence evolution.

A. Resonant case

If the resonance conditions (45) and (46) are satisfied ($\tilde{\delta} = \delta_{09} = 0$) and in the limit $\gamma_f \ll \Gamma, \gamma_m$, we can calculate the variances of the metastable and ground state spins

$$\Delta I_y^2 = \frac{N}{4} \left\{ 1 - \frac{\gamma_m}{\Gamma + \gamma_m} \frac{C}{C + 1} (1 - e^{-2r}) \right\} \quad (49)$$

$$\Delta S_y^2 = \frac{n}{4} \left\{ 1 - \frac{\Gamma}{\Gamma + \gamma_m} \frac{C}{C + 1} (1 - e^{-2r}) \right\} \quad (50)$$

which are plotted as full lines in figure 3.

B. Non-perfectly resonant case

In order to test the robustness of our scheme, let us now concentrate on what happens if the resonance conditions (45) and (46) are only approximatively satisfied. We will focus on the variance of the ground state spin coherence I_{09} .

By adiabatically eliminating the metastable coherence S_{43} one obtains

$$\dot{I}_{09} + [\Gamma_F + ib] I_{09} = f_{09} + \frac{\gamma_m \sqrt{3}}{\gamma_m + \Gamma - i3\tilde{\delta}} \left(f_{43} - \frac{\Omega}{\Delta} f_{47} + i \frac{\Omega g n}{\Delta} \sqrt{\frac{2}{\kappa}} A_{in} \right) \quad (51)$$

The real part in the brackets

$$\Gamma_F = \gamma_f \frac{\Gamma(\gamma_m + \Gamma) + (3\tilde{\delta})^2}{(\gamma_m + \Gamma)^2 + (3\tilde{\delta})^2} \quad (52)$$

is the inverse of the effective time constant for the ground state coherence evolution which would also be the “writing” (or “reading”) time of the quantum memory. $\Gamma_F^{-1} = 2s$ in the example of figure 3 for $\Gamma = 0.1\gamma_m$. It would be proportionally shortened by increasing the

metastable atoms density although Penning collisions prevent in practice metastable atoms densities exceeding 10^{10} - 10^{11} at/cm². The imaginary part in the brackets

$$b = - \left(\gamma_f \frac{3\tilde{\delta}\gamma_m}{(\gamma_m + \Gamma)^2 + (3\tilde{\delta})^2} + \delta_{09} \right) \quad (53)$$

is a light-shift “brought back” to the ground state, which is zero in the resonant case. Equation (51) can be used to calculate the best squeezing (optimized with respect to the transverse spin quadrature) of the ground state coherence: $\Delta I_{best}^2 = \min_{\theta} \Delta I_{\theta}^2$ with $I_{\theta} = I_x \cos \theta + I_y \sin \theta$. We obtain

$$\Delta I_{best}^2 = \frac{N}{4} \left\{ 1 - \frac{\gamma_m}{\Gamma + \gamma_m + (3\tilde{\delta})^2/\Gamma} \frac{C}{C+1} \left[1 - (e^{-2r} + m \sinh(2r)) \right] \right\} \quad (54)$$

where

$$m = 1 - \sqrt{\frac{1}{1 + (b/\Gamma_F)^2}} \quad (55)$$

We show in figure 4 the effect of a frequency mismatch on the normalized spin variances $\Delta \bar{S}_y^2$, $\Delta \bar{I}_y^2$ and the corresponding squeezing transfer efficiencies η_S and η_I

$$\eta_S = \frac{1 - \Delta \bar{S}_y^2}{1 - e^{-2r}} \quad \text{and} \quad \eta_I = \frac{1 - \Delta \bar{I}_y^2}{1 - e^{-2r}} \quad (56)$$

In this example, a frequency mismatch of the order of $\Gamma/3$ in the metastable state or of the order of Γ_F in the ground state affects the efficiency of the squeezing transfer. The condition for the ground state frequency matching (46) imposes stringent requirements on the homogeneity of the magnetic field. Because of the $\sinh(2r)$ in equation (54), the larger the squeezing the worse are the consequences of a mismatch in the condition on $\delta_I = 0$ on the ground state atoms. Physically, if a significant dephasing between the squeezed field and the ground state coherence builds up during the squeezing transfer time, the atoms will see an average between the squeezed and the anti-squeezed quadrature of the field noise. We can easily estimate the required magnetic field homogeneity as follows. Let us introduce the Larmor evolution frequencies in the metastable and ground states: in low field, $\hbar\omega_{\alpha} = \mu_{\alpha}B$ ($\alpha=I,S$) with $\mu_I/h = 3.24\text{kHz/G}$ and $\mu_S/h = 1.87\text{MHz/G}$, and let ΔB be the maximum field difference with respect to the optimal value in the cell volume. For low field, the condition on ΔB to preserve the transfer efficiency reads $\mu_I \Delta B \ll h\Gamma_F$. Since $\Omega^2/\Delta \simeq \Gamma \frac{\Delta}{3\gamma C} \simeq \frac{\mu_S}{h} B$ we get $\frac{\Gamma}{\Gamma_F} \frac{\mu_I}{\mu_S} \frac{\Delta}{3\gamma C} \frac{\Delta B}{B} < 1$ or, in the regime $\Gamma \ll \gamma_m$, $600 \frac{\Delta}{\gamma C} \frac{\Delta B}{B} \ll 1$. With the parameters of figure 3 this gives a condition on the magnetic field inhomogeneity: $\Delta B/B \ll 4 \times 10^{-4}$. In figure




FIG. 4: (a) Normalized ground state spin variance and (b) corresponding quantum transfer efficiency η_I as a function of δ_I/Γ_F (while $\tilde{\delta} = 0$). (c) Normalized metastable state spin variance and (d) corresponding quantum transfer efficiency η_S as a function of $\tilde{\delta}/(\Gamma/3)$ (while $\delta_I = 0$). Symbols: numerical integration of equations (24)-(34). Lines: analytical expression (54). Parameters are the same as in figure 3 and $\Gamma = 0.1\gamma_m$.

5 we calculated the variance of the ground state spin as a function of Γ/γ_m for an increasing inhomogeneity $\Delta B/B$ from zero (dashed line) to 6×10^{-4} . In practice a homogeneity of 100 ppm should be sufficient for the chosen parameters to guarantee that all atoms will be squeezed.

VI. OPTICAL READOUT

A. Outgoing field squeezing

As briefly stressed in [3] the squeezed fluctuations which are stored into the nuclear spins can be retrieved optically in the field exiting the cavity by using the reverse transfer process. Indeed, once the write sequence of the quantum memory has been completed, both the fields




FIG. 5: Normalized “best” variance of the ground state spin as a function of Γ/γ_m (lower x -axis) for an increasing inhomogeneity $\Delta B/B$ from zero to 6×10^{-4} by steps of 1×10^{-4} . On the upper x -axis we show the corresponding homogeneous magnetic field needed to satisfy resonance conditions (45) and (46). Numerical values of parameters are $e^{-2r} = 0.5$, $C = 500$, $\kappa = 100\gamma$, $\Delta = -2000\gamma$, $\gamma = 2 \times 10^7 \text{ s}^{-1}$, $\gamma_m = 5 \times 10^6 \text{ s}^{-1}$, $\gamma_0 = 0$.

and the discharge can be switched off, leaving the atoms in the fundamental state in a spin-squeezed state. After a variable storage time, switching back on only the control field (as well as the discharge) will rapidly put a small fraction of atoms in the metastable state and start the reverse transfer process from the fundamental atoms to the field. The correlations in the ground state will slowly transfer via the metastable state to the intracavity field. This will then result in squeezed fluctuations for the field exiting the cavity, which can be measured by homodyne detection.

More quantitatively, if we still assume that the metastable spin observables and the intra-cavity field adiabatically follow the the ground state spin observables, the evolution equations for the fluctuations of the squeezed component are in the resonant situation and

for $\Gamma \ll \gamma_m$

$$\dot{I}_y(t) = -\Gamma_F I_y(t) + \beta X_{in}(t) + \tilde{f}_y(t) \quad (57)$$

$$X_{out}(t) = \sqrt{2\kappa} X(t) - X_{in}(t) \quad (58)$$

with

$$\beta = \frac{\gamma_m}{\gamma_m + \Gamma} \frac{g_A n \Omega \sqrt{3}}{2\Delta} \sqrt{\frac{2}{\kappa}} \quad (59)$$

$$\tilde{f}_y = \frac{\gamma_m \sqrt{3}}{\gamma_m + \Gamma} \left[\frac{f_{43} - f_{34}}{2i} - \frac{\Omega}{\Delta} \frac{f_{74} - f_{47}}{2i} \right] \quad (60)$$

Denoting by $t = 0$ the start of the readout sequence and by $e^{-2r} = \Delta I_y^2(0)/(N/4)$ the initial squeezing of the ground state nuclear spin, the two-time correlation function of the outgoing field amplitude quadrature can be obtained via (58) after integration of (57)

$$\mathcal{C}(t, t') \equiv \langle X^{out}(t) X^{out}(t') \rangle = \delta(t - t') - 2\Gamma_F \eta_I [1 - e^{-2r}] e^{-\Gamma_F(t+t')} \quad (61)$$

The δ -correlated term corresponds to the vacuum fluctuations contribution, whereas the second term corresponds to a transient squeezing for the outgoing field which is proportional to the initial atomic squeezing. In (61), η_I designates the optimal quantum transfer efficiency in the ground state

$$\eta_I = \frac{\gamma_m}{\gamma_m + \Gamma} \frac{C}{C + 1} \quad (62)$$

The ground state squeezing can be adequately measured by homodyne detection using a temporally matched local oscillator as shown in Refs. [6, 14]. Using a local oscillator with envelope $\mathcal{E}(t)$ the normalized power measured by a Fourier-limited spectrum analyzer integrating over a time T is given by

$$P(t) = \frac{1}{\mathcal{E}(t)^2} \int_{-\pi/T}^{\pi/T} \frac{d\omega}{2\pi} \int_t^{t+T} d\tau \int_t^{t+T} d\tau' e^{-i\omega(\tau-\tau')} \mathcal{E}(\tau) \mathcal{E}(\tau') \mathcal{C}(\tau, \tau') \quad (63)$$

In order to measure the atomic squeezing one has to maximize the temporal overlap between the local oscillator and the field radiated by the atoms: $E(t) \propto e^{-\Gamma_F t}$. For such a local oscillator and for an integration time longer than the readout time Γ_F^{-1} the measured power can be written as the sum of a shot-noise term \mathcal{N} and a time-dependent signal term \mathcal{S} proportional to the initial squeezing:

$$P(t) = \mathcal{N} + \mathcal{S} [1 - e^{-2r}] e^{-2\Gamma_F t} \quad (64)$$

with $\mathcal{S} \simeq \eta_I \mathcal{N}$. The ground state nuclear spin fluctuations can therefore be measured optically with the same efficiency η_I as in the write sequence. However, because of the slow character of the correlation transfer process in the ground state the readout time is as long as the write time.

As expected it is not possible to access the quantum memory faster during the readout than during the write phase. One could think of a faster readout method by transferring the fundamental atoms fluctuations to the metastable atoms and perform the optical readout in the regime $\Gamma \gg \gamma_m$. However, as we showed in section II, starting with a squeezed fundamental spin and first switching on the discharge (without the fields) will transfer very few correlations from the fundamental to the metastable atoms ($\eta_I \ll 1$), and almost no squeezing will be retrieved in the field.

VII. ENTANGLING TWO SEPARATE SAMPLES

A direct and important extension of the previous results is that it is possible to transfer quantum correlations between different light beams to two spatially separated nuclear spins. If one disposes of EPR fields this allows to entangle two separate ensembles [16]. Such EPR atomic states are very useful for quantum information protocols involving the manipulation of continuous variable entanglement, such as atomic teleportation for instance [17].

Let us consider two identical ensembles 1 and 2 illuminated by EPR-correlated vacuum fields $A_{in}^{(i)}$ and coherent control fields Ω_i ($i = 1, 2$). Without loss of generality we assume symmetrical field correlations of the form

$$\Delta^2 X_{in}^{(i)} = \Delta^2 Y_{in}^{(i)} = \cosh(2r) \quad (i = 1, 2) \quad (65)$$

$$\langle X_{in}^{(i)} X_{in}^{(j)} \rangle = -\langle Y_{in}^{(i)} Y_{in}^{(j)} \rangle = \sinh(2r) \quad (i \neq j), \quad (66)$$

i.e. that the amplitude quadratures are correlated and the phase quadratures anti-correlated: $\Delta^2(X_{in}^{(1)} - X_{in}^{(2)}) = \Delta^2(Y_{in}^{(1)} + Y_{in}^{(2)}) = e^{-2r}$. For perfect entanglement ($r = \infty$) these EPR variances vanish. Both spins are initially prepared in a coherent spin state and we assume an equal incident power on both cavities ($\Omega_1 = \Omega_2$). Under the same adiabatic approximations as before, the fluctuations of the transverse spin components satisfy equation of the form (57)

$$\dot{I}_{xi} = -\Gamma_F I_{xi} - \beta Y_{in}^{(i)} + \tilde{f}_{xi}, \quad (67)$$

$$\dot{I}_{yi} = -\Gamma_F I_{yi} + \beta X_{in}^{(i)} + \tilde{f}_{yi}, \quad (68)$$

($i = 1, 2$). Because of the linearity of the coupling in this regime, the EPR atomic nuclear spin operators, $I_{x1} + I_{x2}$ and $I_{y1} - I_{y2}$, are clearly coupled to the EPR field operators

$$\frac{d}{dt}(I_{x1} + I_{x2}) = -\Gamma_F(I_{x1} + I_{x2}) - \beta(Y_{in}^{(1)} + Y_{in}^{(2)}) + \tilde{f}_{x1} + \tilde{f}_{x2} \quad (69)$$

$$\frac{d}{dt}(I_{y1} - I_{y2}) = -\Gamma_F(I_{y1} - I_{y2}) + \beta(X_{in}^{(1)} - X_{in}^{(2)}) + \tilde{f}_{y1} - \tilde{f}_{y2} \quad (70)$$

The amount of EPR-type correlations between the incident fields is given by the half-sum of the EPR variances

$$\mathcal{E}_f = \frac{1}{2} \left[\Delta^2(X_{in}^{(1)} - X_{in}^{(2)}) + \Delta^2(Y_{in}^{(1)} + Y_{in}^{(2)}) \right] = 2e^{-2r} \quad (71)$$

In the Gaussian approximation the entanglement between the nuclear spins can be evaluated using the same quantity (also normalized to 2)

$$\mathcal{E}_I = \frac{2}{N} \left[\Delta^2(I_{x1} + I_{x2}) + \Delta^2(I_{y1} - I_{y2}) \right] \quad (72)$$

It follows from (69-70) that the last two quantities are simply related by

$$\mathcal{E}_I = \eta_I \mathcal{E}_f + 2(1 - \eta_I). \quad (73)$$

Like squeezing entanglement can also be in principle perfectly mapped onto the nuclear spins with an efficiency η_I (62), close to unity in the regime $\Gamma \gg \gamma_m$ and $C \gg 1$. It is interesting to note that in this case the two-spin correlation function of individual spins *inside* one ensemble

$$\mathcal{C}^{(1)} \equiv \overline{\langle I_{x1}^{(i)} I_{x1}^{(j)} \rangle} \simeq \frac{\cosh(2r) - 1}{4N} \quad (74)$$

and the correlation function between two spins belonging to *different* ensembles

$$\mathcal{C}^{(2)} = \overline{\langle I_{x1}^{(i)} I_{x2}^{(j)} \rangle} \simeq \frac{\sinh(2r)}{4N} \quad (75)$$

become approximately equal for a large entanglement - $\mathcal{C}^{(1)} \sim \mathcal{C}^{(2)}$ - which means that one individual spin is about as much correlated with the other spins in its own ensemble as with the spins of the other ensemble.

VIII. THE IMPERFECT POLARIZATION CASE

The nuclear polarization of the sample is defined as

$$P = \frac{\langle I_{00} \rangle - \langle I_{99} \rangle}{\langle I_{00} \rangle + \langle I_{99} \rangle} \quad (76)$$

In practice polarization between 80% and 85% are currently achieved by optical pumping in dilute ^3He samples [18]. If the atoms are prepared in a state which is not fully polarized - $P \neq 1$ - the situation is clearly more complicated than we described in [3] and in the present paper. In particular, equations (24)-(34) and (38) obtained by linearization around the fully polarized state are no longer valid. We did not perform a complete analysis in the $P \neq 1$ case. However, one can have a good idea of the result by using the simplified model of [3] which involves only two metastable sublevels (see figure 6). As in section III, a Raman transition is driven by a coherent control field of Rabi frequency Ω and a squeezed vacuum cavity field:

$$H = H_0 + \hbar \left\{ \Omega S_{31} e^{-i\omega_1 t} + g A S_{32} + \text{h.c.} \right\}. \quad (77)$$

In this toy-model the control field Ω also acts as an optical pumping beam (able to transfer the atoms from sublevel 9 to sublevel 0) and we introduce explicitly a relaxation in the ground state, so that $P \neq 1$ in steady state. Let us introduce for this model the rescaled




FIG. 6: Sublevels 1 and 2 are metastable, level 3 is the excited state, 9 and 0 are the ground state sublevels. We include a relaxation process in the ground state so that $P \neq 1$ in steady state.

coupling constant \tilde{g} , the atomic one-photon detunings Δ_1 and Δ , the two-photon detunings \tilde{d} and δ_I , and two pumping parameters Γ_p and Γ :

$$\Delta_1 = (E_3 - E_1)/\hbar - \omega_1, \quad \Delta = (E_3 - E_2)/\hbar - \omega_2, \quad (78)$$

$$\tilde{\delta} = \Delta_1 - \Delta + \frac{\Omega^2}{\Delta}, \quad \delta_I = (E_0 - E_9)/\hbar - (\omega_1 - \omega_2), \quad (79)$$

$$\Gamma_P = \gamma \frac{\Omega^2}{\Delta^2}, \quad \Gamma = \Gamma_p(1 + C), \quad (80)$$

where γ is the optical coherence decay rate and C is the cooperativity parameter defined by equation (43). For the atomic operators we introduce $S_+ = S_{21}$, $S_- = S_{12}$

$$S_x = (S_- + S_+)/2, \quad S_y = i(S_- - S_+)/2, \quad S_z = (S_{22} + S_{11})/2, \quad (81)$$

and similarly for the ground state operators. In the limit of large one photon detunings the excited state and the optical coherences can be adiabatically eliminated, yielding a set of equations similar to those of Ref. [15] with the addition of metastability exchange. By adiabatically eliminating the field (assumed to be resonant in the cavity) and for $\tilde{\delta} = 0$, $\delta_I = 0$, we obtain:

$$\dot{S}_+ = -(\Gamma_p + \gamma_m)S_+ + \gamma_f I_+ + 2i\tilde{g}AS_+ + f_{S_+} \quad (82)$$

$$\dot{S}_z = -(\Gamma_p + \gamma_m)S_z + \gamma_f I_z + \frac{n\Gamma_p}{2} + i\tilde{g}[A^\dagger S_+ - AS_-] + f_{S_z} \quad (83)$$

$$\dot{A} = -(\kappa + i\Delta_C)A + i\tilde{g}S_+ + \sqrt{2\kappa}A_{in} \quad (84)$$

$$\dot{I}_+ = -(\gamma_f + \Gamma_1)I_+ + \gamma_m S_+ + f_{I_+} \quad (85)$$

$$\dot{I}_z = -(\gamma_f + \Gamma_1)I_z + \gamma_m S_z + f_{I_z} \quad (86)$$

The semiclassical version of equations (82)-(86) has a stationary solution $\langle S_+ \rangle = \langle I_+ \rangle = \langle A_{in} \rangle = 0$ and with

$$\langle S_z \rangle = \frac{\gamma_f + \Gamma_1}{\gamma_m} \langle I_z \rangle P = \frac{1}{1 + \Gamma_1(\Gamma_p + \gamma_m)/(\Gamma_p \gamma_f)} \quad (87)$$

We will have in practice $\Gamma_1 \ll \gamma_f$, meaning that the nuclear polarization in the metastable state $P^* = \langle S_z \rangle / (n/2)$ and the nuclear polarization in the ground state $P = \langle I_z \rangle / (N/2)$ are almost equal. In this toy-model the stationary P is determined by the balance between the decay Γ_1 and the pumping Γ_p . In reality, the atoms will be previously pumped more efficiently with resonant light. When we linearize the equations around the steady state we obtain

$$\dot{S}_+ = -(\tilde{\Gamma} + \gamma_m)S_+ + \frac{ig\Omega n}{\kappa\Delta} P^* \sqrt{2\kappa} A_{in} + \gamma_f I_+ + f_{S_+} \quad (88)$$

$$\dot{I}_+ = -(\gamma_f + \Gamma_1)I_+ + \gamma_m S_+ + f_{I_+} \quad (89)$$

with

$$\tilde{\Gamma} = \Gamma_p(1 + CP^*) \quad (90)$$

Starting from equations (88)-(89) one can proceed as in section V to obtain

$$\Delta \bar{I}_y^2 = \frac{\Delta I_y^2}{NP/4} = \frac{1}{P} + (P^* e^{-2r} - 1)\eta'_I + \frac{\eta'_I}{2\tilde{C}} \frac{P^* - 1}{P} \quad (91)$$

where

$$\tilde{C} = CP^* \quad \text{and} \quad \tilde{\Gamma}_f = \frac{\gamma_f \tilde{\Gamma}}{\tilde{\Gamma} + \gamma_m} \quad (92)$$

$$\eta'_I = \frac{\tilde{C}}{\tilde{C} + 1} \frac{\gamma_m}{\tilde{\Gamma} + \gamma_m} \frac{\tilde{\Gamma}_f}{\tilde{\Gamma}_f + \Gamma_1} \quad (93)$$

For $P^* \simeq P$ and $\tilde{C} \gg 1$, we have finally

$$\Delta \bar{I}_y^2 = \eta'_I e^{-2r} + (1 - \eta'_I)/P \quad (94)$$

Equation (94) shows that the main consequence of having $P \neq 1$ is a rescaling of the cooperativity and the pumping parameter $\tilde{\Gamma}$ and the quantum transfer efficiency η'_I , which are reduced by a factor P . Let us note that, for $P \neq 1$, when no squeezing enters the cavity, the atoms are no longer in a coherent spin state. This shows, however, that strong squeezing transfer is still possible with a non-ideal polarization.

Acknowledgments

Laboratoire Kastler Brossel is UMR 8552 du CNRS, de l'ENS et de l'UPMC. This work was supported by the COVAQIAL European project No. FP6-511004.

IX. APPENDIX

Evolution equations of the density matrix elements under ME collisions are:

$$\begin{aligned} \frac{d}{dt} \rho_{11} &= \gamma_{\text{exc}} \left(-N\rho_{11} + \frac{1}{3}\rho_{99} (\rho_{22} + 3\rho_{11} + 2\rho_{55}) \right) \\ \frac{d}{dt} \rho_{12} &= \gamma_{\text{exc}} \left(-N\rho_{12} + \frac{2}{9}\rho_{99} \left((\rho_{23} + \rho_{56}) \sqrt{3} + 3\rho_{12} \right) + \frac{\sqrt{3}}{9}\rho_{90} (\rho_{22} + 3\rho_{11} + 2\rho_{55}) \right) \end{aligned}$$

$$\begin{aligned}
\frac{d}{dt}\rho_{13} &= \gamma_{\text{exc}} \left(-N\rho_{13} + \frac{1}{3}\rho_{99}(\rho_{13} + \rho_{24}) + \frac{2}{9}\rho_{90}((\rho_{23} + \rho_{56})\sqrt{3} + 3\rho_{12}) \right) \\
\frac{d}{dt}\rho_{14} &= \gamma_{\text{exc}} \left(-N\rho_{14} + \frac{\sqrt{3}}{3}\rho_{90}(\rho_{13} + \rho_{24}) \right) \\
\frac{d}{dt}\rho_{22} &= \gamma_{\text{exc}} \left(-N\rho_{22} + \frac{2}{9}\rho_{99}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) + \frac{2}{9}\rho_{90}(\sqrt{3}\rho_{21} + \rho_{65} + \rho_{32}) \right. \\
&\quad \left. + \frac{2}{9}\rho_{09}(\sqrt{3}\rho_{12} + \rho_{23} + \rho_{56}) + \frac{1}{9}\rho_{00}(\rho_{22} + 3\rho_{11} + 2\rho_{55}) \right) \\
\frac{d}{dt}\rho_{23} &= \gamma_{\text{exc}} \left(-N\rho_{23} + \frac{2}{9}\rho_{99}(\sqrt{3}\rho_{34} + \rho_{56} + \rho_{23}) + \frac{2}{9}\rho_{90}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) \right. \\
&\quad \left. + \frac{\sqrt{3}}{9}\rho_{09}(\rho_{13} + \rho_{24}) + \frac{2}{9}\rho_{00}(\sqrt{3}\rho_{12} + \rho_{23} + \rho_{56}) \right) \\
\frac{d}{dt}\rho_{24} &= \gamma_{\text{exc}} \left(-N\rho_{24} + \frac{2}{9}\rho_{90}(\sqrt{3}(\rho_{23} + \rho_{56}) + 3\rho_{34}) + \frac{1}{3}\rho_{00}(\rho_{13} + \rho_{24}) \right) \\
\frac{d}{dt}\rho_{33} &= \gamma_{\text{exc}} \left(-N\rho_{33} + \frac{1}{9}\rho_{99}(2\rho_{66} + \rho_{33} + 3\rho_{44}) + \frac{2}{9}\rho_{90}(\sqrt{3}\rho_{43} + \rho_{65} + \rho_{32}) \right. \\
&\quad \left. + \frac{2}{9}\rho_{09}(\sqrt{3}\rho_{34} + \rho_{56} + \rho_{23}) + \frac{2}{9}\rho_{00}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) \right) \\
\frac{d}{dt}\rho_{34} &= \gamma_{\text{exc}} \left(-N\rho_{34} + \frac{\sqrt{3}}{9}\rho_{90}(2\rho_{66} + \rho_{33} + 3\rho_{44}) + \frac{2}{9}\rho_{00}(\sqrt{3}(\rho_{23} + \rho_{56}) + 3\rho_{34}) \right) \\
\frac{d}{dt}\rho_{44} &= \gamma_{\text{exc}} \left(-N\rho_{44} + \frac{1}{3}\rho_{00}(2\rho_{66} + \rho_{33} + 3\rho_{44}) \right) \\
\frac{d}{dt}\rho_{55} &= \gamma_{\text{exc}} \left(-N\rho_{55} + \frac{1}{9}\rho_{99}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) - \frac{2}{9}\rho_{90}(\sqrt{3}\rho_{21} + \rho_{65} + \rho_{32}) \right. \\
&\quad \left. - \frac{2}{9}\rho_{09}(\sqrt{3}\rho_{12} + \rho_{23} + \rho_{56}) + \frac{2}{9}\rho_{00}(\rho_{22} + 3\rho_{11} + 2\rho_{55}) \right) \\
\frac{d}{dt}\rho_{56} &= \gamma_{\text{exc}} \left(-N\rho_{56} + \frac{2}{9}\rho_{99}(\sqrt{3}\rho_{34} + \rho_{56} + \rho_{23}) - \frac{1}{9}\rho_{90}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) \right. \\
&\quad \left. - \frac{2\sqrt{3}}{9}\rho_{09}(\rho_{13} + \rho_{24}) + \frac{2}{9}\rho_{00}(\sqrt{3}\rho_{12} + \rho_{23} + \rho_{56}) \right) \\
\frac{d}{dt}\rho_{66} &= \gamma_{\text{exc}} \left(-N\rho_{66} + \frac{2}{9}\rho_{99}(2\rho_{66} + \rho_{33} + 3\rho_{44}) - \frac{2}{9}\rho_{90}(\sqrt{3}\rho_{43} + \rho_{65} + \rho_{32}) \right. \\
&\quad \left. - \frac{2}{9}\rho_{09}(\sqrt{3}\rho_{34} + \rho_{56} + \rho_{23}) + \frac{1}{9}\rho_{00}(2\rho_{22} + \rho_{55} + \rho_{66} + 2\rho_{33}) \right) \\
\frac{d}{dt}\rho_{00} &= \gamma_{\text{exc}} \left(-n\rho_{00} + \frac{1}{3}(3\rho_{44} + \rho_{66} + \rho_{22} + 2\rho_{55} + 2\rho_{33})N \right) \\
\frac{d}{dt}\rho_{09} &= \gamma_{\text{exc}} \left(-n\rho_{09} + \frac{1}{3}N((\rho_{43} + \rho_{21})\sqrt{3} + 2\rho_{32} - \rho_{65}) \right) \\
\frac{d}{dt}\rho_{99} &= \gamma_{\text{exc}} \left(-n\rho_{99} + \frac{1}{3}(\rho_{33} + 2\rho_{22} + 3\rho_{11} + \rho_{55} + 2\rho_{66})N \right)
\end{aligned}$$

-
- [1] W. Heil, H. Humblota, E. Otten, M. Schafera, R. Sarkaua and M. Leduc, Phys. Lett A, **201**, 337 (1995)
- [2] C.Cohen-Tannoudji, J. Dupont-Roc, S. Haroche and F. Laloë, Phys. Rev. Lett. **22**, 758 (1969)
- [3] A. Dantan, G. Reinaudi, A. Sinatra, F. Laloë, E. Giacobino, M. Pinard, Phys. Rev. Lett. **95**, 123002 (2005)
- [4] M. Fleischhauer and M.D. Lukin, Phys. Rev. Lett. **84**, 5094 (2000)
- [5] A.E. Kozhokin, K. Mølmer and E. Polzik, Phys. Rev. A **62**, 033809 (2000)
- [6] A. Dantan and M. Pinard, Phys. Rev. A **69**, 43810 (2004)
- [7] C.H. van der Wal *et al.*, Science **301**, 196 (2003)
- [8] C. Schori, B. Julsgaard, J.L. Sørensen, E.S. Polzik, Phys. Rev. Lett. **89**, 57903 (2002)
- [9] B. Julsgaard, J. Sherson, J.I. Cirac, J. Fiurášek, E.S. Polzik, Nature (London) **432**, 482 (2004)
- [10] B. Julsgaard, A. Kozhokin, E.S. Polzik, Nature (London) **413**, 400 (2001)
- [11] F.D. Colegrove, L.D. Schearer and G.K. Walters, Phys. Rev. **132**, 2561 (1963)
- [12] R.B. Partridge and G.W. Series, Proc. Phys. Soc. **88**, 983 (1966)
- [13] J. Becker *et al.*, Nucl. Instrum. Methods Phys. Res. A **402**, 327 (1998); H. Moller *et al.*, Magn. Reson. Med. **47**, 1029 (2002)
- [14] A. Dantan, J. Cviklinski, M. Pinard, Ph. Grangier, quant-ph/0512175.
- [15] A. Dantan, M. Pinard, V. Josse, N. Nayak, P.R. Berman, Phys. Rev A **67**, 045801 (2003)
- [16] A. Dantan, A. Bramati and M. Pinard, Europhys. Lett. **67**, 881 (2004)
- [17] A. Dantan, N. Treps, A. Bramati, M. Pinard, Phys. Rev. Lett. **95**, 050520 (2005)
- [18] P.-J. Nacher, M. Leduc, J. Physique **46**, 2057 (1985).