



HAL
open science

Nonparametric inference of photon energy distribution from indirect measurements

Éric Moulines, François Roueff, Antoine Souloumiac, Thomas Trigano

► **To cite this version:**

Éric Moulines, François Roueff, Antoine Souloumiac, Thomas Trigano. Nonparametric inference of photon energy distribution from indirect measurements. 2005. hal-00016328v1

HAL Id: hal-00016328

<https://hal.science/hal-00016328v1>

Preprint submitted on 27 Dec 2005 (v1), last revised 12 Jun 2006 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NONPARAMETRIC INFERENCE OF PHOTON ENERGY DISTRIBUTION FROM INDIRECT MEASUREMENTS

E. MOULINES*¹, F. ROUEFF¹, A. SOULOUMIAC² AND T. TRIGANO¹

¹ *ENST Paris, CNRS/LTCl, 34 rue Dareau, 75014 Paris, France.
E-mails: moulines, roueff, trigano@tsi.enst.fr*

² *SSTM/LETS, CEA Saclay, 91191 Gif/Yvette CEDEX, France.
E-mail: antoine.souloumiac@cea.fr*

ABSTRACT. We consider a density estimation problem arising in nuclear physics. Gamma photons are impinging on a semiconductor detector, producing pulses of current. The integral of this pulse is equal to the total amount of charges created by the photon in the detector, which is linearly related to the photon energy. Because the inter-arrival of photons can be shorter than the charge collection time, pulses associated to different photons may overlap leading to a phenomenon known as pileup. The distortions on the photon energy spectrum estimate due to pileup becomes worst when the photon rate increases, making pileup correction techniques a must for high-counting rate experiments. In this paper, we present a novel technique to correct pileup, which extends a method introduced in Hall and Park (2004) for the estimation of the service time from the busy period in M/G/ ∞ models. It is based on a novel formula linking the joint distribution of the energy and duration of the cluster of pulses and the distribution of the energy of the photons. We then assess the performance of this estimator by providing an expression of its integrated square error. A limited Monte-Carlo experiment is presented to illustrate on practical examples the benefits of the pileup correction.

Keywords: indirect observations; marked Poisson processes; nonlinear inverse problems; nonparametric density estimation

1. INTRODUCTION

We consider a problem occurring in nuclear spectroscopy. A radioactive source (a mixture of radionuclides) emits photons which impinge on a semiconductor detector. Photons (X and gamma rays) interact with the semiconductor crystal to produce electron-hole pairs. The migration of these pairs in the semiconductor produce a finite duration pulse of current. Under appropriate experimental conditions (ultra-pure crystal, low temperature), the integral over time of this pulse of current corresponds to the total amount of electron-hole pairs created in the detector, which is proportional to the energy deposited in the semiconductor (see for instance Knoll (1989) or Leo (1994)). In most

classical semiconductor radiation detectors, the pulses amplitudes are recorded and sorted to produce histograms showing, as a function of energy, the number of photons absorbed by the detector. This histogram is used as an estimate of the photon energy distribution (called the *energy spectrum*).

The times between pulses generated by a radiation detector are known as inter-arrival times. The inter-arrival times of pulses are independent of the electrical pulses, and can therefore be shorter than the typical duration of the charge collection, thus creating *clusters* (see Figure 1). In gamma ray spectrometry, this phenomenon is referred to as *pileup*. The phenomenon of pileup causes a distortion of the acquired energy spectrum which becomes more severe as the incoming counting rate increases. This problem has been studied by many authors in the field of nuclear instrumentation since the 1960's (see Bristow (1990) for a detailed review of these early contributions; classical pileup correction techniques are detailed in the ANSI norm (1999)).




FIGURE 1. Illustration of the pile-up phenomenon. (a) : input signal with arrival times T_j , lengths X_j and energies Y_j , $j = n, \dots, n + 2$; (b) : associated on-off process S_t , $S_t = 1$ if the system is busy at time t and $S_t = 0$ otherwise.

In mathematical terms, the problem can be formalized as follows. Denote by $\{T_k, k \geq 1\}$ the sequence of arrival times of the photons, assumed to be the ordered points of an homogeneous Poisson process. The current intensity as a function of time can be modeled as a shot-noise process

$$W(t) \stackrel{\text{def}}{=} \sum_{k \geq 1} F_k(t - T_k), \quad (1)$$

where $\{F_k(s), k \geq 1\}$ are the contributions of each individual photon to the overall intensity. By analogy with queuing models, we call $\{W(t), t \geq 0\}$ the *workload* process. The current pulse $\{F_k(s), k \geq 1\}$ are assumed to be independent copies of a continuous time stochastic process $\{F(s), s \geq 0\}$ whose path has a finite support $[0, X]$ a.s., where X is the pulse duration (the duration of the charge collection). The integral of the pulse $Y \stackrel{\text{def}}{=} \int_0^X F(s) ds$ is equal to the total amount of charge collected for a single photon. Under appropriate experimental condition, this quantity may be shown to be linearly related to the photon energy; for simplicity Y is referred to as the *energy* in the sequel.

The restriction of the workload process to a maximal segment where it is positive (resp. 0) is referred to as a *busy* (resp. *idle*) *period*; in the coverage process literature, these quantities are also referred to as *spacings* and *clumps*. An idle period followed by a busy period is called a *cycle*.

In our experimental setting, the sequence of pulse duration and energy $\{(X_k, Y_k), k \geq 1\}$ is not directly observed. Instead, the only available data are the durations of the busy and idle periods and the total amounts of charge collected on busy periods. Define the on-off process

$$S_t = \sum_{k \geq 1} \mathbb{1}_{[T'_k, T'_k + X'_k)}(t), \quad (2)$$

where $\{T'_k, k \geq 1\}$ is the ordered sequence of busy periods arrivals and $\{X'_k, k \geq 1\}$ the corresponding sequence of durations. We further define, for all $k \geq 1$, $Y'_k \stackrel{\text{def}}{=} \int_{T'_k}^{T'_k + X'_k} W(t) dt$, the total amount of charge of the k -th busy period. Finally we denote by Z_k the duration of the k -th idle period, $Z_1 = T_1$ and, for all $k \geq 2$, $Z_k = T'_k - (T'_{k-1} + X'_{k-1})$. We consider the problem of estimating the distribution of the photon energy Y , having observed n cycles $\{(Z_k, X'_k, Y'_k), k = 1, \dots, n\}$. In the terminology introduced by Pyke (1958), this corresponds to a type II counter.

The problem shares some similarity with service time distribution from busy and idle measurements in a M/G/ ∞ model (see for instance Baccelli and Brémaud (2002)). Note indeed that the M/G/ ∞ model is a particular instance of the above setting, as it corresponds to having $F = \mathbb{1}_{[0, X)}$, so that $X = Y$. There exists a vast literature in

this particular case. Takacs (1962) (see also Hall (1988)) has derived a closed-form relation linking the cumulative distributions functions (cdfs) of the service time X and busy period X' . Bingham and Pitts (1999) derived from this formula an estimator of the service time distribution X , which they apply to the study of biological signals. An alternative estimator has been recently introduced in Hall and Park (2004), in which a kernel-estimator of the probability density function (pdf) of X is derived in a nonparametric framework, together with a bound of the pointwise error.

Although our estimator can be applied to the $M/G/\infty$ framework (thus allowing a comparison with Hall and Park (2004) in this special case), we stress on the fact that we are dealing here simultaneously with durations and energies, without assuming any particular dependence structure between them. Secondly, the main emphasis in the photon problem consists in estimating the distribution of the photon energy and not the distribution of the duration, in sharp contrast with the $M/G/\infty$ problem.

The paper is organized as follows. We give the notations and main assumptions in Section 2, and enounce the basic properties of the model. In Section 3 we present an inversion formula relating the Laplace transform of the cluster duration/energy to the Laplace transform of the density function of interest. We also derive an estimator of this function, which is based on an empirical version of the inversion formula and kernel smoothing. Our main result is presented in Section 4, showing that this estimator achieves standard minimax rates in the sense of the Integrated Squared Error when the pulse duration is almost-surely upper bounded. The study of this error is detailed in Section 5. Some applications and examples are shown in Section 6. Since the present paper is directed towards establishing a theory, practical aspects are not discussed in much details in the present contribution and we refer to Trigano et al. (2005) for a thorough discussion of the implementation and applications to real data. Proofs of the different propositions are presented in appendix.

2. NOTATIONS AND MAIN ASSUMPTIONS

All along the paper, we suppose that

- (H-1) $\{T_k, k \geq 1\}$ is the ordered sequence of the points of a homogeneous Poisson process on the positive half-line with intensity λ .
- (H-2) $\{(X, Y), (X_k, Y_k), k \geq 1\}$ is a sequence of independent and identically distributed $(0, \infty)^2$ -valued random variables with probability distribution denoted by P and independent of $\{T_k, k \geq 1\}$. In addition, $\mathbb{E}[X]$ and $\mathbb{E}[Y]$ are finite.

In other words, $\{(T_k, X_k, Y_k), k \geq 1\}$ is a Poisson point process with control measure $\lambda \text{Leb} \otimes P$, where Leb denotes the Lebesgue measure on the positive half-line. Let us recall a few basic properties satisfied under this assumption by the sequence $\{(Z_k, X'_k, Y'_k), k \geq 1\}$ defined in the introduction. By the lack of memory property of the exponential distribution, the idle periods are independent and identically distributed with common exponential distribution with parameter λ . Moreover they are independent of the busy periods, which also are independent and identically distributed. We denote by (X', Y') a couple having the same distribution as the variables of the sequence $\{(X'_k, Y'_k), k \geq 1\}$ and by P' its probability measure. Using that $\mathbb{E}[X]$ and $\mathbb{E}[Y]$ are finite, it is easily shown that

$$\begin{aligned}\mathbb{E}[X'] &= \{\exp(\lambda \mathbb{E}[X]) - 1\} / \lambda \\ \mathbb{E}[Y'] &= \mathbb{E}[Y] \exp(\lambda \mathbb{E}[X]) .\end{aligned}$$

Our goal is the nonparametric estimation of the distribution of Y ; hence we assume that

(H-3) Y admits a probability density function denoted by m , i.e. $\int_{x>0} P(dx, dy) = m(y) \text{Leb}(dy)$.

As mentioned in Section 1, the marks $\{(X_k, Y_k), k \geq 1\}$ are not directly observed but, instead, we observe the sequence $\{(T'_k, X'_k, Y'_k), k = 1, \dots, n\}$, i.e. the arrival times, duration and integrated energy of the successive busy periods. These quantities are recursively defined as follows. Let $T'_1 = T_1$ and for all $k \geq 2$,

$$T'_k = \inf \left\{ T_i : T_i > \left(T'_{k-1} \vee \max_{j \leq i-1} (T_j + X_j) \right) \right\} ; \quad (3)$$

for all $k \geq 1$,

$$\begin{aligned}X'_k &= \max_{T_i \in [T'_k, T'_{k+1}[} \{T_i + X_i\} - T'_k , \\ Y'_k &= \sum_{i \geq 1} Y_i \mathbb{1}(T'_k \leq T_i < T'_{k+1}) .\end{aligned} \quad (4)$$

Remark 2.1. In this paper, it is assumed that the experiment consists in collecting a number n of cycles. Hence, the total duration of the experiment is equal to $T'_n + X_n$ and is therefore random. A classical renewal argument shows that, as $n \rightarrow \infty$, $(T'_n + X_n)/n$ converges a.s. to the mean duration of a cycle, $1/\lambda + \mathbb{E}[X] = \lambda \exp(\lambda \mathbb{E}[X])$. Another approach, which is more sensible in certain scenarios, is to consider that the total duration of the experiment is given, say equal to \mathbb{T} . In this case, the number of cycles is random, equal to the renewal process of the busy cycles, $N_{\mathbb{T}} = \sum_{k=1}^{\infty} \mathbb{1}\{T'_k \leq \mathbb{T}\}$. As $\mathbb{T} \rightarrow \infty$, the Blackwell theorem shows that $N_{\mathbb{T}}/\mathbb{T} \rightarrow 1/\lambda \exp(\lambda \mathbb{E}[X])$, showing that the asymptotic theory in both cases can be easily related.

3. INVERSION FORMULA AND ESTIMATION

Let \tilde{P} be a probability measure on $\mathbb{R} \times \mathbb{R}$ equipped with the Borel σ -algebra; for all $(s, p) \in \mathbb{C}^+ \times \mathbb{C}^+$, where $\mathbb{C}^+ = \{z \in \mathbb{C}, \operatorname{Re}(z) \geq 0\}$, we define its Laplace transform (or moment generating function) $\mathcal{L}\tilde{P}$ as:

$$\mathcal{L}\tilde{P}(s, p) = \iint e^{-su-pv} \tilde{P}(du, dv) .$$

The following theorem provides a relation between the joint distribution of the individual pulses energies and durations P and the moment-generating function of the distribution of the energies and durations of the busy periods $\mathcal{L}P'$; this key relation will be used to derive an estimator of m .

Theorem 3.1. *Under Assumptions (H-1)–(H-2), for all $(s, p) \in \mathbb{C}^+ \times \mathbb{C}^+$,*

$$\int_{u=0}^{+\infty} e^{-(s+\lambda)u} \{a(u, p) - 1\} du = \frac{\lambda \mathcal{L}P'(s, p)}{s + \lambda} \frac{1}{s + \lambda - \lambda \mathcal{L}P'(s, p)} , \quad (5)$$

where

$$a(u, p) \stackrel{\text{def}}{=} \exp(\lambda \mathbb{E}[e^{-pY}(u - X)_+]) . \quad (6)$$

Proof. See Section D. □

Remark 3.1. Observe that the integral in (5) can be replaced by $\int_{u=-\infty}^{\infty}$ since, in (6), $a(u, p) = 0$ for $u < 0$. Moreover, from (6), we trivially get $|a(u, p)| \leq \exp(\lambda u)$ for $\operatorname{Re}(p) \geq 0$; hence this integral is well defined for $\operatorname{Re}(s) > 0$ and $\operatorname{Re}(p) \geq 0$.

The relation (5) is rather involved and it is perhaps not immediately obvious to see how this relation may yield to an estimator of the distribution of the energy. By logarithmic differentiation with respect to x , (6) implies

$$\frac{\partial}{\partial x} \log a(x, p) = \lambda \mathbb{E}[e^{-pY} \mathbb{1}(X \leq x)] . \quad (7)$$

We consider a kernel function K that integrates to 1 and denote by K^* its Fourier transform, $K^*(\nu) = \int_{-\infty}^{+\infty} K(y) e^{-i\nu y} dy$, so that $K^*(0) = 1$. We further assume that K^* is integrable, so that, for any $y \in \mathbb{R}$,

$$K(y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} K^*(\nu) e^{i\nu y} d\nu .$$

Hence, from (7) and Fubini's theorem, we have, for any bandwidth parameter $h > 0$ and all $y \in \mathbb{R}$,

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{1}{\lambda} \frac{\partial}{\partial x} \log a(x, i\nu) K^*(h\nu) e^{i\nu y} d\nu \\ &= \mathbb{E} \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} K^*(h\nu) e^{i\nu(y-Y)} \mathbb{1}(X \leq x) d\nu \right] \end{aligned} \quad (8)$$

$$= \mathbb{E} \left[\frac{1}{h} K \left(\frac{y-Y}{h} \right) \mathbb{1}(X \leq x) \right]. \quad (9)$$

Taking the limits $x \rightarrow \infty$ and $h \rightarrow 0$ in the previous equation leads to the following explicit inversion formula which will be used to derive our estimator. For any continuity point y of the density m , we have

$$m(y) = \lim_{h \rightarrow 0} \lim_{x \rightarrow +\infty} \left\{ \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{\lambda} \frac{\partial}{\partial x} \log a(x, i\nu) K^*(h\nu) e^{i\nu y} d\nu \right\}. \quad (10)$$

We now observe that for any $p \in \mathbb{C}^+$, the RHS of (5) is integrable on a line $\{c + i\omega, \omega \in \mathbb{R}\}$ where c is an arbitrary positive number. By inverting the Laplace transform, (5) implies that, for all $p \in \mathbb{C}^+$ and $x \in \mathbb{R}_+$,

$$\begin{aligned} a(x, p) &= 1 + \\ & \frac{\lambda}{2\pi} \int_{-\infty}^{+\infty} \frac{\mathcal{L}P'(c + i\omega, p)}{c + i\omega + \lambda} \frac{e^{(c+\lambda+i\omega)x}}{c + i\omega + \lambda - \lambda \mathcal{L}P'(c + i\omega, p)} d\omega. \end{aligned} \quad (11)$$

Our estimator of m is based on (10) and (11) but we need first to estimate λ , the intensity of the underlying Poisson process. Since the idle periods are independent and identically distributed according to an exponential distribution with intensity λ , we use maximum-likelihood estimator based on the durations of the idle periods $\{Z_k, k = 1, \dots, n\}$, namely,

$$\hat{\lambda}_n \stackrel{\text{def}}{=} \left(\frac{1}{n} \sum_{k=1}^n Z_k \right)^{-1}. \quad (12)$$

The function $a(x, i\nu)$ can be estimated from $\{(X'_k, Y'_k), k = 1, \dots, n\}$ by plugging in (11) an estimate of the Laplace transform $\mathcal{L}P'$ of the joint distribution of the busy period duration and energy. More precisely, let \widehat{P}'_n be the associated empirical measure: for any bivariate measurable function g , we denote by

$$\widehat{P}'_n g \stackrel{\text{def}}{=} \iint g(x, y) \widehat{P}'_n(dx, dy) = \frac{1}{n} \sum_{k=1}^n g(X'_k, Y'_k).$$

We consider the following estimator

$$\widehat{a}_n(x, i\nu) = 1 + \frac{\widehat{\lambda}_n}{2\pi} \int_{-\infty}^{+\infty} \frac{\widehat{\lambda}_n \widehat{\mathcal{L}P}'_n(c + i\omega, i\nu)}{c + i\omega + \widehat{\lambda}_n} \frac{e^{(\lambda+c+i\omega)x}}{c + i\omega + \widehat{\lambda}_n - \widehat{\mathcal{L}P}'_n(c + i\omega, i\nu)} d\omega. \quad (13)$$

where

$$\widehat{\mathcal{L}P}'_n(c + i\omega, i\nu) \stackrel{\text{def}}{=} \widehat{\mathcal{L}P}'_n(c + i\omega, i\nu) = \frac{1}{n} \sum_{k=1}^n e^{-(c+i\omega)X'_k - i\nu Y'_k}, \quad (14)$$

In practice, the numerical computation of this integral (and also the one in (21) below) can be done by using efficient numerical packages (we can refer to Gautschi (1997) for an overview of numerical integration methods). Since the integrand is infinitely differentiable and has a modulus decaying as $|\omega|^{-2}$ when $\omega \rightarrow \pm\infty$, the errors in computing this integrals numerically can be made arbitrary small. The numerical error will thus not be taken into account here for brevity.

In order to estimate $\lambda^{-1} \partial \log a / \partial x$, we also need to estimate the partial derivative $\partial a / \partial x$. Because the function $x \mapsto a(x, i\nu)$ (see (11)) is defined as an inverse Fourier transform of an integrable function, it is tempting to estimate its partial derivative simply by multiplying by a factor $\lambda + c + i\omega$ its Fourier transform prior to inversion. This approach however is not directly applicable, because multiplying the integrand by ω in (11) leads to a non absolutely convergent integral. As observed by Hall and Park (2004) in a related problem, it is possible to get rid of this difficulty by finding an explicit expression of the singular part of this function, which can be computed and estimated. Note first that, for any s and p with non-negative real parts, $|\mathcal{L}P'(s, p)| \leq 1$; on the other hand, $\text{Re}(s) > 0$ implies $|\lambda/(s + \lambda)| < 1$. Therefore, for all $(\omega, \nu) \in \mathbb{R} \times \mathbb{R}$,

$$\frac{1}{c + i\omega + \lambda - \lambda \mathcal{L}P'(c + i\omega, i\nu)} = \frac{1}{c + i\omega + \lambda} \sum_{n \geq 0} \left(\frac{\lambda \mathcal{L}P'(c + i\omega, i\nu)}{\lambda + c + i\omega} \right)^n.$$

Using the latter equation, we obtain

$$\begin{aligned} \frac{\lambda \mathcal{L}P'(c + i\omega, i\nu)}{c + i\omega + \lambda} \frac{1}{c + i\omega + \lambda - \lambda \mathcal{L}P'(c + i\omega, i\nu)} \\ = A_1(\omega, i\nu) + A_2(\omega, i\nu) \end{aligned} \quad (15)$$

where we have defined

$$\begin{aligned} A_1(\omega, i\nu) &\stackrel{\text{def}}{=} \frac{\lambda \mathcal{L}P'(c + i\omega, i\nu)}{(c + i\omega + \lambda)^2}, \\ A_2(\omega, i\nu) &\stackrel{\text{def}}{=} \frac{\{\lambda \mathcal{L}P'(c + i\omega, i\nu)\}^2}{(c + i\omega + \lambda)^2} \frac{1}{c + i\omega + \lambda - \lambda \mathcal{L}P'(c + i\omega, i\nu)}. \end{aligned}$$

It is easily seen that the functions $\omega \mapsto A_k(\omega, i\nu)$, $k = 1, 2$ are integrable. Hence we may define, for $k = 1, 2$, and all real numbers x and ν ,

$$a_k(x, i\nu) \stackrel{\text{def}}{=} \frac{1}{2\pi\lambda} \int_{\omega=-\infty}^{\infty} A_k(\omega, i\nu) e^{(\lambda+c+i\omega)x} d\omega \quad (16)$$

and therefore, using (11) and (15), $a(x, i\nu) = 1 + \lambda a_1(x, i\nu) + \lambda a_2(x, i\nu)$, which finally yields

$$\frac{1}{\lambda} \frac{\partial}{\partial x} \log a(x, i\nu) = \frac{1}{a(x, i\nu)} \left[\frac{\partial a_1}{\partial x} + \frac{\partial a_2}{\partial x} \right] (x, i\nu). \quad (17)$$

Recall that the moment generating function of a gamma distribution with shape parameter 2 and scale parameter λ is given by $x \mapsto \lambda^2 / (\lambda - x)^2$. It follows that, for all $u \in \mathbb{R}$,

$$\frac{1}{2\pi} \int_{\omega=-\infty}^{\infty} \frac{e^{(c+i\omega)u}}{(\lambda + c + i\omega)^2} d\omega = u_+ e^{-\lambda u}.$$

Using Fubini's theorem and this equation, we obtain, for all real numbers x and ν ,

$$\begin{aligned} a_1(x, i\nu) &= \frac{1}{2\pi} \int_{\omega=-\infty}^{\infty} \frac{\mathbb{E}[e^{-((c+i\omega)X' + i\nu Y')}] e^{(\lambda+c+i\omega)x}}{(\lambda + c + i\omega)^2} d\omega \\ &= e^{\lambda x} \mathbb{E} \left[e^{-i\nu Y'} (x - X')_+ e^{-\lambda(x-X')} \right] \\ &= \mathbb{E} \left[(x - X')_+ e^{\lambda X' - i\nu Y'} \right], \end{aligned}$$

and, differentiating this latter expression w.r.t. x , we obtain

$$\frac{\partial a_1}{\partial x}(x, i\nu) = \mathbb{E} \left[\mathbb{1}(X' \leq x) e^{\lambda X' - i\nu Y'} \right]. \quad (18)$$

On the other hand, note that $|A_2(\omega, i\nu)| = O(|\omega|^{-3})$ as $\omega \rightarrow \pm\infty$, the derivative of a_2 can (and will) be computed by multiplying the integrand in (16) by $\lambda + c + i\omega$, namely,

$$\begin{aligned} \frac{\partial a_2}{\partial x}(x, i\nu) &= \frac{\lambda}{2\pi} \\ &\times \int_{-\infty}^{+\infty} \frac{\{\mathcal{L}P'(c + i\omega, i\nu)\}^2}{c + i\omega + \lambda} \frac{e^{(\lambda+c+i\omega)x}}{c + i\omega + \lambda - \lambda \mathcal{L}P'(c + i\omega, i\nu)} d\omega. \quad (19) \end{aligned}$$

Eq. (19) and (18) then yield the following estimators for $\partial a_k / \partial x$, $k = 1, 2$,

$$\widehat{I}_{1,n}(x, i\nu) = \frac{1}{n} \sum_{k=1}^n \mathbb{1}(X'_k \leq x) e^{\widehat{\lambda}_n X'_k - i\nu Y'_k} \quad (20)$$

$$\widehat{I}_{2,n}(x, i\nu) = \frac{\widehat{\lambda}_n e^{(c+\widehat{\lambda}_n)x}}{2\pi} \times \int_{-\infty}^{+\infty} \frac{\{\widehat{\mathcal{L}P}'_n(c + i\omega, i\nu)\}^2}{c + i\omega + \widehat{\lambda}_n} \frac{e^{i\omega x}}{c + i\omega + \widehat{\lambda}_n - \widehat{\lambda}_n \widehat{\mathcal{L}P}'_n(c + i\omega, i\nu)} d\omega \quad (21)$$

where $\widehat{\lambda}_n$ and $\widehat{\mathcal{L}P}'_n$ are given respectively by (12) and (14). From (10) and (17), we finally define the following estimator for the energy distribution density function:

$$\widehat{m}_{x,h,n}(y) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \left[\frac{\widehat{I}_{1,n} + \widehat{I}_{2,n}(x, i\nu)}{\widehat{a}_n} \right] K^*(h\nu) e^{i\nu y} d\nu, \quad (22)$$

where \widehat{a}_n , $\widehat{I}_{1,n}$ and $\widehat{I}_{2,n}$ are respectively defined in (13), (20) and (21).

4. MAIN RESULT

We denote respectively by $\|\cdot\|_\infty$, $\|\cdot\|_2$ and $\|\cdot\|_{\mathcal{W}(\beta)}$ the infinite norm, the L^2 -norm and the Sobolev norm of exponent β , that is the norm endowing the Sobolev space

$$\mathcal{W}(\beta) \stackrel{\text{def}}{=} \left\{ g \in L^2(\mathbb{R}) ; \|g\|_{\mathcal{W}(\beta)}^2 \stackrel{\text{def}}{=} \int_{-\infty}^{+\infty} (1 + |\nu|)^{2\beta} |g^*(\nu)|^2 d\nu < \infty \right\},$$

where g^* denotes the Fourier transform of g . Consider the following assumption on the kernel.

(H-4) K^* has a compact support, and there exists constants $C_K > 0$ and $l \geq \beta$. such that for all $\nu \in \mathbb{R}$,

$$|1 - K^*(\nu)| \leq C_K \frac{|\nu|^l}{(1 + |\nu|)^l}.$$

We may now state the main result of this section, which establish the rate of convergence of the integrated square error.

Theorem 4.1. *Let β , C and x be positive numbers. Assume (H-1)–(H-4) and suppose that $X \leq x$ a.s. and $\|m\|_{\mathcal{W}(\beta)} \leq C$. Then there exists $C' > 0$ only depending on K , λ , c , β , x and C such that, for all $M > 0$,*

$$\limsup_{n \rightarrow \infty} \mathbb{P}(n^{\beta/(1+2\beta)} \|\widehat{m}_{x,h,n} - m\|_2 \geq M) \leq C' M^{-2}. \quad (23)$$

where $h_n = n^{-1/(1+2\beta)}$.

Proof. See Section B. □

Remark 4.1. In the application we have considered, the condition $X \leq x$ assumption is always satisfied. Indeed, the pulse duration corresponds to the duration of the charge collection, and therefore to the lifetime of the pairs of electron-holes in the semiconductor detector. This lifetime is always finite and depends primarily of the geometry of the detector.

Remark 4.2. However, the condition $X \leq x$ a.s. can actually be circumvented if, at fixed $x > 0$, one considers $\widehat{m}_{x,h_n,n}$ as an estimator of m_x , defined as the density of the measure $\int_{u=0}^x P(du, dy)$, which is always defined under Assumption (H-3).

Remark 4.3. If X and Y are independent, then $m_x(y) = m(y)\mathbb{P}(X \leq x)$ so that, for all x such that $\mathbb{P}(X \leq x) > 0$, we obtain an estimator of m up to a multiplicative constant.

Remark 4.4. In the $M/G/\infty$ case, i.e. if $X = Y$ a.s., $m_x = m \mathbb{1}_{[0,x]}$. Hence, since

$$\|(\widehat{m}_{x,h_n,n} - m)\mathbb{1}_{[0,x]}\|_2^2 \leq \|(\widehat{m}_{x,h_n,n} - m \mathbb{1}_{[0,x]})\|_2^2,$$

our results apply to the locally integrated error for estimating $m \mathbb{1}_{[0,x]}$. As a comparison, the rate of our estimator is given by the smoothness of $m \mathbb{1}_{[0,x]}$, whereas the rate of the estimator proposed in Hall and Park (2004) for estimating the time service density is given by the smoothness of the pdf of X' (see (Hall and Park, 2004, Eq. (3.7))).

Remark 4.5. The estimators in (23) are functions of $\{(Z_k, X'_k, Y'_k), k = 1, \dots, n\}$, where n is the number of observed cycles. For $t \in \mathbb{R}_+$, denote by \mathcal{N}_t the renewal process associated to the arrivals of the photons, $\mathcal{N}_t \stackrel{\text{def}}{=} \sum_{k=1}^{\infty} \mathbb{1}\{T_k \leq t\}$. The number of arrivals during n cycles is equal to $\tilde{n} = \mathcal{N}_{T'_n + X'_n}$ and is therefore random. As n tends to infinity, $(T'_n + X'_n)/n$ converges almost surely to the mean of the cycle duration, $\exp(\lambda\mathbb{E}[X])/\lambda$ and it can be easily shown that the n -th return to an idle period (that is, $T'_n + X'_n$) is a stopping time with respect to the natural history of \mathcal{N}_t . Therefore, by the Blackwell theorem, $\mathcal{N}_{T'_n + X'_n}/(T'_n + X'_n)$ converges to λ . Therefore, $\tilde{n}/n = \mathcal{N}_{T'_n + X'_n}/n$ converges almost surely to $\exp(\lambda\mathbb{E}[X])$. It is well known that the minimax integrated rate for estimating m from $\{Y_k, k = 1, \dots, \tilde{n}\}$ with m in a β -Sobolev ball is $\tilde{n}^{1/(1+2\beta)}$, the only non-standard feature being that the density estimator is calculated by using a random number of data, which does not alter the density's estimator first-order property. Since \tilde{n}/n converges almost surely to a constant, Theorem 4.1 shows that the rates achieved by our estimator is the minimax integrated rate.

5. DECOMPOSITION OF THE ERROR

We give in this section theoretical results for the proposed estimators by introducing auxiliary variables, which will be used in the proof of

the main theorem. For any positive numbers W , x and $\tilde{\lambda}$, define

$$\widehat{\Delta}_n(W) \stackrel{\text{def}}{=} \sup_{(\omega, \nu) \in [-W, W]^2} |\mathcal{L}P'(c + i\omega, i\nu) - \widehat{\mathcal{L}P}'_n(c + i\omega, i\nu)| ;$$

$$\widehat{E}_n(W; x, \tilde{\lambda}) \stackrel{\text{def}}{=} \sup_{\nu \in [-W, W]} \left| \int \mathbb{1}_{[0, x]}(u) e^{\tilde{\lambda}(u-x)} e^{-i\nu y} (P' - \widehat{P}'_n)(du, dy) \right| .$$

Proposition 5.1 provides bounds for the random variables $\widehat{\Delta}_n$ and \widehat{E}_n .

Proposition 5.1. *Assume (H-1)–(H-2). Then $M_1 \stackrel{\text{def}}{=} \mathbb{E}(\max\{X', Y'\})$ is finite and the following inequalities hold for all $\varepsilon > 0$, $r > 0$ and $W > 1$:*

$$\mathbb{P}(|\widehat{\Delta}_n(W)| \geq \varepsilon) \leq \frac{4rM_1}{\varepsilon} + \left(1 + \frac{W}{r}\right)^2 \exp\left(-\frac{n\varepsilon^2}{16}\right) ; \quad (24)$$

$$\sup_{x, \tilde{\lambda} > 0} \mathbb{P}(|\widehat{E}_n(W; x, \tilde{\lambda})| \geq \varepsilon) \leq \frac{4rM_1}{\varepsilon} + \left(1 + \frac{W}{r}\right) \exp\left(-\frac{n\varepsilon^2}{16}\right) . \quad (25)$$

Proof. See Appendix C □

Since our estimate depends on $\hat{\lambda}_n$ and $\widehat{\mathcal{L}P}'_n$, we introduce auxiliary functions to exhibit both dependencies. Define the following functions depending on reals h , x , $\tilde{\lambda}$ and on any probability measure \tilde{P} :

$$\tilde{a}(x, i\nu; \tilde{\lambda}, \tilde{P}) \stackrel{\text{def}}{=} 1 + \frac{\tilde{\lambda} e^{(c+\tilde{\lambda})x}}{2\pi} \int_{-\infty}^{+\infty} \frac{\mathcal{L}\tilde{P}(c + i\omega, i\nu)}{c + i\omega + \tilde{\lambda}} \frac{e^{i\omega x}}{c + i\omega + \tilde{\lambda} - \tilde{\lambda}\mathcal{L}\tilde{P}(c + i\omega, i\nu)} d\omega \quad (26)$$

$$\tilde{I}_1(x, i\nu; \tilde{\lambda}, \tilde{P}) \stackrel{\text{def}}{=} \iint_{\mathbb{R}_+^2} \mathbb{1}_{\{u \leq x\}} e^{\tilde{\lambda}u - i\nu v} \tilde{P}(du, dv)$$

$$\tilde{I}_2(x, i\nu; \tilde{\lambda}, \tilde{P}) \stackrel{\text{def}}{=} \frac{\tilde{\lambda} e^{(\tilde{\lambda}+c)x}}{2\pi} \int_{-\infty}^{+\infty} \frac{(\mathcal{L}\tilde{P}(c + i\omega, i\nu))^2}{c + i\omega + \tilde{\lambda}} \frac{e^{i\omega x}}{c + i\omega + \tilde{\lambda} - \tilde{\lambda}\mathcal{L}\tilde{P}(c + i\omega, i\nu)} d\omega$$

and define

$$\tilde{m}(y; x, h, \tilde{\lambda}, \tilde{P}) \stackrel{\text{def}}{=} \frac{1}{2\pi} \int_{-\infty}^{+\infty} \left[\frac{\tilde{I}_1 + \tilde{I}_2}{\tilde{a}}(x, i\nu; \tilde{\lambda}, \tilde{P}) \right] K^*(h\nu) e^{i\nu y} d\nu \quad (27)$$

whenever the integral is well defined. Hence, by (6), (13), (19), (18), (20) and (21), for $i = 1, 2$,

$$\tilde{a}(x, i\nu; \lambda, P') = a(x, i\nu) \quad \text{and} \quad \tilde{I}_i(x, i\nu; \lambda, P') = \frac{\partial a_i}{\partial x}(x, i\nu) , \quad (28)$$

$$\tilde{a}(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) = \hat{a}_n(x, i\nu) \quad \text{and} \quad \tilde{I}_i(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) = \hat{I}_{i,n}(x, i\nu) \quad (29)$$

and $\widehat{m}_{x,h,n}(y) = \widetilde{m}(y; x, h, \widehat{\lambda}_n, \widehat{P}'_n)$. Now define

$$b_1(y) \stackrel{\text{def}}{=} m(y) - \mathbb{E} \left[\frac{1}{h} K \left(\frac{y - Y}{h} \right) \right] ; \quad (30)$$

$$b_2(y) \stackrel{\text{def}}{=} \mathbb{E} \left[\frac{1}{h} K \left(\frac{y - Y}{h} \right) \right] - \widetilde{m}(y; x, h, \lambda, P') ; \quad (31)$$

$$V_1(y) \stackrel{\text{def}}{=} \widetilde{m}(y; x, h, \lambda, P') - \widetilde{m}(y; x, h, \widehat{\lambda}_n, P') ; \quad (32)$$

$$V_2(y) \stackrel{\text{def}}{=} \widetilde{m}(y; x, h, \widehat{\lambda}_n, P') - \widehat{m}_{x,h,n}(y) \quad (33)$$

so that, by definition,

$$\widehat{m}_{x,h,n} - m = b_1 + b_2 + V_1 + V_2 . \quad (34)$$

In this decomposition, b_1 and b_2 are deterministic functions and V_1, V_2 are random processes. We now provide bounds for these quantities in the L^2 sense.

Theorem 5.1. *Let β, x and h be positive numbers and n be a positive integer. Assume (H-1)–(H-4). If $m \in \mathcal{W}(\beta)$, then we have*

$$\|b_1\|_2^2 \leq C_K^2 h^{2\beta} \|m\|_{\mathcal{W}(\beta)}^2 ; \quad (35)$$

$$\|b_2\|_2^2 \leq \|K\|_2^2 h^{-1} \mathbb{P}[X > x] . \quad (36)$$

Moreover, there exist positive constants M and η only depending on c and λ such that the two following assertions hold.

(i) We have

$$\|V_1\|_2^2 \leq M^2 \|K\|_2^2 (1+x)^2 h^{-1} e^{4(c+2\lambda)x} (\widehat{\lambda}_n - \lambda)^2 \quad (37)$$

on the event

$$E_1 \stackrel{\text{def}}{=} \left\{ |\widehat{\lambda}_n - \lambda| \leq \eta (1+x)^{-1} e^{-(c+2\lambda)x} \right\} . \quad (38)$$

(ii) For all $W \geq 1$ such that $[-Wh, Wh]$ contains the support of K^* , we have

$$\|V_2\|_2^2 \leq M^2 \|K\|_2^2 h^{-1} e^{4(c+2\lambda)x} \left[\widehat{\Delta}_n(W) + W^{-1} + \widehat{E}_n(W; x, \widehat{\lambda}_n) \right]^2 \quad (39)$$

on the event E_1 intersected with the event

$$E_2 \stackrel{\text{def}}{=} \left\{ \widehat{\Delta}_n(W) + W^{-1} \leq \eta e^{-(c+2\lambda)x} \right\} . \quad (40)$$

Proof. See Appendix A. \square

In this result, b_1 is the usual bias in kernel nonparametric estimation; b_2 is a non-usual bias term which only vanishes when X is bounded, it correspond to the fact that the limit $x \rightarrow \infty$ is not attained in (10); the fluctuation term V_1 accounts for the error in the estimation of λ by $\widehat{\lambda}_n$ and is of order $h^{-1}\sqrt{n}$ for fixed x and V_2 accounts for the

error in the estimation of $\mathcal{L}P'$ by $\widehat{\mathcal{L}P}'_n$ and, by using Proposition 5.1, it can be shown to be “almost” of order $h^{-1}\sqrt{n}$ for W chosen to diverge quickly enough with respect to n . The events E_1 and E_2 have probability tending to 1 as n tends to infinity; they are induced by the fraction present in the definition (22) of the estimator as they primarily avoid the denominator to get close to zero.

We now give a result on the consistency of our estimator, and also on a rate of convergence, based on Theorem 5.1 and Proposition 5.1 by imposing a superexponential tail for X .

Corollary 5.1. *Let $\beta > 0$ and $\gamma > 1$. Assume (H-1)–(H-4) and suppose that $m \in \mathcal{W}(\beta)$ and $\mathbb{P}[X > x] = O(e^{-|x|^\gamma})$. Then, for all $\epsilon > 0$, as $n \rightarrow +\infty$,*

$$\|m - \widehat{m}_{x_n, h_n, n}\|_2^2 = O_{\mathbb{P}}(n^{\epsilon-2\beta/(1+2\beta)}) \quad , \quad (41)$$

where $h_n \asymp n^{-1/(1+2\beta)}$ and $x_n \asymp (\log n)^{\gamma'}$ with $\gamma' \in (\gamma^{-1}, 1)$.

Proof. We set $W_n \stackrel{\text{def}}{=} n$. By Proposition 5.1, we get by choosing $\epsilon = C(\log(n)/n)^{1/2}$ and $r = n^{-1/2}$:

$$\mathbb{P}(|\widehat{\Delta}_n(W_n)| \geq C(\log(n)/n)^{1/2}) \leq \frac{4}{C} \log^{-1/2}(n) + (1 + \sqrt{n})^2 n^{-C/16}$$

which tends to 0 as $n \rightarrow \infty$ for $C > 32$. Hence,

$$|\widehat{\Delta}_n(W_n)| = O_{\mathbb{P}}\{(\log(n)/n)^{1/2}\} .$$

Similarly, because $\widehat{\lambda}_n$ is independent of $\{(X'_k, Y'_k), k = 1, \dots, n\}$, one has that $|\widehat{E}_n(W_n; x_n, \widehat{\lambda}_n)| = O_{\mathbb{P}}\{\log(n)/n\}^{1/2}$. For any $\delta_1 \geq 0$, $\delta_2 > 0$ and $\epsilon > 0$, one has that

$$x_n^{\delta_1} \exp(\delta_2 x_n) = o(n^\epsilon) . \quad (42)$$

Since $\widehat{\lambda}_n = \lambda + O_{\mathbb{P}}(n^{-1/2})$ and $|\widehat{\Delta}_n(W_n) + W_n^{-1}| = O_{\mathbb{P}}((\log(n)/n)^{1/2})$, E_1 and E_2 have probability tending to one, so that, applying again (42) and the bounds of Theorem 5.1 finally give, for all $\epsilon > 0$,

$$\|V_i\| = O_{\mathbb{P}}((h_n n)^{\epsilon-1/2}) \quad , \quad i = 1, 2 .$$

Now using the superexponential tail assumption for X , we have $\mathbb{P}(X > x_n) = O(\exp\{-\log^{\gamma'}(n)\}) = o(n^\epsilon)$ for all $\epsilon > 0$ and the result follows. \square

As seen from (41), the estimator almost achieves the standard non-parametric minimax rate $n^{-\beta/(1+2\beta)}$ that one would obtain by observing $\{(X_k, Y_k), k = 1, \dots, n\}$ directly. If X is bounded, then the rate can be made more precise as in Theorem 4.1: by taking x equal to an upper bound for X (so that $b_2 = 0$) and $h_n \asymp n^{-1/(1+2\beta)}$, one easily gets from the above proof that

$$\|m - \widehat{m}_{x, h_n, n}\|_2^2 = O_{\mathbb{P}}(\log(n)n^{-2\beta/(1+2\beta)}) \quad , \quad (43)$$

thus a loss of $\log(n)$ in comparison with the claimed rate. This $\log(n)$ can in fact be removed as shown in Appendix B.

6. APPLICATIONS — DISCUSSION

The present paper is directed towards the construction of an estimator and deriving elements of its asymptotic theory. We will therefore satisfy ourselves by providing simple examples and will refer the reader to Trigano et al. (2005) for an in-depth discussion of the selection of the tuning parameters (*e.g.* the kernel bandwidth, the truncation bound, etc) and the analysis of many different data sets.

We first consider a simulated data set. Samples are drawn from the following the density

$$f(x, y) = \mathcal{N}_{20,3}(x) \times (0.6\mathcal{N}_{100,6}(y) + 0.4\mathcal{N}_{130,9}(y)) , \quad (44)$$

where $\mathcal{N}_{a,b}$ denotes the gaussian distribution of mean a and standard deviation b truncated to \mathbb{R}_+ ; The intensity of the Poisson process is set to $\lambda = 0.04$; Figure 2 shows the true density and a kernel estimate of the marginal of the piled-up distribution g , based on 10^5 samples. Figure 3 displays the difference between the true and the estimated




FIGURE 2. dashed line: marginal density of the energy for the pdf (44); solide line: kernel density estimator of the pileup observations based on 10^5 samples.

density, obtained using the kernel bandwidth $h = 2$ and the upper bound $x = 80$. The pileup correction is very effective, the estimated energy distribution $\hat{m}_{x,h,n}$ captures most of the important features of

the original distribution m . Note that the second sharp mode of the original density is well recovered after the pileup correction, whereas it is severely distorted in absence of pileup correction. The fake modes at 200 and 230 are totally removed after pileup correction.




FIGURE 3. dashed line: marginal density of the energy for the pdf (44); solid line: estimator $\hat{m}_{x,h,n}$.

n	ISE	c	ISE	x	ISE
		1	$2.232 \cdot 10^{-2}$	20	$5.417 \cdot 10^{-3}$
1000	$4.760 \cdot 10^{-3}$	0.1	$3.852 \cdot 10^{-4}$	40	$1.905 \cdot 10^{-4}$
5000	$1.089 \cdot 10^{-3}$	0.01	$4.002 \cdot 10^{-4}$	60	$3.836 \cdot 10^{-4}$
10000	$3.852 \cdot 10^{-4}$	0.001	$4.348 \cdot 10^{-4}$	80	$5.100 \cdot 10^{-4}$
20000	$2.042 \cdot 10^{-4}$	0.0001	$3.852 \cdot 10^{-4}$	100	$2.229 \cdot 10^{-2}$
		0.00001	$4.426 \cdot 10^{-4}$	$\max_{i \leq n} X'_i$	$2.231 \cdot 10^{-2}$

TABLE 1. Integrated Square Error Monte-Carlo estimates as n , c or x varies.

We now present some results using a model of the energy distribution of the Cesium 137 radionuclide (including Compton effect). We draw $n = 500000$ samples of (X, Y) using the adaptive rejection sampling algorithm; the number of samples seems may appear to be large, but such large number are commonly used in nuclear spectrometry, especially when active sources are measured. Figure 4 illustrates the differences between the distribution of the observations when the counting

rate λ is low (dashed curve) (that is when the piled-up phenomenon is negligible), an distribution of the same element when the counting rate is high (solid curve) ; we can remark that the Compton continuum (which is the smooth part of the density on the left of the spike) is also distorted, since electrical pulses generated by Compton photons are also susceptible to pile-up.






FIGURE 4. Illustration of ideal and piled-up energy spectra of the Cs 137 element

Figure 5 illustrates the behaviour of our estimator and compares it with the ideal energy spectrum. We observe good adequation between both densities and pile-up correction in the histograms.

We now discuss on the choice of the parameters c , x and h . It appears that the parameter c has little influence, provided it is not chosen too large to avoid numerical issues. This is hardly surprising, since the Bromwich integral used to compute the inverse Laplace transform does not theoretically depend on the choice of c (see e.g. Doetsch (1974)). The choice of the parameter x is more delicate, when the impulse durations are not upper-bounded, and this choice must be examined using the upper bounds of Theorem 5.1. On the one hand, choosing x too small introduces a bias in (22), since the control of the bias term b_2 is not guaranteed in that case; on the other hand, taking x too large does not ensure that the variance term V_1 and V_2 are controlled, since in this case the conditions (38) and (40) may not be satisfied. However, in our practical applications, an approximate bound of X can be



(a)



(b)

FIGURE 5. Energy spectrum of the Cs 137 element — (a) Estimate $\hat{m}_{x,h,n}$; (b) Ideal probability density function

determined and the number of samples n is generally large enough for allowing to choose this bound for x . In standard nonparametric estimation, there are several ways of choosing a data-driven bandwidth parameter. It is not yet clear how these methods can be adapted to this non-standard density estimation scenario, except in special cases. For instance, when X and Y are independent, it is easily seen that Y' is an infinite mixture of the auto-convolutions of the density m and thus has a density with similar smoothness (this can be seen in Figures 2 and 4 where X and Y have been drawn independently). A possible approach would then consist in using an automatic bandwidth selector (such that cross validation) on the observations $\{Y'_k, k = 1, \dots, n\}$, and use the obtained optimal bandwidth for the estimator $\hat{m}_{x,h,n}$. This type of ideas and other discussions on practical applications of these results are discussed in the companion paper Trigano et al. (2005).

APPENDIX A. PROOF OF THEOREM 5.1

Bound for b_1 . Observe that b_1 is the usual bias in non-parametric kernel estimation. The bound of the integrated error is classically given, for density in a Sobolev space, by

$$\|b_1\|_2^2 = \int_{-\infty}^{\infty} |1 - K^*(h\nu)|^2 |m^*(\nu)|^2 d\nu \leq C_K^2 h^{2\beta} \|m\|_{W(\beta)}^2,$$

which shows (35).

Bound for b_2 . By (9), (17), (27) and (28), we find

$$b_2(y) = \mathbb{E} \left[\frac{1}{h} K \left(\frac{y - Y}{h} \right) \mathbb{1}(X > x) \right].$$

An application of the Cauchy-Schwarz Inequality yields (36).

Bound for V_1 . We will show below that there exist positive constants M and η such that, on E_1 (as defined in (38)),

$$\sup_{\nu \in \mathbb{R}} \left| \frac{\partial}{\partial \tilde{\lambda}} \left[\frac{\tilde{I}_1 + \tilde{I}_2}{\tilde{a}} \right] (x, i\nu; \hat{\lambda}_n, P') \right| \leq M (1 + x) e^{(2c+4\lambda)x}. \quad (45)$$

Using (27) and (32), the Parseval Theorem and the latter relation imply

$$\|V_1\|_2^2 \leq \int_{-\infty}^{\infty} \left| M (1 + x) e^{(2c+4\lambda)x} (\hat{\lambda}_n - \lambda) \right|^2 |K^*(h\nu)|^2 d\nu$$

on the event E_1 , which yields (37). Hence it remains to show (45).

First observe that, by definition of \tilde{I}_1 , one gets trivially, for all $\tilde{\lambda} > 0$,

$$|\tilde{I}_1(x, i\nu; \tilde{\lambda}, P')| \leq e^{\tilde{\lambda}x} \quad \text{and} \quad \left| \frac{\partial}{\partial \tilde{\lambda}} \tilde{I}_1(x, i\nu; \tilde{\lambda}, P') \right| \leq x e^{\tilde{\lambda}x}. \quad (46)$$

Inserting (6) into (28), we get, for all $\nu \in \mathbb{R}$,

$$|\tilde{a}(x, i\nu; \lambda, P')| = \exp(\lambda \mathbb{E}[\cos(\nu Y) (x - X)_+]) \in [e^{-\lambda x}, e^{\lambda x}]. \quad (47)$$

Let $\eta_0 > 0$ to be chosen later. From (26), Lemma E.1 shows that, there exists a constant M_1 only depending on λ , c and η_0 such that, for all $\tilde{\lambda}$ in $[0, \lambda + \eta_0]$ and $\nu \in \mathbb{R}$,

$$\begin{aligned} & \left| (\tilde{a}(x, i\nu; \tilde{\lambda}, P') - 1)e^{-(\tilde{\lambda}+c)x} - (\tilde{a}(x, i\nu; \lambda, P') - 1)e^{-(\lambda+c)x} \right| \\ & \leq M_1 |\tilde{\lambda} - \lambda|. \end{aligned} \quad (48)$$

From (47) and (48) and since, for all real y , $|1 - e^y| \leq |y|e^{|y|}$, we get, for all $\tilde{\lambda}$ in $[0, \lambda + \eta_0]$ and $\nu \in \mathbb{R}$,

$$\begin{aligned} |\tilde{a}|(x, i\nu; \tilde{\lambda}, P') & \geq |\tilde{a}|(x, i\nu; \lambda, P') e^{(\tilde{\lambda}-\lambda)x} - |e^{(\tilde{\lambda}-\lambda)x} - 1| \\ & \quad - M_1 |\tilde{\lambda} - \lambda| e^{(\tilde{\lambda}+c)x}, \end{aligned}$$

hence

$$|\tilde{a}|(x, i\nu; \tilde{\lambda}, P') \geq e^{(\tilde{\lambda}-2\lambda)x} - [M_1 e^{(\tilde{\lambda}+c)x} + x e^{|\tilde{\lambda}-\lambda|x}] |\tilde{\lambda} - \lambda|.$$

Note that, taking $\eta_0 = c$ and $M'_1 = M_1 \vee 1$, the term between brackets is at most $M'_1 e^{(\tilde{\lambda}-2\lambda)x} (1+x) e^{(c+2\lambda)x}$ for $\tilde{\lambda} \in [\lambda - \eta_0, \lambda + \eta_0]$ so that we get, on the event E_1 with $\eta \leq \eta_1 \stackrel{\text{def}}{=} \{\eta_0 \wedge (M'_1)^{-1}/2\}$,

$$|\tilde{a}|(x, i\nu; \hat{\lambda}_n, P') \geq \frac{1}{2} e^{(\hat{\lambda}_n-2\lambda)x}. \quad (49)$$

From (26) and using similar bounds as in Lemma E.1, one easily shows that, for some constant M_2 only depending on λ , c and η_1 , for all $\tilde{\lambda} \in \mathbb{R}$ such that $|\lambda - \tilde{\lambda}| \leq \eta_1$,

$$\begin{aligned} |\tilde{a}(x, i\nu; \tilde{\lambda}, P')| & \leq M_2 e^{(\tilde{\lambda}+c)x} \quad \text{and} \\ \left| \frac{\partial \tilde{a}}{\partial \tilde{\lambda}}(x, i\nu; \tilde{\lambda}, P') \right| & \leq M_2 (1+x) e^{(\tilde{\lambda}+c)x}, \end{aligned} \quad (50)$$

$$\begin{aligned} |\tilde{I}_2(x, i\nu; \tilde{\lambda}, P')| & \leq M_2 e^{(\tilde{\lambda}+c)x} \quad \text{and} \\ \left| \frac{\partial \tilde{I}_2}{\partial \tilde{\lambda}}(x, i\nu; \tilde{\lambda}, P') \right| & \leq M_2 (1+x) e^{(\tilde{\lambda}+c)x}. \end{aligned} \quad (51)$$

Collecting (46), (49) and the two last displayed bounds shows that (45) holds on E_1 , for any $\eta \leq \eta_1$.

Bound for V_2 . Since the support of K^* is included in $[-Wh, Wh]$, By Parseval Theorem, (27) and (33), the claimed bound is implied by

$$\begin{aligned} & \sup_{|\nu| \leq W} \left| \frac{\tilde{I}_1 + \tilde{I}_2}{\tilde{a}}(x, i\nu; \hat{\lambda}_n, P') - \frac{\tilde{I}_1 + \tilde{I}_2}{\tilde{a}}(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) \right| \\ & \leq M e^{2(c+2\lambda)x} \left[\widehat{\Delta}_n(W) + W^{-1} + \widehat{E}_n(W; x, \hat{\lambda}_n) \right], \end{aligned} \quad (52)$$

which we now show. Using (26), we may write

$$\begin{aligned} & |\tilde{a}(x, i\nu; \tilde{\lambda}, \tilde{P}) - \tilde{a}(x, i\nu; \tilde{\lambda}, P')| \\ &= \left| \frac{\tilde{\lambda} e^{(c+\tilde{\lambda})x}}{2\pi} \int_{-\infty}^{+\infty} [\Psi_{\tilde{z}}(\tilde{\lambda}, \omega) - \Psi_z(\tilde{\lambda}, \omega)] d\omega \right|, \end{aligned} \quad (53)$$

where Ψ is defined in Lemma E.1, and where the complex functions \tilde{z} and z are defined as

$$z(\omega) \stackrel{\text{def}}{=} (e^{i\omega x} \mathcal{L}P'(c + i\omega, i\nu); \mathcal{L}P'(c + i\omega, i\nu))$$

and

$$\tilde{z}(\omega) \stackrel{\text{def}}{=} (e^{i\omega x} \mathcal{L}\tilde{P}(c + i\omega, i\nu); \mathcal{L}\tilde{P}(c + i\omega, i\nu)).$$

Using (53) and assertion (ii) of Lemma E.1, there exists $M_1 > 0$ such that, for all $\tilde{\lambda} \leq \lambda + \eta_1$,

$$\begin{aligned} & \sup_{|\nu| \leq W} |\tilde{a}(x, i\nu; \tilde{\lambda}, P') - \tilde{a}(x, i\nu; \tilde{\lambda}, \widehat{P}'_n)| \\ & \leq M_1 e^{(c+\tilde{\lambda})x} \left(\widehat{\Delta}_n(W) + \frac{1}{W} \right). \end{aligned} \quad (54)$$

It is also clear that for all $\tilde{\lambda} \leq \lambda + \eta_1$,

$$\sup_{|\nu| \leq W} |\tilde{I}_1(x, i\nu; \tilde{\lambda}, \widehat{P}'_n) - \tilde{I}_1(x, i\nu; \tilde{\lambda}, P')| \leq e^{\tilde{\lambda}x} \widehat{E}_n(W; \tilde{\lambda}, x) \quad (55)$$

and

$$\begin{aligned} & |\tilde{I}_2(x, i\nu; \tilde{\lambda}, \tilde{P}) - \tilde{I}_2(x, i\nu; \tilde{\lambda}, P')| \\ &= \left| \frac{e^{(c+\tilde{\lambda})x}}{2\pi} \int_{-\infty}^{+\infty} [\Psi_{\tilde{z}}(\tilde{\lambda}, \omega) - \Psi_z(\tilde{\lambda}, \omega)] d\omega \right|, \end{aligned}$$

with

$$z(\omega) \stackrel{\text{def}}{=} (e^{i\omega x} (\mathcal{L}P'(c + i\omega, i\nu))^2; \mathcal{L}P'(c + i\omega, i\nu))$$

and

$$\tilde{z}(\omega) \stackrel{\text{def}}{=} (e^{i\omega x} (\mathcal{L}\tilde{P}(c + i\omega, i\nu))^2; \mathcal{L}\tilde{P}(c + i\omega, i\nu)).$$

Consequently, using assertion (ii) of Lemma E.1, we have for all $\tilde{\lambda} \leq \lambda + \eta_1$,

$$\begin{aligned} & \sup_{|\nu| \leq W} |\tilde{I}_2(x, i\nu; \tilde{\lambda}, \widehat{P}'_n) - \tilde{I}_2(x, i\nu; \tilde{\lambda}, P')| \\ & \leq M_2 e^{(\tilde{\lambda}+c)x} \left(\widehat{\Delta}_n(W) + \frac{1}{W} \right). \end{aligned} \quad (56)$$

We now derive a lower bound for $\hat{a}_n(x, i\nu) = a(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n)$; By (54), we get

$$\inf_{|\nu| \leq W} |\hat{a}|(x, i\nu) \geq \inf_{|\nu| \leq W} |\tilde{a}|(x, i\nu; \hat{\lambda}_n, P') - M_1 e^{(c+\hat{\lambda}_n)x} \left(\widehat{\Delta}_n(W) + \frac{1}{W} \right)$$

Recall that E_1 and E_2 are defined in (38) and (40) respectively. Using (49), which holds on E_1 for any $\eta \leq \eta_1$, we get, on $E_1 \cap E_2$,

$$\inf_{|\nu| \leq W} \left| \tilde{a}(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) \hat{a}_n(x, i\nu) \right| \geq \frac{1}{2} \left[\frac{1}{2} - M_1 \eta \right] e^{(2\hat{\lambda}_n - 4\lambda)x} . \quad (57)$$

Hence we set $\eta \stackrel{\text{def}}{=} (4M_1)^{-1} \wedge \eta_1$, so that the term between brackets is at least $1/4$. Finally, using that, for all complex number x, y, z, x', y', z' ,

$$\frac{x+y}{z} - \frac{x'+y'}{z'} = \frac{(z'-z)(x+y) + z(x-x') + z(y-y')}{zz'}$$

and collecting (46), (50), (51), (54), (55), (56) and (57) leads to (52).

APPENDIX B. PROOF OF THE THEOREM 4.1

In this section we denote by η_i, M_i and $C_i, i = 0, 1, 2, \dots$ some positive constants only depending on $\|m\|_{\mathcal{W}(\beta)}, K, \lambda, c$ and x . We will also use the notations introduced Section A. As shown in this section, we have $\|b_1\|_2^2 \leq C_K^2 \|m\|_{\mathcal{W}(\beta)}^2 h^{2\beta}$ and since $\mathbb{P}(X > x) = 0$, we have $b_2 = 0$.

By (6), because $\mathbb{E}[X] < \infty$, it is easily seen that $|a(x, i\nu)| \geq e^{-\lambda x}$. This can be used in the ratio appearing in (22) to lower bound \hat{a}_n for n large by using that $\hat{a}_n(x, i\nu)$ converges to $a(x, i\nu)$. However this will not allow bounds of the ratio in the mean square sense. For obtaining mean square error estimates, we consider the following modified estimator which (artificially) circumvent this difficulty. Let $\eta_0 > \lambda$ and denote by A_n the set

$$A_n \stackrel{\text{def}}{=} \{\hat{\lambda}_n \leq \eta_0\} \cap \left\{ \inf_{h_n \nu \in \text{Supp}(K^*)} |\hat{a}_n|(x, i\nu) \geq \frac{1}{5} \exp(-\hat{\lambda}_n x) \right\} ,$$

where $\text{Supp}(K^*)$ denotes the (compact) support of K^* . Define

$$\check{m}_{x,h,n}(y) = \mathbb{1}_{A_n} \hat{m}_{x,h,n}(y) . \quad (58)$$

We will show that

$$\sup_{n \geq 1} n^{2\beta/(1+2\beta)} \mathbb{E} \|\check{m}_{x,h,n} - m\|_2^2 \leq C_0 . \quad (59)$$

Let V_3 be the random process

$$V_3(y) \stackrel{\text{def}}{=} \check{m}(y; x, h_n, \lambda, P') - \hat{m}_{x,h,n}(y) ,$$

so that

$$\|\check{m}_{x,h,n} - m\|_2^2 \leq \|b_1\|_2^2 + \|m\|_2^2 \mathbb{1}_{A_n^c} + \|V_3\|_2^2 \mathbb{1}_{A_n} .$$

We will show that there exist $C_1 > 0$ such that, for n large enough,

$$\mathbb{P}(A_n^c) \leq C_1 n^{-1} ; \quad (60)$$

$$\mathbb{E}[\|V_3\|_2^2 \mathbb{1}_{A_n}] \leq C_1 (h_n n)^{-1} . \quad (61)$$

Since $\|b_1\|_2^2 \leq C_K^2 \|m\|_{\mathcal{W}(\beta)}^2 h^{2\beta}$ and $\|m\|_2 \leq \|m\|_{\mathcal{W}(\beta)}$, the three last displays yield the bound (59). The bound (23) then follows by writing

$$\begin{aligned} & \mathbb{P}(n^{\beta/(1+2\beta)} \|\widehat{m}_{x,h_n,n} - m\|_2 \geq M) \\ & \leq \mathbb{P}(n^{\beta/(1+2\beta)} \|\check{m}_{x,h_n,n} - m\|_2 \geq M) + \mathbb{P}(A_n^c) \leq C_0 M^{-2} + C_1 n^{-1}, \end{aligned}$$

where we applied the Markov Inequality, (59) and (60). It now remains to show (60) and (61).

Proof of the bound (60). We set $W_n \stackrel{\text{def}}{=} n$, so that for n large enough, $h_n \nu \in \text{Supp}(K^*)$ implies $|\nu| \leq W_n$. As in (57), we have, on $E_1 \cap E_2$,

$$\inf_{|\nu| \leq W_n} |\widehat{a}_n|(x, i\nu) \geq \frac{1}{4} e^{(\widehat{\lambda}_n - 2\lambda)x}.$$

Hence the intersection of $\{\widehat{\lambda}_n \leq \eta_0\}$, E_1 , E_2 and $\{\exp((\widehat{\lambda}_n - 2\lambda)x)/4 \geq \exp(-\widehat{\lambda}_n x)/5\}$ is included in A_n . Since the last inequality and E_2 both contain $|\widehat{\lambda}_n - \lambda| \leq \eta_2$ for $\eta_2 > 0$ small enough, we get

$$\mathbb{P}(A_n^c) \leq \mathbb{P}(\widehat{\lambda}_n > \eta_0) + \mathbb{P}(|\widehat{\lambda}_n - \lambda| > \eta_2) + \mathbb{P}(\widehat{\Delta}_n(n) + n^{-1} > \eta_3).$$

Clearly the two first probabilities in the RHS are $O(n^{-1})$. For n large enough, the last probability is less than $\mathbb{P}(\widehat{\Delta}_n(n) > \eta_3/2)$, which is $o(n^{-1})$ by applying Proposition 5.1, say with $r = n^{-2}$. We thus get (60) for n large enough.

Proof of the bound (61). By (27), (28) and (29), V_3 is defined as the inverse Fourier transform of

$$V_3^*(\nu) = K^*(h_n \nu) \left[\frac{\partial_x a_1 + \partial_x a_2}{a} - \frac{\widehat{I}_{1,n} + \widehat{I}_{2,n}}{\widehat{a}_n} \right] (x, i\nu),$$

where $\partial_x a_i$ is a shorthand notation for $\partial a_i / \partial x$. Using that

$$\begin{aligned} & \left| \frac{\partial_x a_1 + \partial_x a_2}{a} - \frac{\widehat{I}_{1,n} + \widehat{I}_{2,n}}{\widehat{a}_n} \right| \leq \\ & \frac{1}{|\widehat{a}_n|} \left[\sum_{i=1,2} |\partial_x a_i - \widehat{I}_{i,n}| + \left| \frac{\partial_x a_1 + \partial_x a_2}{a} \right| |\widehat{a}_n - a| \right], \end{aligned}$$

we obtain that, on the set A_n defined above, for all $\nu \in \mathbb{R}$,

$$|V_3^*(\nu)| \leq 5 |K^*(h_n \nu)| \left[\mathcal{E}_{1,n} + \mathcal{E}_{2,n} + \mathcal{E}_n \left| \frac{\partial_x a_1 + \partial_x a_2}{a} \right| (x, i\nu) \right],$$

where, for $i = 1, 2$, we define $\mathcal{E}_{i,n} \stackrel{\text{def}}{=} e^{\widehat{\lambda}_n x} |\partial_x a_i - \widehat{I}_{i,n}| (x, i\nu)$ and $\mathcal{E}_n \stackrel{\text{def}}{=} e^{\widehat{\lambda}_n x} |\widehat{a}_n - a| (x, i\nu)$. Multiplying by $\mathbb{1}_{A_n}$, taking the expectation and

applying the Parseval Theorem yield

$$\begin{aligned} & \mathbb{E}[\|V_3\|_2^2 \mathbb{1}_{A_n}] \\ & \leq C_2 \left[h_n^{-1} \|K\|_2^2 \sum_{i=1}^2 \sup_{\nu \in \mathbb{R}} \mathbb{E}[\mathbb{1}_{A_n} \mathcal{E}_{i,n}^2] + M_1^2 \sup_{\nu \in \mathbb{R}} \mathbb{E}[\mathbb{1}_{A_n} \mathcal{E}_n] \right], \quad (62) \end{aligned}$$

where, by (9) and (17) and Parseval's theorem,

$$M_1^2 \stackrel{\text{def}}{=} \int_{-\infty}^{\infty} |K^*(h_n \nu)|^2 \left| \frac{\partial_x a_1 + \partial_x a_2}{a} \right|^2 (x, i\nu) d\nu \leq \|K^*\|_{\infty} \|m\|_2^2.$$

By (18) and (20), we have

$$\begin{aligned} \left| \partial_x a_1 - \widehat{I_{1,n}} \right| (x, i\nu) & \leq \left| \partial_x a_1(x, i\nu) - \tilde{I}_1(x, i\nu; \hat{\lambda}_n, P') \right| \\ & \quad + \left| \tilde{I}_1(x, i\nu; \hat{\lambda}_n, P') - \tilde{I}_1(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) \right|. \end{aligned}$$

Using this decomposition in $\mathcal{E}_{1,n}$, the independence of $\hat{\lambda}_n$, (X'_k, Y'_k) , $k = 1, \dots, n$, the fact that $\text{Var}(\mathbb{1}(X' \leq x) e^{\hat{\lambda} X' - i\nu Y'}) \leq 2e^{2\hat{\lambda}x}$ and the bound $\hat{\lambda}_n \leq \eta_0$ on A_n , we get

$$\mathbb{E}[\mathbb{1}_{A_n} \mathcal{E}_{1,n}^2] \leq M_2 \left\{ \mathbb{E} \left[\mathbb{1}_{A_n} \left| \partial_x a_1(x, i\nu) - \tilde{I}_1(x, i\nu; \hat{\lambda}_n, P') \right|^2 \right] + n^{-1} \right\}$$

Using (46) and the mean value theorem for bounding the first expectation shows that the first term is $O(1/n)$ and thus

$$\sup_{\nu \in \mathbb{R}} \mathbb{E}[\mathcal{E}_{1,n}^2] \leq C_3 n^{-1} \quad (63)$$

By (19) and (21), we have

$$\begin{aligned} \left| \partial_x a_2 - \widehat{I_{2,n}} \right| (x, i\nu) & \leq \left| \partial_x a_2(x, i\nu) - \tilde{I}_2(x, i\nu; \hat{\lambda}_n, P') \right| \\ & \quad + \left| \tilde{I}_2(x, i\nu; \hat{\lambda}_n, P') - \tilde{I}_2(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) \right|. \end{aligned}$$

Using (28) and (51), by the mean value Theorem, we get

$$\sup_{\nu \in \mathbb{R}} \mathbb{E} \left[\mathbb{1}_{A_n} e^{2\hat{\lambda}_n x} \left| \partial_x a_2(x, i\nu) - \tilde{I}_2(x, i\nu; \hat{\lambda}_n, P') \right|^2 \right] \leq M_3 n^{-1}$$

Using (79) in the proof of Lemma E.1, for all $\nu \in \mathbb{R}$, on the set $\{\hat{\lambda}_n \leq \eta_0\}$,

$$\begin{aligned} & \left| \tilde{I}_2(x, i\nu; \hat{\lambda}_n, P') - \tilde{I}_2(x, i\nu; \hat{\lambda}_n, \widehat{P}'_n) \right| \\ & \leq M_4 \int_{-\infty}^{\infty} g(\omega) |\mathcal{L}P' - \widehat{\mathcal{L}P}'_n|(c + i\omega, i\nu) d\omega, \end{aligned}$$

where g is an integrable function only depending on c and λ . Inserting the three last bounds in the definition of $\mathcal{E}_{2,n}$, we obtain

$$\sup_{\nu \in \mathbb{R}} \mathbb{E}[\mathbb{1}_{A_n} \mathcal{E}_{2,n}^2] \leq C_4 n^{-1}.$$

Comparing (11) with (19) and (13) with (21), one easily sees that similar argument applies for bounding \mathcal{E}_n on the set A_n , giving

$$\sup_{\nu \in \mathbb{R}} \mathbb{E}[\mathbb{1}_{A_n} \mathcal{E}_n^2] \leq C_5 n^{-1}.$$

Inserting (63) and the two last displays into (62) shows (61).

APPENDIX C. PROOF OF PROPOSITION 5.1

We have $M_1 \leq \mathbb{E}[X'] + \mathbb{E}[Y'] < \infty$. Denote by $\mathbf{z}_i \stackrel{\text{def}}{=} (\omega_i, \nu_i)$, $i = 1, 2$ and define the function $L_{\mathbf{z}}(x, y) \stackrel{\text{def}}{=} e^{-(c+i\omega)x - i\nu y}$; we get,

$$|L_{\mathbf{z}_1}(x, y) - L_{\mathbf{z}_2}(x, y)| \leq g(x, y) |\mathbf{z}_1 - \mathbf{z}_2|_1, \quad (x, y) \in \mathbb{R}_+^2 \quad (64)$$

where $g(x, y) \stackrel{\text{def}}{=} \max(x, y)$ and $|\mathbf{z}_1 - \mathbf{z}_2|_1 \stackrel{\text{def}}{=} |\omega_1 - \omega_2| + |\nu_1 - \nu_2|$. Note that

$$\widehat{\Delta}_n(W) = \sup_{\mathbf{z} \in [-W, W]^2} |\widehat{P}'_n L_{\mathbf{z}} - P' L_{\mathbf{z}}|.$$

Let $N \stackrel{\text{def}}{=} \lceil W/r \rceil^2$, where $\lceil x \rceil$ denotes the unique integer in $[x, x+1)$. Then there exists a net $\{\mathbf{z}_k\}_{1 \leq k \leq N}$ so that

$$[-W, W]^2 \subset \bigcup_{k=1}^N C(\mathbf{z}_k, r),$$

where $C(\mathbf{z}_k, r) \stackrel{\text{def}}{=} \{\mathbf{z} \in \mathbb{R}^2 : |\mathbf{z} - \mathbf{z}_k|_1 \leq r\}$. Using this covering in the above expression of $\widehat{\Delta}_n$, we get

$$\widehat{\Delta}_n(W) \leq \max_{1 \leq k \leq N} \left\{ \sup_{\mathbf{z} \in C(\mathbf{z}_k, r)} |\widehat{P}'_n L_{\mathbf{z}} - P' L_{\mathbf{z}}| \right\}. \quad (65)$$

Using (64) for bounding each term in the max of (65), we get

$$\begin{aligned} & \sup_{\mathbf{z} \in C(\mathbf{z}_k, r)} |\widehat{P}'_n L_{\mathbf{z}} - P' L_{\mathbf{z}}| \quad (66) \\ &= \sup_{\mathbf{z} \in C(\mathbf{z}_k, r)} |\widehat{P}'_n(L_{\mathbf{z}} - L_{\mathbf{z}_k}) + P'(L_{\mathbf{z}_k} - L_{\mathbf{z}}) + (\widehat{P}'_n L_{\mathbf{z}_k} - P' L_{\mathbf{z}_k})| \\ &\leq r(\widehat{P}'_n g + P' g) + |(\widehat{P}'_n - P') L_{\mathbf{z}_k}|. \quad (67) \end{aligned}$$

Inserting this bound in (65), for proving (24), it is now sufficient to bound $\mathbb{P}(r[\widehat{P}'_n + P'](g) \geq \varepsilon/2)$ and $\mathbb{P}(\max_{1 \leq k \leq N} |(\widehat{P}'_n - P') L_{\mathbf{z}_k}| \geq \varepsilon/2)$.

Using that $P'g = M_1$ and Markov's inequality, we get, if $rM_1 \leq \varepsilon/4$,

$$\begin{aligned} & \mathbb{P}\left(r[\widehat{P}'_n + P']g \geq \frac{\varepsilon}{2}\right) \\ &= \mathbb{P}\left(\frac{r\widehat{P}'_ng}{\varepsilon/2 - rM_1} \geq 1\right) \leq \frac{rM_1}{\varepsilon/2 - rM_1} \leq \frac{4rM_1}{\varepsilon}, \end{aligned} \quad (68)$$

where, in the last inequality, we used that $1/(x-1) \leq 2/x$ for all $x \geq 2$ with $x = \varepsilon/(2rM_1)$. Since the LHS is at most 1 and the RHS is more than one when $rM_1 > \varepsilon/4$, this inequality holds in all cases, yielding the first term in the RHS of (24). We now consider the second term in the RHS of (24). Since the $\{(X'_k, Y'_k), k \geq 1\}$ are independent and identically distributed and $|L_{\mathbf{z}}| \leq 1$, by using Hoeffding's inequality (see e.g. Appendix 6 of Van Der Waart and Wellner (1996)), we get, for all \mathbf{z} ,

$$\mathbb{P}\left(\max_{1 \leq k \leq N} |(\widehat{P}'_n - P'_n)L_{\mathbf{z}}| \geq \frac{\varepsilon}{2}\right) \leq 4N \exp\left(-\frac{n\varepsilon^2}{16}\right).$$

The proof is concluded by using that $N \leq (W/r + 1)^2$.

To prove inequality (25), let now L_{ν} be defined as $L_{\nu}(\mathbf{u}; x, \tilde{\lambda}) \stackrel{\text{def}}{=} \mathbb{1}_{[0,x]}(u)e^{\tilde{\lambda}(u-x)}e^{-i\nu y}$ with $\mathbf{u} = (u, y)$; same calculations can be done, yielding, for all positive x and $\tilde{\lambda}$,

$$\begin{aligned} |L_{\nu_1}(\mathbf{u}; x, \tilde{\lambda}) - L_{\nu_2}(\mathbf{u}; x, \tilde{\lambda})| &\leq \\ &\mathbb{1}_{[0,x]}(u)e^{\tilde{\lambda}(u-x)}|e^{-i\nu_1 y} - e^{-i\nu_2 y}| \leq g(\mathbf{u})|\nu_1 - \nu_2|, \end{aligned}$$

where the function g is here defined as $g((u, y)) \stackrel{\text{def}}{=} |y|$. Using that $P'g \leq M_1$ and $|L_{\nu}| \leq 1$, inequality (25) stems along the same lines as above.

APPENDIX D. PROOF OF THE THEOREM 3.1

Denote by \bar{E}_t the integrated workload at time t , that is:

$$\bar{E}_t \stackrel{\text{def}}{=} \int_0^t W(u)du, \quad (69)$$

where $\{W(t), t \geq 0\}$ is the workload process given in (1). Recall that $\{S_t, t \geq 0\}$ denotes the on-off process equal to 0 in idle periods and equal to 1 in busy periods (see (2)). Define by $\rho(x, y)$ the probability:

$$\rho(x, y) = \mathbb{P}(S_x = 0, \bar{E}_x \leq y). \quad (70)$$

In a first step, we calculate the Laplace transform $\mathcal{L}\rho$ of ρ using the renewal process of the idle and busy periods. Note that this renewal process is stationary. Define by $\{R_n, n \geq 1\}$ the successive time instants of the end of the busy periods and by $\{A_n, n \geq 1\}$ the integrated

workload at the end of the busy periods,

$$R_n \stackrel{\text{def}}{=} \sum_{k=1}^n (X'_k + Z_{k-1}) \quad \text{and} \quad A_n \stackrel{\text{def}}{=} \sum_{k=1}^n Y'_k, \quad n \geq 1, \quad (71)$$

where we have set $R_0 \stackrel{\text{def}}{=} 0$ and $A_0 \stackrel{\text{def}}{=} 0$.

Proposition D.1. *Under Assumption (H-1)-(H-2), for any $(s, p) \in \mathbb{C}^2$ such that $\text{Re}(s) > 0$ and $\text{Re}(p) > 0$,*

$$\mathcal{L}\rho(s, p) = \frac{1}{s + \lambda - \lambda \mathcal{L}P'(s, p)} \times \frac{\mathcal{L}P'(s, p)}{p(s + \lambda)} + \frac{1}{p(s + \lambda)}.$$

Proof. The proof is based on classical renewal arguments (see for instance Coccozza-Thivent (1997)) and the fact that for all integer k , the idle period Z_k is distributed according to an exponential distribution with scale parameter λ , \mathcal{E}_λ . Note that the event $\{S_x = 0, \bar{E}_x \leq y\}$ may be decomposed as

$$\begin{aligned} & \{S_x = 0, \bar{E}_x \leq y\} \\ &= \{x < T'_1\} \cup \left(\bigcup_{n \geq 1} \left\{ T'_n + X'_n \leq x < T'_{n+1}, \sum_{k=1}^n Y'_k \leq y \right\} \right) \\ &= \{x < T'_1\} \cup \left(\bigcup_{n \geq 1} \{R_n \leq x < R_n + Z_n, A_n \leq y\} \right), \end{aligned} \quad (72)$$

where A_n and R_n are defined in (71). Consequently :

$$\rho(x, y) - e^{-\lambda x} = \lambda \int_0^{+\infty} \sum_{n \geq 1} \mathbb{P}(x - u < R_n \leq x, A_n \leq y) e^{-\lambda u} du,$$

and since

$$\begin{aligned} & \int_0^{+\infty} \lambda \mathbb{P}(x - u < R_n \leq x, A_n \leq y) e^{-\lambda u} du \\ &= \mathbb{P}(R_n \leq x, A_n \leq y) - \lambda \int_0^{+\infty} \mathbb{P}(R_n \leq u - x, A_n \leq y) e^{-\lambda u} du, \end{aligned}$$

the proof follows from the identity

$$\int_0^\infty \int_0^\infty \mathbb{P}(R_n \leq x, A_n \leq y) e^{-sx} e^{-py} dx dy = \frac{1}{sp} \left(\frac{\lambda}{s + \lambda} \mathcal{L}P'(s, p) \right)^n.$$

□

We will now derive another expression for $\mathcal{L}\rho$, using standard properties of the Poisson process.

Proposition D.2. *Under Assumption (H-1)-(H-2), for any $(s, p) \in \mathbb{C}^2$ such that $\operatorname{Re}(s) > 0$ and $\operatorname{Re}(p) > 0$,*

$$\begin{aligned} \mathcal{L}\rho(s, p) &= \frac{1}{p(s + \lambda)} \\ &+ \frac{1}{p} \int_0^{+\infty} e^{-sx} e^{-\lambda x} \left[\exp \left(\lambda \int_0^{\infty} k(x, v) e^{-pv} dv \right) - 1 \right] dx . \end{aligned}$$

Proof. Denote by $\{\mathcal{N}_t, t \geq 0\}$ the counting process associated to the homogeneous Poisson process $\{T_k, k \geq 0\}$ of the arrivals, more explicitly $\mathcal{N}_t = \sum_{n=1}^{\infty} \mathbb{1}\{T_n \leq t\}$. By conditioning the event $\{S_x = 0, \bar{E}_x \leq y\}$ on the event $\{\mathcal{N}_x = n\}$,

$$\begin{aligned} \rho(x, y) &= e^{-\lambda x} + \\ &\sum_{n \geq 1} \mathbb{P}(\mathcal{N}_x = n) \mathbb{P} \left(\{T_i + X_i \leq x\}_{i=1}^n, \sum_{k=1}^n Y_k \leq y \mid \mathcal{N}_x = n \right) . \end{aligned} \quad (73)$$

The conditional distribution of the arrival times (T_1, \dots, T_n) given $\{\mathcal{N}_x = n\}$ is equal to the distribution of the order statistics of n independent and identically distributed uniform random variables on $[0, x]$; hence, for any n -tuple (x_1, \dots, x_n) of positive real numbers,

$$\begin{aligned} \mathbb{P}(T_1 \leq x_1, \dots, T_n \leq x_n \mid \mathcal{N}_x = n) \\ = \mathbb{P}(U_{(1)} \leq x_1, \dots, U_{(n)} \leq x_n) , \end{aligned} \quad (74)$$

where $\{U_k\}_{k=1}^n$ are independent and identically distributed random variables uniformly distributed on $[0, x]$ and $U_{(1)} \leq \dots \leq U_{(n)}$ are the order statistics. Therefore, (73) and (74) imply that

$$\begin{aligned} A &\stackrel{\text{def}}{=} \mathbb{P} \left(\{T_i + X_i \leq x\}_{i=1}^n, \sum_{k=1}^n Y_k \leq y \mid \mathcal{N}_x = n \right) \\ &= \frac{1}{x^n} \int \cdots \int \prod_{k=1}^n \mathbb{1}\{u_k + x_k \leq x\} \mathbb{1} \left\{ \sum_{k=1}^n y_k \leq y \right\} \prod_{k=1}^n P(dx_k, dy_k) du_k , \end{aligned}$$

since the latter integral is invariant by permuting the indexes. An application of the Fubini theorem leads to

$$A = \frac{1}{x^n} \int \cdots \int \mathbb{1} \left\{ \sum_{k=1}^n y_k \leq y \right\} \prod_{k=1}^n \kappa(x, dy_k) ,$$

where $\kappa(x, dy)$ is the probability kernel defined by

$$\kappa(x, dy) \stackrel{\text{def}}{=} \int (x - u) \mathbb{1}\{u \leq x\} P(du, dy) . \quad (75)$$

We obtain, for any p such that $\operatorname{Re}(p) > 0$,

$$\begin{aligned} \int_0^\infty \rho(x, v) e^{-pv} dv &= \frac{e^{-\lambda x}}{p} + \frac{1}{p} \sum_{n \geq 1} \frac{\lambda^n}{n!} e^{-\lambda x} \left[\int_0^\infty \kappa(x, dv) e^{-pv} \right]^n \\ &= \frac{e^{-\lambda x}}{p} + \frac{e^{-\lambda x}}{p} \left[\exp \left(\lambda \int_0^\infty \kappa(x, dv) e^{-pv} \right) - 1 \right], \end{aligned}$$

and hence

$$\begin{aligned} \mathcal{L}\rho(s, p) &= \frac{1}{p(s + \lambda)} \\ &\quad + \frac{1}{p} \int_0^{+\infty} e^{-su} e^{-\lambda u} \left[\exp \left(\lambda \int_0^\infty e^{-pv} \kappa(u, dv) \right) - 1 \right] du. \end{aligned}$$

□

The proof of Theorem 3.1 is then a direct consequence of Proposition D.1 and Proposition D.2 and the fact that

$$a(x, p) = \exp \left(\lambda \int_0^\infty e^{-pv} \kappa(x, dv) \right)$$

The result is extrapolated on the line $\operatorname{Re}(p) = 0$ by continuity in p at fixed s such that $\operatorname{Re}(s) > 0$.

APPENDIX E. TECHNICAL LEMMAS

Lemma E.1. *Let $c > 0$ and $\eta_0 > 0$. for any complex valued functions z_1 and z_2 satisfying*

$$\sup_{\omega \in \mathbb{R}, i=1,2} |z_i(\omega)| \leq 1, \quad (76)$$

let $z = (z_1, z_2)$ and denote by Ψ_z the function defined on $\mathbb{R}_+ \times \mathbb{R}$ by

$$\Psi_z(\tilde{\lambda}, \omega) \stackrel{\text{def}}{=} \frac{z_1(\omega)}{(c + i\omega + \tilde{\lambda})(c + i\omega + \tilde{\lambda} - \tilde{\lambda}z_2(\omega))}.$$

Then the following assertions hold :

- (i) *The function $\tilde{\lambda} \mapsto \int_{-\infty}^{+\infty} \Psi_z(\tilde{\lambda}, \omega) d\omega$ is continuously differentiable on \mathbb{R}_+ and its derivative is bounded independently of z over $\tilde{\lambda} \in [0, \eta_0]$.*
- (ii) *There exists $K > 0$ only depending on c and η_0 such as, for any $W \geq 1$ or $W = \infty$ and any function $\tilde{z} = (\tilde{z}_1, \tilde{z}_2)$ also satisfying (76),*

$$\begin{aligned} \sup_{\tilde{\lambda} \in [0, \eta_0]} \left| \int_{-\infty}^{+\infty} \Psi_z(\tilde{\lambda}, \omega) d\omega - \int_{-\infty}^{+\infty} \Psi_{\tilde{z}}(\tilde{\lambda}, \omega) d\omega \right| \\ \leq K \left(\max_{i=1,2} \sup_{\omega \in [-W, W]} |z_i(\omega) - \tilde{z}_i(\omega)| + \frac{1}{W} \right). \end{aligned}$$

Proof. For all ω in \mathbb{R} and $\tilde{\lambda} \in [0, \eta_0]$, by using (76), we obtain

$$\left| \partial_{\tilde{\lambda}} \Psi(\tilde{\lambda}, \omega) \right| \leq \frac{3(c + |\omega|) + 4\eta_0}{(c^2 + \omega^2) \left(\sqrt{(c + \tilde{\lambda})^2 + \omega^2} - \tilde{\lambda} \right)^2}. \quad (77)$$

From Jensen inequality, one can easily show that for all α in $[0, 1]$ and $\tilde{\lambda} \in [0, \eta_0]$,

$$\sqrt{(c + \tilde{\lambda})^2 + \omega^2} \geq (c + \tilde{\lambda})\sqrt{\alpha} + |\omega|\sqrt{1 - \alpha}. \quad (78)$$

Choosing α close enough to 1 so that $(\sqrt{\alpha} - 1)\eta_0 + c\sqrt{\alpha} > 0$, (77) and (78) yields to

$$\left| \partial_{\tilde{\lambda}} \Psi(\tilde{\lambda}, \omega) \right| \leq \frac{3(c + |\omega|) + 4\eta_0}{(c^2 + \omega^2) \left((\sqrt{\alpha} - 1)\eta_0 + c\sqrt{\alpha} + |\omega|\sqrt{1 - \alpha} \right)^2},$$

which is valid independently of $\tilde{\lambda}$ in $[0, \eta_0]$, and whose RHS is integrable over ω in \mathbb{R} , hence (i). For showing (ii), observe that for all $\tilde{\lambda}$ in $[0, \eta_0]$

$$\begin{aligned} & \Psi_z(\tilde{\lambda}, \omega) - \Psi_{\tilde{z}}(\tilde{\lambda}, \omega) \\ &= \frac{(z_1(\omega) - \tilde{z}_1(\omega))}{(c + i\omega + \tilde{\lambda} - \tilde{\lambda}z_2(\omega))(c + i\omega + \tilde{\lambda} - \tilde{\lambda}\tilde{z}_2(\omega))} \\ &+ \frac{\tilde{\lambda}(\tilde{z}_1(\omega)z_2(\omega) - \tilde{z}_2(\omega)z_1(\omega))}{(c + i\omega + \tilde{\lambda})(c + i\omega + \tilde{\lambda} - \tilde{\lambda}z_2(\omega))(c + i\omega + \tilde{\lambda} - \tilde{\lambda}\tilde{z}_2(\omega))}. \end{aligned}$$

Using again (77) and since, using (76), $|\tilde{z}_1z_2 - \tilde{z}_2z_1| = |\tilde{z}_1(z_2 - \tilde{z}_2) - \tilde{z}_2(z_1 - \tilde{z}_1)| \leq |z_2 - \tilde{z}_2| + |z_1 - \tilde{z}_1|$, for α chosen as above and for each $\tilde{\lambda}$ in $[0, \eta_0]$, we have

$$\begin{aligned} & |\Psi_z(\tilde{\lambda}, \omega) - \Psi_{\tilde{z}}(\tilde{\lambda}, \omega)| \leq \\ & \frac{(c + |\omega| + 3\eta_0)}{\sqrt{c^2 + \omega^2} \left((\sqrt{\alpha} - 1)\eta_0 + c\sqrt{\alpha} + |\omega|\sqrt{1 - \alpha} \right)^2} \max_{i=1,2} |z_i(\omega) - \tilde{z}_i(\omega)|. \end{aligned} \quad (79)$$

Observe that in the RHS above, the fraction is integrable over $\omega \in \mathbb{R}$ and is equivalent to $[(1 - \alpha)|\omega|]^{-2}$ as $|\omega| \rightarrow \infty$. Consequently, there exists constants K_1 and K_2 depending only on c and η_0 such as

$$\begin{aligned} & \int_{-W}^{+W} |\Psi_z(\tilde{\lambda}, \omega) - \Psi_{\tilde{z}}(\tilde{\lambda}, \omega)| d\omega \\ & \leq K_1 \max_{i=1,2} \sup_{\omega \in [-W, W]} |z_i(\omega) - \tilde{z}_i(\omega)| \frac{1}{\sqrt{1 - \alpha}(c\sqrt{\alpha} + (\sqrt{\alpha} - 1)\eta_0)} \end{aligned}$$

and, since $\max_{i=1,2} |z_i(\omega) - \tilde{z}_i(\omega)| \leq 2$,

$$\int_{[-W, W]^c} |\Psi_z(\tilde{\lambda}, \omega) - \Psi_{\tilde{z}}(\tilde{\lambda}, \omega)| d\omega \leq K_2 W^{-1},$$

hence (ii). □

REFERENCES

- ANSI NORM (1999). *American National Standard for Calibration and Use of Germanium Spectrometers for the Measurement of Gamma-Ray Emission Rates of Radionuclides. ANSI N42.14-1999*. American National Standards Institute.
- BACCELLI, F. and BRÉMAUD, P. (2002). *Elements of Queueing Theory*. Springer.
- BINGHAM, N. H. and PITTS, S. M. (1999). Non-Parametric Estimation for the M/G/ ∞ Queue. *Ann. Inst. Statist. Math.* **51** 71–97.
- BRISTOW, Q. (1990). *Theoretical and experimental investigations of coincidences in Poisson distributed pulse trains and spectral distortion caused by pulse pileup*. Ph.D. thesis, Carleton University, Ottawa, Canada.
- COCOZZA-THIVENT, C. (1997). *Processus Stochastiques et Fiabilité des Systèmes*. Springer Verlag.
- DOETSCH, G. (1974). *Introduction to the Theory and the Application of the Laplace Transform*. Springer-Verlag.
- GAUTSCHI, W. (1997). *Numerical Analysis: an Introduction*. 1st ed. Birkhäuser.
- HALL, P. (1988). *Introduction to the Theory of Coverage Processes*. Wiley.
- HALL, P. and PARK, J. (2004). Nonparametric Inference about Service Time Distribution from Indirect Measurements. *Jour. Roy. Statist. Soc.* **66** 861–875.
- KNOLL, G. (1989). *Radiation Detection and Measurement*. 2nd ed. Wiley.
- LEO, W. R. (1994). *Techniques for Nuclear and Particle Physics Experiments: A How-To Approach*. Springer Verlag.
- PYKE, R. (1958). On renewal processes related to type I and type II counter models. *Ann. Math. Statist.* **29** 737–754.
- TAKACS, L. (1962). *Introduction to the Theory of Queues*. Oxford University Press.
- TRIGANO, T., MONTAGU, T., MOULINES, E., ROUEFF, F. and SOULOUMIAC, A. (2005). Analytical Pileup Correction Method for HPGe Detectors. *submitted* .
- VAN DER WAART, A. W. and WELLNER, J. (1996). *Weak Convergence and Empirical Processes with Applications to Statistics*. Springer Verlag.