



HAL
open science

New method to derive transport properties including diffusion in a two-temperature plasma

Vincent Rat, Pascal André, Jacques Aubreton, Marie-Françoise Elchinger,
Pierre Fauchais, André Lefort

► **To cite this version:**

Vincent Rat, Pascal André, Jacques Aubreton, Marie-Françoise Elchinger, Pierre Fauchais, et al..
New method to derive transport properties including diffusion in a two-temperature plasma. 15
International Symposium on Plasma Chemistry, 2001, France. pp.819-824. hal-00019934

HAL Id: hal-00019934

<https://hal.science/hal-00019934v1>

Submitted on 2 Mar 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New method to derive transport properties including diffusion in a two-temperature plasma

V. Rat¹, P. André², J. Aubreton¹, M.F. Elchinger¹, P. Fauchais¹, and A. Lefort²

¹ SPCTS University of Limoges, 123 av. A. Thomas, 87060 Limoges cedex, France

² LAEPT Blaise Pascal University, 24 av. des Landais, 63177 Aubière cedex, France

Abstract

A new derivation of transport properties in a two-temperature plasma has been performed. The electron kinetic temperature T_e is supposed to be different from that of heavy species T_h . The resolution of the Boltzmann's equation, thanks to the Chapman-Enskog method, is used to calculate transport coefficients and it allows to the generalisation of bracket integrals out of thermal equilibrium. Two-temperature diffusion coefficients are defined and the obtained results are presented for an atmospheric two-temperature argon plasma.

1. Introduction

Recent works [1] have shown that the simplified theory of transport properties out of thermal equilibrium introduced by Devoto [2] and Bonnefoi [3] neglecting the coupling between electrons and heavy species, and very often used in modeling, leads to unphysical results. Thus, two-temperature transport coefficients have been derived [4] without the Bonnefoi's assumptions starting from the Boltzmann's equation. The electron temperature T_e is supposed to be different from that of heavy species T_h . Only elastic processes are considered in a collision dominated plasma. Firstly, is presented the principle of the resolution of the Boltzmann's equation in a two-temperature model. Secondly, the modifications of this resolution, with respect to equilibrium, give rise to a generalisation of bracket integrals giving transport coefficients. The differences with thermal equilibrium are then highlighted. Lastly, two-temperature diffusion coefficients resulting of this derivation are displayed as function of the non-equilibrium parameter $\theta=T_e/T_h$.

2. Resolution of the Boltzmann equation

The distribution function f_i of the i th species is the solution of the integro-differential equation of Boltzmann [5]:

$$\frac{Df_i}{Dt} = \sum_{j=1}^N \iiint (f'_i f'_j - f_i f_j) g b d b d \epsilon d \bar{v}_j \quad (1)$$

f'_i is the distribution function after the elastic collision of the i th species, g is the relative velocity of the species i and j , b and ϵ are respectively the impact parameter and the incidence azimuthal angle.

It has been assumed that the zero-order approximation function is Maxwellian at T_e for electrons and T_h for heavy species. The distribution function of the i th species, solution of (1), is approximated by a Maxwellian distribution function $f_i^{(0)}$ perturbed by a slowly time and space dependent first-order perturbation function Φ_i of the i th species with the temperature T_i :

$$f_i = f_i^{(0)}(1 + \Phi_i) \quad (2)$$

Inserting (2) into (1), it can be shown that:

$$\frac{Df_i^{(0)}}{Dt} = I_i^{(0)} + \sum_{j=1}^N \iiint f_i^{(0)} f_j^{(0)} ((\Phi_i' + \Phi_j') K_i - \Phi_i - \Phi_j) \mathcal{E} \mathcal{B} d\mathcal{B} d\mathcal{E} d\vec{v}_j \quad (3)$$

$I_i^{(0)}$ represents the zero-order approximation of the Chapman-Enskog's expansion and does not vanish for two colliding particles with different temperatures:

$$I_i^{(0)} = \sum_{j=1}^N \iiint (f_i^{(0)} f_j^{(0)} - f_i^{(0)} f_j^{(0)}) \mathcal{E} \mathcal{B} d\mathcal{B} d\mathcal{E} d\vec{v}_j \quad (4)$$

A term $K_i(W_i, \theta_{ij})$, taking into account the thermal non-equilibrium when electrons and heavy species collide, has been introduced as follows:

$$f_i^{(0)} f_j^{(0)} = f_i^{(0)} f_j^{(0)} K_i(W_i, \theta_{ij}) \quad \forall i, j \in [1; N] \quad (5)$$

where $K_i(W_i, \theta_{ij}) = \exp(-(W_i'^2 - W_i^2)(1 - \theta_{ij}))$ (6)

with $\theta_{ij} = \frac{T_i}{T_j}$ (7)

and W_i' is the reduced velocity after collision such as $\bar{W}_i = \left(\frac{m_i}{2kT_i}\right)^{1/2} \bar{V}_i$ where $\bar{V}_i = \vec{v}_i - \vec{v}_0$. \vec{v}_i and \vec{v}_0 are respectively the velocity of the i th species and flow velocity.

The introduction of K_i allows the definition of bracket integrals to be generalised out of thermal equilibrium.

The calculation of $\frac{Df_i^{(0)}}{Dt}$ and $\frac{Df_i^{(0)}}{Dt}$ ($i \geq 2$) is obtained using the equations of change

[4]. The transport terms in equations (3), show that transport phenomena are due to a new gradient $\bar{V} \ln \theta$, which characterizes the temperature difference between electrons and heavy species, the heavy species temperature gradient, the velocity gradient, external forces (forced diffusion), the concentration and pressure gradients and a term acting on the hydrostatic pressure to the first-order approximation of Sonine polynomials.

Following the linear form of transport terms, it can be supposed that the first-order perturbation function can be written as:

$$\Phi_i = -\bar{A}_i \cdot \bar{\nabla} \ln T_h - \bar{B}_i : \bar{\nabla} \bar{v}_0 + \sum_{j=1}^N \bar{C}_i^j \cdot \bar{d}_j + D_i Q_i^{(0)} + \sum_{j=1}^N \bar{E}_i^j \cdot g_j \bar{\nabla} \ln \theta - \bar{F}_i \cdot \bar{\nabla} \ln \theta \quad (8)$$

The diffusion forces are written as follows for *electrons*:

$$\bar{d}_1 = \frac{\rho_1}{\rho} \sum_{j=1}^N n_j \bar{F}_j - n_1 \bar{F}_1 + \left(\frac{x_1 \theta}{D} - \frac{\rho_1}{\rho} \right) \bar{\nabla} p + \frac{\theta p}{D^2} \bar{\nabla} x_1 \quad (9)$$

and
$$g_1 = \frac{x_1 p (1 - x_1)}{D^2} \quad (10)$$

For *heavy species*, it is written:

$$\bar{d}_i = \frac{\rho_i}{\rho} \sum_{j=1}^N n_j \bar{F}_j - n_i \bar{F}_i + \left(\frac{x_i}{D} - \frac{\rho_i}{\rho} \right) \bar{\nabla} p + \frac{p}{D} \bar{\nabla} x_i - \frac{x_i (\theta - 1) p}{D^2} \bar{\nabla} x_i \quad (11)$$

and
$$g_i = -\frac{x_i x_1 p}{D^2} \quad (12)$$

ρ_i , \bar{F}_i , x_i and p are respectively the density, an external force and the molar fraction of the i th species and the total pressure.

$Q_i^{(0)}$ corresponds to the exchanged kinetic energy between electrons and heavy species during collisions. It can be shown:

$$Q_i^{(0)} = 4k_B n_1 (T_h - T_c) \left(\frac{8k_B T_c}{\pi m_1} \right)^{1/2} \sum_{j=2}^N n_j \frac{m_1}{m_j} \bar{Q}_{ij}^{(1,1)} \quad (13)$$

k_B being the Boltzmann constant.

The unknowns \bar{A}_i , \bar{B}_i , \bar{C}_i^j , D_i , \bar{E}_i^j and \bar{F}_i are determined by assuming that they can be splitted on the Sonine polynomial basis [5] which allows the introduction of systems of linear equations following the approximation order.

The introduction of $\bar{\nabla} \ln \theta$ prevents from considering the uncoupling between electrons and heavy species in the resolution of systems of linear equations which gives the transport coefficients. As a result, two-temperature diffusion coefficients not only include ordinary and thermal diffusion but also new diffusion coefficients due to the temperature difference ($\bar{\nabla} \ln \theta$) between electrons and heavy species [4].

3. Bracket integrals

Transport coefficients results in the resolution of systems of linear equations. Each element q_{ij}^{mp} (m and p represents the order of approximation of Sonine polynomials) of the considered matrix depends on bracket integrals. The latter are expressed as a linear combination of collision integrals which account for the interaction between the colliding species.

For example, in the first-order approximation of Sonine polynomials, the following bracket integral can be written:

$$\left[\bar{W}_i S_{3/2}^1(w_i^2) \bar{W}_j S_{3/2}^1(w_j^2) \right]_{ij} = C \sum_{r\ell} A_{11r\ell} \Omega_{ij}^{(r,\ell)} \quad (14)$$

where C depends on m_i , m_j , T_i and T_j .

$\Omega_{ij}^{(r,\ell)}$, i and j being the colliding species, is the integral of the transport cross section over a Maxwellian distribution function. Table 1 gives the values of the coefficients $A_{11r\ell}$ at equilibrium ($\theta=1.0$) and for different values of θ calculated for a collision between an electron and a heavy species, that is for a mass ratio of 10^{-5} .

Table 1: Calculation of $A_{11r\ell}$ for different values of θ and with a mass ratio $m_j/m_i=10^{-5}$.

θ	1.00	1.25	1.50	2.00	3.00
A_{1111}	-13.75	-8.50	-5.00	-0.63	-3.75
A_{1121}	5.00	3.30	3.00	4.50	11.00
A_{1122}	2.00	0.75	-0.50	-3.00	-8.00
A_{1131}	-1.00	-1.40	-2.00	-3.50	-7.00
A_{1132}	0.00	0.50	1.00	2.00	4.00

Firstly, it is shown that our results strictly lead to those of Hirschfelder et al [5]. Secondly, the coefficients $A_{11r\ell}$ are drastically changed following the applied thermal non-equilibrium. Lastly, coefficients such as A_{1132} do not vanish when the non-equilibrium parameter θ is different from unity. As a result, collision integrals $\Omega_{ij}^{(r,\ell)}$ with indexes (r, ℓ) higher than at thermal equilibrium should be taken into account out of thermal equilibrium.

4. Two-temperature diffusion coefficients

Two-temperature diffusion coefficients are displayed for a two-temperature argon plasma. Calculations are performed at atmospheric pressure and the two-temperature plasma composition is obtained using the Saha equation for ionization of Van de Sanden et al [6].

Electrons, Ar and Ar^+ species have been considered.

First, it has to be noted that the rate of convergence of diffusion coefficients are similar to that of thermal equilibrium that is the second-order approximation is sufficient for diffusion coefficients implying electrons. However, for argon, around 5000 K (the electron temperature) the third-order approximation is required because of the Ramsauer effect of the momentum cross section.


The two-temperature diffusion coefficients D_{eAr} and D_{Ar^+e} are respectively displayed in figure 1(a) and 1(b) as a function the electron temperature from 1000 up to 20000 for different values of θ (between 1 and 3). The calculation have been performed to the third-order approximation. When ionisation is weak, namely below 6000 K, two-temperature ordinary diffusion coefficients increase with temperature.

For D_{eAr} , the effect of temperature is dominant with respect to the collision integral $\bar{Q}_{eAr}^{(1,1)}$ which increases slowly with T_e . However, between 4500 and about 6000 K, the effect of the collision integral $\bar{Q}_{eAr}^{(1,1)}$ overcomes the effect of temperature but, at these temperatures the binary approximation is not valid anymore. As a result, the Ar^+ species has to be taken into account and the effect of the charge transfer collision integral $\bar{Q}_{ArAr^+}^{(1,1)}$ contributes to increase


ordinary diffusion coefficients since it decreases with temperature faster than $\bar{Q}_{eAr}^{(1,1)}$ increases (for example, $\bar{Q}_{ArAr^+}^{(1,1)} / \bar{Q}_{eAr}^{(1,1)}$ is about 55 at 5000 K). The same evolution for D_{Ar^+e} is observed, the charge transfer collision integral being dominant at low temperature.

When ionization becomes significant around 6000 K, D_{eAr} and D_{Ar^+e} decreases with temperature because the charged-charged collision integral $\bar{Q}_{eAr^+}^{(1,1)}$ counterbalances the effect of temperature.

Besides, it is observed in figure 1(a) and 1(b) that two-temperature diffusion coefficients decrease with the non-equilibrium parameter θ . Figure 2 depicts the influence of the non equilibrium parameter θ on diffusion coefficients and that it can be written that at low temperature $D_{eAr} = \theta D_{Ar^+e}$.



(a)



(b)

Figure 1: Two-temperature diffusion coefficient D_{eAr} (a) and D_{Ar^+e} (b) as function of electron temperature T_e and θ .




Figure 2: Ratio D_{eAr}/D_{Are} as function of electron temperature T_e and θ .

5. Conclusion

A new method of derivation of transport properties in a two-temperature plasma has been performed. Two-temperature transport coefficients have been derived starting from the Boltzmann's equation. The latter is solved by using the well-known Chapman-Enskog method which has been adapted to thermal non-equilibrium plasmas. It has been assumed that the distribution function of species, solution of the Boltzmann's equation, is a Maxwellian, at T_e for electrons and T_h for heavy species, perturbed by a slowly time and space dependent first-order perturbation function. The introduction of the gradient $\bar{\nabla}\theta$ allows to maintain the coupling between electrons and heavy species in the calculation of two-temperature transport coefficients. Two-temperature diffusion coefficients in an atmospheric argon plasma were calculated between electrons and heavy species, contrarily to what happens when using the simplified theory of transport coefficients of Devoto[2] and Bonnefoi [3].

References

- [1] V. Rat, J. Aubreton, M.F. Elchinger, and P. Fauchais, *Plasma Chem. Plasma Process.* **21** (2001), 355-364.
- [2] R.S. Devoto, *Phys. Fluids* **10** (1967), 2105-2112.
- [3] C. Bonnefoi, State Thesis, University of Limoges, France (1983).
- [4] V. Rat, P. André, J. Aubreton, M.F. Elchinger, P. Fauchais, and A. Lefort, to be published, *Phys. Rev. E*.
- [5] J.O. Hirschfelder, C.F. Curtis, and R.B. Bird, *Molecular Theory of Gases and Liquids*, 2nd ed., Wiley, New York (1964).
- [6] M.C.M. Van de Sanden, P.P.J.M. Schram, A.G. Peeters, J.A.M. van der Mullen and G.M. Kroesen, *Physical Review A*, **40** (1989), 5273-5276.