



# Stability in Discrete Tomography: Some Positive Results

Sara Brunetti, Alain Daurat

## ► To cite this version:

Sara Brunetti, Alain Daurat. Stability in Discrete Tomography: Some Positive Results. Discrete Applied Mathematics, 2005, 147, pp.207-226. 10.1016/j.dam.2004.09.012 . hal-00023085

**HAL Id: hal-00023085**

**<https://hal.science/hal-00023085>**

Submitted on 19 Apr 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Stability in Discrete Tomography: Some Positive Results

Sara Brunetti <sup>a</sup> and Alain Daurat <sup>b</sup>

<sup>a</sup>*Dipartimento di Scienze Matematiche e Informatiche, Università di Siena, Via  
del Capitano 15, 53100, Siena, Italy*

<sup>b</sup>*LSIT UMR 7005 CNRS-ULP, Pôle API, Boulevard Sébastien Brant, 67400  
Illkirch-Graffenstaden, France*

---

## Abstract

The problem of reconstructing finite subsets of the integer lattice from X-rays has been studied in discrete mathematics and applied in several fields like image processing, data security, electron microscopy. In this paper we focus on the stability of the reconstruction problem for some special lattice sets. First we prove that if the sets are additive, then a stability result holds for very small errors. Then, we study the stability of reconstructing convex sets from both an experimental and a theoretical point of view. Numerical experiments are conducted by using linear programming that support the conjecture that convex sets are additive with respect to a set of suitable directions, and consequently the reconstruction problem is stable. The theoretical investigation provides a stability result for lattice sets. It is used to prove the following property: if a sequence of lattice convex sets have X-rays in suitable directions which converge to X-rays of a convex body, then it converges to this convex body.

*Key words:* Discrete Tomography, Stability, Linear Programming, Additivity, Convexity

---

## 1 Introduction

A *lattice set* is a non-empty finite subset of the integer lattice  $\mathbb{Z}^2$ . A *lattice direction* is a direction directed by a vector in  $\mathbb{Z}^2 \setminus \{0\}$ , and it can also be given by an equation  $p(x, y) = ax + by$  with  $a, b \in \mathbb{Z}$ . Further, the *X-ray* of a lattice

---

*Email addresses:* sara.brunetti@unisi.it (Sara Brunetti),  
daurat@dpt-info.u-strasbg.fr (Alain Daurat).

set  $E$  in a lattice direction  $p$  is the function  $X_p E$  giving the number of points in  $E$  on each line parallel to this direction, formally  $X_p E(k) = \text{card}(\{M \in E : p(M) = k\})$ . Discrete Tomography is the area of mathematics and computer science that deals with the inverse problem of reconstructing lattice sets from a finite set of X-rays. An overview on this subject highlighting the applications, the mathematical foundations, and the algorithms in Discrete Tomography is provided by the book [16].

In this paper we focus on the stability of the reconstruction problem.

Informally, a problem is stable if a small perturbation of the data does not change the corresponding solutions too much. Therefore, the stability problem is of main importance in practical applications where the X-rays are possibly affected by errors. For instance, in electron microscopy, techniques that enable to count the number of atoms lying in a line up to an error of  $\pm 1$  are known [14]. So, in case of instability, the reconstructed set can be quite different from the original one even if the error on the data is small. In [1] the authors prove that when  $m > 2$ , the two sets can be even disjoint, permitting an error of  $2(m-1)$  on the X-rays. First, we show in Remark 6 that to obtain a stability result even with a very small error on the data the requirement of uniqueness for the sets is not enough. To this goal, we shall consider the reconstruction of lattice sets with some additional constraints.

In Section 3 we treat the stability of reconstructing additive sets. This class of sets was first introduced by Fishburn et al. in [10]. Here we just recall to the reader that additivity implies uniqueness, whereas the converse is not true. Additionally, the notion of additivity should be regarded as a property of the solutions of the linear program associated to the reconstruction problem. We prove that if the sets are additive, then a stability result holds (Proposition 7).

In Section 4 we study the stability of reconstructing convex sets from both an experimental and a theoretical point of view. In the former, we use linear programming to deal with this problem. Experimental results suggest the conjecture that for the set of directions  $\{x, y, 2x + y, -x + 2y\}$ , convex sets are additive. This would imply that the results of Section 2 may hold to convex sets so giving a stability result that corresponds to the continuous case where the reconstruction problem for convex bodies is well-posed ([22]). In the latter, the theoretical result (Proposition 18) confirms stability for convex sets by exploiting the result in [22]. The last proposition of this paper allows to use Discrete Tomography to solve Hammer's X-ray problem which is reconstructing a continuous convex body from continuous X-rays. More precisely we prove that a convex body is arbitrarily near lattice convex sets on the condition that the X-rays of the lattice convex sets are near enough the ones of the convex body. In practice it means that we can reconstruct a convex body

from discrete X-rays to any precision if the resolution and the error about the X-ray can be as small as wanted. This result is very linked to other tries to approximate the convex body from a sequence of discrete objects reconstructed from the X-rays (see [19]).

## 2 The Problem

The reconstruction problem is the task of determining any lattice set having the given X-rays. Stability concerns how sensitive is the problem to noisy data. Hence one can ask whether a perturbation of the data correspond solutions that are close. To study the problem we define a measure for the error on the X-rays and one for the distance of two solutions. Let  $\mathcal{D}$  be a set of  $m$  prescribed lattice directions with  $m \geq 2$  and let  $E, F$  be lattice sets:

$$DX_{\mathcal{D}}(E, F) = \max_{p \in \mathcal{D}} \sum_{k \in \mathbb{Z}} |X_p E(k) - X_p F(k)|$$

and

$$\text{card}(E \triangle F) = \text{card}((E \setminus F) \cup (F \setminus E)).$$

The formulation of the problem that we consider is the following:

**Problem 1** *Let  $E$  be known. Determine  $F$  maximizing  $\text{card}(E \triangle F)$ , with the constraint that  $DX_{\mathcal{D}}(E, F)$  is given.*

Let us introduce some definitions that we need in the following.

**Definition 2** *A lattice set  $E$  is additive with respect to  $\mathcal{D}$ , or  $\mathcal{D}$ -additive, if there is a function  $e$  which gives a value  $e_p(k)$  for each line  $p = k$  parallel to a direction  $p$  of  $\mathcal{D}$  such that for all  $M$  in  $\mathbb{Z}^2$ :*

$$M \in E \text{ if and only if } \sum_{p \in \mathcal{D}} e_p(p(M)) > 0.$$

This definition introduced by Fishburn et al. can be better understood with linear programming: a lattice set  $E$  is additive if it is the unique solution of the linear programming problem which looks for a fuzzy set which has the same X-rays than  $E$ .

**Definition 3** *A lattice set  $E$  is unique with respect to  $\mathcal{D}$ , or  $\mathcal{D}$ -unique, if  $F \subset \mathbb{Z}^2$  and  $X_p E = X_p F$  for any  $p \in \mathcal{D}$  imply  $E = F$ .*

There is an intimate relationship between these two definitions: every  $\mathcal{D}$ -additive set is  $\mathcal{D}$ -unique and the converse is true if  $m = 2$  (see[10]).

As a last remark we recall that if  $p$  and  $q$  are two directions, then a  $p$ -line does not always intersect a  $q$ -line. Indeed  $\mathbb{Z}^2$  can be split in  $\det(p, q)$   $pq$ -lattices such that in each  $pq$ -lattice a  $p$ -line intersects with any  $q$ -line. Precisely a  $pq$ -lattice has the form:

$$L_i^{pq} = \{M : p(M) = i \pmod{\det(p, q)} \text{ and } q(M) = \kappa i \pmod{\det(p, q)}\}$$

where  $\kappa$  only depends on the directions  $p$  and  $q$  (see for example [7]). Moreover we denote by  $\langle i, j \rangle_{pq}$  the point  $M$  such that  $p(M) = i$  and  $q(M) = j$ . Notice that this point is in  $\mathbb{Z}^2$  only if  $p = i$  and  $q = j$  are in the same  $pq$ -lattice.

### 3 Stability for Additive Sets

In this section we study the stability of reconstructing  $\mathcal{D}$ -additive sets. We begin to study Problem 1 with  $E$  and  $F$  verifying the constraint  $DX_{\mathcal{D}}(E, F) \leq 1$ .

In the first two lemmas additivity is not required.

The condition  $DX_{\mathcal{D}}(E, F) \leq 1$  permits the X-rays of the two sets to differ by one in at most a line for each direction. Then,  $p \in \mathcal{D}$  and an integer  $k_p$  exists such that  $|X_p E(k_p) - X_p F(k_p)| = 1$  and  $X_p E(k) = X_p F(k)$  for  $k \neq k_p$ .

**Lemma 4** *If  $p \in \mathcal{D}$  and an integer  $k_p$  exist such that  $|X_p E(k_p) - X_p F(k_p)| = 1$ , then for every  $q \in \mathcal{D}$  there is an integer  $k_q$  such that  $|X_q F(k_q) - X_q E(k_q)| = 1$  and  $\langle k_p, k_q \rangle_{pq} \in \mathbb{Z}^2$ .*

**PROOF.** Let  $L_i^{pq}$  be the  $pq$ -lattice containing the line  $p = k_p$ , or equivalently  $k_p \in p(L_i^{pq})$ . Suppose that  $X_p F(k_p) - X_p E(k_p) = +1$ . Thus, we have that

$$\sum_{k \in p(L_i^{pq})} X_p F(k) = 1 + \sum_{k \in p(L_i^{pq})} X_p E(k).$$

Using the consistency of the X-rays for  $F$  and  $E$ , the previous identity leads to the following

$$\sum_{k \in q(L_i^{pq})} X_q F(k) = 1 + \sum_{k \in q(L_i^{pq})} X_q E(k),$$

for all  $q$  in  $\mathcal{D}$ . From this, the thesis easily follows.  $\square$

In the next lemma we show that all the lines with error 1 have a common point and this point is in  $\mathbb{Z}^2$ . In the following, we assume that  $\text{card}(F) > \text{card}(E)$  and for any  $p \in \mathcal{D}$  the integer  $k_p$  is as in the previous lemma.

**Lemma 5** *If  $DX_{\mathcal{D}}(E, F) = 1$ , then a point  $W \in \mathbb{Z}^2$  exists such that*

$$X_p F(k) = X_p E(k) + 1, \text{ if } k = p(W)$$

$$X_p F(k) = X_p E(k), \quad \text{otherwise}$$

*for all the directions  $p$  in  $\mathcal{D}$ .*

**PROOF.** Let  $p, q$  and  $r$  be directions in  $\mathcal{D}$  and suppose that  $A = \langle k_p, k_q \rangle_{pq}$ ,  $B = \langle k_p, k_r \rangle_{pr}$ ,  $C = \langle k_q, k_r \rangle_{qr}$  are three distinct points. Let  $a, b$  be such that  $r = ap + bq$ . Thus, summing up we can write:

$$\sum_{M \in F} r(M) = a \sum_{M \in F} p(M) + b \sum_{M \in F} q(M)$$

and by grouping line by line we obtain:

$$\sum_k k X_r F(k) = a \sum_k k X_p F(k) + b \sum_k k X_q F(k).$$

We can exhibit the corresponding identity for the set  $E$ . As a result of the difference of these two identities we obtain that  $k_r = ak_p + bk_q$  and so  $r(A) = r(B) = r(C)$ . Thus, the three points  $A, B$  and  $C$  coincide and the claim is proved.  $\square$

**Remark 6** *Given any three lattice directions we can construct two sets  $E, F$  in such a way that they are (non-additive) sets of uniqueness. (We do not give the proof for reasons of space limit and we refer the reader to [10]). Figure 1 illustrates two such sets verifying the constraint  $DX_{\mathcal{D}}(E, F) = 1$ . Since they are disjoint, Proposition 7 does not hold for  $\mathcal{D}$ -unique sets.*

Since uniqueness is not sufficient to have stable solutions for the reconstruction problem, we suppose that  $E$  and  $F$  are  $\mathcal{D}$ -additive, that is  $E = \{M : e(M) > 0\}$  and  $F = \{M : f(M) > 0\}$ .

**Proposition 7** *Let  $E$  and  $F$  be  $\mathcal{D}$ -additive lattice sets. If  $DX_{\mathcal{D}}(E, F) = 1$ , then  $\text{card}(E \triangle F) = 1$ .*

**PROOF.** Let  $W$  be as in Lemma 5. At first suppose that  $W \notin E$  and let  $E' = E \cup \{W\}$ . For each direction  $p$  in  $\mathcal{D}$  we have that  $X_p E' = X_p F$ . Finally, since additivity of  $F$  implies uniqueness of  $F$ , we conclude that  $F = E \cup \{W\}$ . On the contrary, if  $W \in E$  we study the following:

$$\Phi_E = \sum_{M \in \mathbb{Z}^2} \sum_{p \in \mathcal{D}} e_p(p(M))(1_E(M) - 1_F(M)).$$




Fig. 1.  $E$  and  $F$  are non-additive sets of uniqueness such that  $DX_{\mathcal{D}}(E, F) = 1$  and  $E \cap F = \emptyset$ .

Rewriting it as

$$\sum_{M \in E} \sum_{p \in \mathcal{D}} e_p(p(M))(1_E(M) - 1_F(M)) + \sum_{M \notin E} \sum_{p \in \mathcal{D}} e_p(p(M))(1_E(M) - 1_F(M)),$$

we notice that  $\Phi_E > 0$ , because the additivity of  $E$  implies that if  $M$  is in  $E$ , then  $e(M) > 0$  and  $1_E(M) = 1$  holds, and otherwise  $e(M) \leq 0$  and  $1_E(M) = 0$ . We can also explicit the terms  $X_p E$  and  $X_p F$  in  $\Phi_E$  so obtaining that

$$\Phi_E = \sum_{k \notin p(W)} \sum_{p \in \mathcal{D}} e_p(k)(X_p E(k) - X_p F(k)) + \sum_{p \in \mathcal{D}} e_p(p(W))(X_p E(p(W)) - X_p F(p(W)))$$

that is strictly less than zero.  $\square$

**Remark 8** *Let us notice that in the proof, additivity for  $F$  and just uniqueness for  $E$  are needed.*

If we consider the case where the error is larger than 1, we have instability even when the error is just equal to 2, if the number of lattice directions is larger than 2. More in detail, the instability follows from the result of [1, Theorem 1] because the sets constructed in the proof of [1] are actually  $\mathcal{D}$ -additive. Therefore we can restate it as follows:

**Proposition 9** (see [1]) *For any  $n$  and a set  $\mathcal{D}$  of  $m \geq 3$  directions there exist  $E$  and  $F$   $\mathcal{D}$ -additive such that  $\text{card}(E) = \text{card}(F) \geq n$ ,  $DX_{\mathcal{D}}(E, F) = 2$  and  $E \cap F = \emptyset$ .*

## 4 Stability for Convex Sets

In this section we study Problem 1 for convex sets from both an experimental and a theoretical point of view.

Any *convex set* is the intersection of a convex polygon and the digital plane  $\mathbb{Z}^2$ . The definition of convex set pass through that of convex polygon, and this can be used to determine results for the discrete case from the continuous case. In this way, convex sets are uniquely determined by their X-rays taken in a suitable set of directions [11], and this set of directions distinguishes convex bodies [13]. So in the “continuous” plane an analogous result holds, and additionally the reconstruction problem is stable [22]. Moreover we notice that there is a connection between additive sets and convex sets, since an euclidean ball is additive with respect to two orthogonal directions ([9]).

Experiments support the conjecture that convex sets are additive for a suitable set of directions, and indeed they accord with Proposition 7. The experimental results are better for a larger error thanks to the property of convexity.

In the second part, we conduct a theoretical study that confirms stability for convex sets.

### 4.1 Experimental results

In this section we experimentally study the stability of the reconstruction of convex sets via linear programming. Our experiments support the suspect that the results in the continuous have a correspondence in the “digital” plane.

Actually we consider in this section a class of lattice sets which is more general than the convex sets [5].

For each point  $M = (x_M, y_M) \in \mathbb{Z}^2$  the four quadrants around  $M$  are defined by the following formulas:

$$\begin{aligned} R_0(M) &= \{(x, y) \in \mathbb{Z}^2 / x \leq x_M \text{ and } y \leq y_M\}, \\ R_1(M) &= \{(x, y) \in \mathbb{Z}^2 / x \geq x_M \text{ and } y \leq y_M\}, \\ R_2(M) &= \{(x, y) \in \mathbb{Z}^2 / x \geq x_M \text{ and } y \geq y_M\}, \\ R_3(M) &= \{(x, y) \in \mathbb{Z}^2 / x \leq x_M \text{ and } y \geq y_M\}. \end{aligned}$$

**Definition 10** *A lattice set  $E$  is Q-convex if and only if for each  $M \notin E$  there exists  $i \in \{0, 1, 2, 3\}$  such that  $R_i(M) \cap E = \emptyset$ .*



An example of Q-convex set is given on the left part of Figure 2.

We have generated 184 Q-convex sets of semi-perimeter from 4 to 370 using an uniform generator ([4], inspired from [17]). Then their X-rays in the set of directions  $\mathcal{D} = \{x, y, 2x + y, -x + 2y\}$  have been computed. (These directions have been chosen because the X-rays along them uniquely determine the convex sets ([11]) and they contain the horizontal and vertical directions). We then used these X-rays and any error  $e \in \{0, 1, 2, 3\}$  as input data in the following linear-program:

Maximizing

$$\sum_{(i,j) \in E} (1 - v_{i,j}) + \sum_{(i,j) \notin E} v_{i,j} \quad (1)$$

such that

$$\sum_{p(i,j)=k} v_{i,j} = X_p E(k) + er_{p,k}^+ - er_{p,k}^- \quad (2)$$

$$\sum_k er_{p,k}^+ + er_{p,k}^- \leq e \quad (3)$$

$$0 \leq v_{i,j} \leq 1, er_{p,k}^+ \geq 0, er_{p,k}^- \geq 0 \quad (4)$$

We solved the linear program with the software soplex which implements the simplex algorithm ([23]). Notice that solving this problem with  $v_{i,j} \in \mathbb{Z}$  permits to exactly find the maximum of  $\text{card}(E \triangle F)$  where  $F$  describes *all* the lattice sets such that  $DX_{\mathcal{D}}(E, F) \leq e$ . Unfortunately, integer-linear-program is an NP-hard problem, and hence we solved the *relaxed* problem where the unknown variables can be fractional: this computation provides an upper bound to  $\text{card}(E \triangle F)$ . Figure 2 illustrates (on the right side-hand) a solution of the linear programming for  $\text{card}(E) = 200$  and  $e = 3$ . The different grey-scale colors of the squares correspond to different values of  $v_{i,j}$ .

The complete results are summarized in Figures 3 and 4. In Figure 3 the upper bound to  $\text{card}(E \triangle F)$  is divided by  $\text{card}(E)$ , so that each value gives an upper bound to the relative distance from a given set. Moreover the black squares show the values of the maximum of the quantity (1) when the constraints (2),(3) are replaced by  $X_p E(k) - 1 \leq \sum_{p(i,j)=k} v_{i,j} \leq X_p E(k) + 1$ : these values give an upper bound to  $\text{card}(E \triangle F)$  when  $DX'_{\mathcal{D}}(E, F) = \max_{p \in \mathcal{D}} \max_{k \in \mathbb{Z}} |X_p E(k) - X_p F(k)| = 1$ .

These experimental results bring the following comments:

- If  $DX_{\mathcal{D}}(E, F) = 0$ , then we always found a null relative distance. In other words, according to our experiments every Q-convex set is  $\mathcal{D}$ -additive. In fact this property was first conjectured by L. Thorens ([21]) (with additivity replaced by uniqueness), and can be seen as a variant of Conjecture 4.6 of [2] and Theorem 5.7 of [11]. We can set out the conjecture as follows:




Fig. 2. A Q-convex set  $E$  and the corresponding extremal values of  $v_{i,j}$  for  $e = 3$ . In this case we have  $\text{card}(E) = 200$  and  $\sum_{(i,j) \in E} (1 - v_{i,j}) + \sum_{(i,j) \in E^c} v_{i,j} = 33.7$ .




Fig. 3. An upper bound to  $\frac{\text{card}(E \Delta F)}{\text{card}(E)}$  for the Q-convex generated sets. (Only 40 % of the 184 generated sets have been represented for readability)

**Conjecture 11** *If  $\mathcal{D}$  is a set of directions which contains  $\{x, y\}$ , such that all the directions are not in the same quadrant and they uniquely determine the convex sets, then every Q-convex set is  $\mathcal{D}$ -additive.*

Notice that the property about the quadrants is necessary because there is a counter-example with  $\mathcal{D} = \{x, y, x + y, x + 5y\}$ .

- For a fixed error  $e$ , the relative distance looks to converge to zero as  $\text{card}(E)$

grows. If we divide by  $\sqrt{\text{card}(E)}$  instead of  $\text{card}(E)$ , this ratio seems to be bounded so that in *average*  $\text{card}(E \triangle F) = O(\sqrt{\text{card}(E)})$  according to our experiments (see Figure 4). It must be noticed that in the case  $e = 1$ , the maximum error for lattice sets is always 1 for the generated cases according to the result of Proposition 7. Since the theoretical result holds for additive sets, the experiments could be interpreted as a further evidence of the conjecture.

- If  $DX'_D(E, F) = 1$ , then the relative distance does not seem to converge to zero, but the computed values are only upper bounds, that is, we do not know if the fractional values mirror instability or they are just an artifact introduced by relaxing the integral constraints of the problem. In the former case, the reconstruction of convex sets would not be applied easily in the continuous world (as in medical imaging), because a rounding error of the measurements can always be of  $\pm 1$ .




Fig. 4. An upper bound to  $\frac{\text{card}(E \triangle F)}{\sqrt{\text{card}(E)}}$  for the 184 generated Q-convex sets

## 4.2 Theoretical results

In this section we first exploit a stability result for convex bodies [22] to deal with the corresponding problem for convex lattice sets and then we use this result to choose that it is possible to reconstruct convex bodies from X-rays, by the intermediate of lattice convex sets and Discrete Tomography.

#### 4.2.1 Preliminaries

A *convex body* is a compact convex subset of  $\mathbb{R}^2$  with non-empty interior. We denote the set of all the convex bodies by  $\mathcal{K}_*$ . The X-ray  $X_p U$  of the convex body  $U$  in direction  $p$  is the function giving the length of each chord of  $U$  parallel to  $p$ . More precisely  $X_p U(\alpha)$  is the length of the intersection of  $U$  with the line  $p = \alpha$ . The *Steiner symmetral*  $S_p(U)$  of  $U$  in direction  $p$  is the closure of the union of all open segments on lines parallel to  $p$  of the same length as  $X_p U$  centered about a fixed line orthogonal to  $p$ . So the Steiner symmetral  $S_p(U)$  and the X-ray  $X_p U$  contains exactly the same information.

**Definition 12** *A set of direction  $\mathcal{D}$  is a Gardner-McMullen set of directions if any convex body is characterized by all its X-rays in the directions of  $\mathcal{D}$ .*

We recall a result of [12, Proposition 6.1, Theorem 4.5]:

**Theorem 13 (Gardner-Gritzmann)** *A set  $\mathcal{D} = \{p_1, p_2, p_3, p_4\}$  of four lattice directions is a Gardner-McMullen set of directions if and only if the cross-ratio of the directions arranged in order of increasing angle with the positive  $x$ -axis is not in  $\{\frac{4}{3}, \frac{3}{2}, 2, 3, 4\}$ .*

**Example 14** *This theorem implies that the set  $\mathcal{D} = \{x, y, 2x + y, -x + 2y\}$  is a Gardner-Gritzmann set of directions.*

In the following we suppose that  $\mathcal{D} = \{p_1, p_2, p_3, p_4\}$  is a Gardner-McMullen set of four directions. So the mapping  $\mu : \mathcal{K}_* \mapsto (S_{p_1} U, S_{p_2} U, S_{p_3} U, S_{p_4} U)$  is injective.

Let  $\mathcal{K}_*$  be endowed with Nikodym's distance:

$$d_N(U, V) = m(U \Delta V),$$

where  $m(U)$  denotes the Lebesgue measure on  $\mathbb{R}^2$ . Now are we in place to state the stability result for convex bodies (see Theorem of [22, section 3.1]):

*if  $\mathcal{K}_*$  is endowed with the topology induced by the Nikodym's distance,  $\mu$  is continuous and continuously invertible from  $\mu(\mathcal{K}_*)$ .*

We shall reformulate this theorem. Consider the map  $\sigma_{\mathcal{D}} : U \mapsto (X_p U)_{p \in \mathcal{D}}$ ; if  $\mathcal{D}$  is a Gardner-McMullen set of direction, then  $\sigma_{\mathcal{D}}(U)$  is injective. Let  $\mathcal{X}_{\mathcal{D}}$  be the range of  $\sigma_{\mathcal{D}}$ . We endow  $\mathcal{X}_{\mathcal{D}}$  with the following distance:

$$d_X((f_p)_{p \in \mathcal{D}}, (g_p)_{p \in \mathcal{D}}) = \max_{p \in \mathcal{D}} \int_{-\infty}^{+\infty} \frac{|f_p(\alpha) - g_p(\alpha)|}{\sqrt{a_p^2 + b_p^2}} d\alpha,$$

where  $(f_p)_{p \in \mathcal{D}}, (g_p)_{p \in \mathcal{D}}$  are in  $\mathcal{X}_{\mathcal{D}}$ ,  $a_p$  and  $b_p$  are defined by  $p(x, y) = a_p x + b_p y$ . (Notice that each integral in the definition of this distance corresponds exactly

to the Nikodym distance if X-rays are considered as Steiner symmetral.) We also use the notation  $d_{\mathcal{X}_{\mathcal{D}}}(U, V) = d_X(\sigma_{\mathcal{D}}(U), \sigma_{\mathcal{D}}(V))$ .

The Theorem of [22, section 3.1] can be rewritten as follows:

**Theorem 15 (Volčič)** *Let  $\mathcal{D}$  be a Gardner-McMullen set of four lattice directions the inverse  $\sigma_{\mathcal{D}}^{-1}$  of the function  $\sigma_{\mathcal{D}}$  is a continuous function from  $\mathcal{X}_{\mathcal{D}}$  to  $\mathcal{K}_*$ .*

For any bounded set  $E \subset \mathbb{R}^2$  we define  $R_{\max}(E) = \max_{M \in E} \|M\|$  where  $\|\cdot\|$  is the euclidean norm. The set  $\mathcal{K}_{\varepsilon}^1 = \{U \in \mathcal{K}_* : R_{\max}(U) \leq 1 \text{ and } m(U) \geq \varepsilon\}$  is a compact subset of  $\mathcal{K}_*$ ; it follows that  $\sigma_{\mathcal{D}}(\mathcal{K}_{\varepsilon}^1)$  is a compact subset of  $\mathcal{X}_{\mathcal{D}}$  and so the function  $\sigma_{\mathcal{D}}^{-1}$  restricted to  $\sigma_{\mathcal{D}}(\mathcal{K}_{\varepsilon}^1)$  is uniformly continuous. So we can give a more precise formulation of the previous theorem:

**Corollary 16** *Let  $\mathcal{D}$  be a Gardner-McMullen set of four lattice directions. For any  $\varepsilon > 0$  there exists  $\eta > 0$  such that any  $U, V \in \mathcal{K}_{\varepsilon}^1$  satisfy:*

$$d_{\mathcal{X}_{\mathcal{D}}}(U, V) < \eta \implies d_N(U, V) < \varepsilon.$$

#### 4.2.2 A stability result for lattice convex sets

In this section  $\mathcal{D}$  is a Gardner-McMullen set of four lattice directions. Since a Gardner-McMullen set of lattice direction uniquely determines convex lattice sets [11], we use the result enunciated in Corollary 16 to get a stability result for convex lattice sets.

At first we need a lemma which is a direct consequence of Pick's theorem. We recall that a lattice polygon is a polygon whose vertexes are in  $\mathbb{Z}^2$ , and a simple polygon is a polygon whose edges have a non-empty intersection only if they are consecutive.

**Lemma 17** *If  $P \subset \mathbb{R}^2$  is simple lattice polygon, then  $\text{card}(P \cap \mathbb{Z}^2) \leq 2m(P) + 2$ .*

**PROOF.** By Pick's theorem [20,15] we have  $m(P) = i + \frac{b}{2} - 1$  where  $i$  is the number of lattice points which are in the interior of  $P$  and  $b$  is the number of lattice points which are in the border of  $P$ .

So  $\text{card}(P \cap \mathbb{Z}^2) = i + b \leq 2i + b = 2(m(P) + 1)$ .  $\square$

In the sequel each direction  $p$  of  $\mathcal{D}$  has the form  $p(x, y) = a_px + b_py$  with  $a_p, b_p$  integer.

**Proposition 18** *For any  $\varepsilon > 0$  and  $K > 1$ , there exists  $\eta > 0, M > 0$  such that any lattice convex non-segment sets  $E$  and  $F$  such that  $\frac{\text{card}(E)}{(R_{\max}(E))^2}, \frac{\text{card}(F)}{(R_{\max}(F))^2} \geq \varepsilon$ ,  $R_{\max}(E), R_{\max}(F) \geq M$ ,  $\frac{1}{K} \leq \frac{R_{\max}(E)}{R_{\max}(F)} \leq K$  satisfy:*

$$\frac{DX(E, F)}{(\max(R_{\max}(E), R_{\max}(F)))^2} < \eta \implies \frac{\text{card}(E \triangle F)}{(\max(R_{\max}(E), R_{\max}(F)))^2} < \varepsilon + \frac{17}{\max(R_{\max}(E), R_{\max}(F))}$$

**PROOF.** We define  $\varepsilon_c = \min(\frac{\varepsilon}{4K^2}, \frac{\varepsilon}{2})$ . Let  $\eta_c$  given by Corollary 16 applied to  $\varepsilon_c$ . We take  $M$  such that  $\frac{6}{M} \leq \frac{\eta_c}{2}$ ,  $\frac{1}{(KM)^2} \leq \varepsilon_c$  and  $M \geq 8$ . So we suppose that  $E$  and  $F$  are sets which satisfy the conditions of the propositions.

Let us consider sets  $E_c = \frac{1}{N}\text{conv}(E)$ ,  $F_c = \frac{1}{N}\text{conv}(F)$  and the number  $N = \max(R_{\max}(E), R_{\max}(F))$ .

The sets  $E_c$  and  $F_c$  are convex polygons of  $\mathbb{R}^2$ , and since they are not segments,  $E_c$  and  $F_c$  are convex bodies. Additionally, they are simple *lattice* polygons being their vertices in  $(\mathbb{Z}/N)^2$ . By Lemma 17 applied to  $P = NE_c$  we have  $m(E_c) \geq \frac{1}{N^2}(\frac{\text{card}(E)}{2} - 1)$ . So:

$$\begin{aligned} m(E_c) &\geq \frac{1}{N^2}(\frac{\text{card}(E)}{2} - 1) \\ &\geq \frac{\text{card}(E)}{2(KR_{\max}(E))^2} - \frac{1}{(KR_{\max}(E))^2} \\ &\geq \frac{\varepsilon}{2K^2} - \frac{1}{(KM)^2} \\ &\geq 2\varepsilon_c - \varepsilon_c = \varepsilon_c \end{aligned}$$

Similarly  $m(F_c) \geq \varepsilon_c$ . So  $E_c, F_c \in \mathcal{K}_\varepsilon^1$ .

Now we suppose that  $\frac{DX(E, F)}{N^2} < \eta$  with  $\eta = \frac{\eta_c}{4}$  and we estimate  $d_{\mathcal{X}_D}(E_c, F_c)$ . We have that:

$$\begin{aligned} \frac{X_p E(n) - 1}{N} &\leq X_p E_c(\frac{n}{N}) \leq \frac{X_p E(n) + 1}{N} \\ \frac{X_p F(n) - 1}{N} &\leq X_p F_c(\frac{n}{N}) \leq \frac{X_p F(n) + 1}{N} \end{aligned}$$

so that

$$|X_p E_c(\frac{n}{N}) - X_p F_c(\frac{n}{N})| \leq \frac{|X_p E(n) - X_p F(n)| + 2}{N}.$$

Since

$$|X_p E_c(\alpha) - X_p F_c(\alpha)| \leq \max(|X_p E_c(\frac{\lfloor \alpha N \rfloor}{N}) - X_p F_c(\frac{\lfloor \alpha N \rfloor}{N})|, |X_p E_c(\frac{\lceil \alpha N \rceil}{N}) - X_p F_c(\frac{\lceil \alpha N \rceil}{N})|)$$

we get

$$\begin{aligned} \int_{-\infty}^{+\infty} |X_p E_c(\alpha) - X_p F_c(\alpha)| d\alpha &\leq \frac{2}{N} \sum_{n=\lceil -N\sqrt{a_p^2+b_p^2} \rceil}^{\lfloor N\sqrt{a_p^2+b_p^2} \rfloor} \frac{|X_p E(n) - X_p F(n)| + 2}{N} \\ &= \frac{2}{N^2} \sum_{n=-\infty}^{+\infty} |X_p E(n) - X_p F(n)| + \frac{2(2N\sqrt{a_p^2+b_p^2} + 1)}{N^2}. \end{aligned}$$

Finally,  $a_p$  and  $b_p$  are integer and, so  $\sqrt{a_p^2 + b_p^2} \geq 1$ , and we conclude that

$$\begin{aligned} d_{\mathcal{X}_D}(E_c, F_c) &\leq \frac{2}{N^2} DX(E, F) + \frac{6}{N} \\ &\leq 2\eta + \frac{6}{M} \\ &\leq \frac{\eta_c}{2} + \frac{\eta_c}{2} = \eta_c. \end{aligned}$$

By Corollary 16 we have that  $d_N(E_c, F_c) \leq \varepsilon_c \leq \frac{\varepsilon}{2}$ .

The symmetric difference  $E_c \triangle F_c$  is the union of components of  $E_c \setminus F_c$  and of  $F_c \setminus E_c$ . Let  $C_j$  denotes the closure of the  $j$ th component and  $E_c \triangle F_c = \cup_{j=1}^k C_j$ . Each component  $C_j$  is a simple polygon  $(S_0 A_0 A_1 A_2 \dots A_l S_1 B_0 B_1 B_2 B_m)$  where  $A_0, \dots, A_l$  are consecutive vertexes of  $E_c$ ,  $B_0, \dots, B_m$  are consecutive vertexes of  $F_c$ , and  $S_0, S_1$  are intersection of an edge of  $E_c$  and an edge of  $F_c$ .

We consider the polygon  $C'_i$  union of the three following polygons:

- $\text{conv}((S_0 A_0 B_0) \cap (\mathbb{Z}/N)^2)$
- the polygon  $(A_0 A_1 \dots A_l B_0 B_1 \dots B_l)$
- $\text{conv}((S_1 A_l B_m) \cap (\mathbb{Z}/N)^2)$

This polygon is included in  $C_i$  and is a simple polygon whose vertexes are all in  $(\mathbb{Z}/N)^2$ , so by Lemma 17 we have  $\text{card}(C'_i \cap (\mathbb{Z}/N)^2) = \text{card}(C_i \cap (\mathbb{Z}/N)^2) \leq 2m(C'_i) + 2 \leq 2m(C_i) + 2$ .

So finally

$$\text{card}(E \triangle F) = \sum_{j=1}^k \text{card}(C_j \cap (\mathbb{Z}/N)^2) \leq \sum_{j=1}^k (2m(C_j) + 2) = 2d_N(E_c, F_c) + 2k.$$

The vertexes of  $E_c$  and  $F_c$  are in  $\mathbb{Z}/N^2 \cap [-1, 1]^2$  so the polygons  $E_c$  and  $F_c$  have less than  $2(2N + 1)$  vertexes. Each component  $C_i$  contains at least one

vertex of  $E_c$  or one vertex of  $F_c$  so  $k \leq 4(2N + 1)$ . Moreover  $d_N(E_c, F_c) \leq \frac{\varepsilon}{2}$  so  $\text{card}(E \triangle F) \leq \varepsilon N^2 + 8(2N + 1) = \varepsilon N^2 + 16N + 8$ . We have supposed that  $N \geq M \geq 8$  so finally  $\text{card}(E \triangle F) \leq \varepsilon N^2 + 17N$ .  $\square$

This upper bound overestimates the symmetric difference because we actually count also points of the border of  $E_c \cap F_c$  and the number of components is less than the number of vertexes of  $E_c$  and  $F_c$ , that in turn is less than  $4N$ .

#### 4.2.3 Reconstruction of a convex body from noisy discrete X-rays

In this section we always suppose that  $\mathcal{D}$  is a Gardner-McMullen set of directions. If  $F$  is a convex body then we know that it is completely determined by its continuous X-rays in  $\mathcal{D}$ . Anyway it does not give an algorithm to reconstruct  $F$  from its X-rays. The aim of this section is to use previous Proposition to show that it is possible, in theory, to reconstruct  $F$  by using Discrete Tomography. For this we fix an integer  $n$ , and we suppose that we have a lattice convex set  $E_n$  which could be seen as an approximation of  $F$  to the resolution  $\frac{1}{n}$ . But as we do not know  $F$ , the only assertions about  $E_n$  is the nearness of the discrete X-rays of  $E_n$  and the continuous X-rays of  $F$ . Proposition 20 will show that assertions which only consider the X-rays of  $E_n$  exist such that the set  $E_n$  converges, in a certain sense, to  $F$  when  $n$  tends to infinity.

If  $p = ax + by$  is a direction,  $\|p\|$  designs  $\sqrt{a^2 + b^2}$ . We start with an easy lemma which will be useful in the following:

**Lemma 19** *If  $E$  is any bounded subset of  $\mathbb{R}^2$  and  $p, p'$  are two directions then*

$$\begin{aligned} \max\left(\frac{|\alpha_1|}{\|p\|}, \frac{|\alpha_2|}{\|p\|}, \frac{|\alpha'_1|}{\|p'\|}, \frac{|\alpha'_2|}{\|p'\|}\right) &\leq R_{\max}(E) \\ &\leq \max(\|\langle \alpha_1, \alpha'_1 \rangle_{pp'}\|, \|\langle \alpha_1, \alpha'_2 \rangle_{pp'}\|, \|\langle \alpha_2, \alpha'_1 \rangle_{pp'}\|, \|\langle \alpha_2, \alpha'_2 \rangle_{pp'}\|) \end{aligned}$$

where  $\alpha_1 = \inf_{z \in E} p(z)$ ,  $\alpha_2 = \sup_{z \in E} p(z)$ ,  $\alpha'_1 = \inf_{z \in E} p'(z)$ ,  $\alpha'_2 = \sup_{z \in E} p'(z)$ .

**Proposition 20** *Let  $F$  be a convex body, and  $(E_n)_{n \in \mathbb{N}}$  a sequence of non-segment convex lattice sets such that for any  $p \in \mathcal{D}$  there hold:*

$$\frac{1}{n} \max\{k \in \mathbb{Z} : X_p E_n(k) \neq 0\} \xrightarrow{n \rightarrow \infty} \sup\{\alpha \in \mathbb{R} : X_p F(\alpha) \neq 0\} \quad (5)$$

$$\frac{1}{n} \min\{k \in \mathbb{Z} : X_p E_n(k) \neq 0\} \xrightarrow{n \rightarrow \infty} \inf\{\alpha \in \mathbb{R} : X_p F(\alpha) \neq 0\} \quad (6)$$

$$\frac{1}{n} \max_{p \in \mathcal{D}} \sum_{k \in \mathbb{Z}} \left| \frac{X_p E_n(k)}{n} - X_p F\left(\frac{k}{n}\right) \right| \xrightarrow{n \rightarrow \infty} 0 \quad (7)$$



then

$$\frac{1}{n^2} \text{card}(E_n \Delta (nF \cap \mathbb{Z}^2)) \xrightarrow{n \rightarrow \infty} 0.$$

**PROOF.** Let  $(F_n)_{n \in \mathbb{N}}$  be the sequence of convex lattice sets, defined by  $F_n = nF \cap \mathbb{Z}^2$ . To prove this proposition we are going to show that an integer  $N$  exists such that for  $n > N$  the sets  $E_n$  and  $F_n$  verify the conditions of Proposition 18. The thesis follows by applying the proposition.

At first we derive some conditions we need to our goal.

Since  $\frac{\text{card}(F_n)}{n^2} \xrightarrow{n \rightarrow \infty} m(F)$  and  $\frac{R_{\max}(F_n)}{n} \xrightarrow{n \rightarrow \infty} R_{\max}(F)$ , it follows that

$$\frac{\text{card}(F_n)}{(R_{\max}(F_n))^2} \xrightarrow{n \rightarrow \infty} \frac{m(F)}{(R_{\max}(F))^2} > 0. \quad (8)$$

We have

$$\frac{X_p F_n(k) - 1}{n} \leq X_p F\left(\frac{k}{n}\right) \leq \frac{X_p F_n(k) + 1}{n},$$

and, hence by condition (7)

$$\frac{1}{n^2} DX(E_n, F_n) \xrightarrow{n \rightarrow \infty} 0. \quad (9)$$

As a consequence of this and  $|\text{card}(E_n) - \text{card}(F_n)| \leq DX(E_n, F_n)$ , we obtain that  $\frac{\text{card}(E_n) - \text{card}(F_n)}{n^2} \xrightarrow{n \rightarrow \infty} 0$ .

We choose arbitrarily two directions  $p, p'$  of  $\mathcal{D}$ . Let  $\alpha_1 = \min_{z \in F} p(z)$ ,  $\alpha_2 = \max_{z \in F} p(z)$ ,  $\alpha'_1 = \min_{z \in F} p'(z)$ ,  $\alpha'_2 = \max_{z \in F} p'(z)$ . By Lemma 19 applied to  $E_n$ , the conditions (5),(6) and the continuity of the function  $(\alpha, \alpha') \rightarrow \langle \alpha, \alpha' \rangle_{pp'}$ , there exists an integer  $N_1$  such that for  $n > N_1$  we have:  $M_1 \leq \frac{R_{\max}(E_n)}{n} \leq M_2$  with  $M_1 = \frac{1}{2} \max(\frac{|\alpha_1|}{\|p\|}, \frac{|\alpha_2|}{\|p\|}, \frac{|\alpha'_1|}{\|p'\|}, \frac{|\alpha'_2|}{\|p'\|})$  and  $M_2 = 2 \max(\|\langle \alpha_1, \alpha'_1 \rangle_{pp'}\|, \|\langle \alpha_1, \alpha'_2 \rangle_{pp'}\|, \|\langle \alpha_2, \alpha'_1 \rangle_{pp'}\|, \|\langle \alpha_2, \alpha'_2 \rangle_{pp'}\|)$ . Thus, by this and the previous deduction, we get

$$\frac{\text{card}(E_n)}{(R_{\max}(E_n))^2} \geq \frac{\text{card}(E_n)}{(nM_2)^2} \xrightarrow{n \rightarrow \infty} \frac{m(F)}{(M_2)^2} > 0. \quad (10)$$

Moreover, an integer  $N_2 > N_1$  exists such that for  $n > N_2$  there holds:  $\frac{M_1}{2R_{\max}(F)} \leq \frac{R_{\max}(E_n)}{R_{\max}(F_n)} \leq \frac{2M_2}{R_{\max}(F)}$ .

Now we are going to use these properties to show that we can apply Proposition 18 to  $E_n$  and  $F_n$ . To prove the thesis, we have to find, for any  $\varepsilon > 0$ , an  $N$  such that  $\frac{1}{n^2} \text{card}(E_n \Delta (nF \cap \mathbb{Z}^2)) \leq \varepsilon$  for  $n > N$ . Let  $K = \max(\frac{2M_2}{R_{\max}(F)}, \frac{2R_{\max}(F)}{M_1})$

and  $\varepsilon' = \frac{\varepsilon}{2(2KR_{\max}(F))^2}$ . Without loss of generality let us suppose that  $0 < \varepsilon' < \frac{m(F)}{2(R_{\max}(F))^2}, \frac{m(F)}{2(M_2)^2}$ ,

- We have that  $\frac{1}{K} \leq \frac{R_{\max}(E_n)}{R_{\max}(F_n)} \leq K$  for  $n > N_2$ .
- By the conditions (8) and (10) there exists an integer  $N_3$  such that  $\frac{\text{card}(E_n)}{(R_{\max}(E_n))^2}, \frac{\text{card}(F_n)}{(R_{\max}(F_n))^2} \geq \min(\frac{m(F)}{2(R_{\max}(F))^2}, \frac{m(F)}{2(M_2)^2}) \geq \varepsilon'$ . Hence  $\frac{\text{card}(E_n)}{(R_{\max}(E_n))^2}, \frac{\text{card}(F_n)}{(R_{\max}(F_n))^2} \geq \varepsilon'$ .
- For any fixed  $M > 0$  there exists an integer  $N_3$  such that for any  $n > N_3$  we have  $R_{\max}(E_n) \geq M$  and  $R_{\max}(F_n) \geq M$ .
- For any fixed  $\eta > 0$ , by property (9), an integer  $N_4$  exists such that  $\frac{1}{n^2}DX(E_n, F_n) < \eta(\frac{R_{\max}(F)}{2K})^2$  for any  $n > N_4$ .
- There exists an integer  $N_5$  such that  $\frac{R_{\max}(F_n)}{n} \geq \frac{R_{\max}(F)}{2}$  for  $n > N_5$ .

So, for any  $n > N = \max(N_2, N_3, N_4, N_5)$ , the sets  $E_n$  and  $F_n$  satisfy the conditions of Proposition 18 (with  $\varepsilon'$  instead of  $\varepsilon$  and  $\eta$  and  $M$  chosen as in the proposition). Therefore we have:

$$\begin{aligned} \frac{DX(E_n, F_n)}{(\max(R_{\max}(E_n), R_{\max}(F_n)))^2} < \eta \implies \\ \frac{\text{card}(E_n \Delta F_n)}{(\max(R_{\max}(E_n), R_{\max}(F_n)))^2} < \varepsilon' + \frac{17}{\max(R_{\max}(E_n), R_{\max}(F_n))}. \end{aligned}$$

We have  $\max(R_{\max}(E_n), R_{\max}(F_n)) \geq \frac{R_{\max}(F_n)}{K} \geq \frac{nR_{\max}(F)}{2K}$  and  $\max(R_{\max}(E_n), R_{\max}(F_n)) \leq KR_{\max}(F_n) \leq 2KnR_{\max}(F)$ .

So  $\frac{DX(E_n, F_n)}{(\max(R_{\max}(E_n), R_{\max}(F_n)))^2} \leq \frac{n^2\eta(\frac{R_{\max}(F)}{2K})^2}{(\frac{nR_{\max}(F)}{2K})^2} = \eta$ , then

$$\begin{aligned} \text{card}(E_n \Delta F_n) &\leq \varepsilon'(\max(R_{\max}(E_n), R_{\max}(F_n)))^2 + 17\max(R_{\max}(E_n), R_{\max}(F_n)) \\ &\leq \varepsilon'(2KnR_{\max}(F))^2 + 17(2KnR_{\max}(F)) \\ &\leq n^2\frac{\varepsilon}{2} + 34KR_{\max}(F)n \\ &\leq n^2\frac{\varepsilon}{2} + n^2\frac{\varepsilon}{2} \quad \text{for } n > N_6 = \frac{68KR_{\max}(F)}{\varepsilon} \\ &= \varepsilon n^2 \end{aligned}$$

□

**Remark 21** *This last proposition gives a positive result for the reconstruction of a convex body from X-rays. Nevertheless it does not really give a concrete algorithm to reconstruct this convex body for at least two reasons:*

- *the result is not quantitative: the difference  $\text{card}(E_n \Delta \text{conv}(nF \cap \mathbb{Z}^2))$  is not bounded by a precise function on the X-rays errors.*

- *when the resolution is fixed, we have to reconstruct a lattice convex set from approximative X-rays. Algorithms are known in the exact case ([5]) or for more general classes than lattice convex sets ([3,6]) but not exactly in this case.*

## Acknowledgements

Acknowledgments are due to Mohamed Tajine for his questions about convergence, partially answered by Proposition 20.

## References

- [1] A. Alpers, P. Grizmann, and L. Thorens. Stability and instability in discrete tomography. In *Dagstuhl Seminar: Digital and Image Geometry 2000*, volume 2243 of *Lecture Notes in Comp. Sci.*, pages 175–186. Springer, 2001.
- [2] E. Barcucci, A. Del Lungo, M. Nivat, and R. Pinzani. X-rays characterizing some classes of discrete sets. *Linear Algebra Appl.*, 339:3–21, 2001.
- [3] Y. Boufkhad, O. Dubois, and M. Nivat. Reconstructing (h,v)-convex 2-dimensional patterns of objects from approximate horizontal and vertical projections. *Theoret. Comput. Sci.*, 290(3):1647–1664, 2003.
- [4] S. Brunetti and A. Daurat. Generating Q-convex sets. in preparation.
- [5] S. Brunetti and A. Daurat. An algorithm reconstructing convex lattice sets. *Theoret. Comput. Sci.*, 304(1-3):35–57, 2003.
- [6] S. Brunetti, A. Daurat, and A. Del Lungo. Approximate X-rays reconstruction of special lattice sets. *Pure Math. Appl.*, 11(3):409–425, 2000.
- [7] A. Daurat, A. Del Lungo, and M. Nivat. Median points of discrete sets according to a linear distance. *Disc. and Comp. Geom.*, 23:465–483, 2000.
- [8] A. Del Lungo and M. Nivat. Reconstruction of connected sets from two projections. In *Discrete tomography*, Appl. Numer. Harmon. Anal., pages 163–188. Birkhäuser Boston, 1999.
- [9] P. C. Fishburn, J. C. Lagarias, J. A. Reeds, and L. A. Shepp. Sets uniquely determined by projections on axes. I. Continuous case. *SIAM J. Appl. Math.*, 50(1):288–306, 1990.
- [10] P. C. Fishburn and L. A. Shepp. Sets of uniqueness and additivity in integer lattices. In *Discrete tomography*, Appl. Numer. Harmon. Anal., pages 35–58. Birkhäuser, 1999.

- [11] R. J. Gardner and P. Gritzmann. Discrete tomography: determination of finite sets by X-rays. *Trans. Amer. Math. Soc.*, 349(6):2271–2295, 1997.
- [12] R. J. Gardner and P. Gritzmann. Discrete tomography: Determination of finite sets by X-rays. *Trans. Am. Math. Soc.*, 349(6):2271–2295, 1997.
- [13] R. J. Gardner and P. McMullen. On Hammer’s X-ray problem. *J. Lond. Math. Soc., II. Ser.*, 21:171–175, 1980.
- [14] P. Gritzmann, S. de Vries, and M. Wiegmann. Approximating binary images from discrete X-rays. *SIAM J. Optim.*, 11(2):522–546, 2000.
- [15] B. Grünbaum and G. C. Shephard. Pick’s theorem. *Amer. Math. Monthly*, 100(2):150–161, 1993.
- [16] G. T. Herman and A. Kuba, editors. *Discrete tomography*. Appl. Numer. Harmon. Anal. Birkhäuser, 1999.
- [17] W. Hochstättler, M. Loeb, and C. Moll. Generating convex polyominoes at random. *Discrete Math.*, 153(1-3):165–176, 1996.
- [18] A. Kuba and G. T. Herman. Discrete tomography: a historical overview. In *Discrete tomography*, Appl. Numer. Harmon. Anal., pages 3–34. Birkhäuser Boston, Boston, MA, 1999.
- [19] D. Kölzow, A. Kuba, and A. Volčič. An algorithm for reconstructing convex bodies from their projections. *Disc. and Comp. Geom.*, 4:205–237, 1989.
- [20] G. A. Pick. Geometrisches zur zahlenlehre. *Lotos (Prag)*, 19:311–319, 1899.
- [21] L. Thorens. Personal communication, 2000.
- [22] A. Volčič. Well-posedness of the Gardner-McMullen reconstruction problem. In *Measure theory, Oberwolfach 1983 (Oberwolfach, 1983)*, volume 1089 of *Lecture Notes in Math.*, pages 199–210. Springer, 1984.
- [23] R. Wunderling. *Paralleler und Objektorientierter Simplex-Algorithmus*. PhD thesis, ZIB technical report TR 96-09, 1996.