



**HAL**  
open science

## Evaluation of long-term tropospheric NO<sub>2</sub> data obtained by GOME over East Asia in 1996-2002

H. Irie, K. Sudo, H. Akimoto, A. Richter, J.-P. Burrows, T. Wagner, M. Wenig, S. Beirle, Y. Kondo, V.-P. Sinyakov, et al.

### ► To cite this version:

H. Irie, K. Sudo, H. Akimoto, A. Richter, J.-P. Burrows, et al.. Evaluation of long-term tropospheric NO<sub>2</sub> data obtained by GOME over East Asia in 1996-2002. *Geophysical Research Letters*, 2005, 32, pp.L11810. 10.1029/2005GL022770 . hal-00068749

**HAL Id: hal-00068749**

**<https://hal.science/hal-00068749>**

Submitted on 21 Jan 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Evaluation of long-term tropospheric NO<sub>2</sub> data obtained by GOME over East Asia in 1996–2002

H. Irie,<sup>1</sup> K. Sudo,<sup>1</sup> H. Akimoto,<sup>1</sup> A. Richter,<sup>2</sup> J. P. Burrows,<sup>2</sup> T. Wagner,<sup>3</sup> M. Wenig,<sup>4</sup> S. Beirle,<sup>3</sup> Y. Kondo,<sup>5</sup> V. P. Sinyakov,<sup>6</sup> and F. Goutail<sup>7</sup>

Received 21 February 2005; revised 14 April 2005; accepted 16 May 2005; published 14 June 2005.

[1] Long-term tropospheric nitrogen dioxide (NO<sub>2</sub>) column data obtained by the Global Ozone Monitoring Experiment (GOME) (*G*-NO<sub>2</sub>) are evaluated to confirm the trends found in tropospheric NO<sub>2</sub> abundances over East Asia between 1996 and 2002. For three locations in Central and East Asia, the *G*-NO<sub>2</sub> values are compared with tropospheric columns estimated from coincident observations of total NO<sub>2</sub> by ground-based UV/visible spectrometers and stratospheric NO<sub>2</sub> by satellite solar occultation sensors (*E*-NO<sub>2</sub>). The comparisons show a slight linear drift in *G*-NO<sub>2</sub> data from 1996 to 2002. However, it is much smaller than the standard deviation of the differences between *G*-NO<sub>2</sub> and *E*-NO<sub>2</sub> and much smaller than the increasing trends in NO<sub>2</sub> seen by GOME over the industrial areas of China, demonstrating the validity of the trends estimated using the GOME data. **Citation:** Irie, H., et al. (2005), Evaluation of long-term tropospheric NO<sub>2</sub> data obtained by GOME over East Asia in 1996–2002, *Geophys. Res. Lett.*, 32, L11810, doi:10.1029/2005GL022770.

### 1. Introduction

[2] It is well known that nitrogen oxides (NO<sub>x</sub> = NO + NO<sub>2</sub>) play a crucial role in the Earth's atmosphere. As a result of rapid economic growth in East Asian countries, an increase in the anthropogenic emissions of NO<sub>x</sub> from these countries is thought to be occurring, due mainly to rising consumption of fossil fuels [Akimoto, 2003]. While tropospheric NO<sub>x</sub> abundance should respond to the changes in emission from adjacent regions, its temporal evolution over East Asia has not yet been quantified due to the lack of continuous long-term observations by a single instrument. The Global Ozone Monitoring Experiment (GOME) onboard the European Remote Sensing-2 (ERS-2) satellite measured tropospheric NO<sub>2</sub> column abundances over East Asia from July 1995 until June 2003 [Leue et al., 2001;

Richter and Burrows, 2002; Martin et al., 2002, 2003; Beirle et al., 2004; A. Richter et al., Significant increase in nitrogen dioxide levels over China observed from space, submitted to *Nature*, 2005, hereinafter referred to as Richter et al., submitted manuscript, 2005]. Recently, Richter et al. (submitted manuscript, 2005) have combined the GOME tropospheric NO<sub>2</sub> column data (*G*-NO<sub>2</sub>) with those of the Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY) to show a highly significant increase in NO<sub>2</sub> of ~50% over the industrial areas of China (30°–40°N and 110°–123°E) between 1996 and 2004. In Europe from 1996 to 2001, *G*-NO<sub>2</sub> has shown a reduction consistent with the pronounced reduction of NO<sub>x</sub> emissions (Richter et al., submitted manuscript, 2005). *G*-NO<sub>2</sub> has been compared to independent measurements over Europe and the U.S. [e.g., Heland et al., 2002; Petritoli et al., 2004; Martin et al., 2004], and good agreement was found. For East Asia, however, despite a large uncertainty in emission estimates, no validation comparison has been made so far. In the present study, the long-term *G*-NO<sub>2</sub> data obtained in Central and East Asia are evaluated by comparing monthly-mean *G*-NO<sub>2</sub> tropospheric column data with those estimated from long-term observations by ground-based UV/vis. zenith-sky spectrometers and satellite-borne solar occultation sensors (*E*-NO<sub>2</sub>). The comparisons support the results of Richter et al. (submitted manuscript, 2005) indicating a significant increase in NO<sub>2</sub> over the industrial areas of China between 1996 and 2002.

### 2. GOME Tropospheric NO<sub>2</sub> Column (*G*-NO<sub>2</sub>)

[3] The GOME instrument, a nadir-viewing UV/vis. spectrometer, was launched aboard the ERS-2 satellite in April 1995 [Burrows et al., 1999]. GOME observations with a spatial resolution of 40 km (in latitude) × 320 km (in longitude) are made three times across the flight track using a scanning mirror, achieving nearly global coverage within 3 days. The satellite crosses the equator at 10:30 am local time in the descending node, so that GOME observations at northern mid-latitudes occur around 11:00 am. For the *G*-NO<sub>2</sub> tropospheric column data used in this study, the retrieval technique was described in detail by Richter and Burrows [2002] and recent improvements were given by Richter et al. (submitted manuscript, 2005). To minimize the shielding effects of clouds on the retrieval, we used GOME data with a cloud fraction less than 0.2, based on a cloud retrieval algorithm (FRESCO) [Koelemeijer et al., 2001]. It has been shown that *G*-NO<sub>2</sub> tropospheric column data was tightly correlated with surface NO<sub>2</sub> concentration in highly-polluted regions (tropospheric NO<sub>2</sub> reaching 200 × 10<sup>14</sup> cm<sup>-2</sup>) in Italy throughout the year 2000 under optimized comparison conditions (well-mixed boundary layer

<sup>1</sup>Frontier Research Center for Global Change, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan.

<sup>2</sup>Institute of Environmental Physics, University of Bremen, Bremen, Germany.


<sup>3</sup>Institut für Umweltphysik, Heidelberg University, Heidelberg, Germany.

<sup>4</sup>NASA Goddard Space Flight Center, Atmospheric Chemistry and Dynamics Branch, Greenbelt, Maryland, USA.

<sup>5</sup>Research Center for Advanced Science and Technology, University of Tokyo, Tokyo, Japan.

<sup>6</sup>Kyrgyz National University, Bishkek, Kyrgyzstan.

<sup>7</sup>Service d'Aéronomie du CNRS, Verrières le Buisson, France.



**Figure 1.** Time series of NO<sub>2</sub>(tot) (black) and NO<sub>2</sub>(str) (red) values for Issyk-Kul at sunrise (SR). The SAGE II- and HALOE-based NO<sub>2</sub>(str) values have been unified for each month.

and low cloud coverage) [Petritoli *et al.*, 2004], indicating the sensitivity of the GOME observations to NO<sub>2</sub> near the surface. The present study uses monthly-averaged  $G\text{-NO}_2$  data derived on a  $0.5^\circ$  (in latitude)  $\times$   $0.5^\circ$  (in longitude) grid. We focus on the years 1996–2002, when nearly complete seasonal cycles were observed by GOME. Uncertainty in  $G\text{-NO}_2$  arises primarily due to errors in the a-priori NO<sub>2</sub> profiles used and the tropospheric air mass factor calculations. Uncertainty in a single  $G\text{-NO}_2$  value was estimated to be  $\sim 15 \times 10^{14} \text{ cm}^{-2}$  [Richter and Burrows, 2002] or 35–60% under highly-polluted conditions [Boersma *et al.*, 2004], suggesting that there is much smaller uncertainty in the monthly-averaged  $G\text{-NO}_2$  data.

### 3. Estimate of Tropospheric NO<sub>2</sub> Column ( $E\text{-NO}_2$ )

[4] To estimate  $E\text{-NO}_2$  values for the period 1996–2002, we combine observations by ground-based UV/vis. zenith-sky spectrometers and satellite-borne solar occultation sensors (Stratospheric Aerosol and Gas Experiment II (SAGE II) and Halogen Occultation Experiment (HALOE)) with the Chemical Atmospheric general circulation model for Study of Atmospheric Environment and Radiative forcing (CHASER) [Sudo *et al.*, 2002].  $E\text{-NO}_2$  is defined as:

$$E\text{-NO}_2 = (\text{NO}_2(\text{tot}) - \text{NO}_2(\text{str})) \times F,$$


where NO<sub>2</sub>(tot) is the total (troposphere + stratosphere) NO<sub>2</sub> vertical column derived from a UV/vis. spectrometer at a local time of sunrise (SR) or sunset (SS), NO<sub>2</sub>(str) is the stratospheric NO<sub>2</sub> vertical column derived from SAGE II (ver. 6.2) and HALOE (ver. 19) observations at SR or SS, and  $F$  is a diurnal correction factor defined as the ratio of the modeled tropospheric NO<sub>2</sub> column at the time of the GOME measurements to that at SR or SS.

[5] For estimating  $E\text{-NO}_2$  values, we focus on three sites, (1) Moshiri (44.4°N, 142.3°E, 200 m asl), Japan, (2) Issyk-Kul (42.6°N, 77.0°E, 1650 m asl), Kyrgyzstan, and (3) Zhigansk (66.8°N, 123.4°E, 50 m asl), Russia, where UV/vis. spectrometer observations at SR and SS were made as part of the Network for the Detection of Stratospheric Change (NDSC) during the 1996–2002 period. These sites are located in or closer to East Asia than other UV/vis. spectrometer observation sites available in the NDSC database for the period. We used the differential optical absorption spectroscopy (DOAS) technique and air mass factor for ground-based observations (AMF) to derive NO<sub>2</sub>(tot) at SR and SS from the spectra measured. Constant AMF values of 17.0, 16.5, and 15.86 over 1996–2002 were used for Moshiri, Issyk-Kul, and Zhigansk, respectively.

These AMF values were taken from the NDSC database. It should be noted that the use of constant AMF values may contribute to the uncertainty in NO<sub>2</sub>(tot), because AMF can vary depending on season. It should also be noted that an additional uncertainty in NO<sub>2</sub>(tot) may arise depending on season, because the sensitivity of the observations at SR and SS to the troposphere is much less than that to the stratosphere and varies according to the vertical profile of NO<sub>2</sub>. Moreover, the retrievals of NO<sub>2</sub>(tot) for the three sites have not been performed in a fully consistent manner, including the same NO<sub>2</sub> cross sections and a radiative transfer model. The effect of these uncertainties on the  $G\text{-NO}_2/E\text{-NO}_2$  comparison is discussed in the results and discussion section below.

[6] We next selected stratospheric NO<sub>2</sub> vertical profiles observed by SAGE II/HALOE at SR or SS within  $2^\circ$  latitude, 500 km distance, and 1 day of the ground-based UV/vis. spectrometer observations. For every coincident stratospheric observation, the NO<sub>2</sub>(str) value and its uncertainty were calculated by integrating the vertical profiles of observed NO<sub>2</sub> concentrations and their errors, respectively, for altitudes between 18 and 50 km. For both SAGE II and HALOE data, integrations including altitudes below 18 km sometimes led to NO<sub>2</sub>(str) values that were larger than NO<sub>2</sub>(tot), especially for Zhigansk, which is characterized by a lower pollution level than Moshiri and Issyk-Kul. At lower altitudes, the relative errors in the SAGE II and HALOE NO<sub>2</sub> concentration data increase and exceed 100% below 18 and 14 km, respectively, even in the summer when stratospheric NO<sub>2</sub> abundances are largest. This suggests that the discrepancy could be accounted for by the SAGE II and HALOE NO<sub>2</sub> concentration profile errors. The summertime NO<sub>2</sub> amounts between the tropopause and 18 km were calculated assuming that the NO<sub>2</sub> concentration decreased monotonically from 18 km to the tropopause, where the concentration was set to zero. The calculated NO<sub>2</sub> amounts ( $\sim 4$ ,  $\sim 4$ , and  $\sim 10 \times 10^{14} \text{ cm}^{-2}$ , for Moshiri, Issyk-Kul and Zhigansk, respectively) are considered as the maximum uncertainty in NO<sub>2</sub>(str) due to exclusion of NO<sub>2</sub> between the tropopause and 18 km.

[7] For each ground-based measurement site at SR and SS, the above calculations give two sets of NO<sub>2</sub>(str) stratospheric column values, one from SAGE II and one from HALOE. To check consistency between these two sets, we used the criteria of a 2-hour difference and a 500-km distance between the SAGE II and HALOE observations to identify coincident SAGE II- and HALOE-based column data within  $2^\circ$  latitude of the ground-based measurement locations. Comparisons between the two sets of NO<sub>2</sub>(str) columns showed that the median values of the differences (SAGE II minus HALOE) ( $\pm 67\%$  range) were  $0.0 (+3.3/-2.7) \times 10^{14}$  and  $6.6 (+2.2/-5.0) \times 10^{14} \text{ cm}^{-2}$  for local SR and SS, respectively. For local SS, the median of the relative differences was 20% (+13%/–17%). Since the NO<sub>2</sub> data obtained by SAGE II at SS could be biased [Bracher *et al.*, 2005], the SAGE II NO<sub>2</sub>(str) stratospheric column values at SS were adjusted to agree with the HALOE values at SS. The tropospheric NO<sub>2</sub> columns at SR or SS were then estimated by subtracting the SAGE II- or HALOE-based NO<sub>2</sub>(str) values from the corresponding NO<sub>2</sub>(tot) values. For example, the estimated NO<sub>2</sub>(str) and NO<sub>2</sub>(tot) values for Issyk-Kul are shown in Figure 1. In most of the cases over



**Figure 2.** Time series of  $G\text{-NO}_2$  (red) and  $E\text{-NO}_2$  (black) values for (a) Moshiri, (b) Issyk-Kul, and (c) Zhigansk. For each site, the mean difference ( $G\text{-NO}_2$  minus  $E\text{-NO}_2$ ) ( $\pm$ standard deviation) is also shown. (d) The differences between  $G\text{-NO}_2$  and  $E\text{-NO}_2$  are all shown. The line is the linear least-square fit to the data.

the three sites,  $\text{NO}_2(\text{tot})$  is greater than  $\text{NO}_2(\text{str})$ , indicating some contribution of tropospheric  $\text{NO}_2$  to  $\text{NO}_2(\text{tot})$ .

[8] To derive  $F$  values, we used a global tropospheric chemistry model, CHASER [Sudo *et al.*, 2002], with 32 vertical layers (surface to  $\sim 5$  hPa) and T42 horizontal resolution ( $\sim 2.8^\circ$  latitude  $\times$   $\sim 2.8^\circ$  longitude). CHASER includes 53 chemical species and 139 reactions, including gas, liquid, and heterogeneous reactions, and takes into account  $\text{O}_x\text{-NO}_x\text{-HO}_x\text{-CH}_4\text{-CO}$  chemistry and the oxidation of non-methane hydrocarbons. For each of the ground-based observation sites used in this study, we calculated daily  $F$  values ( $\sim 0.5\text{--}0.6$  on average) by dividing the modeled tropospheric  $\text{NO}_2$  column at the times of the GOME measurements by that at SR or SS. One- $\sigma$  standard deviations (s.d.) of the monthly-averaged  $F$  values ( $\sim 13\text{--}24\%$ ) are assumed to represent the uncertainty in the daily  $F$  value. After multiplying the  $F$  values with the above-estimated tropospheric  $\text{NO}_2$  column abundances for both SR and SS, the  $E\text{-NO}_2$  values were estimated and averaged for each month, yielding monthly-mean  $E\text{-NO}_2$  values for each of the ground-based observation sites.


#### 4. Results and Discussion

[9] The comparisons between the monthly-mean  $G\text{-NO}_2$  and  $E\text{-NO}_2$  tropospheric column amounts over Moshiri, Issyk-Kul, and Zhigansk are shown in Figures 2a–2c, respectively. Error bars for the  $G\text{-NO}_2$  values represent  $1\sigma$  s.d. of the monthly-mean  $G\text{-NO}_2$  values over a  $2^\circ$  (in latitude)  $\times$   $2^\circ$  (in longitude) region centered on the ground-based measurement sites. Error bars for  $E\text{-NO}_2$  represent the uncertainties in  $\text{NO}_2(\text{str})$  and  $F$ , as described above. As shown in the figures, most of the  $G\text{-NO}_2$  values agree with the  $E\text{-NO}_2$  values to within their combined error ranges. The differences are nearly independent of season, suggesting that the uncertainty in  $E\text{-NO}_2$ , due to the use of

constant AMF values and the seasonal change in sensitivity of the UV/vis. spectrometer observations to the troposphere, is smaller than the errors shown in the figures. Over 1996–2002, the mean differences ( $G\text{-NO}_2$  minus  $E\text{-NO}_2$ ) ( $\pm 1\sigma$  s.d.) were  $-2 (\pm 6) \times 10^{14}$ ,  $1 (\pm 6) \times 10^{14}$ , and  $4 (\pm 4) \times 10^{14}$   $\text{cm}^{-2}$ , for Moshiri, Issyk-Kul, and Zhigansk, respectively. Thus, since no significant difference in the  $G\text{-NO}_2/E\text{-NO}_2$  comparison results was found among the three locations, we next averaged all the differences between  $G\text{-NO}_2$  and  $E\text{-NO}_2$  values for all the locations. The mean difference was estimated to be  $0.6 \times 10^{14}$   $\text{cm}^{-2}$ , with a  $1\sigma$  s.d. of  $6 \times 10^{14}$   $\text{cm}^{-2}$ , suggesting that a  $G\text{-NO}_2$  variation exceeding  $6 \times 10^{14}$   $\text{cm}^{-2}$  can be interpreted as being statistically meaningful over East Asia. We note that the values of  $E\text{-NO}_2$  could be affected systematically by the errors in AMF and the weak sensitivity of UV/vis. spectrometer observations to the troposphere. However, the result may suggest that the systematic bias in the retrieved  $G\text{-NO}_2$  data is less than  $6 \times 10^{14}$   $\text{cm}^{-2}$ , if one considers previous validation studies made by Heland *et al.* [2002] and Martin *et al.* [2004]. They estimated the differences between  $G\text{-NO}_2$  and the tropospheric  $\text{NO}_2$  columns derived from aircraft observations to be less than  $7 \times 10^{14}$   $\text{cm}^{-2}$  over Austria in spring of 2001 [Heland *et al.*, 2002] and less than  $6 \times 10^{14}$   $\text{cm}^{-2}$  on average over eastern Texas and the southeast U.S. in the summers of 1999 and 2000 [Martin *et al.*, 2004].

[10] These previous comparisons have been made under polluted conditions (tropospheric  $\text{NO}_2$  columns reaching  $60 \times 10^{14}$   $\text{cm}^{-2}$ ), whereas relatively clean sites have been used in the present study, due to the lack of ground-based UV/vis. spectrometer observation at polluted sites in East Asia. In addition, the UV/vis. spectrometer observations in East Asia were available only at SR and SS. Observations at SR and SS are generally less sensitive to tropospheric  $\text{NO}_2$  than those in daytime. Thus, a ground-based UV/vis. spectrometer observation in polluted regions in daytime is highly desirable over East Asia to perform precise validation of future observations by nadir-viewing satellite instruments.

[11] The magnitude of the additional errors in  $E\text{-NO}_2$  depends mainly on season and is almost constant for the same seasons. Therefore, we can assess a possible long-term drift in the  $G\text{-NO}_2$  data over years 1996–2002 by



**Figure 3.** Time series of the mean  $G\text{-NO}_2$  values in industrial areas of China ( $30^\circ\text{--}40^\circ\text{N}$  and  $110^\circ\text{--}123^\circ\text{E}$ ) for winter (December–February) (blue), spring (March–May) (green), summer (June–August) (red), and fall (September–November) (black). Error bars represent standard deviations of  $G\text{-NO}_2$  over the region. The lines are the linear least-square fits to the data.

investigating the year-to-year variations in the differences between  $G\text{-NO}_2$  and  $E\text{-NO}_2$  for each season. Since the differences are nearly independent of season as described above, all the differences are plotted in Figure 2d. We find that the differences tend to decrease, at a rate of  $\sim 0.7 \times 10^{14} \text{ cm}^{-2} \text{ year}^{-1}$  on average (Figure 2d). However, this trend is much smaller than the  $1\sigma$  s.d. of all the differences estimated above and is negative, showing a counter trend towards the reported increasing trend in the industrial areas of China ( $30^\circ\text{--}40^\circ\text{N}$  and  $110^\circ\text{--}123^\circ\text{E}$ ) (Richter et al., submitted manuscript, 2005). Since insignificant, negative trends were seen in the time series of the  $G\text{-NO}_2/E\text{-NO}_2$  differences at each site (not shown), the result is not affected by the different retrieval procedures employed at each site.

[12] We take into account the drift in  $G\text{-NO}_2$  and estimate the linear trend in tropospheric NO<sub>2</sub> amounts over the industrial areas of China for each season (Figure 3). The most significant increasing trend in NO<sub>2</sub> occurred in wintertime ( $\sim 9.0 \times 10^{14} \text{ cm}^{-2} \text{ year}^{-1}$ ) with a correlation coefficient ( $R^2$ ) of 0.93. As the observed trends are much greater than the estimated drift, these results are very similar to those of Richter et al. (submitted manuscript, 2005) and confirm their trend analysis. Accounting for the drift estimated above, the mean relative trends from 1996 to 2002 are estimated to be 8, 6, 8, and 7%  $\text{year}^{-1}$  for winter, spring, summer, and fall, respectively. Thus, a continuous increase in tropospheric NO<sub>2</sub> abundances likely occurred in the industrial regions of China between 1996 and 2002, at a rate of  $7 \pm 1\% \text{ year}^{-1}$ . This trend may be greater than that over the whole country, but is likely to represent the qualitative tendency over China. This additional information from GOME will be useful to reduce uncertainty in bottom-up emission estimates for China.

## 5. Conclusions

[13] To evaluate the long-term tropospheric NO<sub>2</sub> column data obtained by GOME ( $G\text{-NO}_2$ ) in East Asia, we used tropospheric NO<sub>2</sub> abundances estimated based on observations by ground-based UV/vis. spectrometer and satellite-borne solar occultation sensors (SAGE II and HALOE) ( $E\text{-NO}_2$ ) for three locations in Central and East Asia (Moshiri in Japan, Issyk-Kul in Kyrgyzstan, and Zhigansk in Russia). The mean differences between  $G\text{-NO}_2$  and  $E\text{-NO}_2$  were  $(0.6 \pm 6.0) \times 10^{14} \text{ cm}^{-2}$ , suggesting that there is no significant bias in the  $G\text{-NO}_2$  data in East Asia. For making more precise comparisons, however, ground-based observations in polluted regions during daytime would be more suitable than those used in this study. The comparisons between  $G\text{-NO}_2$  and  $E\text{-NO}_2$  show that a slight drift in  $G\text{-NO}_2$  data occurred, at an average rate of  $-0.7 \times 10^{14} \text{ cm}^{-2} \text{ year}^{-1}$  from 1996 to 2002. However, this is much smaller than the observed increases in NO<sub>2</sub> over the industrial areas of China ( $30^\circ\text{--}40^\circ\text{N}$  and  $110^\circ\text{--}123^\circ\text{E}$ ), confirming the trend estimates made here and by Richter et al. (submitted manuscript, 2005). This multi-year  $G\text{-NO}_2$  data showing a large increasing trend of  $7 \pm 1\% \text{ year}^{-1}$  over the region will be useful for reducing the uncertainty in bottom-up emission estimates for China.

[14] **Acknowledgments.** The authors thank ESA for launching the ERS-2 satellite. We appreciate helpful discussions with O. Wild, M. Naja, and E. Bucsele. HI is grateful to the satellite groups in Bremen and Heidelberg Universities for their excellent retrievals. The ground-based UV/vis. observation data were obtained from <http://www.ndsc.ncep.noaa.gov>. The SAGE II and HALOE data were obtained from the NASA/LaRC data center.

## References

- Akimoto, H. (2003), Global air quality and pollution, *Science*, *302*, 1716–1719.
- Beirle, S., U. Platt, M. Wenig, and T. Wagner (2004), Highly resolved global distribution of tropospheric NO<sub>2</sub> using GOME narrow swath mode data, *Atmos. Chem. Phys.*, *4*, 1913–1924.
- Boersma, K. F., H. J. Eskes, and E. J. Brinksma (2004), Error analysis for tropospheric NO<sub>2</sub> retrieval from space, *J. Geophys. Res.*, *109*, D04311, doi:10.1029/2003JD003962.
- Bracher, A., M. Sinnhuber, A. Rozanov, and J. P. Burrows (2005), Using a photochemical model for the validation of NO<sub>2</sub> satellite measurements at different solar zenith angles, *Atmos. Chem. Phys.*, *5*, 393–408.
- Burrows, J. P., et al. (1999), The Global Ozone Monitoring Experiment (GOME): Mission concept and first scientific results, *J. Atmos. Sci.*, *56*(2), 151–175.
- Heland, J., H. Schlager, A. Richter, and J. P. Burrows (2002), First comparison of tropospheric NO<sub>2</sub> column densities retrieved from GOME measurements and in situ aircraft profile measurements, *Geophys. Res. Lett.*, *29*(20), 1983, doi:10.1029/2002GL015528.
- Koelemeijer, R. B. A., P. Stammes, J. W. Hovenier, and J. F. de Haan (2001), A fast method for retrieval of cloud parameters using oxygen A band measurements from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, *106*(D4), 3475–3490.
- Leue, C., M. Wenig, T. Wagner, O. Klimm, U. Platt, and B. Jähne (2001), Quantitative analysis of NO<sub>x</sub> emissions from Global Ozone Monitoring Experiment satellite image sequences, *J. Geophys. Res.*, *106*(D6), 5493–5505.
- Martin, R. V., et al. (2002), An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.*, *107*(D20), 4437, doi:10.1029/2001JD001027.
- Martin, R. V., D. J. Jacob, K. Chance, T. P. Kurosu, P. I. Palmer, and M. J. Evans (2003), Global inventory of nitrogen oxide emissions constrained by space-based observations of NO<sub>2</sub> columns, *J. Geophys. Res.*, *108*(D17), 4537, doi:10.1029/2003JD003453.
- Martin, R. V., D. D. Parrish, T. B. Ryerson, D. K. Nicks Jr., K. Chance, T. P. Kurosu, D. J. Jacob, E. D. Sturges, A. Fried, and B. P. Wert (2004), Evaluation of GOME satellite measurements of tropospheric NO<sub>2</sub> and HCHO using regional data from aircraft campaigns in the southeastern United States, *J. Geophys. Res.*, *109*, D24307, doi:10.1029/2004JD004869.
- Petritoli, A., P. Bonasoni, G. Giovanelli, F. Ravegnani, I. Kostadinov, D. Bortoli, A. Weiss, D. Schaub, A. Richter, and F. Fortezza (2004), First comparison between ground-based and satellite-borne measurements of tropospheric nitrogen dioxide in the Po basin, *J. Geophys. Res.*, *109*, D15307, doi:10.1029/2004JD004547.
- Richter, A., and J. P. Burrows (2002), Tropospheric NO<sub>2</sub> from GOME measurements, *Adv. Space Res.*, *29*, 1673–1683.
- Sudo, K., M. Takahashi, J. Kurokawa, and H. Akimoto (2002), CHASER: A global chemical model of the troposphere—1. Model description, *J. Geophys. Res.*, *107*(D17), 4339, doi:10.1029/2001JD001113.

H. Akimoto, H. Irie, and K. Sudo, Frontier Research Center for Global Change, Japan Agency for Marine-Earth Science and Technology, 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Kanagawa 236-0001, Japan. (irie@jamstec.go.jp)

S. Beirle and T. Wagner, Institut für Umweltphysik, Heidelberg University, D-69120 Heidelberg, Germany.

J. P. Burrows and A. Richter, Institute of Environmental Physics, University of Bremen, D-28334 Bremen, Germany.

F. Goutail, Service d'Aéronomie du CNRS, BP 3, Verrières le Buisson 91371, France.

Y. Kondo, Research Center for Advanced Science and Technology, University of Tokyo, 4-6-1, Komaba, Meguro, Tokyo 153-8904, Japan.

V. P. Sinyakov, Kyrgyz National University, 547, Frunze St., 720033 Bishkek, Kyrgyzstan.

M. Wenig, NASA Goddard Space Flight Center, Code 916: The Atmospheric Chemistry and Dynamics Branch, Greenbelt, MD 20771, USA.