



HAL
open science

About the Trimmed and the Poincare-Dulac normal form of diffeomorphisms

Jacky Cresson, Jasmin Raissy

► **To cite this version:**

Jacky Cresson, Jasmin Raissy. About the Trimmed and the Poincare-Dulac normal form of diffeomorphisms. 2006. hal-00076843

HAL Id: hal-00076843

<https://hal.science/hal-00076843>

Preprint submitted on 28 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ABOUT THE TRIMMED AND THE POINCARÉ-DULAC NORMAL FORM OF DIFFEOMORPHISMS

by

Jacky CRESSON & Jasmin RAISSY

Abstract. — We study two particular continuous prenormal forms as defined by Jean Ecalle and Bruno Vallet for local analytic diffeomorphism of \mathbb{C}^{ν} : the Trimmed form and the Poincaré-Dulac normal form. We first give a self-contained introduction to the mould formalism of Jean Ecalle. We provide a dictionary between moulds and the classical Lie algebraic formalism using non-commutative formal power series. We then give full proofs and details for results announced by J. Ecalle and B. Vallet about the Trimmed form of diffeomorphisms. We then discuss a mould approach to the classical Poincaré-Dulac normal form of diffeomorphisms. We discuss the universal character of moulds taking place in normalization problems.

Contents

1. Introduction.....	2
2. Diffeomorphisms, automorphisms and continuous prenormalization.....	3
3. Moulds and prenormalization.....	5
3.1. Reminder about moulds.....	5
3.1.1. Moulds and non-commutative formal power series.....	5
3.1.2. Moulds algebra.....	6
3.1.3. Composition of moulds.....	7
3.1.4. Exponential and logarithm of moulds.....	8
3.1.5. A technical lemma.....	9
3.2. Prenormalization.....	10
3.3. Universality of moulds and prenormalization.....	12
4. The Trimmed form.....	13
4.1. Cancelling non-resonant terms.....	13
4.1.1. Around the Baker-Campbell-Hausdorff formula.....	13
4.1.2. The simplified form and the moulds dem^{\bullet} and Dem^{\bullet}	15
4.2. The Trimmed form.....	16
4.2.1. The Trimmed form up to order r	16
4.2.2. The moulds ${}_r\text{sem}^{\bullet}$ and ${}_r\text{Sem}^{\bullet}$	17
4.2.3. The Trimmed form.....	18
4.2.4. The mould Trem^{\bullet}	19
4.3. About Ecalle-Vallet results.....	20
4.3.1. Relation between the alphabets $\mathcal{B}_{\mathcal{A}(F)}$ and $\mathcal{D}_{\mathcal{A}(F)}$	20
4.3.2. The mould dem^{\bullet}	20
5. The Poincaré-Dulac normal form.....	21
5.1. Homogeneous components and the Trimmed form.....	22
5.2. The Poincaré normal form of order r	23
5.3. The Poincaré-Dulac normal form.....	24

Appendix A. About the Baker-Campbell-Hausdorff formula.....	25
Appendix B. Proof of theorem 3.....	25
References.....	27

1. Introduction

In this paper we study the set of *local analytic resonant diffeomorphisms* of \mathbb{C}^ν using the theory of *continuous prenormalization* developed by J. Ecalle ([6],[7]). We assume that diffeomorphisms have a diagonalizable linear part and we work in a chart where the linear part is diagonal. The diffeomorphism is called in this case in *prepared form*. Let f be a diffeomorphism in prepared form. Roughly speaking a *prenormal form* of f is a diffeomorphism f_{pren} , conjugated to f , of the form $f_{\text{pren}} = f_{\text{lin}} + f_{\text{rem}}$ where f_{rem} is made of resonant terms. A *normal form* is a prenormal form containing the minimal number of resonant terms, with formal invariants as coefficients. Although a normal form can be considered as the simplest prenormal form, it is not in general possible to compute it. Even if an *algorithmic* procedure can be obtained [2], its exact shape is related to the vanishing of certain quantities depending polynomially on the Taylor coefficients of the diffeomorphisms. This can not be decided by a computer.

We look for *calculable* prenormal forms, *i.e.* prenormal forms which can be obtained using a procedure which is *algorithmic* and *implementable*. As an example of such prenormal forms, we study *continuous prenormal forms* as defined by J. Ecalle [6].

We mainly focus on two particular continuous prenormal forms, one introduced by J. Ecalle and B. Vallet [7] called the *Trimmed form* and the classical *Poincaré-Dulac normal form*. The framework of continuous prenormalization is the *mould formalism* developed by J. Ecalle since 1970. We provide a self-contained introduction to this formalism, omitting some aspects which will not be used in this paper. We refer to ([3],[4]) for more details.

The Trimmed form is first studied. We give complete proofs for results which are announced by J. Ecalle and B. Vallet [7] with (or without) a sketch of proof. In particular, we give all the details for the computations of the different moulds associated to the Trimmed form. We also give *closed* formulae for these moulds using a different initial alphabet.

The *Poincaré-Dulac normal form* is then discussed in the mould framework and compared to the Trimmed form. We obtain two universal moulds Poin^\bullet and Dulac^\bullet . These two universal moulds are associated to the Poincaré normalization procedure and the Poincaré-Dulac normal form. It seems impossible to obtain such objects using the existing methods of perturbation theory. The mould formalism provides a direct and algorithmic way to capture the universal features of a normalization procedure.

2. Diffeomorphisms, automorphisms and continuous prenormalization

We consider local analytic diffeomorphisms of \mathbb{C}^ν with 0 as a fixed point and a *diagonalizable* linear part. We work in a given analytic chart where the linear part is assumed to be in diagonal form. In such a case, the diffeomorphism is called in *prepared form* by J. Ecalle.

Let $f : \mathbb{C}^\nu \rightarrow \mathbb{C}^\nu$, $\nu \in \mathbb{N}$ defined by

$$(2.1) \quad f(x_1, \dots, x_\nu) = (e^{\lambda_1} x_1, \dots, e^{\lambda_\nu} x_\nu) + h(x_1, \dots, x_\nu),$$

with $f(0) = 0$, and $h = (h_1, \dots, h_\nu)$, $h_i \in \mathbb{C}\{x\}$ for all $i = 1, \dots, \nu$. We denote by f_{lin} the linear part of f , *i.e.* $f_{\text{lin}}(x_1, \dots, x_\nu) = (e^{\lambda_1} x_1, \dots, e^{\lambda_\nu} x_\nu)$.

J. Ecalle looks for the *substitution operator* associated to f , denoted by F and defined by

$$(2.2) \quad F : \begin{array}{ccc} \mathbb{C}\{x\} & \rightarrow & \mathbb{C}\{x\}, \\ \phi & \mapsto & \phi \circ f, \end{array}$$

where \circ is the usual composition of functions.

As f is a diffeomorphism, the substitution operator F is an automorphism of $(\mathbb{C}\{x\}, \cdot)$ where \cdot is the usual product of functions on $\mathbb{C}\{x\}$, *i.e.* for all $\phi, \psi \in \mathbb{C}\{x\}$, we have

$$(2.3) \quad F(\phi \cdot \psi) = F\phi \cdot F\psi,$$

and $F^{-1}(\phi) = \phi \circ f^{-1}$.

J. Ecalle uses the following result, which is a direct consequence of the Taylor expansion theorem:

Lemma 1. — *Let f be an analytic diffeomorphism of \mathbb{C}^ν in prepared form and F its associated substitution operator. There exist a decomposition of F as*

$$(2.4) \quad F = F_{\text{lin}} \left(\text{Id} + \sum_{n \in A(F)} B_n \right),$$

where $A(F)$ is an infinite set of indices $n \in \mathbb{Z}^\nu$, F_{lin} the substitution operator associated to f_{lin} , and for all $n \in A(F)$, B_n is a homogeneous differential operator of degree n , i.e. for all $m \in \mathbb{N}^\nu$,

$$(2.5) \quad B_n(x^m) = \beta_{n,m} x^{n+m}, \quad \beta_{n,m} \in \mathbb{C}.$$

In the following, we work essentially with the substitution operator F . In order to simplify our statements, we call *diffeo*(s) the automorphism F associated to a given diffeomorphism f .

Definition 1. — *Let F and F_{conj} be two local analytic diffeos of \mathbb{C}^ν . The diffeo F_{conj} is called conjugated to F if there exists a change of variables h of C^ν such that the associated substitution operator denoted by Θ satisfies*

$$(2.6) \quad F_{\text{conj}} = \Theta \cdot F \cdot \Theta^{-1}.$$

The substitution operator Θ is called the *normalizator* in the following. When the change of variables h is of class formal, C^k or C^ω , we speak of a formal, C^k or analytic normalization.

Definition 2. — *Let F be an analytic diffeo of \mathbb{C}^ν in prepared form. A prenormal form for F , denoted by F_{pran} , is an automorphism of $\mathbb{C}\{x\}$ conjugated to F such that*

$$(2.7) \quad F_{\text{pran}} \cdot F_{\text{lin}} = F_{\text{lin}} \cdot F_{\text{pran}}.$$

J. Ecalle has introduced in [6] and extensively studied in [7] a very particular class of prenormal forms called *continuous prenormal forms*.

Definition 3. — *Let F be a diffeo of \mathbb{C}^ν in prepared form given by*

$$F = F_{\text{lin}} \left(\text{Id} + \sum_{n \in A(F)} B_n \right).$$

A continuous prenormal form F_{pren} is an automorphism of $\mathbb{C}\{x\}$ of the form

$$(2.8) \quad F_{\text{pren}} = F_{\text{lin}} \left(\sum_{\mathbf{n} \in A(F)^*} \text{Pren}^{\mathbf{n}} B_{\mathbf{n}} \right),$$

where $A(F)^*$ is the set of sequences $\mathbf{n} = (n_1, \dots, n_r)$, $n_i \in A(F)$, $r \geq 0$, $\text{Pren}^{\mathbf{n}} \in \mathbb{C}$ satisfying

$$(2.9) \quad \text{Pren}^{\mathbf{n}} = 0 \quad \text{if} \quad \|\mathbf{n}\| \neq 0,$$

with $\|\mathbf{n}\| = n_1 + \dots + n_r$ for all $\mathbf{n} \in A(F)^*$, $\mathbf{n} = n_1 \dots n_r$, and $B_{\mathbf{n}} = B_{n_1} \dots B_{n_r}$ with the usual composition of differential operators.

These forms are calculable using the formalism of *moulds* developed by J. Ecalle since 1970.

3. Moulds and prenormalization

3.1. Reminder about moulds. — We provide a self-contained introduction to the formalism of moulds and we refer to the articles of J. Ecalle or to the lectures ([3],[4]) for more details.

3.1.1. Moulds and non-commutative formal power series. — We denote by A an alphabet, finite or not. A letter of A is denoted by a . Let A^* denotes the set of *words* constructed on A , i.e. the sequences $a_1 \dots a_r$, $r \geq 0$, with $a_i \in A$, with the convention that for $r = 0$ we have the *empty-word* denoted by \emptyset . We denote with bold letter \mathbf{a} a word of A^* . We have a natural action on A^* provided by the usual *concatenation* of two words $\mathbf{a}, \mathbf{b} \in A^*$, which glues the words \mathbf{a} to \mathbf{b} , i.e. \mathbf{ab} .

Definition 4. — Let \mathbb{K} be a ring (or a field) and A a given alphabet. A \mathbb{K} -valued mould on A is a map from A^* to \mathbb{K} , denoted by M^\bullet .

The evaluation of M^\bullet on a word $\mathbf{a} \in A^*$ is denoted by $M^{\mathbf{a}}$

As an example, we define a \mathbb{C} -valued mould on $A(F)$ by

$$(3.1) \quad \begin{array}{ccc} \text{Pren}^\bullet : & A(F)^* & \longrightarrow \mathbb{C} \\ & \mathbf{n} & \longmapsto \text{Pren}^{\mathbf{n}}. \end{array}$$

The mould Pren^\bullet is obtained collecting the coefficients of a formal power serie $\sum_{\mathbf{n} \in A(F)^*} \text{Pren}^{\mathbf{n}} B_{\mathbf{n}}$. There exist a one-to-one correspondence between moulds and formal

power series.

For $r \geq 0$, we denote by A_r^* the set of words of length r , with the convention that $A_0^* = \{\emptyset\}$. We denote by $\mathbb{K}\langle A \rangle$ the set of finite \mathbb{K} -linear combinations of elements of A^* , *i.e.* *non-commutative* polynomials on A with coefficients in K , and by $\mathbb{K}_r\langle A \rangle$ the set of \mathbb{K} -linear combination of elements of A_r^* , *i.e.* the set of non-commutative homogeneous polynomials of degree r . We have a natural *graduation* on $\mathbb{K}\langle A \rangle$ by the length of words:

$$(3.2) \quad \mathbb{K}\langle A \rangle = \bigoplus_{r=0}^{\infty} \mathbb{K}_r\langle A \rangle.$$

The completion of $\mathbb{K}\langle A \rangle$ with respect to the graduation by length denoted by $\mathbb{K}\langle\langle A \rangle\rangle$ is the set of formal power series with coefficients in \mathbb{K} . An element of $\mathbb{K}\langle\langle A \rangle\rangle$ is denoted by

$$(3.3) \quad \sum_{\mathbf{a} \in A^*} M^{\mathbf{a}} \mathbf{a}, \quad M^{\mathbf{a}} \in \mathbb{K},$$

where this sum must be understood as

$$(3.4) \quad \sum_{r \geq 0} \left(\sum_{\mathbf{a} \in A_r^*} M^{\mathbf{a}} \mathbf{a} \right).$$

Let M^\bullet be a \mathbb{K} -valued mould on A , its generating serie denoted by Φ_M belongs to $\mathbb{K}\langle\langle A \rangle\rangle$ and is defined by

$$(3.5) \quad \Phi_M = \sum_{\mathbf{a} \in A^*} M^{\mathbf{a}} \mathbf{a},$$

or in a condensed way as $\sum_{\bullet} M^{\bullet} \bullet$. This correspondence provide a *one-to-one mapping* from the set of \mathbb{K} -valued moulds on A denoted by $\mathcal{M}_{\mathbb{K}}(A)$ and $\mathbb{K}\langle\langle A \rangle\rangle$.

3.1.2. Moulds algebra. — The set of moulds $\mathcal{M}_{\mathbb{K}}(A)$ inherits a *structure of algebra* from $\mathbb{K}\langle\langle A \rangle\rangle$. The sum and product of two moulds M^\bullet and N^\bullet is denoted by $M^\bullet + N^\bullet$ and $M^\bullet \cdot N^\bullet$ respectively and defined by

$$(3.6) \quad \begin{aligned} (M^\bullet + N^\bullet)^{\mathbf{a}} &= M^{\mathbf{a}} + N^{\mathbf{a}}, \\ (M^\bullet \cdot N^\bullet)^{\mathbf{a}} &= \sum_{\mathbf{a}^1 \mathbf{a}^2 = \mathbf{a}} M^{\mathbf{a}^1} N^{\mathbf{a}^2}, \end{aligned}$$

for all $\mathbf{a} \in A^*$ where the sum corresponds to all the partition of \mathbf{a} as a concatenation of two words \mathbf{a}^1 and \mathbf{a}^2 of A^* .

The neutral element for the mould product is denoted by 1^\bullet and defined by

$$(3.7) \quad 1^\bullet = \begin{cases} 1 & \text{if } \bullet = \emptyset, \\ 0 & \text{otherwise,} \end{cases}$$

Let M^\bullet be a mould. We denote by \mathbb{M}^\bullet the inverse of M^\bullet for the mould product when it exists, *i.e.* the solution of the mould equation:

$$(3.8) \quad M^\bullet \cdot \mathbb{M}^\bullet = \mathbb{M}^\bullet \cdot M^\bullet = 1^\bullet.$$

3.1.3. Composition of moulds. — Assuming that A possesses a *semi-group* structure, we can define a non-commutative version of the classical operation of *substitution* of formal power series.

We denote by \star an internal law on A , such that (A, \star) is a semi-group. We denote by $\|\cdot\|_\star$ the mapping from A^* to A defined by

$$(3.9) \quad \begin{aligned} \|\cdot\|_\star: \quad A^* &\longrightarrow A, \\ \mathbf{a} = a_1 \dots a_r &\longmapsto a_1 \star \dots \star a_r. \end{aligned}$$

The \star will be omitted when clear from the context.

The set $\mathbb{K}\langle\langle A \rangle\rangle$ is graded by $\|\cdot\|_\star$. A *homogeneous component* of degree $a \in A$ of a non-commutative serie $\Phi_M = \sum_{\mathbf{a} \in A^*} M^{\mathbf{a}} \mathbf{a}$ is the quantity

$$(3.10) \quad \Phi_M^a = \sum_{\substack{\mathbf{a} \in A^*, \\ \|\mathbf{a}\|_\star = a}} M^{\mathbf{a}} \mathbf{a}.$$

We have by definition

$$(3.11) \quad \Phi_M = \sum_{a \in A} \Phi_M^a.$$

Definition 5 (Composition). — Let (A, \star) be a semi-group structure. Let M^\bullet and N^\bullet be two moulds on $\mathcal{M}_{\mathbb{K}}(A)$ and Φ_M, Φ_N their associated generating series. The substitution of Φ_N in Φ_M , denoted by $\Phi_M \circ \Phi_N$ is defined by

$$(3.12) \quad \Phi_M \circ \Phi_N = \sum_{\mathbf{a} \in A^*} M^{\mathbf{a}} \Phi_N^{\mathbf{a}},$$

where $\Phi_N^{\mathbf{a}}$ is given by $\Phi_N^{a_1} \dots \Phi_N^{a_r}$ for $\mathbf{a} = a_1 \dots a_r$.

We denote by $M^\bullet \circ N^\bullet$ the mould of $\mathcal{M}_{\mathbb{K}}(A)$ such that

$$(3.13) \quad \Phi_M \circ \Phi_N = \sum_{\mathbf{a} \in A^*} (M^\bullet \circ N^\bullet)^{\mathbf{a}} \mathbf{a}.$$

Equation (3.13) define a natural operation on moulds denoted \circ and called *composition*. Using $\|\cdot\|_\star$ we can give a closed formula for the composition of two moulds.

Lemma 2. — *Let (A, \star) be a semi-group and M^\bullet, N^\bullet be two moulds of $\mathcal{M}_{\mathbb{K}}(A)$. We have for all $\mathbf{a} \in A^*$,*

$$(3.14) \quad (M^\bullet \circ N^\bullet)^{\mathbf{a}} = \sum_{k=1}^{l(\mathbf{a})} \sum_{\mathbf{a}^1 \dots \mathbf{a}^k \bar{\star} \mathbf{a}} M^{\|\mathbf{a}^1\|_\star \dots \|\mathbf{a}^k\|_\star} N^{\mathbf{a}^1} \dots N^{\mathbf{a}^k},$$

where $\mathbf{a}^1 \dots \mathbf{a}^k \bar{\star} \mathbf{a}$ denotes all the partitions of \mathbf{a} such that $\mathbf{a}^i \neq \emptyset$, $i = 1, \dots, k$.

Proof. — Equation (3.12) is equivalent to

$$(3.15) \quad \Phi_M \circ \Phi_N = \sum_{r \geq 0} \sum_{\mathbf{b} = b_1 \dots b_r \in A_r^*} M^{b_1 \dots b_r} \left(\sum_{\mathbf{a}^1 \in A^*, \|\mathbf{a}^1\|_\star = b_1} N^{\mathbf{a}^1} \right) \dots \left(\sum_{\mathbf{a}^r \in A^*, \|\mathbf{a}^r\|_\star = b_r} N^{\mathbf{a}^r} \right).$$

Let $\mathbf{a} \in A^*$ be a given word of A^* . Each partition of \mathbf{a} of the form $\mathbf{a} = \mathbf{a}^1 \dots \mathbf{a}^k$, $k = 1, \dots, l(\mathbf{a})$, occurs in the sum (3.15) with a coefficient given by

$$(3.16) \quad M^{b_1 \dots b_r} N^{\mathbf{a}^1} \dots N^{\mathbf{a}^k},$$

where $b_i = \|\mathbf{a}^i\|_\star$. Collecting all these coefficients, we obtain the formula (3.14) for the coefficient of \mathbf{a} in $\Phi_M \circ \Phi_N$. \square

The neutral element for the mould composition is denoted by \mathbf{I}^\bullet and defined by

$$(3.17) \quad \mathbf{I}^\bullet = \begin{cases} 1 & \text{if } l(\bullet) = 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $l(\bullet)$ denotes the length of a word of A^* .

3.1.4. Exponential and logarithm of moulds. — We denote by $(\mathbb{K}\langle\langle A \rangle\rangle)_*$ the set of formal power series without a constant term. We define the *exponential* of an element $x \in (\mathbb{K}\langle\langle A \rangle\rangle)_*$, denoted by $\exp(x)$ using the classical formula

$$(3.18) \quad \exp(x) = \sum_{n \geq 0} \frac{x^n}{n!}.$$

The *logarithm* of an element $1 + x \in 1 + (\mathbb{K}\langle\langle A \rangle\rangle)_*$ is denoted by $\log(1 + x)$ and defined by

$$(3.19) \quad \log(1 + x) = \sum_{n \geq 0} (-1)^{n+1} \frac{x^n}{n!}.$$

These two applications have their natural counterpart in $\mathcal{M}_{\mathbb{K}}(A)$.

Definition 6. — Let M^\bullet be a mould of $\mathcal{M}_{\mathbb{K}}(A)$ and Φ_M the associated generating serie. Assume that $\exp(\Phi_M)$ is defined. We denote by $\text{Exp}M^\bullet$ the mould satisfying the equality

$$(3.20) \quad \exp\left(\sum_{\bullet} M^\bullet\right) = \sum_{\bullet} \text{Exp}M^\bullet.$$

Simple computations lead to the following direct definition of Exp on moulds:

$$(3.21) \quad \text{Exp}M^\bullet = \sum_{n \geq 0} \frac{[M^\bullet]_{(\times n)}}{n!},$$

where $[M^\bullet]_{(\times n)}$, $n \in \mathbb{N}$, stands for

$$(3.22) \quad [M^\bullet]_{(\times n)} = \underbrace{M^\bullet \cdots M^\bullet}_{n \text{ times}}.$$

The same procedure can be applied to define the logarithm of a mould.

Definition 7. — Let M^\bullet be a mould of $\mathcal{M}_{\mathbb{K}}(A)$ and Φ_M the associated generating serie. Assume that $\log(1 + \Phi_M)$ is defined. We denote by $\text{Log}M^\bullet$ the mould satisfying the equality

$$(3.23) \quad \log\left(1 + \sum_{\bullet} M^\bullet\right) = \sum_{\bullet} \text{Log}M^\bullet.$$

A direct definition of Log is then given by

$$(3.24) \quad \text{Log}M^\bullet = \sum_{n \geq 0} (-1)^{n+1} \frac{[M^\bullet]_{(\times n)}}{n!}.$$

As \exp and \log satisfy $\exp \circ \log = \log \circ \exp = 1$, we have

$$(3.25) \quad \text{Exp}(\text{Log}M^\bullet) = \text{Log}(\text{Exp}M^\bullet) = 1^\bullet.$$

3.1.5. A technical lemma. — In this section, we derive simple results for the exponential and logarithm of moulds with non-zero components only on words of length 1.

Lemma 3. — Let us denote by Z^\bullet a mould of $\mathcal{M}_{\mathbb{K}}(A)$ such that $Z^\bullet = 0$ for all \bullet of length different from 1. For all $\mathbf{a} \in A^*$, $r \geq 1$, we have

$$(3.26) \quad [Z^\bullet]_{\times r}^{\mathbf{a}} = \begin{cases} Z^{a_1} \cdots Z^{a_r} & \text{if } l(\mathbf{a}) = r, \mathbf{a} = a_1 \cdots a_r, \\ 0 & \text{otherwise.} \end{cases}$$

$$(3.27) \quad [\text{Exp}Z^\bullet]^{\mathbf{a}} = 1^{\mathbf{a}} + \frac{1}{l(\mathbf{a})!} [Z^\bullet]_{(\times l(\mathbf{a}))}^{\mathbf{a}},$$

$$(3.28) \quad [\text{Log}Z^\bullet]^{\mathbf{a}} = \frac{(-1)^{l(\mathbf{a})+1}}{l(\mathbf{a})!} [Z^\bullet]_{(\times l(\mathbf{a}))}^{\mathbf{a}}.$$

Proof. — We first remark that equations (3.27) and (3.28) easily follow from equation (3.26).

The proof of equation (3.26) is done by induction on r . Formula (3.26) is trivially true for $r = 1$. Assume that formula (3.26) is true for $r \geq 1$. By definition, we have

$$(3.29) \quad [Z^\bullet]_{(\times r+1)} = Z^\bullet \times [Z^\bullet]_{(\times r)}.$$

Let $\mathbf{a} = a\mathbf{b}$, then by assumption on Z^\bullet we obtain

$$(3.30) \quad [Z^\bullet]_{(\times r+1)}^{a\mathbf{b}} = Z^a [Z^\bullet]_{(\times r)}^{\mathbf{b}}.$$

As the mould $[Z^\bullet]_{(\times r)}$ is non-trivial only on words of length r , we deduce that the mould $[Z^\bullet]_{(\times r+1)}$ is non-trivial only on words of length $r + 1$.

Moreover, using the fact that $[Z^\bullet]_{(\times r)}^{a_1 \dots a_r} = Z^{a_1} \dots Z^{a_r}$ for all $a_i \in A$, we also deduce that $[Z^\bullet]_{(\times r+1)}^{a_1 \dots a_{r+1}} = Z^{a_1} \dots Z^{a_{r+1}}$. This concludes the proof. \square

3.2. Prenormalization. — Let F be a diffeo in prepared form given by

$$F = \text{F}_{\text{lin}} \left(\text{Id} + \sum_{n \in A(F)} B_n \right).$$

Let Φ_Θ be an automorphism of $\mathbb{C}\{x\}$ of the form

$$(3.31) \quad \Phi_\Theta = \sum_{\mathbf{n} \in A(F)^*} \Theta^{\mathbf{n}} B_{\mathbf{n}},$$

where $\Theta^{\mathbf{n}} \in \mathbb{C}$ for all $\mathbf{n} \in A(F)^*$, *i.e.* $\Phi_\Theta \in \mathbb{C}\langle\langle \mathbf{B} \rangle\rangle$, where $\mathbf{B} = \{B_n\}_{n \in A(F)}$ and $\Theta^\bullet \in \mathcal{M}_{\mathbb{C}}(A(F))$.

Using the moulds 1^\bullet and \mathbf{I}^\bullet we write $\text{Id} + \sum_{n \in A(F)} B_n$ as an element of $\mathbb{C}\langle\langle \mathbf{B} \rangle\rangle$:

$$(3.32) \quad \text{Id} + \sum_{n \in A(F)} B_n = \sum_{\bullet} (1^\bullet + \mathbf{I}^\bullet) B_{\bullet}.$$

We assume that F is conjugated to an automorphism F_{conj} via Φ_Θ . Equation (2.6) is then given by

$$(3.33) \quad F_{\text{conj}} = \Phi_\Theta \cdot F \cdot \Phi_\Theta^{-1}.$$

The automorphism F_{conj} can be written as

$$(3.34) \quad F_{\text{conj}} = \text{F}_{\text{lin}} \left(\sum_{\bullet} C^\bullet B_{\bullet} \right).$$

Equation (3.33) is then equivalent to

$$(3.35) \quad F_{\text{lin}} \left(\sum_{\bullet} C^{\bullet} B_{\bullet} \right) = \left(\sum_{\bullet} \Theta^{\bullet} B_{\bullet} \right) F_{\text{lin}} \left(\sum_{\bullet} (1^{\bullet} + \Gamma^{\bullet}) B_{\bullet} \right) \left(\sum_{\bullet} \Theta B_{\bullet} \right),$$

where Θ is such that $\Theta \cdot \Theta^{\bullet} = \Theta^{\bullet} \cdot \Theta = 1^{\bullet}$, i.e. $\Phi_{\Theta}^{-1} = \sum_{\bullet} \Theta B_{\bullet}$.

In order to explicit C^{\bullet} we need to understand the action of a formal power serie of $\mathbb{C}\langle\langle \mathbf{B} \rangle\rangle$ on F_{lin} . We have the following fundamental lemma:

Lemma 4. — Let $M^{\bullet} \in \mathcal{M}_{\mathbb{C}}(A(F))$. We have

$$(3.36) \quad \left(\sum_{\bullet} M^{\bullet} B_{\bullet} \right) F_{\text{lin}} = F_{\text{lin}} \left(\sum_{\bullet} e^{\Delta} (M^{\bullet})^{\bullet} B_{\bullet} \right),$$

where e^{Δ} is a map from $\mathcal{M}_{\mathbb{C}}(A(F))$ to $\mathcal{M}_{\mathbb{C}}(A(F))$ defined by

$$(3.37) \quad e^{\Delta} (M^{\bullet})^{\mathbf{n}} = e^{-\lambda \cdot \|\mathbf{n}\|} M^{\mathbf{n}} \text{ for all } \mathbf{n} \in A(F)^*.$$

Proof. — Let $B_{\mathbf{n}} = B_{n_1 \dots n_r}$ such that $B_{n_i}(x^m) = \beta_m^{n_i} x^{m+n_i}$, $\beta_m^{n_i} \in \mathbb{C}$, $i = 1, \dots, r$, for all $m \in \mathbb{N}^{\nu}$. We have

$$(3.38) \quad B_{\mathbf{n}}(x^m) = \beta_{m+n_r+\dots+n_2}^{n_1} \beta_{m+n_r+\dots+n_3}^{n_2} \dots \beta_m^{n_r} x^{m+n_1+\dots+n_r}.$$

As $F_{\text{lin}}(x^m) = e^{\lambda \cdot m} x^m$ we obtain

$$(3.39) \quad \begin{aligned} B_{\mathbf{n}}(F_{\text{lin}}(x^m)) &= e^{\lambda \cdot m} B_{\mathbf{n}}(x^m), \\ &= e^{-\lambda \cdot (n_1+\dots+n_r)} e^{\lambda \cdot (m+n_1+\dots+n_r)} B_{\mathbf{n}}(x^m), \\ &= e^{-\lambda \cdot (n_1+\dots+n_r)} F_{\text{lin}}(B_{\mathbf{n}}(x^m)), \\ &= F_{\text{lin}}(e^{-\lambda \cdot (n_1+\dots+n_r)} B_{\mathbf{n}}(x^m)). \end{aligned}$$

This concludes the proof. □

Next lemma gives an explicit formula to compute the mould C^{\bullet} assuming that the mould Θ^{\bullet} is known.

Lemma 5. — Equation (3.35) is equivalent to the mould equation

$$(3.40) \quad C^{\bullet} = e^{\Delta} (\Theta^{\bullet}) \cdot (1^{\bullet} + \Gamma^{\bullet}) \cdot \Theta.$$

Proof. — Using lemma 4, we have

$$\begin{aligned}
 (3.41) \quad F_{\text{lin}} \left(\sum_{\bullet} C^{\bullet} B_{\bullet} \right) &= \left(\sum_{\bullet} \Theta^{\bullet} B_{\bullet} \right) F_{\text{lin}} \left(\sum_{\bullet} (1^{\bullet} + \Gamma^{\bullet}) B_{\bullet} \right) \left(\sum_{\bullet} \ominus B_{\bullet} \right), \\
 &= F_{\text{lin}} \left(\sum_{\bullet} e^{\Delta} (\Theta^{\bullet}) B_{\bullet} \right) \left(\sum_{\bullet} (1^{\bullet} + \Gamma^{\bullet}) B_{\bullet} \right) \left(\sum_{\bullet} \ominus B_{\bullet} \right), \\
 &= F_{\text{lin}} \left(\sum_{\bullet} (e^{\Delta} (\Theta^{\bullet}) \cdot (1^{\bullet} + \Gamma^{\bullet}) \cdot \ominus) B_{\bullet} \right).
 \end{aligned}$$

This concludes the proof. \square

As a consequence, choosing carefully the normalizer Φ_{Θ} , we can obtain an inductive expression for the mould of normalization C^{\bullet} .

We will give explicit formulae for C^{\bullet} using specific moulds for Θ^{\bullet} in the next section.

3.3. Universality of moulds and prenormalization. — Lemma 5 gives an important feature of the mould formalism in the context of continuous prenormalization. Formula (3.40) is valid whatever is the underlying alphabet $A(F)$. We then obtain a *universal* object underlying the prenormalization problem which is studied.

For example, in the context of *linearization*, i.e. $F_{\text{conj}} = F_{\text{lin}}$, the universal mould of linearization which defined the linearizing change of variables is given as follow (see [3] Chap. III for more details):

Theorem 1. — Let $\mathbf{L} = \{L_r\}_{r \geq 1}$, $r \in \mathbb{N}$, be the set of \mathbb{C} -valued functions $L_r : \mathbb{C}^r \rightarrow \mathbb{C}$ defined by

$$(3.42) \quad L_r(x_1, \dots, x_r) = [(e^{-(x_1 + \dots + x_r)} - 1) (e^{-(x_2 + \dots + x_r)} - 1) \dots (e^{-x_r} - 1)]^{-1},$$

for all $(x_1, \dots, x_r) \in \mathbb{C}^r \setminus S_r$ where the singular set S_r is given by

$$(3.43) \quad S_r = \{x_r = 0\} \cup \{x_r + x_{r-1} = 0\} \cup \dots \cup \{x_1 + \dots + x_r = 0\}.$$

If F possesses a non-resonant linear part λ , the mould of formal linearization is given for all $\mathbf{n} \in A(F)^*$, $\mathbf{n} = n_1, \dots, n_r$, by

$$(3.44) \quad \Theta^{n_1 \dots n_r} = L_r(\omega_1, \dots, \omega_r),$$

where $\omega_i = n_i \cdot \lambda$ for $i = 1, \dots, r$.

This result can not be obtained using other existing formalisms. Of course, anybody knows that an expression like (3.44) is the important quantity entering the linearization problem. However, the previous result associates universal coefficients from which one can compute the desired linearization map for a given particular diffeo F by posing

$$\Phi_{\Theta} = \sum_{\mathbf{n} \in \mathcal{A}(F)^*} \Theta^{\mathbf{n}} B_{\mathbf{n}}.$$

4. The Trimmed form

In this section, we give complete proofs of results concerning the *Trimmed form* defined by J. Ecalle and B. Vallet announced in [7] with a sketch of proof and without explicit computations.

4.1. Cancelling non-resonant terms. — In this section, we give a mould approach to the classical problem of cancellation of non-resonant terms.

4.1.1. Around the Baker-Campbell-Hausdorff formula. — Let F be a diffeo in prepared form given by (2.4). The operator $\text{Id} + \sum_{n \in \mathcal{A}(F)} B_n$ is an automorphism of $\mathbb{C}\{x\}$ which can be viewed as the exponential of a vector field, *i.e.*

$$(4.1) \quad \text{Id} + \sum_{n \in \mathcal{A}(F)} B_n = \exp \left(\sum_{m \in \mathcal{A}(F)} D_m \right),$$

where D_n is a homogeneous differential operator of degree m and order 1, *i.e.* a derivation on $\mathbb{C}\{x\}$, $m = (m_1, \dots, m_\nu) \in \mathbb{Z}^\nu$, with all $m_i \in \mathbb{N}$, $i = 1, \dots, \nu$ except at most one which can be -1 , and $\mathcal{A}(F)$ the set of degrees coming in the decomposition.

We look for an automorphism given by the exponential of a vector field \mathbf{V} given by

$$(4.2) \quad \mathbf{V} = \sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{dem}^{\mathbf{n}} B_{\mathbf{n}},$$

or equivalently given on the alphabet $\mathcal{A}(F)^*$ by

$$(4.3) \quad \mathbf{V} = \sum_{\mathbf{m} \in \mathcal{A}(F)^*} \text{Dem}^{\mathbf{m}} D_{\mathbf{m}},$$

where $V_{\mathbf{m}=m_1 \dots m_r} = V_{m_1} V_{m_2} \dots V_{m_r}$ with the usual composition of differential operators.

The action of $\exp \mathbf{V}$ on F is given by

$$(4.4) \quad \exp \mathbf{V} \cdot F \cdot \exp(-\mathbf{V})$$

Equation (4.4) can be analyzed using the moulds expression of \mathbf{V} and F with respect to the alphabet $\mathcal{A}(F)$. We have the following lemma:

Lemma 6. — Equation (4.4) is equal to

$$(4.5) \quad \exp \mathbf{V} \cdot F \cdot \exp(-\mathbf{V}) = F_{\text{lin}} \exp \left(\tilde{\mathbf{V}} + \mathbf{D} - \mathbf{V} + \dots \right),$$

where the \dots stands for a formal power serie beginning with words of length at least 2, and \mathbf{D} and $\tilde{\mathbf{V}}$ are vector fields defined by $\mathbf{D} = \sum_{m \in \mathcal{A}(F)} D_m$ and

$$(4.6) \quad \tilde{\mathbf{V}} = \sum_{\mathbf{m} \in \mathcal{A}(F)^*} e^{-\lambda \cdot \|\mathbf{m}\|} \text{Dem}^{\mathbf{m}} D_{\mathbf{m}},$$

respectively.

Proof. — Using the Baker-Campbell-Hausdorff formula (BCH_2), we obtain

$$(4.7) \quad \begin{aligned} \exp \mathbf{D} \cdot \exp(-\mathbf{V}) &= \exp(\mathbf{D} \star (-\mathbf{V})), \\ &= \exp(\mathbf{D} - \mathbf{V} + \text{h.o.t.}), \end{aligned}$$

where h.o.t. stands for *higher order terms*.

Using lemma 4, we have

$$(4.8) \quad \exp \mathbf{V} \cdot F_{\text{lin}} = F_{\text{lin}} \cdot \exp \tilde{\mathbf{V}},$$

where $\tilde{\mathbf{V}}$ is given by

$$(4.9) \quad \tilde{\mathbf{V}} = \sum_{\mathbf{m} \in \mathcal{A}(F)^*} e^{-\lambda \cdot \|\mathbf{m}\|} \text{Dem}^{\mathbf{m}} D_{\mathbf{m}}.$$

As a consequence, applying again (BCH_2) we obtain

$$(4.10) \quad \begin{aligned} \exp \tilde{\mathbf{V}} \cdot \exp(\mathbf{D} \star (-\mathbf{V})) &= \exp \left(\tilde{\mathbf{V}} \star (\mathbf{D} \star (-\mathbf{V})) \right), \\ &= \exp \left(\tilde{\mathbf{V}} + \mathbf{D} - \mathbf{V} + \dots \right), \end{aligned}$$

where the \dots stand for a formal power serie beginning with words of length at least 2. This concludes the proof. \square

4.1.2. *The simplified form and the moulds dem^\bullet and Dem^\bullet .* — The main consequence of lemma 6 is that we can cancel the non-resonant terms of \mathbf{D} using a simple vector field \mathbf{V} .

Theorem 2 (Simplified form). — *Let \mathbf{V} be the vector field defined by the mould*

$$(4.11) \quad \text{Dem}^\bullet = \begin{cases} \frac{I^{\mathbf{m}}}{1 - e^{|\mathbf{m}| \lambda}} & \text{for } \mathbf{m} \in \mathcal{A}(F)^* \setminus \mathcal{R}(F), \\ 0 & \text{otherwise,} \end{cases}$$

where $\mathcal{R}(F)$ is the set of resonant words of $\mathcal{A}(F)^*$, i.e. $\mathbf{m} \in \mathcal{R}(F)$ if and only if $\mathbf{m} \cdot \lambda = 0$. We denote by dem^\bullet the associated mould on $\mathcal{M}_{\mathbb{C}}(\mathcal{A}(F))$, i.e.

$$(4.12) \quad \mathbf{V} = \sum_{\bullet} \text{Dem}^\bullet \mathbf{D}_{\bullet} = \sum_{\bullet} \text{dem}^\bullet \mathbf{B}_{\bullet}.$$

We call *simplified form of F* and we denote by F_{Sem} the automorphism obtained from F under the action of $\exp \mathbf{V}$. We have

$$(4.13) \quad \begin{aligned} F_{\text{Sem}} &= F_{\text{lin}} \left(\sum_{\mathbf{m} \in \mathcal{A}(F)^*} \text{Sem}^{\mathbf{m}} D_{\mathbf{m}} \right), \\ &= F_{\text{lin}} \left(\sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{sem}^{\mathbf{n}} B_{\mathbf{n}} \right) \end{aligned}$$

with the mould Sem^\bullet given by

$$(4.14) \quad \text{Sem}^\bullet = e^{\Delta} (\text{Exp}(\text{Dem}^\bullet)) \cdot \text{Exp}(I^\bullet) \cdot \text{Exp}(-\text{Dem}^\bullet),$$

and the mould sem^\bullet given by

$$(4.15) \quad \text{sem}^\bullet = e^{\Delta} (\text{Exp}(\text{dem}^\bullet)) \cdot (1^\bullet + I^\bullet) \cdot \text{Exp}(-\text{dem}^\bullet).$$

Proof. — We have $F_{\text{Sem}} = \exp \mathbf{V} \cdot F \cdot \exp(-\mathbf{V})$ with $\mathbf{V} = \sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{dem}^{\mathbf{n}} B_{\mathbf{n}}$. As a consequence, we have $\exp \mathbf{V} = \sum_{\mathbf{n} \in \mathcal{A}(F)^*} (\text{Exp} \text{dem}^\bullet)^{\mathbf{n}} B_{\mathbf{n}}$ and the formula for sem^\bullet follows from lemma 5 using $\Theta^\bullet = \text{Exp}(\text{dem}^\bullet)$.

For Sem^\bullet , we first use lemma 4 to obtain

$$(4.16) \quad \exp \mathbf{V} F_{\text{lin}} = F_{\text{lin}} \left(\sum_{\mathbf{m} \in \mathcal{A}(F)^*} [e^{\Delta} (\text{Exp} \text{Dem}^\bullet)]^{\mathbf{n}} D_{\mathbf{m}} \right).$$

As a consequence, the conjugacy equation is equivalent to

$$(4.17) \quad \begin{aligned} F_{\text{Sem}} &= \exp \mathbf{V} \cdot F \cdot \exp(-\mathbf{V}), \\ &= F_{\text{lin}} \left(\sum_{\bullet} e^{\Delta} (\text{ExpDem}^{\bullet} \text{D}_{\bullet}) \right) \left(\sum_{\bullet} \text{ExpI}^{\bullet} \text{D}_{\bullet} \right) \left(\sum_{\bullet} \text{Exp}(-\text{Dem}^{\bullet}) \text{D}_{\bullet} \right), \\ &= F_{\text{lin}} \left(\sum_{\bullet} [e^{\Delta} (\text{ExpDem}^{\bullet}) \cdot \text{ExpI}^{\bullet} \cdot \text{Exp}(-\text{Dem}^{\bullet})]^{\bullet} \text{D}_{\bullet} \right). \end{aligned}$$

This concludes the proof. \square

The mould Sem^{\bullet} can be compute explicitly. We first introduce some convenient notations:

Let $\mathbf{m} = m_1 \dots m_r$ be a word of length r , $r \geq 1$. We denote by $\mathbf{m}^{\leq i}$ and $\mathbf{m}^{> i}$ the word

$$(4.18) \quad \mathbf{m}^{\leq i} = m_1 \dots m_i, \quad \mathbf{m}^{> i} = m_{i+1} \dots m_r.$$

Moreover we denote by $d(\mathbf{m})$ the index of the last m_i in $\mathbf{m} = m_1 \dots m_r$ such that $\lambda.m_i = 0$, and we denote by $q(\mathbf{m})$ the first index just before of the first zero $\omega_j = \lambda.m_j$.

Theorem 3. — For all $\mathbf{m} \in \mathcal{A}(F)^*$, we have

$$(4.19) \quad \begin{aligned} \text{Sem}^{\mathbf{m}} &= \frac{(-1)^{l(\mathbf{m})}}{l(\mathbf{m})!} [\text{Dem}^{\bullet}]_{(\times l(\mathbf{m}))}^{\mathbf{m}} + \frac{1}{l(\mathbf{m})!} + \sum_{j=d(\mathbf{m})+1}^{l(\mathbf{m})+1} \frac{(-1)^{l(\mathbf{m}^{> j})} [\text{Dem}^{\bullet}]_{(\times l(\mathbf{m}^{\geq j}))}^{\mathbf{m}^{\geq j}}}{l(\mathbf{m}^{< j})! l(\mathbf{m}^{\geq j})!} + e^{-\lambda.\|\mathbf{m}\|} \mathbf{1}^{\mathbf{m}} \\ &+ \sum_{i=1}^{q(\mathbf{m}) \wedge (l(\mathbf{m})-1)} \frac{e^{-\lambda.\|\mathbf{m}^{\leq i}\|}}{l(\mathbf{m}^{\leq i})!} [\text{Dem}^{\bullet}]_{(\times l(\mathbf{m}^{\leq i}))}^{\mathbf{m}^{\leq i}} \times \\ &\left(\frac{(-1)^{l(\mathbf{m})}}{l(\mathbf{m})!} [\text{Dem}^{\bullet}]_{(\times l(\mathbf{m}^{> i}))}^{\mathbf{m}^{> i}} + \frac{1}{l(\mathbf{m}^{> i})!} + \sum_{j=d(\mathbf{m}^{> i})+1}^{l(\mathbf{m}^{> i})+1} \frac{(-1)^{l(\mathbf{m}^{> j}) \geq j} [\text{Dem}^{\bullet}]_{(\times l((\mathbf{m}^{> i})^{\geq j}))}^{(\mathbf{m}^{> i})^{\geq j}}}{l((\mathbf{m}^{> i})^{< j})! l((\mathbf{m}^{> i})^{\geq j})!} \right). \end{aligned}$$

The proof is done in appendix B.

4.2. The Trimmed form. — The Trimmed form is constructed by induction applying successively the previous simplification scheme to remove non-resonant terms of higher and higher degrees. The mould formalism allows us to explicit some particular moulds underlying this construction as well as algorithmic and explicit formulae for some of them.

4.2.1. The Trimmed form up to order r . — We can use the simplification procedure previously defined inductively in order to cancel non-resonant terms of higher and higher degrees.

Definition 8 (Trimmed form up to order r). — Let $r \in \mathbb{N}$, the Trimmed form up to order r is defined as F_{Sem}^r obtained from F after r successive simplifications, i.e.

$$(4.20) \quad F = F_{\text{Sem}}^0 \xrightarrow{\text{Simp}^1} F_{\text{Sem}}^1 \xrightarrow{\text{Simp}^2} \dots \xrightarrow{\text{Simp}^r} F_{\text{Sem}}^r,$$

where Simp^i is the automorphism of simplification defined by

$$(4.21) \quad \text{Simp}^i = \exp(\mathbf{V}_i),$$

with \mathbf{V}_i the vector fields associated to the mould Dem^\bullet on the alphabet $\mathcal{A}(F_{\text{Sem}}^{i-1})$ associated to F_{Sem}^{i-1} .

Using theorem 2, we deduce the following useful result:

Theorem 4. — For all $r \in \mathbb{N}$, the Trimmed form up to order r denoted F_{Sem}^r possesses a mould expansion, i.e. there exists moulds denoted by ${}_r\text{Sem}^\bullet \in \mathcal{M}_{\mathbb{C}}(\mathcal{A}(F))$ and ${}_r\text{sem}^\bullet \in \mathcal{M}_{\mathbb{C}}(\mathcal{A}(F))$ such that

$$(4.22) \quad F_{\text{sem}}^r = F_{\text{lin}} \left(\sum_{\bullet} {}_r\text{Sem}^\bullet \mathbf{D}_\bullet \right) = F_{\text{lin}} \left(\sum_{\bullet} {}_r\text{sem}^\bullet \mathbf{B}_\bullet \right).$$

Despite its moulds expansion, the Trimmed form up to order r is *not* a prenormal form as it remains non-resonant terms for sequences of length $l \geq r + 1$.

4.2.2. The moulds ${}_r\text{sem}^\bullet$ and ${}_r\text{Sem}^\bullet$. — The mould ${}_r\text{sem}^\bullet$ has a simple expression in function of sem^\bullet .

Lemma 7. — For all $r \in \mathbb{N}$, we have

$$(4.23) \quad {}_r\text{sem}^\bullet = \underbrace{\text{sem}^\bullet \circ \dots \circ \text{sem}^\bullet}_{r \text{ times}}.$$

Proof. — The simplification procedure can be written as follows:

$$(4.24) \quad \sum_{\bullet} \mathbf{I}^\bullet \mathbf{B}_\bullet \longmapsto \sum_{\bullet} \text{sem}^\bullet \mathbf{B}_\bullet.$$

Iterating this mapping we go from step i to $i + 1$

$$(4.25) \quad \sum_{\bullet} {}_i\text{sem}^\bullet \mathbf{B}_\bullet = \sum_{\bullet} \mathbf{I}^\bullet {}_{i+1}\mathbf{B}_\bullet \longmapsto \sum_{\bullet} {}_{i+1}\text{sem}^\bullet \mathbf{B}_\bullet = \sum_{\bullet} \text{sem}^\bullet {}_{i+1}\mathbf{B}_\bullet,$$

where $\sum_{\bullet} \mathbf{I}^\bullet {}_{i+1}\mathbf{B}_\bullet$ denotes the homogeneous decomposition constructed on F_{Sem}^i .

By definition of the composition for moulds we have

$$(4.26) \quad \sum_{\bullet} \text{sem}^\bullet {}_{i+1}\mathbf{B}_\bullet = \sum_{\bullet} (\text{sem}^\bullet \circ {}_i\text{sem}^\bullet) \mathbf{B}_\bullet,$$

from which we deduce the recursive relation

$$(4.27) \quad {}_{i+1}\text{sem}^\bullet = \text{sem}^\bullet \circ {}_i\text{sem}^\bullet.$$

We conclude by induction on i . □

For the mould ${}_r\text{Sem}^\bullet$ we have a more complicated formula:

Lemma 8. — *For all $r \in \mathbb{N}$, we have*

$$(4.28) \quad \text{Log}[_r\text{Sem}^\bullet] = \underbrace{\text{Log}(\text{Sem}^\bullet) \circ \dots \circ \text{Log}(\text{Sem}^\bullet)}_{r \text{ times}}.$$

The fact that we must take the Log of Sem^\bullet instead of Sem^\bullet is related to the fact that the alphabet of derivation ${}_{i+1}\text{D}_\bullet$ constructed at step i from F_{sem}^i is not related to $\sum_\bullet {}_i\text{Sem}^\bullet\text{D}_\bullet$ but to its logarithm.

Proof. — The simplification procedure can be written as follows:

$$(4.29) \quad \exp\left(\sum_\bullet \text{I}^\bullet\text{D}_\bullet\right) \mapsto \sum_\bullet \text{Sem}^\bullet\text{D}_\bullet = \exp\left(\sum_\bullet \text{Log}(\text{Sem}^\bullet)\text{D}_\bullet\right).$$

Iterating this mapping we go from step i to $i + 1$

$$(4.30) \quad \begin{aligned} \exp\left(\sum_\bullet \text{Log}[_i\text{Sem}^\bullet]\text{D}_\bullet\right) &= \exp\left(\sum_\bullet \text{I}^\bullet{}_{i+1}\text{D}_\bullet\right) \\ &\downarrow \\ \exp\left(\sum_\bullet \text{Log}[_{i+1}\text{Sem}^\bullet]\text{D}_\bullet\right) &= \exp\left(\sum_\bullet \text{Log}(\text{Sem}^\bullet)_{i+1}\text{D}_\bullet\right), \end{aligned}$$

where $\sum_\bullet \text{I}^\bullet{}_{i+1}\text{D}_\bullet$ denotes the homogeneous decomposition constructed on $\sum_\bullet \text{Log}[_i\text{Sem}^\bullet]\text{D}_\bullet$.

By definition of the composition of moulds, we deduce that

$$(4.31) \quad \text{Log}[_{i+1}\text{Sem}^\bullet] = \text{Log}(\text{Sem}^\bullet) \circ \text{Log}[_i\text{Sem}^\bullet].$$

We conclude the proof by induction on i . □

4.2.3. The Trimmed form. —

Definition 9. — *The Trimmed form of F is the limit of the simplification procedure.*

Theorem 5. — *The Trimmed form is a continuous prenormal form given by*

$$(4.32) \quad \begin{aligned} F_{\text{Trem}} &= F_{\text{lin}} \left(\sum_{\mathbf{m} \in \mathcal{A}(F)^*} \text{Trem}^{\mathbf{m}} D_{\mathbf{m}} \right), \\ &= F_{\text{lin}} \left(\sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{trem}^{\mathbf{n}} B_{\mathbf{n}} \right) \end{aligned}$$

with the moulds Trem^\bullet and trem^\bullet defined by

$$(4.33) \quad \begin{aligned} \text{Trem}^\bullet - 1^\bullet &= \text{limstat}_{r \rightarrow \infty} [\text{Sem}^\bullet - 1^\bullet]^{(or)}, \\ \text{trem}^\bullet - 1^\bullet &= \text{limstat}_{r \rightarrow \infty} [\text{sem}^\bullet - 1^\bullet]^{(or)}, \end{aligned}$$

where limstat is the stationary limit.

The proof is a direct consequence of the simplification procedure.

Remark 1. — *Following ([7] §.7) we have divergence and resurgence of the simplification procedure. This is not the case when working directly with the diffeomorphism instead of its associated automorphism of substitution. However, this problem can be avoided (see [7] p.8).*

4.2.4. *The mould Trem^\bullet .* — We can compute the mould Trem^\bullet using a simple remark. By definition, we have the following identities

$$(4.34) \quad \text{Trem}^\bullet = \text{Sem}^\bullet \circ \text{Trem}^\bullet,$$

$$(4.35) \quad \text{Trem}^\bullet = \text{Trem}^\bullet \circ \text{Sem}^\bullet.$$

Using the first equation and the definition of composition for moulds we obtain for all $\mathbf{m} \in \mathcal{A}(F)^*$

$$(4.36) \quad \text{Trem}^{\mathbf{m}} = \text{Sem}^{|\mathbf{m}|} \text{Trem}^{\mathbf{m}} + \text{s.l.},$$

where s.l denotes terms which depend on Trem^\bullet for words with a strictly *short length* than $l(\mathbf{m})$.

The mould Trem^\bullet takes non-trivial values only on resonant words, *i.e.* $\mathbf{m} \in \mathcal{A}(F)^*$ such that $\|\mathbf{m}\| \cdot \lambda = 0$. However, the mould Sem^\bullet is equal to 1 on resonant words of length 1. As a consequence, equation (4.36) can not be used to compute the mould Trem^\bullet by induction on the length of words.

Using equation (4.35) we obtain

$$(4.37) \quad \text{Trem}^{\mathbf{m}} = \text{Trem}^{\mathbf{m}} \text{Sem}^{m_1} \dots \text{Sem}^{m_r} + \text{s.l.}$$

4.3. About Ecalle-Vallet results. — All our computations have been done in the alphabet $\mathcal{D}_{\mathcal{A}(F)}$. However, J. Ecalle and B. Vallet [7] use the initial alphabet $\mathcal{B}_{\mathcal{A}(F)}$ to formulate their results. In order to compare our approach, we first give a simple formula connecting the two alphabets. We then discuss the essential differences between the moulds dem^\bullet , sem^\bullet and trem^\bullet with the moulds Dem^\bullet , Sem^\bullet , and Trem^\bullet . The main point is that contrary to our moulds, Ecalle-Vallet moulds can not be expressed via closed formulae, except for dem^\bullet .

4.3.1. Relation between the alphabets $\mathcal{B}_{\mathcal{A}(F)}$ and $\mathcal{D}_{\mathcal{A}(F)}$. — By definition, we have the identity

$$(4.38) \quad 1 + \sum_{n \in \mathcal{A}(F)} B_n = \exp \left(\sum_{m \in \mathcal{A}(F)} D_m \right).$$

Using the logarithm, we obtain

$$(4.39) \quad \log \left(1 + \sum_{n \in \mathcal{A}(F)} B_n \right) = \sum_{m \in \mathcal{A}(F)} D_m.$$

As $\sum_{n \in \mathcal{A}(F)} B_n = \sum_{\mathbf{n} \in \mathcal{A}^*(F)} \mathbf{I}^{\mathbf{n}} B_{\mathbf{n}}$, we have

$$(4.40) \quad \sum_{\mathbf{n} \in \mathcal{A}^*(F)} (\text{Log} \mathbf{I}^\bullet)^{\mathbf{n}} B_{\mathbf{n}} = \sum_{m \in \mathcal{A}(F)} D_m.$$

We finally deduce the following relation between $\mathcal{D}_{\mathcal{A}(F)}$ and $\mathcal{B}_{\mathcal{A}(F)}$:

Lemma 9. — *For all $D_m \in \mathcal{D}_{\mathcal{A}(F)}$, we have*

$$(4.41) \quad D_m = \sum_{\mathbf{n} \in \mathcal{A}(F)^*, \|\mathbf{n}\|=m} (\text{Log} \mathbf{I}^\bullet)^{\mathbf{n}} B_{\mathbf{n}}.$$

The proof is based on the fact that a differential operator $B_{\mathbf{n}}$ is of order $\|\mathbf{n}\|$.

4.3.2. The mould dem^\bullet . — By definition, we have the identity

$$(4.42) \quad \sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{dem}^{\mathbf{n}} B_{\mathbf{n}} = \sum_{m \in \mathcal{A}(F) \setminus \mathcal{R}_{\mathcal{A}(F)}} \frac{D_m}{1 - e^{m \cdot \lambda}}.$$

Using lemma 9, we deduce:

Lemma 10. — *The mould dem^\bullet of $\mathcal{M}_{\mathbb{C}}(A(F))$ is defined for all $\mathbf{n} \in A(F)^*$ by*

$$(4.43) \quad \text{dem}^{\mathbf{n}} = \frac{(-1)^{l(\mathbf{n})+1}}{l(\mathbf{n})!} \frac{1}{1 - e^{\|\mathbf{n}\|.\lambda}} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}))}^{\mathbf{n}} \mathbf{1}_{N(F)}(\mathbf{n}),$$

where $N(F) = \{\mathbf{n} \in A(F)^*, \|\mathbf{n}\|.\lambda \neq 0\}$ is the set of non-resonant words of $A(F)^*$ and $\mathbf{1}_J$ is the indicatrice of the set J , i.e. $\mathbf{1}_J(x)$ is equal to 1 if $x \in J$, 0 otherwise.

This mould is defined directly by Ecalle-Vallet without any details (see [7], p.30).

Proof. — Equation (4.42) can be rewritten as

$$(4.44) \quad \sum_{\mathbf{n} \in A(F)^*} \text{dem}^{\mathbf{n}} B_{\mathbf{n}} = \sum_{m \in \mathcal{A}(F)} \frac{D_m}{1 - e^{m.\lambda}} \mathbf{1}_{\{m.\lambda \neq 0\}}(m).$$

Using lemma 9, we have

$$(4.45) \quad \begin{aligned} \sum_{m \in \mathcal{A}(F)} \frac{D_m}{1 - e^{m.\lambda}} \mathbf{1}_{\{m.\lambda \neq 0\}}(m) &= \sum_{m \in \mathcal{A}(F)} \sum_{\substack{\mathbf{n} \in A(F)^*, \\ \|\mathbf{n}\|=m}} \frac{(\text{Log} \mathbf{I}^\bullet)^{\mathbf{n}}}{1 - e^{m.\lambda}} B_{\mathbf{n}}, \\ &= \sum_{\mathbf{n} \in A(F)^*} \frac{(\text{Log} \mathbf{I}^\bullet)^{\mathbf{n}}}{1 - e^{\|\mathbf{n}\|.\lambda}} \mathbf{1}_{N(F)} B_{\mathbf{n}}, \end{aligned}$$

using the fact that

$$(4.46) \quad \bigcup_{m \in \mathcal{A}(F)} \{\mathbf{n} \in A(F)^*, \|\mathbf{n}\|=m\} = A(F)^*,$$

by assumption.

Using lemma 3 for the mould \mathbf{I}^\bullet , we obtain for all $\mathbf{n} \in A(F)^*$

$$(4.47) \quad \text{Log} \mathbf{I}^{\mathbf{n}} = \frac{(-1)^{l(\mathbf{n})+1}}{l(\mathbf{n})!} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}))}^{\mathbf{n}}.$$

Replacing $\text{Log} \mathbf{I}^\bullet$ by its expression in equation (4.45) we conclude the proof. \square

5. The Poincaré-Dulac normal form

The Trimmed form is constructed using cancellation of non-resonant terms as the classical Poincaré-Dulac normal form. However, these two prenormal forms do not coincide in general. We introduce the universal mould associated to the Poincaré-Dulac normal form and the universal mould of the associated cancellation procedure. The difference between the two procedures lies in the treatment of the homogeneous components of the diffeomorphism. For a classical approach to the Poincaré-Dulac normal form we refer to ([1] §.B p.178).

5.1. Homogeneous components and the Trimmed form. — We keep the notations introduced in §.4.1. In order to discuss the cancellation of non-resonant terms, we must write our prepared form as follows:

$$(5.1) \quad \text{Id} + \sum_{n \in \mathcal{A}(F)} B_n = \exp \mathbf{D} = \exp \left(\sum_{m \in \mathcal{A}(F)} D_m \right) = \exp \left(\sum_{k \geq 1} \mathbf{D}_k \right),$$

where

$$(5.2) \quad \mathbf{D}_k = \sum_{n \in \mathcal{A}(F), |n|=k} D_n,$$

denotes the homogeneous component of degree k of the vector field \mathbf{D} .

For a given vector field \mathbf{D} we introduce the following *degree of resonance*, denoted by K :

$$(5.3) \quad K = \min_{k \geq 1} \{N_k \neq \emptyset\},$$

where N_k denotes the set of non-resonant letters $m \in \mathcal{A}(F)$ of degree k , *i.e.*

$$(5.4) \quad N_k = \{m \in \mathcal{A}(F) \mid |m| = k, m \cdot \lambda = 0\}.$$

As a consequence, we have

$$(5.5) \quad \mathbf{D} = \sum_{1 \leq k < K} \mathbf{D}_k + \mathbf{D}_K + \sum_{k > K} \mathbf{D}_k.$$

The first sum up to order $K - 1$ is made of resonant terms. The first non-resonant terms belong to \mathbf{D}_K .

The field \mathbf{V} introduced in §.4.1.2 cancel the non-resonant terms of degree K but introduces several other terms in the homogeneous components of degree $> K$ which can be non-resonant. As a consequence, even if the field \mathbf{V} is constructed in order to cancel *all* the non-resonant terms of the vector field \mathbf{D} we have an effective cancellation only for the components of degree K .

As a consequence, the vector field \mathbf{V} must be modified in order to cancel *only* non-resonant terms of degree K .

Theorem 6 (Poincaré normalization procedure). — *Let \mathbf{S} be the vector field defined by the mould*

$$(5.6) \quad \text{Den}^\bullet = \begin{cases} \frac{1}{1 - e^{m.\lambda}} & \text{for } m \in \mathbb{N}_K(F), \\ 0 & \text{otherwise,} \end{cases}$$

We denote by den^\bullet the associated mould on $\mathcal{M}_\mathbb{C}(A(F))$, i.e.

$$(5.7) \quad \mathbf{S} = \sum_{\bullet} \text{Den}^\bullet \mathbf{D}_\bullet = \sum_{\bullet} \text{den}^\bullet \mathbf{B}_\bullet.$$

We call simplified form of F and we denote by F_{Poin} the automorphism obtained from F under the action of $\exp \mathbf{S}$. We have

$$(5.8) \quad \begin{aligned} F_{\text{Poin}} &= F_{\text{lin}} \left(\sum_{\mathbf{m} \in \mathcal{A}(F)^*} \text{Poin}^{\mathbf{m}} D_{\mathbf{m}} \right), \\ &= F_{\text{lin}} \left(\sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{poin}^{\mathbf{n}} B_{\mathbf{n}} \right) \end{aligned}$$

with the mould Poin^\bullet given by

$$(5.9) \quad \text{Poin}^\bullet = e^\Delta (\text{Exp}(\text{Den}^\bullet)) \cdot \text{Exp}(\mathbf{I}^\bullet) \cdot \text{Exp}(-\text{Den}^\bullet),$$

and the mould poin^\bullet given by

$$(5.10) \quad \text{poin}^\bullet = e^\Delta (\text{Exp}(\text{den}^\bullet)) \cdot (\mathbf{1}^\bullet + \mathbf{I}^\bullet) \cdot \text{Exp}(-\text{den}^\bullet).$$

The proof is exactly the same as those of theorem 2.

5.2. The Poincaré normal form of order r . — We apply the Poincaré normalization procedure inductively in order to cancel non-resonant terms in homogeneous components of higher and higher degree.

Definition 10 (Poincaré normal form up to order r). — *Let $r \in \mathbb{N}$, the Poincaré normal form up to order r is defined as F_{Poin}^r obtained from F after r successive simplifications, i.e.*

$$(5.11) \quad F = F_{\text{Poin}}^0 \xrightarrow{\text{Simp}^1} F_{\text{Poin}}^1 \xrightarrow{\text{Simp}^2} \dots \xrightarrow{\text{Simp}^r} F_{\text{Poin}}^r,$$

where Simp^i is the automorphism of simplification defined by

$$(5.12) \quad \text{Simp}^i = \exp(\mathbf{S}_i),$$

with \mathbf{S}_i the vector fields associated to the mould Den^\bullet on the alphabet $\mathcal{A}(F_{\text{Poin}}^{i-1})$ associated to F_{Poin}^{i-1} .

Using theorem 6, we obtain:

Theorem 7. — For all $r \in \mathbb{N}$, the Poincaré normal form up to order r denoted F_{Poin}^r possesses a mould expansion, i.e. there exist moulds denoted by ${}_r\text{Poin}^\bullet \in \mathcal{M}_{\mathbb{C}}(\mathcal{A}(F))$ and ${}_r\text{poin}^\bullet \in \mathcal{M}_{\mathbb{C}}(A(F))$ such that

$$(5.13) \quad F_{\text{Poin}}^r = F_{\text{lin}} \left(\sum_{\bullet} {}_r\text{Poin}^\bullet D_{\bullet} \right) = F_{\text{lin}} \left(\sum_{\bullet} {}_r\text{poin}^\bullet B_{\bullet} \right).$$

As for the moulds ${}_r\text{sem}^\bullet$ and ${}_r\text{Sem}^\bullet$, we have explicit inductive formulae to compute the moulds ${}_r\text{poin}^\bullet$ and ${}_r\text{Poin}^\bullet$ using only poin^\bullet and Poin^\bullet .

5.3. The Poincaré-Dulac normal form. — The mould formulation of the Poincaré-Dulac normal form is:

Definition 11. — The Poincaré-Dulac normal form of F is the limit of the Poincaré normalization procedure.

Theorem 8. — The Poincaré-Dulac normal form is a continuous prenormal form given by

$$(5.14) \quad \begin{aligned} F_{\text{Dulac}} &= F_{\text{lin}} \left(\sum_{\mathbf{m} \in \mathcal{A}(F)^*} \text{Dulac}^{\mathbf{m}} D_{\mathbf{m}} \right), \\ &= F_{\text{lin}} \left(\sum_{\mathbf{n} \in \mathcal{A}(F)^*} \text{dulac}^{\mathbf{n}} B_{\mathbf{n}} \right) \end{aligned}$$

with the moulds Dulac^\bullet and dulac^\bullet defined by

$$(5.15) \quad \begin{aligned} \text{Dulac}^\bullet - 1^\bullet &= \lim_{\text{stat}_{r \rightarrow \infty}} [\text{Poin}^\bullet - 1^\bullet]^{(or)}, \\ \text{dulac}^\bullet - 1^\bullet &= \lim_{\text{stat}_{r \rightarrow \infty}} [\text{poin}^\bullet - 1^\bullet]^{(or)}, \end{aligned}$$

where \lim_{stat} is the stationary limit.

The mould Dulac^\bullet (or dulac^\bullet) is the *universal* part of the Poincaré-Dulac normal form as it does not depends on the exact values of the coefficients coming in the Taylor expansion of the diffeomorphism. It seems impossible to characterize such kind of object without using moulds.

Appendix A

About the Baker-Campbell-Hausdorff formula

The Baker-Campbell-Hausdorff formula covers at least two formulae which are of interest for the computation of continuous prenormal forms for vector fields and diffeomorphisms.

Let A and B be two linear operators. We denote by

$$\exp A = \sum_{k \geq 0} \frac{A^k}{k!},$$

where $A^k = A \circ \dots \circ A$, k times. The Baker-Campbell-Hausdorff formula is given by

$$(\exp A) \cdot B \cdot (\exp(-A)) = \sum_{m \geq 0} \frac{B_m}{m!}, \quad (BCH_1)$$

where

$$(A.1) \quad B_m = [A, B]_m = [A[A, \dots, [A, B]] \dots],$$

with the convention that $B_0 = B$.

A consequence of this formula is

$$\exp A \exp B = \exp(A \star B), \quad (BCH_2)$$

where

$$(A.2) \quad A \star B = A + B + \frac{1}{2}[A, B] + \frac{1}{12}[A, [A, B]] - \frac{1}{12}[B, [A, B]] + \dots$$

Appendix B

Proof of theorem 3

In order to compute the mould Sem^\bullet , we first compute $\text{ExpI}^\bullet \cdot \text{Exp}(-\text{Dem}^\bullet)$. We have

$$(B.1) \quad \begin{aligned} (\text{ExpI}^\bullet \cdot \text{Exp}(-\text{Dem}^\bullet))^{\mathbf{n}} &= \sum_{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n}} (\text{ExpI}^\bullet)^{\mathbf{n}^1} \text{Exp}(-\text{Dem}^\bullet)^{\mathbf{n}^2}, \\ &= \sum_{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n}} \left(1^{\mathbf{n}^1} + \frac{1}{l(\mathbf{n}^1)!} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}^1))}^{\mathbf{n}^1} \right) \left(1^{\mathbf{n}^2} + \frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^2)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} \right), \\ &= \sum_{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n}} \left(1^{\mathbf{n}^1} 1^{\mathbf{n}^2} + 1^{\mathbf{n}^1} \frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^2)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} \right. \\ &\quad \left. + 1^{\mathbf{n}^2} \frac{1}{l(\mathbf{n}^1)!} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}^1))}^{\mathbf{n}^1} + \frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^1)! l(\mathbf{n}^2)!} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}^1))}^{\mathbf{n}^1} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} \right). \end{aligned}$$

It is clear that $(\text{Exp} \mathbf{I}^\bullet \cdot \text{Exp}(-\text{Dem}^\bullet))^\emptyset = 1$. If $l(\mathbf{n}) \geq 1$ we have

$$\begin{aligned}
(\text{Exp} \mathbf{I}^\bullet \cdot \text{Exp}(-\text{Dem}^\bullet))^\mathbf{n} &= \frac{(-1)^{l(\mathbf{n})}}{l(\mathbf{n})!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}))}^\mathbf{n} + \frac{1}{l(\mathbf{n})!} [\mathbf{I}^\bullet]_{(\times l(\mathbf{n}))}^\mathbf{n} \\
&\quad + \sum_{\substack{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n} \\ \mathbf{n}^1 \neq \emptyset}} \left(\frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^1)! l(\mathbf{n}^2)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} \right), \\
\text{(B.2)} \qquad \qquad \qquad &= \frac{(-1)^{l(\mathbf{n})}}{l(\mathbf{n})!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}))}^\mathbf{n} + \frac{1}{l(\mathbf{n})!} \\
&\quad + \sum_{j=d(\mathbf{n})+1}^{l(\mathbf{n})+1} \frac{(-1)^{l(\mathbf{n}^{\geq j})} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^{\geq j}))}^{\mathbf{n}^{\geq j}}}{l(\mathbf{n}^{< j})! l(\mathbf{n}^{\geq j})!}.
\end{aligned}$$

Now we can compute Sem^\bullet .

$$\begin{aligned}
\text{(B.3)} \qquad \text{Sem}^\mathbf{n} &= (e^\Delta (\text{Exp}(\text{Dem}^\bullet)) \cdot \text{Exp}(\mathbf{I}^\bullet) \cdot \text{Exp}(-\text{Dem}^\bullet))^\mathbf{n}, \\
&= \sum_{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n}} (e^\Delta \text{Exp}(\text{Dem}^\bullet))^{\mathbf{n}^1} \left(\frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^2)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} \right. \\
&\quad \left. + \frac{1}{l(\mathbf{n}^2)!} + \sum_{j=d(\mathbf{n}^2)+1}^{l(\mathbf{n}^2)+1} \frac{(-1)^{l((\mathbf{n}^2)^{\geq j})} [\text{Dem}^\bullet]_{(\times l((\mathbf{n}^2)^{\geq j}))}^{(\mathbf{n}^2)^{\geq j}}}{l((\mathbf{n}^2)^{< j})! l((\mathbf{n}^2)^{\geq j})!} \right), \\
&= \sum_{\mathbf{n}^1 \mathbf{n}^2 = \mathbf{n}} e^{-\lambda \cdot \|\mathbf{n}^1\|} \left(1^{\mathbf{n}^1} + \frac{1}{l(\mathbf{n}^1)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^1))}^{\mathbf{n}^1} \right) \times \\
&\quad \left(\frac{(-1)^{l(\mathbf{n}^2)}}{l(\mathbf{n}^2)!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^2))}^{\mathbf{n}^2} + \frac{1}{l(\mathbf{n}^2)!} + \sum_{j=d(\mathbf{n}^2)+1}^{l(\mathbf{n}^2)+1} \frac{(-1)^{l((\mathbf{n}^2)^{\geq j})} [\text{Dem}^\bullet]_{(\times l((\mathbf{n}^2)^{\geq j}))}^{(\mathbf{n}^2)^{\geq j}}}{l((\mathbf{n}^2)^{< j})! l((\mathbf{n}^2)^{\geq j})!} \right), \\
&= \frac{(-1)^{l(\mathbf{n})}}{l(\mathbf{n})!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}))}^\mathbf{n} + \frac{1}{l(\mathbf{n})!} + \sum_{j=d(\mathbf{n})+1}^{l(\mathbf{n})+1} \frac{(-1)^{l(\mathbf{n}^{\geq j})} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^{\geq j}))}^{\mathbf{n}^{\geq j}}}{l(\mathbf{n}^{< j})! l(\mathbf{n}^{\geq j})!} \\
&\quad + e^{-\lambda \cdot \|\mathbf{n}\|} 1^\mathbf{n} + \sum_{i=1}^{q(\mathbf{n}) \wedge (l(\mathbf{n})-1)} \frac{e^{-\lambda \cdot \|\mathbf{n}^{\leq i}\|}}{l(\mathbf{n}^{\leq i})!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^{\leq i}))}^{\mathbf{n}^{\leq i}} \times \\
&\quad \left(\frac{(-1)^{l(\mathbf{n}^{> i})}}{l(\mathbf{n}^{> i})!} [\text{Dem}^\bullet]_{(\times l(\mathbf{n}^{> i}))}^{\mathbf{n}^{> i}} + \frac{1}{l(\mathbf{n}^{> i})!} + \sum_{j=d(\mathbf{n}^{> i})+1}^{l(\mathbf{n}^{> i})+1} \frac{(-1)^{l(\mathbf{n}^{> i})^{\geq j}} [\text{Dem}^\bullet]_{(\times l((\mathbf{n}^{> i})^{\geq j}))}^{(\mathbf{n}^{> i})^{\geq j}}}{l((\mathbf{n}^{> i})^{< j})! l((\mathbf{n}^{> i})^{\geq j})!} \right).
\end{aligned}$$

This concludes the proof.

References

- [1] V.I. Arnold, *Chapitres supplémentaires de la théorie des équations différentielles ordinaires*, Ed. Librairie du Globe, Paris (1996).
- [2] A. Baider, Unique normal forms for vector fields and Hamiltonian, *J. Diff. Equ.* 78 (1989), 33-52.
- [3] J. Cresson, *Mould calculus and normalization of vector fields and diffeomorphisms*, Lectures at the University of Pisa, Prépublications de l'I.H.É.S. (2006) 27.p.
- [4] J. Cresson, *Calcul Moulien*, Prépublications de l'I.H.É.S. 06/22 (2006) 98.p.
- [5] Ecalle J., Singularités non abordables par la géométrie, *Ann. Inst. Fourier*, 42 (1-2), 1992, 73-164.
- [6] Ecalle J., Schlomiuk D., The nilpotent and distinguished form of resonant vector fields or diffeomorphisms, *Ann. Inst. Fourier* 43 5 (1993) 1407-1483.
- [7] J. Ecalle, B. Vallet, Prenormalization, correction, and linearization of resonant vector fields or diffeomorphisms, *Prepublication d'Orsay* (1995) 101.p.
- [8] J. Ecalle, B. Vallet, Correction an linearization of resonant vector fields and diffeomorphisms, *Math. Z.* 229 (1998) 249-318.

Jacky CRESSON

Université de Pau et des Pays de l'Adour,
Laboratoire de Mathématiques appliquées de Pau, CNRS UMR 5142
jacky.cresson@univ-pau.fr

and

Institut des Hautes Études Scientifiques (I.H.É.S.)
Le Bois-Marie, 35 Route de Chartres, F-91440 Bures sur Yvette, France.
cresson@ihes.fr

Jasmin RAISSY
University of Pisa

raissy@mail.dm.unipi.it

JACKY CRESSON
JASMIN RAISSY