



**HAL**  
open science

## Spectroscopic Evidence for Gas Phase Formation of Successive Beta-turns in a Three-Residue Peptide Chain

W. Chin, I. Compagnon, J. P. Dognon, C. Canuel, F. Piuzzi, I. Dimicoli, G. von Helden, G. Meijer, M. Mons

► **To cite this version:**

W. Chin, I. Compagnon, J. P. Dognon, C. Canuel, F. Piuzzi, et al.. Spectroscopic Evidence for Gas Phase Formation of Successive Beta-turns in a Three-Residue Peptide Chain. *Journal of the American Chemical Society*, 2005, 127, pp.1388-1389. 10.1021/ja042860b . hal-00081949

**HAL Id: hal-00081949**

**<https://hal.science/hal-00081949>**

Submitted on 7 Sep 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Spectroscopic Evidence for Gas Phase Formation of Successive $\beta$ -turns in a Three-Residue Peptide Chain

Wutharath Chin,<sup>†</sup> Isabelle Compagnon,<sup>‡</sup> Jean-Pierre Dognon,<sup>†</sup> Clélia Canuel,<sup>†</sup> François Piuze,<sup>†</sup> Iliana Dimicoli,<sup>†</sup> Gert von Helden,<sup>§</sup> Gerard Meijer,<sup>§</sup> and Michel Mons<sup>\*†</sup>

*Contribution from Laboratoire Francis Perrin (URA CEA CNRS 2453), Service des Photons, Atomes et Molécules, Centre d'Etudes de Saclay, Bât. 522, 91191 Gif-sur-Yvette Cedex, France and FOM Institute for Plasma Physics Rijnhuizen, Edisonbaan 14, 3439 MN Nieuwegein, The Netherlands*

Author E-mail : mmons@cea.fr

Gas phase studies of small biomolecules<sup>1-3</sup> and peptides<sup>4-9</sup> have gained a considerable interest in the recent years as they provide experimental counterparts to the numerous high level calculations on short peptide chains,<sup>10-15</sup> documenting the competition between secondary structures such as  $\beta$ -strands,  $\beta$ -turns,  $2_7$  ribbons,  $\alpha$ - and  $3_{10}$  helices. In the present communication, we report an optical spectroscopic study in the gas phase on the three-residue peptide chain, Ac-Phe-Gly-Gly-NH<sub>2</sub>, (Ac=acetyl). The conformational analysis carried out from the IR spectroscopy is backed by high-level theoretical calculations, which enables a structural assignment in terms of intramolecular H-bonding. In particular, we provide evidence for the spontaneous formation of successive entangled  $\beta$ -turns in the gas phase.

In the experiment, molecules are laser-desorbed and cooled in a supersonic expansion.<sup>16,17</sup> UV spectra (Fig. S1) are recorded in the absorption region of the phenylalanine UV chromophore using mass-selective resonant two-photon ionization (R2PI). At least five conformers, distinguishable in the UV spectrum, are populated in the jet (labeled A1-A4 and B in Fig. S1).

The IR spectrum of each conformer in the amide A region (3  $\mu$ m) has been obtained in Saclay by the IR/UV double resonance technique<sup>1,2</sup> using an OPO (1  $\text{cm}^{-1}$  resolution) as an IR source.<sup>8</sup> The IR spectra (Fig. 1a) of the five conformers A1-A4 and B of Ac-Phe-Gly-Gly-NH<sub>2</sub> show resolved absorption bands assigned to NH stretching modes of the molecule. Some bands are intense, broad and significantly red-shifted compared to the range of free or nearly free NH's (3420 - 3550  $\text{cm}^{-1}$ ).<sup>8</sup> Those characteristics are unambiguous signatures of intramolecular H-bonding. The presence of IR bands in an intermediate region (3420-3450  $\text{cm}^{-1}$ ) indicates weaker interactions of the corresponding NH groups, as for example in C<sub>5</sub> close contacts in  $\beta$  strand-like ( $\beta_L$ ) conformations or in NH-aromatic interactions.<sup>5,6,8</sup> These basic considerations indicate that conformers A1-A4 in Fig. 1 contain strong intramolecular H-bonds that involve at least two NH moieties. The spectrum of conformer B is qualitatively different. It exhibits two narrow and intense bands that are moderately red-shifted (at 3380 and 3392  $\text{cm}^{-1}$ ), indicating that two medium-strength H-bonds are present. The amide I-II (5-8  $\mu$ m) absorption spectrum of this conformer (Fig. 2) was recorded using the same IR/UV double resonance technique, employing the free electron laser FELIX<sup>17,18</sup> (15  $\text{cm}^{-1}$  resolution) as the IR source. In contrast to the amide A region, the amide I region exhibits only one partially resolved feature.

The here proposed H-bonding assignment of these five conformers are consistent with recently reported IR spectra of shorter peptides.<sup>19</sup> The main conformer of Ac-Gly-Phe-NH<sub>2</sub>, assigned to a double  $\gamma$ -fold structure, shows two broad C<sub>7</sub> bands

in the 3300-3370  $\text{cm}^{-1}$  region, bearing a striking resemblance to the red part of the present A1-A4 spectra (Fig. 1a). On the other hand, the minor conformers of Ac-Gly-Phe-NH<sub>2</sub> and Ac-Phe-Gly-NH<sub>2</sub>, assigned to  $\beta$ -turns,<sup>18,19</sup> exhibit a C<sub>10</sub> H-bond signature at 3388 and 3387  $\text{cm}^{-1}$  respectively. This is very close to the slightly red shifted bands observed for conformer B.


This first analysis provides evidence for a large propensity of the Ac-Phe-Gly-Gly-NH<sub>2</sub> tripeptide chain to form an extended intramolecular H-bonding network in the gas phase, presumably containing at least two strong C<sub>7</sub> H-bonds (conformers A1-A4). Interestingly, however, the most intense feature observed in the UV spectrum (conformer B; Fig. S1) exhibits a radically different IR pattern, which might be due to  $\beta$ -turns.

Focusing onto this conformer B, quantum mechanical geometry optimizations (DFT B3LYP/6-31+G(d)) have been carried out for a set of selected conformations that exhibit only two H-bonds (Fig. 1b; Table S1). Considered were: structures with (1) two C<sub>10</sub> bonds (successive entangled  $\beta$ -turns), (2) one C<sub>10</sub> and one C<sub>7</sub> bond (successive entangled  $\beta$ -turns and  $\gamma$ -turns in C<sub>10</sub>-C<sub>7</sub> as well as C<sub>7</sub>-C<sub>10</sub> order), (3) one central C<sub>7</sub>  $\gamma$ -turn accompanied by either a parallel C<sub>13</sub> or an antiparallel C<sub>11</sub> bond, and (4) having a  $\beta$  strand-like ( $\beta_L$ ; C<sub>5</sub>) conformation on Phe (very stable in the protected amino acid<sup>6,8</sup>) followed by two successive  $\gamma$ -folds (C<sub>7</sub>-C<sub>7</sub>). For comparison, two remarkable secondary structures have also been considered: (5) a triple C<sub>7</sub> structure ( $2_7$  ribbon) and (6) a triple C<sub>5</sub> structure ( $\beta$ -strand). The orientation (anti, gauche+ and gauche-) of the Phe side-chain relative to the backbone was chosen either arbitrarily or, when possible, to allow stabilizing NH- $\pi$  interactions. In structures (1-2), among the four canonical types of  $\beta$ -turns, types I or II were chosen for the Phe-Gly part, as they have been shown to be the most stable forms in Ac-Phe-Gly-NH<sub>2</sub>.<sup>19</sup> For the Gly-Gly part, all the four types were *a priori* considered, but steric considerations led to retain only I-I and II-I' combinations.  $3_{10}$  helices (which also belong to family (1)) were found to converge to I-I forms. For  $\gamma$ -turns on Gly residue (families (2) and (5)), both inverse or direct, i.e.,  $\gamma_L$  or  $\gamma_D$  respectively, were considered, in contrast to  $\gamma$ -turns on Phe for which  $\gamma_L$  are much more stable.<sup>8</sup> Figure 1c displays calculated absorption spectra in the amide A region for selected conformations characteristic of the six families described above. A large variety of red shifts and hence of H-bond strengths is found and the calculated spectra can be compared to the experimental absorption pattern of conformer B. This comparison (Fig. 1c and Table S2) suggests that the central  $\gamma$ -fold (3),  $2_7$ -ribbon (5) and  $\beta$ -strand (6) secondary structures can be rejected, although one should remark that in the case of the A1-A4 conformers, the  $2_7$ -ribbon family accounts for the observed red-shifts, in agreement with the qualitative analysis. For a further analysis of the spectrum of conformer B, only conformations of


<sup>†</sup> CEA Saclay

<sup>‡</sup> FOM Institute for Plasmaphysics Rijnhuizen.

<sup>§</sup> Present address: Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195 Berlin, Germany



**Figure 1.** a) IR/UV double resonance spectrum in the amide A region of five conformers of Ac-Phe-Gly-Gly-NH<sub>2</sub> labeled A1-A4 and B. The spectral gap in the 3470-3500 cm<sup>-1</sup> region is due to an undesired absorption in the crystal of the IR source. b) H-bonding schemes in a series of conformations exhibiting 2 H-bonds (labeled (1) to (4)), three H-bonds (5) or none (6); c) DFT B3LYP/6-31+G(d) calculated stick spectra (scaling factor 0.960) of selected conformations (see text and Table S1) for comparison with the spectrum of conformer B.



**Figure 2.** Amide I and II IR spectrum of conformer B of Ac-Phe-Gly-Gly-NH<sub>2</sub> (lower bold trace and dots) and DFT B3LYP/6-31+G(d) calculated spectra (narrow lines) of selected conformations (see text; Table S1) obtained by convoluting the stick spectra by a 15 cm<sup>-1</sup> width gaussian-like function to account for the experimental line widths. Scaling factors of 0.980 and 0.970 have been used in the amide I and II regions respectively.

families (1), (2) and (4) are considered, as the agreement between their amide A spectra and the experiment is within the expected range (usually better than 15 cm<sup>-1</sup>).<sup>8,19</sup> The best agreement is found for structures exhibiting *two successive beta-turns, of type II and I' respectively*, and a Phe side chain in either (a) or (g-) orientation. This backbone structural assignment has been cross-checked against spectroscopic data from the amide I/II regions as well as energetic considerations.

Calculations (Fig. 2, Table S2) show that the amide I absorption spectrum is sensitive to both H-bonding and Phe side-chain orientation. Comparison to the experiment allows us to reject conformations possessing one or two  $\gamma$ -folds (Families (2) and (4)), as their spectra have too large amide I splittings. We also can distinguish between very similar structures, like II(a)-I' and II(g-)-I', or II(a)-I' and II(a)- $\gamma_D$  (Fig. S2). The best fit to the experimental amide I and II spectrum (Fig. 2) is provided by the double  $\beta$ -turn type II(a)-type I' conformation.

This assignment is further supported by refined energetic calculations (MP2/6-31+G(d)// B3LYP/6-31+G(d); Table S1): the type II(a)-type I' conformation is the most stable form among all the conformations here considered.

As a conclusion, for the first time for a peptide chain model of this size, a detailed conformational analysis has been carried out, revealing the intrinsic folding properties of the chain. The abundance of the double  $\beta$ -turn form observed indicates a significant evolution of the folding preferences by sequence lengthening. Indeed, a previous study revealed that a Phe-Gly chain preferentially adopts an extended  $\beta_L$ - $\gamma_L$  conformation; the type II  $\beta$ -turn conformation had only minor abundance.<sup>19</sup> This propensity is reversed in the present longer peptide, presumably because of the rather high stability of the  $\beta$ -turns in the Gly-Gly part, as suggested by recent high level calculations.<sup>15,20</sup>

Finally, one can remark that nascent 3<sub>10</sub> helices (type I-type I conformations, Fig. S2) are less stable than type II-type I' conformations, probably because the central Gly residue strongly favors the type II  $\beta$ -turn of the Phe-Gly part of the chain. The absence of the 3<sub>10</sub> helix conformation should probably be viewed as a specific effect of the Gly residue. This result might be of importance since the Gly residue, which is less demanding in terms of calculation facility, is generally considered as a case study and a popular model in theoretical studies.


**Acknowledgment.** This work was supported by the Stichting voor Fundamenteel Onderzoek der Materie (FOM) in providing the required beam time on FELIX and by the FP6 "Structuring the European Research Area" Program of the European Community. We also wish to thank the FELIX staff for their skilful assistance.

**Supporting Information Available:** UV spectrum of Ac-Phe-Gly-Gly-NH<sub>2</sub> (Fig. S1); Structural and energetic parameters (Table S1), calculated IR data (Table S2) and relevant DFT optimized structures (Fig. S2).

## References

- (1) Zwier, T. S. *J. Phys. Chem. A* **2001**, *105*, 8827.
- (2) Robertson, E. G.; Simons, J. P. *Phys. Chem. Chem. Phys.* **2001**, *3*, 1.
- (3) Jarrold, M. F. *Annu. Rev. Phys. Chem.* **2000**, *51*, 179.
- (4) Cohen, R.; Brauer, B.; Nir, E.; Grace, L.; de Vries, M. S. *J. Phys. Chem. A* **2000**, *104*, 6351.
- (5) Dian, B. C.; Longarte, A.; Zwier, T. S. *Science* **2002**, *296*, 2369.
- (6) Unterberg, C.; Gerlach, A.; Schrader, T.; Gerhards, M. *J. Chem. Phys.* **2003**, *118*, 8296.
- (7) Hünig, I.; Kleinermanns, K. *Phys. Chem. Chem. Phys.* **2004**, *6*, 2650.
- (8) Chin, W.; Mons, M.; Dognon, J.-P.; Piuze, F.; Tardivel, B.; Dimicoli, I. *Phys. Chem. Chem. Phys.* **2004**, *6*, 2700.
- (9) Kapota, C.; Lemaire, J.; Maitre, P.; Ohanessian, G. *J. Am. Chem. Soc.* **2004**, *126*, 1836.
- (10) Topol, I. A.; Burt, S. K.; Derety, E.; Tang, T. H.; Perczel, A.; Rashin, A.; Csizmadia, I. G. *J. Am. Chem. Soc.* **2001**, *123*, 6054.
- (11) Wu, Y. D.; Zhao, Y. L. *J. Am. Chem. Soc.* **2001**, *123*, 5313.
- (12) Improta, R.; Barone, V.; Kudin, K. N.; Scuseria, G. E. *J. Am. Chem. Soc.* **2001**, *123*, 3311.
- (13) Wiczorek, R.; Dannenberg, J. J. *J. Am. Chem. Soc.* **2003**, *125*, 14065.
- (14) Wiczorek, R.; Dannenberg, J. J. *J. Am. Chem. Soc.* **2003**, *125*, 8124.
- (15) Jalkanen, K. J.; Elstner, M.; Suhai, S. *J. Mol. Struct. (Theochem)* **2004**, *675*, 61.
- (16) Piuze, F.; Dimicoli, I.; Mons, M.; Tardivel, B.; Zhao, Q. *Chem. Phys. Lett.* **2000**, *320*, 282.
- (17) Bakker, J. M.; Aleese, L. M.; Meijer, G.; von Helden, G. *Phys. Rev. Lett.* **2003**, *91*, 203003.
- (18) Chin, W.; Dognon, J.-P.; Canuel, C.; Piuze, F.; Dimicoli, I.; Mons, M.; Compagnon, I.; von Helden, G.; Meijer, G. *J. Chem. Phys.*, in press.
- (19) Chin, W.; Dognon, J.-P.; Piuze, F.; Tardivel, B.; Dimicoli, I.; Mons, M. *J. Am. Chem. Soc.*, in press.
- (20) Möhle, K.; Gussmann, M.; Hofmann, H. *J. Comput. Chem.* **1997**, *18*, 1415.

## Table of Content Artwork



We report the first gas phase spectroscopic study of a three-residue model of a peptide chain, Ac-Phe-Gly-Gly-NH<sub>2</sub> (Ac=acetyl), using the IR/UV double resonance technique. The existence of at least five different conformers under supersonic expansion conditions is established, most of them exhibiting rather strong intramolecular H-bonds. One of the most populated conformers, however, exhibits a different H-bonding network characterized by two weak H-bonds. Comparison of the amide A and I/II experimental data with Density Functional Theory calculations carried out on a series of selected conformations enables us to assign this conformer to two successive β-turns along the peptide chain; the two H-bonds being of C<sub>10</sub> type, i.e., each of them closing a 10-atom ring in the molecule. The corresponding form is found to be more stable than the 3<sub>10</sub> helix secondary structure (not observed), presumably because of specific effects due to the glycine residues.