



HAL
open science

Explicit spectral gap estimates for the linearized Boltzmann and Landau operators with hard potentials

Céline Baranger, Clément Mouhot

► **To cite this version:**

Céline Baranger, Clément Mouhot. Explicit spectral gap estimates for the linearized Boltzmann and Landau operators with hard potentials. *Revista Matemática Iberoamericana*, 2005, 21, pp.819-841. hal-00087219

HAL Id: hal-00087219

<https://hal.science/hal-00087219>

Submitted on 21 Jul 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Explicit spectral gap estimates for the linearized Boltzmann and Landau operators with hard potentials

Céline Baranger and Clément Mouhot

July 21, 2006

Abstract

This paper deals with explicit spectral gap estimates for the linearized Boltzmann operator with hard potentials (and hard spheres). We prove that it can be reduced to the Maxwellian case, for which explicit estimates are already known. Such a method is constructive, does not rely on Weyl's Theorem and thus does not require Grad's splitting. The more physical idea of the proof is to use geometrical properties of the whole collision operator. In a second part, we use the fact that the Landau operator can be expressed as the limit of the Boltzmann operator as collisions become grazing in order to deduce explicit spectral gap estimates for the linearized Landau operator with hard potentials.

Mathematics Subject Classification (2000): 76P05 Rarefied gas flows, Boltzmann equation [See also 82B40, 82C40, 82D05].

Keywords: spectral gap, Boltzmann linearized operator, Landau linearized operator, geometrical properties, explicit, grazing collision limit, hard potentials.

Contents

1	Introduction	1
2	The Boltzmann linearized operator	8
3	The Landau linearized operator	15

1 Introduction

This paper is devoted to the study of the spectral properties of the linearized Boltzmann and Landau collision operators with hard potentials. In this work we shall obtain new quantitative estimates on the spectral gap of these operators. Before we explain our

methods and results in more details, let us introduce the problem in a precise way. The Boltzmann equation describes the behavior of a dilute gas when the only interactions taken into account are binary elastic collisions. It reads in \mathbb{R}^N ($N \geq 2$)

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = Q^{\mathcal{B}o}(f, f),$$

where $f(t, x, v)$ stands for the time-dependent distribution function of density of particles in the phase space. The N -dimensional Boltzmann collision operator Q is a quadratic operator, which is local in (t, x) . The time and position are only parameters and therefore shall not be written in the sequel: the estimates proven in this paper are all local in (t, x) . Thus it acts on $f(v)$ by

$$Q^{\mathcal{B}o}(f, f)(v) = \int_{v_* \in \mathbb{R}^N} \int_{\sigma \in \mathbb{S}^{N-1}} B(\cos \theta, |v - v_*|) [f'_* f' - f_* f] d\sigma dv_*$$

where we have used the shorthands $f = f(v)$, $f_* = f(v_*)$, $f' = f(v')$, $f'_* = f(v'_*)$. The velocities are given by

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma, \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma.$$

The collision kernel B is a non-negative function which only depends on $|v - v_*|$ and $\cos \theta = k \cdot \sigma$ where $k = (v - v_*)/|v - v_*|$.

Consider the collision operator obtained by the linearization process around the Maxwellian *global* equilibrium state denoted by M

$$L^{\mathcal{B}o}h(v) = \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} B(\cos \theta, |v - v_*|) M(v_*) [h'_* + h' - h_* - h] d\sigma dv_*,$$

where $f = M(1+h)$ and $M(v) = e^{-|v|^2}$. Notice that for the associated linearized equation, the entropy is the $L^2(M)$ norm of h and thus by differentiating, the entropy production is

$$\langle h, L^{\mathcal{B}o}h \rangle_{L^2(M)} = -\frac{1}{4} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} B(\cos \theta, |v - v_*|) [h'_* + h' - h_* - h]^2 M M_* d\sigma dv_* dv.$$

This quantity is non-positive. At the level of the linearized equation, this corresponds to the first part of Boltzmann's H -theorem, and it implies that the spectrum of $L^{\mathcal{B}o}$ in $L^2(M)$ is non-positive. Let us denote $D^{\mathcal{B}o}(h) = -\langle h, L^{\mathcal{B}o}h \rangle_{L^2(M)}$. We shall call this quantity “(linearized) entropy dissipation functional” by analogy with the nonlinear case.

In the case of long-distance interaction, the collisions occur mostly for very small deviation angle θ . In the case of the Coulomb potential, for which the Boltzmann collision operator is meaningless (see [18, Annex I, Appendix A]), one has to replace it by the Landau collision operator

$$Q^{\mathcal{L}a}(f, f)(v) = \nabla_v \cdot \left(\int_{v_* \in \mathbb{R}^N} \mathbf{a}(v - v_*) [f_*(\nabla f) - f(\nabla f)_*] dv_* \right),$$

with $\mathbf{a}(z) = |z|^2 \Phi(z) \Pi_{z^\perp}$, where Π_{z^\perp} is the orthogonal projection onto z^\perp , i.e

$$(\Pi_{z^\perp})_{i,j} = \delta_{i,j} - \frac{z_i z_j}{|z|^2}.$$

This operator is used for instance in models of plasma in the case of a Coulomb potential, i.e a gas of (partially or totally) ionized particles (for more details see [16] and the references therein). Applying the same linearization process than for the Boltzmann operator (around the same global equilibrium M), we define the linearized Landau operator

$$L^{\mathcal{L}^a} h(v) = M(v)^{-1} \nabla_v \cdot \left(\int_{v_* \in \mathbb{R}^N} \mathbf{a}(v - v_*) [(\nabla h) - (\nabla h)_*] M M_* dv_* \right),$$

and the (linearized) Landau entropy dissipation functional

$$\begin{aligned} D^{\mathcal{L}^a}(h) &= - \langle h, L^{\mathcal{L}^a} h \rangle_{L^2(M)} \\ &= \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Phi(v - v_*) |v - v_*|^2 \|\Pi_{(v-v_*)^\perp} [(\nabla h) - (\nabla h)_*]\|^2 M M_* dv_* dv \end{aligned}$$

which is also non-positive. It implies that the spectrum of $L^{\mathcal{L}^a}$ in $L^2(M)$ is non-positive.

Let us now write down our assumptions for the collision kernel B :

- B is a tensorial product

$$B = b(\cos \theta) \Phi(|v - v_*|), \quad (1.1)$$

where Φ and b are non-negative functions (this is the case for instance for collision kernels deriving from interaction potentials behaving like inverse-power laws).

- The kinetic part Φ is bounded from below at infinity, i.e

$$\exists R \geq 0, c_\Phi > 0 \mid \forall r \geq R, \Phi(r) \geq c_\Phi. \quad (1.2)$$

This assumption holds for hard potentials (and hard spheres).

- The angular part b satisfies

$$c_b = \inf_{\sigma_1, \sigma_2 \in \mathbb{S}^{N-1}} \int_{\sigma_3 \in \mathbb{S}^{N-1}} \min\{b(\sigma_1 \cdot \sigma_3), b(\sigma_2 \cdot \sigma_3)\} d\sigma_3 > 0. \quad (1.3)$$

This covers all the physical cases.

Remarks: 1. Notice that there is no b left in $Q^{\mathcal{L}^a}$ and $L^{\mathcal{L}^a}$ but the function Φ is definitely the same in both Landau and Boltzmann operators. Therefore the assumptions on the

Landau operator reduce to (1.2). Thus we deal with the so-called “hard potentials” case for the Landau operator, which excludes the Coulomb potential.

2. The assumption that B is a tensorial product is made for a sake of simplicity. Indeed, one could easily adapt the proofs in section 2 to relax this assumption. The price to pay would be a more technical condition on the collision kernel B .

The spectral properties of the linearized Boltzmann and Landau operators have been extensively studied. In particular, there are of crucial interest for perturbative approach issues. For instance, the convergence to equilibrium has been studied in this context, as well as the hydrodynamical limit (see [12]).

On the one hand, for hard potentials, the existence of a spectral gap as soon as the kinetic part of the collision kernel is bounded from below at infinity is a classical result, which can be traced back unto Grad himself. The only method was up to now to work under the assumption of Grad’s angular cutoff, and to apply Weyl’s Theorem to L^B , written as a compact perturbation of a multiplication operator (a very clear presentation of this proof can be found in [5]). The picture of the spectrum obtained for the operator (under Grad’s cutoff assumption) is described by figures 1 and 2 (see [4]).




Figure 1: Spectrum of the collision operator for strictly hard potential with angular cutoff




Figure 2: Spectrum of the collision operator for Maxwell’s molecules with angular cutoff

A similar method has been applied to the Landau linearized operator with hard potential in [7].

On the other hand, for the particular case of Maxwellian molecules (for $L^{\mathcal{B}}$), a complete and explicit diagonalisation has been obtained first by symmetry arguments in [19], and then by Fourier methods in [1]. The spectral gap for the “over-Maxwellian” collision kernel of the Landau linearized operator (i.e collision kernels which are bounded from below by one for Maxwellian molecules) can be derived from results in [11], by a linearization process. Notice also that in the case of the so-called Kac’s equation, an explicit entropy production estimate, based on a cancellation method, was given in [10]; this method can be linearized in order to give explicit spectral gap estimates for “over-quadratic” linearized Kac’s operator (for which the physical meaning is not clear!). Nevertheless we did not manage to adapt this strategy to the Boltzmann operator with hard potentials. Notice however that Wennberg [20] gave an extension of the very first entropy estimates of Desvillettes [8] to allow for hard and soft potentials. His idea has some similarities with ours: to avoid the region in $\mathbb{R}^N \times \mathbb{R}^N$ where $\Phi(|v - v_*|)$ is small.

A specific study of the spectral properties of the linearized operator was made for *non-cutoff* hard potentials in [15]. Nevertheless this article was critically reviewed some years later in [14]. Also another specific study for “radial cutoff potentials” was done in [6].

Finally notice that it is proved in [2] that the Boltzmann linearized operator with soft potential has no spectral gap. The resulting spectrum is described in figure 3 (see [4]).




Figure 3: Spectrum of the collision operator for soft potentials with angular cutoff

But if one allows a loss on the algebraic weight of the norm, it was proved in [13] a “degenerated spectral gap” result of the form

$$\|L^{\mathcal{B}o}h\|_{L^2(M)} \geq C \|h\|_{L^2_\gamma(M)} \quad \forall h \perp \{1; v; |v|^2\},$$

where $\gamma < 0$ is the power of the kernel Φ . It is based on inequalities proven in [2] and Weyl’s Theorem.

However, the perturbative method has drawbacks, all coming from the fact that it does not rely on a physical argument. First it is not explicit, that is the width of the spectral gap is not known, which is problematic when one wants to obtain quantitative estimates of convergence to equilibrium. Secondly it gives no information about how this spectral

gap is sensitive to the perturbation of the collision kernel. Finally approaches based on Weyl's Theorem rely strongly on Grad's cutoff assumption via "Grad's splitting", which means to deal separately with the gain and the loss part of the collision operator.

Our method is geometrical and based on a physical argument. It gives explicit estimates and deals with the whole operator, with or without angular cutoff. Up to our knowledge, as far as spectral gaps are considered, it covers all the results of the above-mentioned articles dealing with hard potentials, with or without angular cutoff. We think likely that this geometrical method could also be adapted to give explicit versions of "degenerated spectral gap" results in the case of soft potentials, even if up to now we did not manage to do it.

We now state our main theorems:

Theorem 1.1 (The Boltzmann linearized operator). *Under the assumptions (1.1), (1.2), (1.3), the Boltzmann entropy dissipation functional $D^{\mathcal{B}_0}$ with $B = \Phi b$ satisfies, for all $h \in L^2(M)$*

$$D^{\mathcal{B}_0}(h) \geq C_{\Phi, b}^{\mathcal{B}_0} D_0^{\mathcal{B}_0}(h), \quad (1.4)$$

where $D_0^{\mathcal{B}_0}(h)$ stands for the entropy dissipation functional with $B_0 \equiv 1$ and

$$C_{\Phi, b}^{\mathcal{B}_0} = \left(\frac{c_\Phi c_b e^{-4R^2}}{32 |\mathbb{S}^{N-1}|} \right)$$

with R, c_Φ, c_b being defined in (1.2), (1.3).

As a consequence we deduce quantitative estimates on the spectral gap of the linearized Boltzmann operator, namely for all $h \in L^2(M)$ orthogonal in $L^2(M)$ to 1, v and $|v|^2$, we have

$$D^{\mathcal{B}_0}(h) \geq C_{\Phi, b}^{\mathcal{B}_0} |\lambda_0^{\mathcal{B}_0}| \|h\|_{L^2(M)}^2. \quad (1.5)$$

Here $\lambda_0^{\mathcal{B}_0}$ is the first non-zero eigenvalue of the linearized Boltzmann operator with $B_0 \equiv 1$ (that is, for Maxwellian molecules with no angular dependence, sometimes called pseudo-Maxwellian molecules) which equals in dimension 3 (see [1])

$$\lambda_0^{\mathcal{B}_0} = -\pi \int_0^\pi \sin^3 \theta d\theta = -\frac{4\pi}{3}.$$

Remark: As an application of this theorem, let us give explicit formulas for the spectral gap $S_\gamma^{\mathcal{B}_0}$ of the Boltzmann linearized operator with $b \geq 1$ and $\Phi(z) = |z|^\gamma$, $\gamma > 0$, in dimension 3. Then $c_b \geq |\mathbb{S}^2|$ and for any given R we can take $c_\Phi = R^\gamma$. Thus we get

$$S_\gamma^{\mathcal{B}_0} \geq \left(\frac{R^\gamma e^{-4R^2}}{32} \right) \frac{4\pi}{3}$$

for any $R > 0$. An easy computation leads to the lower bound

$$S_\gamma^{\mathcal{B}_0} \geq \frac{\pi (\gamma/8)^{\gamma/2} e^{-\gamma/2}}{24}$$

by optimizing the free parameter R .

Theorem 1.2 (The Landau linearized operator). *Under assumptions (1.2), the Landau entropy dissipation functional $D^{\mathcal{L}^a}$ with collision kernel Φ satisfies, for all $h \in L^2(M)$*

$$D^{\mathcal{L}^a}(h) \geq C_{\Phi}^{\mathcal{L}^a} D_0^{\mathcal{L}^a}(h) \quad (1.6)$$

where $D_0^{\mathcal{L}^a}(h)$ stands for the Landau entropy dissipation functional with $\Phi_0 \equiv 1$ and

$$C_{\Phi}^{\mathcal{L}^a} = \left(\frac{c_{\Phi} \beta_R}{8 \alpha_N} \right)$$

with

$$\alpha_N = \int_{\mathbb{R}^{N-1}} e^{-|V|^2} dV, \quad \beta_R = \int_{\{V \in \mathbb{R}^{N-1} \mid |V| \geq 2R\}} e^{-|V|^2} dV.$$

As a consequence we deduce quantitative estimates on the spectral gap of the linearized Landau operator, namely for all $h \in L^2(M)$ orthogonal in $L^2(M)$ to $1, v$ and $|v|^2$, we have

$$D^{\mathcal{L}^a}(h) \geq C_{\Phi}^{\mathcal{L}^a} |\lambda_0^{\mathcal{L}^a}| \|h\|_{L^2(M)}^2. \quad (1.7)$$

Here $\lambda_0^{\mathcal{L}^a}$ is the first non-zero eigenvalue of the linearized Landau operator with $\Phi_0 \equiv 1$ (that is, for Maxwellian molecules).

Moreover in dimension 3, by grazing collisions limit, we can estimate $\lambda_0^{\mathcal{L}^a}$ thanks to the explicit formula on the spectral gap of the Boltzmann linearized operator for Maxwellian molecules

$$|\lambda_0^{\mathcal{L}^a}| \geq 2\pi. \quad (1.8)$$

Remarks: 1. As for the Boltzmann linearized operator, we can deduce from this theorem an explicit formula for a lower bound on the spectral gap $S_{\gamma}^{\mathcal{L}^a}$ for the Landau linearized operator with hard potentials $\Phi(z) = |z|^{\gamma}$, $\gamma > 0$, in dimension 3. We get

$$S_{\gamma}^{\mathcal{L}^a} \geq \left(\frac{R^{\gamma} e^{-4R^2}}{8} \right) 2\pi$$

for any $R > 0$. An easy computation leads to the lower bound

$$S_{\gamma}^{\mathcal{L}^a} \geq \frac{\pi (\gamma/8)^{\gamma/2} e^{-\gamma/2}}{4}$$

by optimizing the free parameter R .

2. The modulus of the first non-zero eigenvalue of the Landau linearized operator for Maxwellian molecules is estimated here by grazing collisions limit. Other methods would have been the linearization of entropy estimates in [11], or to use the decomposition (established in [17]) of the Landau operator for Maxwellian molecules into a Fokker-Planck

part (for which spectral gap is already known) and a spherical diffusion process, which can only increase the spectral gap; and then to linearize the estimate thus obtained.

3. More generally, it is likely that an explicit spectral gap for the Landau linearized operator with hard potentials could be directly computed by existing methods even if up to our knowledge this is the first explicit formula. But Theorem 1.2 is stronger: it says that the property proved on the Boltzmann operator with hard potentials, namely “cancellations for small relative velocities can be neglected as far as entropy production is concerned”, remains true for the Landau linearized operator with hard potentials.

The idea of the proof is to reduce the case of hard potentials (in the generalized sense (1.2)) to the Maxwellian case. The difficulty is to deal with the cancellations of the kinetic collision kernel Φ on the diagonal $v = v_*$.

The starting point is the following inequality which is a corollary of [3, Theorem 2.4]

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} |\xi(x) - \xi(y)|^2 |x - y|^\gamma M(x) M(y) dx dy \\ \geq K_\gamma \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} |\xi(x) - \xi(y)|^2 M(x) M(y) dx dy \end{aligned} \quad (1.9)$$

for $\gamma \geq 0$, ξ some function, and

$$K_\gamma = \frac{1}{4 \int_{\mathbb{R}^N} M} \inf_{x, y \in \mathbb{R}^N} \int_{\mathbb{R}^N} \min \{|x - z|^\gamma, |z - y|^\gamma\} M(z) dz.$$

It was first suggested by Villani [16, Chap. 5, section 1.4], in the context of the study of entropy-entropy dissipation inequalities for the Landau equation with hard potentials, that this inequality could allow to prove that hard potentials reduce to the Maxwellian case as far as convergence to equilibrium is concerned.

The proof of (1.9) relies strongly on the existence of a “triangular inequality” for some function $F(x, y)$ integrated: in (1.9), the function F is simply $|\xi(x) - \xi(y)|^2$ which satisfies

$$F(x, y) \leq 2F(x, z) + 2F(z, y).$$

The main difficulty is hence to obtain such a “triangular inequality” adapted to our case for the Boltzmann linearized operator. It will be discussed in details in section 2 together with the proof of Theorem 1.1. Section 3 will be devoted to the Landau linearized operator: using results of section 2, we will prove Theorem 1.2 thanks to a grazing collision limit.

2 The Boltzmann linearized operator

In this section, we present the proof of inequality (1.4) in Theorem 1.1. In order to “avoid” the diagonal $v \sim v_*$ where Φ is not uniformly bounded from below, we use the following argument: performing a collision with small relative velocity (i.e. for a small

$|v - v_*|$) is the same than performing two collisions with great relative velocity, provided that the pre- and post-collisional velocities are the same. One could summarize the situation in this way: when a collision with small relative velocity occurs, at the same time, two collisions with great relative velocity occur, which give the same pre- and post-collisional velocities, and which produce *at least the same amount of entropy*.

Before proving (1.4), let us begin with a preliminary lemma dealing with the angular part of the collision kernel. This lemma is based on the same geometrical idea as the one we shall use for the treatment of the cancellations of Φ : the introduction of some well-chosen intermediate collision. This first step is made for the sake of simplicity: we show that in the sequel of this section one can set $b \equiv 1$ without restriction. It makes the proof clearer, and simplifies somehow the constants.

Let us denote from now on

$$k(v, v_*, v', v'_*) = [h(v) + h(v_*) - h(v') - h(v'_*)]^2.$$

Lemma 2.1 (Homogeneization of the angular collision kernel b). *Under the assumptions (1.1), (1.2), (1.3), for all $h \in L^2(M)$,*

$$D^{\mathcal{B}o}(h) \geq \frac{c_b}{4|\mathbb{S}^{N-1}|} D_1^{\mathcal{B}o}(h) \tag{2.1}$$

where $D_1^{\mathcal{B}o}$ denotes the entropy dissipation functional with $B = \Phi(|v - v_*|)$ instead of $B = \Phi(|v - v_*|) b(\theta)$.

Remark: This lemma allows to bound from below the entropy dissipation functional by one with an “uniform angular collision kernel”, i.e a constant c_b , even when b is not bounded from below by a positive number uniformly on the sphere. Notice for instance that the condition $c_b > 0$ is satisfied for b having only finite number of 0.

Proof of Lemma 2.1. First, we write down an appropriate representation of the operator. The functional $D^{\mathcal{B}o}$ reads in “ σ -representation”

$$D^{\mathcal{B}o}(h) = \frac{1}{4} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} \Phi(|v - v_*|) b \left(\frac{v - v_*}{|v - v_*|} \cdot \sigma_1 \right) M M_* k(v, v_*, v', v'_*) d\sigma_1 dv_* dv.$$

(for the classical representations of the Boltzmann operator we refer to [16]). Then keeping σ_1 fixed we do the change of variable $(v, v_*) \rightarrow (\frac{v+v_*}{2}, \frac{v-v_*}{2})$, whose jacobian is $(-1/2)^N$. Let us denote $\Omega = \frac{v+v_*}{2}$ and $\Omega' = \frac{v-v_*}{2}$. We obtain

$$D^{\mathcal{B}o}(h) = \frac{2^N}{4} \int_{\Omega \in \mathbb{R}^N} \int_{\Omega' \in \mathbb{R}^N} \int_{\mathbb{S}^{N-1}} \Phi(2|\Omega'|) b \left(\frac{\Omega'}{|\Omega'|} \cdot \sigma_1 \right) k(\Omega + \Omega', \Omega - \Omega', \Omega + |\Omega'| \sigma_1, \Omega - |\Omega'| \sigma_1) e^{-2|\Omega|^2 - 2|\Omega'|^2} d\sigma_1 d\Omega d\Omega'$$

(recall that $|\Omega|^2 + |\Omega'|^2 = \frac{|v|^2 + |v_*|^2}{2}$).

We now write Ω' in spherical coordinates $\Omega' = r \sigma_2$, the other variables being kept fixed, and use Fubini's Theorem

$$D^{\mathcal{B}o}(h) = \frac{2^N}{4} \int_{\Omega \in \mathbb{R}^N} \int_{r \in \mathbb{R}_+} r^{N-1} \Phi(2r) e^{-2|\Omega|^2 - 2r^2} \int_{\sigma_1 \in \mathbb{S}^{N-1}} \int_{\sigma_2 \in \mathbb{S}^{N-1}} b(\sigma_1 \cdot \sigma_2) k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_1, \Omega - r\sigma_1) d\sigma_1 d\sigma_2 dr d\Omega.$$

Now we apply a geometrical idea that we shall also use below in the treatment of cancellations of Φ : namely we add a third artificial variable. Let us thus introduce two collision points u and u_* on the sphere of center Ω and radius r (see figure 4) and replace the collision “ (v, v_*) gives (v', v'_*) ” by the two collisions “ (v, v_*) gives (u, u_*) ” and “ (u, u_*) gives (v', v'_*) ”.




Figure 4: Introduction of an intermediate collision

Then, we shall use the following “triangular” inequality on the collision points:

$$\left[(h(v) + h(v_*)) - (h(v') + h(v'_*)) \right]^2 \leq 2 \left[(h(v) + h(v_*)) - (h(u) + h(u_*)) \right]^2 + 2 \left[(h(u) + h(u_*)) - (h(v') + h(v'_*)) \right]^2. \quad (2.2)$$

So let us add a third “blind” variable σ_3 on the sphere

$$D^{\mathcal{B}o}(h) = \frac{2^N}{4 |\mathbb{S}^{N-1}|} \int_{\Omega \in \mathbb{R}^N} \int_{r \in \mathbb{R}_+} r^{N-1} \Phi(2r) e^{-2|\Omega|^2 - 2r^2} \int_{\sigma_1 \in \mathbb{S}^{N-1}} \int_{\sigma_2 \in \mathbb{S}^{N-1}} \int_{\sigma_3 \in \mathbb{S}^{N-1}} b(\sigma_1 \cdot \sigma_2) k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_1, \Omega - r\sigma_1) d\sigma_1 d\sigma_2 d\sigma_3 dr d\Omega.$$

As variables σ_1 , σ_2 and σ_3 are equivalent, one can change the “blind” variable into either

σ_1 or σ_2 and compute the mean to get

$$\begin{aligned} D^{\mathcal{B}o}(h) &= \frac{2^N}{4 |\mathbb{S}^{N-1}|} \int_{\Omega \in \mathbb{R}^N} \int_{r \in \mathbb{R}_+} r^{N-1} \Phi(2r) e^{-2|\Omega|^2 - 2r^2} \int_{\sigma_1 \in \mathbb{S}^{N-1}} \int_{\sigma_2 \in \mathbb{S}^{N-1}} \int_{\sigma_3 \in \mathbb{S}^{N-1}} \\ &\quad \frac{1}{2} \left[b(\sigma_1 \cdot \sigma_3) k(\Omega + r\sigma_3, \Omega - r\sigma_3, \Omega + r\sigma_1, \Omega - r\sigma_1) \right. \\ &\quad \left. + b(\sigma_2 \cdot \sigma_3) k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_3, \Omega - r\sigma_3) \right] d\sigma_1 d\sigma_2 d\sigma_3 dr d\Omega, \end{aligned}$$

which yields

$$\begin{aligned} D^{\mathcal{B}o}(h) &\geq \frac{2^N}{4 |\mathbb{S}^{N-1}|} \int_{\Omega \in \mathbb{R}^N} \int_{r \in \mathbb{R}_+} \int_{\sigma_1 \in \mathbb{S}^{N-1}} \int_{\sigma_2 \in \mathbb{S}^{N-1}} \int_{\sigma_3 \in \mathbb{S}^{N-1}} \Phi(2r) \\ &\quad \frac{1}{2} \min\{b(\sigma_1 \cdot \sigma_3), b(\sigma_2 \cdot \sigma_3)\} \left[k(\Omega + r\sigma_3, \Omega - r\sigma_3, \Omega + r\sigma_1, \Omega - r\sigma_1) \right. \\ &\quad \left. + k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_3, \Omega - r\sigma_3) \right] e^{-2|\Omega|^2 - 2r^2} d\sigma_1 d\sigma_2 d\sigma_3 dr d\Omega, \end{aligned}$$

The triangular inequality needed on k is

$$\begin{aligned} k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_1, \Omega - r\sigma_1) &\leq 2k(\Omega + r\sigma_3, \Omega - r\sigma_3, \Omega + r\sigma_1, \Omega - r\sigma_1) \\ &\quad + 2k(\Omega + r\sigma_2, \Omega - r\sigma_2, \Omega + r\sigma_3, \Omega - r\sigma_3) \end{aligned}$$

and follows from (2.2). Thus if one sets

$$c_b = \inf_{\sigma_1, \sigma_2 \in \mathbb{S}^{N-1}} \int_{\sigma_3 \in \mathbb{S}^{N-1}} \min\{b(\sigma_1 \cdot \sigma_3), b(\sigma_2 \cdot \sigma_3)\} d\sigma_3,$$

one obtains (going back to the classical representation)

$$\begin{aligned} D^{\mathcal{B}o}(h) &\geq \frac{c_b}{4 |\mathbb{S}^{N-1}|} \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Phi(|v - v_*|) M M_* k(v, v_*, v', v'_*) d\sigma dv_* dv \\ &\geq \frac{c_b}{4 |\mathbb{S}^{N-1}|} D_1^{\mathcal{B}o}(h) \end{aligned}$$

which concludes the proof. \square

Lemma 2.2 (Treatment of the cancellations of Φ). *Under the assumptions (1.2) on Φ , for all $h \in L^2(M)$*

$$D_1^{\mathcal{B}o}(h) \geq \left(\frac{c_\Phi e^{-4R^2}}{8} \right) D_0^{\mathcal{B}o}(h) \quad (2.3)$$

where $D_1^{\mathcal{B}o}$ is the entropy dissipation functional with $B = \Phi(|v - v_*|)$ and $D_0^{\mathcal{B}o}$ is the entropy dissipation functional with $B = 1$.

Proof of Lemma 2.2. We assume here that $b \equiv 1$. Lemma 2.1 indeed shows that this is no restriction modulo a factor $c_b/(4|\mathbb{S}^{N-1}|)$. Let us consider the so-called “ ω -representation” (see [16] again): the vector σ integrated on the sphere becomes $\omega = \frac{v'-v}{|v'-v|}$ and the change of variable changes the angular kernel into

$$\tilde{b}(\theta) = 2^{N-1} \sin^{N-2} \left(\frac{\theta}{2} \right).$$

where $\cos \theta = 2(k \cdot \omega)^2 - 1$ with $k = (v - v_*)/|v - v_*|$. The operator $D_1^{\mathcal{B}o}(h)$ thus becomes

$$D_1^{\mathcal{B}o}(h) = \frac{1}{4} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-1}} \Phi(|v - v_*|) \tilde{b}(\theta) M M_* k(v, v_*, v', v'_*) d\omega dv dv_*.$$

where the velocities v', v'_* are given by

$$v' = v - (v - v_*)\omega, \quad v'_* = v_* + (v - v_*)\omega.$$

Then keeping ω fixed we do the following change of variable

$$v = r_1\omega + V_1, \quad v_* = r_2\omega + V_2$$

with $V_1, V_2 \in \omega^\perp$. The Jacobian of the change of variable is 1 since the decompositions are orthogonal. Finally we obtain the following representation

$$D_1^{\mathcal{B}o}(h) = \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} e^{-r_1^2 - r_2^2} \Phi \left(\sqrt{|r_2 - r_1|^2 + |V_2 - V_1|^2} \right) \tilde{b}(\theta) k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) dr_1 dr_2 dV_2 dV_1 d\omega.$$

Assume that Φ is non-decreasing. This is no restriction since $\Phi \geq \tilde{\Phi}$, with

$$\tilde{\Phi}(r) = \inf_{r' \geq r} \Phi(r'),$$

and $\tilde{\Phi}$ satisfies assumption (1.2) with the same constant as Φ . This monotonicity yields

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} e^{-r_1^2 - r_2^2} \Phi(|r_2 - r_1|) \tilde{b}(\theta) k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) dr_1 dr_2 dV_2 dV_1 d\omega.$$

We now introduce two collision points u and u_* (see figure 5) and replace the collision “ (v, v_*) gives (v', v'_*) ” by the two collisions “ (v, u_*) gives (v', u) ” and “ (u, v_*) gives (u_*, v') ”.

Then, we shall use the following “triangular” inequality on the collision points:

$$\begin{aligned} [(h(v) + h(v_*)) - (h(v') + h(v'_*))]^2 &\leq 2[(h(v) + h(u_*)) - (h(u) + h(v'_*))]^2 \\ &\quad + 2[(h(u) + h(v_*)) - (h(v') + h(u_*))]^2. \end{aligned}$$




Figure 5: Introduction of an intermediate collision

Recall that $\int_{\mathbb{R}} e^{-r^2} dr = \sqrt{\pi}$. Let us add a third artificial integration variable r_3 on \mathbb{R}

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{4\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \int_{r_3 \in \mathbb{R}} \Phi(|r_2 - r_1|) \tilde{b}(\theta_{1,2}) k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-r_1^2 - r_2^2 - r_3^2} dr_1 dr_2 dr_3 dV_1 dV_2 d\omega.$$

From now on, indexes of θ denote the points which are chosen to compute the angle. Now we rename r_1, r_2, r_3 first in r_1, r_3, r_2 , secondly in r_3, r_2, r_1 and we take the mean of these two quantities. We get

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{8\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \int_{r_3 \in \mathbb{R}} e^{-r_1^2 - r_2^2 - r_3^2} \left[\tilde{b}(\theta_{1,3}) \Phi(|r_3 - r_1|) k(r_1\omega + V_1, r_3\omega + V_2, r_3\omega + V_1, r_1\omega + V_2) + \tilde{b}(\theta_{2,3}) \Phi(|r_2 - r_3|) k(r_3\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_3\omega + V_2) \right] dr_1 dr_2 dr_3 dV_1 dV_2 d\omega.$$

Then,

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{8\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \int_{r_3 \in \mathbb{R}} \min \{ \tilde{b}(\theta_{1,3}) \Phi(|r_3 - r_1|), \tilde{b}(\theta_{2,3}) \Phi(|r_2 - r_3|) \} \left[k(r_1\omega + V_1, r_3\omega + V_2, r_3\omega + V_1, r_1\omega + V_2) + k(r_3\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_3\omega + V_2) \right] e^{-r_1^2 - r_2^2 - r_3^2} dr_1 dr_2 dr_3 dV_1 dV_2 d\omega. \quad (2.4)$$

Now we use the following triangular inequality above-mentioned which means translated

on k

$$k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) \leq 2k(r_1\omega + V_1, r_3\omega + V_2, r_3\omega + V_1, r_1\omega + V_2) \\ + 2k(r_3\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_3\omega + V_2).$$

Plugging it in (2.4), we get

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{16\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \int_{r_3 \in \mathbb{R}} \\ \min \{ \tilde{b}(\theta_{1,3}) \Phi(|r_3 - r_1|), \tilde{b}(\theta_{2,3}) \Phi(|r_2 - r_3|) \} \\ k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-r_1^2 - r_2^2 - r_3^2} dr_1 dr_2 dr_3 dV_1 dV_2 d\omega,$$

which yields

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{16\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \\ \left(\int_{r_3 \in \mathbb{R}} \min \{ \tilde{b}(\theta_{1,3}) \Phi(|r_3 - r_1|), \tilde{b}(\theta_{2,3}) \Phi(|r_2 - r_3|) \} e^{-r_3^2} dr_3 \right) \\ k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-r_1^2 - r_2^2} dr_1 dr_2 dV_1 dV_2 d\omega.$$

We now restrict the domain of integration for r_3 to the set

$$\mathcal{D}_{r_1, r_2} = \{ r_3 \in \mathbb{R} \mid |r_3 - r_1| \geq |r_1 - r_2| \text{ and } |r_2 - r_3| \geq |r_1 - r_2| \}.$$

Since \tilde{b} is non-decreasing, and

$$\cos \theta = \frac{|V_1 - V_2|^2 - |r_1 - r_2|^2}{|V_1 - V_2|^2 + |r_1 - r_2|^2}$$

which is non-increasing with respect to $|r_1 - r_2|$ when V_1, V_2 are kept frozen, it is easy to check that on this domain we have $\theta_{1,3} \geq \theta_{1,2}$ and $\theta_{2,3} \geq \theta_{1,2}$ and thus $\tilde{b}(\theta_{1,3}) \geq \tilde{b}(\theta_{1,2})$ and $\tilde{b}(\theta_{2,3}) \geq \tilde{b}(\theta_{1,2})$. Therefore we get

$$D_1^{\mathcal{B}o}(h) \geq \frac{1}{16\sqrt{\pi}} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} \\ \left(\int_{r_3 \in \mathcal{D}_{r_1, r_2}} \min \{ \Phi(|r_3 - r_1|), \Phi(|r_2 - r_3|) \} e^{-r_3^2} dr_3 \right) \tilde{b}(\theta_{1,2}) \\ k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-r_1^2 - r_2^2} dr_1 dr_2 dV_1 dV_2 d\omega.$$

Under assumption (1.2), an easy computation leads to

$$\left(\int_{r_3 \in \mathcal{D}_{r_1, r_2}} \min \{ \Phi(|r_3 - r_1|), \Phi(|r_2 - r_3|) \} e^{-|r_3|^2} dr_3 \right) \geq c_\Phi \sqrt{\pi} e^{-4R^2} > 0$$

as soon as $|r_1 - r_2| \leq R$, i.e

$$\left(\int_{r_3 \in \mathcal{D}_{r_1, r_2}} \min \{ \Phi(|r_3 - r_1|), \Phi(|r_2 - r_3|) \} e^{-|r_3|^2} dr_3 \right) \geq c_\Phi \sqrt{\pi} e^{-4R^2} 1_{|r_1 - r_2| \leq R}.$$

By taking the mean of this estimate and the one obtained by replacing Φ by its bound from below $c_\Phi 1_{r \geq R}$, we deduce that

$$D_1^{\mathcal{B}o}(h) \geq \min \left(\frac{c_\Phi e^{-4R^2}}{8}, \frac{c_\Phi}{2} \right) \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} e^{-|V_1|^2 - |V_2|^2} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) \tilde{b}(\theta) e^{-r_1^2 - r_2^2} dr_1 dr_2 dV_1 dV_2 d\omega.$$

If we now go back to the classical representation and simplify the minimum, we obtain

$$\begin{aligned} D_1^{\mathcal{B}o}(h) &\geq \left(\frac{c_\Phi e^{-4R^2}}{8} \right) \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} M M_* k(v, v_*, v', v'_*) d\sigma dv dv_* \\ &= \left(\frac{c_\Phi e^{-4R^2}}{8} \right) D_0^{\mathcal{B}o}(h) \end{aligned}$$

which concludes the proof of the lemma. \square

The proof of Theorem 1.1 is a straightforward consequence of inequalities (2.1) and (2.3).

3 The Landau linearized operator

We now prove Theorem 1.2. The idea here is to take the grazing collisions limit in some inequalities on the Boltzmann linearized operator obtained thanks to the geometrical method used in Section 2. In fact, the most natural idea would have been to look for a geometrical property on the Landau linearized operator similar to the triangular inequality used for the Boltzmann linearized operator. But as collision circles reduce to lines in the grazing limit, the triangular inequality becomes trivial, and thus does not seem sufficient to apply the method of section 2. It could be linked to the fact that in the grazing collisions limit one loses some information on the geometry of the collision.

The problem that has to be tackled is to keep track of the angular collision kernel b . In fact we need it only for particular b , namely

$$b_\varepsilon(\theta) = \frac{j_\varepsilon(\theta)}{\varepsilon^2 \sin^{N-2} \frac{\theta}{2}} \quad (3.1)$$

where $j_\varepsilon(\theta) = j(\theta/\varepsilon)/\varepsilon$ is a sequence of mollifiers (approximating $\delta_{\theta=0}$) with compact support in $[0, \pi/2]$ and non-increasing on this interval. It is easy to see that $\tilde{b}_\varepsilon = 2^{N-1} \sin^{N-2} \left(\frac{\theta}{2} \right) b_\varepsilon$ is also non-increasing on $[0, \pi]$. Following the same strategy as in Lemma 2.2 but keeping track of the angular part of the collision kernel, one obtains

Lemma 3.1. Under the assumptions (1.1), (1.2), (1.3), plus the assumption that $\tilde{b} = 2^{N-1} \sin^{N-2}(\frac{\theta}{2}) b$ is non-increasing, one gets for all $h \in L^2(M)$

$$D_{b,\Phi}^{\mathcal{B}_o}(h) \geq \left(\frac{c_\Phi \beta_R}{8\alpha_N} \right) D_{b,1}^{\mathcal{B}_o}(h) \quad (3.2)$$

with

$$\alpha_N = \int_{\mathbb{R}^{N-1}} e^{-|V|^2} dV, \quad \beta_R = \int_{\{V \in \mathbb{R}^{N-1} \mid |V| \geq 2R\}} e^{-|V|^2} dV.$$

Here $D_{b,\Phi}^{\mathcal{B}_o}$ stands for the entropy dissipation functional with $B = \Phi b$ and $D_{b,1}^{\mathcal{B}_o}$ stands for the entropy dissipation functional with $B = b$.

Proof of Lemma 3.1. The geometrical idea of Lemma 2.2 can be applied to the variables V_1, V_2 . Let us thus introduce two collision points u and u_* (see figure 6) and replace the collision “ (v, v_*) gives (v', v'_*) ” by the two collisions “ (v, u_*) gives (v', u) ” and “ (u, v_*) gives (u_*, v'_*) ”.




Figure 6: Introduction of an intermediate collision

Then, we shall use the following “triangular” inequality on the collision points:

$$\begin{aligned} [(h(v) + h(v_*)) - (h(v') + h(v'_*))]^2 &\leq 2[(h(v) + h(u_*)) - (h(v') + h(u))]^2 \\ &\quad + 2[(h(u) + h(v_*)) - (h(u_*) + h(v'_*))]^2. \end{aligned}$$

Now we introduce an artificial third variable V_3 on ω^\perp . Let us denote

$$\alpha_N = \int_{\mathbb{R}^{N-1}} e^{-|V|^2} dV$$

By inverting either V_1 and V_3 or V_2 and V_3 , taking the mean, and using the “triangular” inequality above-mentioned we get

$$D_{b,\Phi}^{\mathcal{B}_o}(h) \geq \frac{1}{16\alpha_N} \int_{\mathbb{S}^{N-1}} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} e^{-r_1^2 - r_2^2} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} \left(\int_{V_3 \in \omega^\perp} \min \left\{ \tilde{b}(\theta_{1,3}) \Phi(|V_3 - V_1|), \tilde{b}(\theta_{2,3}) \Phi(|V_2 - V_3|) \right\} e^{-|V_3|^2} dV_3 \right) k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-|V_1|^2 - |V_2|^2} dr_1 dr_2 dV_1 dV_2 d\omega.$$

Let us now restrict the integration along V_3 to the domain

$$\mathcal{D}_{V_1, V_2} = \{V_3 \mid |V_3 - V_1| \geq |V_1 - V_2| \text{ and } |V_2 - V_3| \geq |V_1 - V_2|\}.$$

Then since the expression

$$\cos \theta = \frac{|V_1 - V_2|^2 - |r_1 - r_2|^2}{|V_1 - V_2|^2 + |r_1 - r_2|^2}$$

is non-decreasing according to $|V_1 - V_2|$ when r_1, r_2 are kept frozen, and \tilde{b} is non-increasing, we get $\theta_{1,3} \leq \theta_{1,2}$ and $\theta_{2,3} \leq \theta_{1,2}$ (see figure 6) and so $\tilde{b}(\theta_{1,3}) \geq \tilde{b}(\theta_{1,2})$ and $\tilde{b}(\theta_{2,3}) \geq \tilde{b}(\theta_{1,2})$. Consequently

$$D_{b,\Phi}^{\mathcal{B}_o}(h) \geq \frac{1}{16\alpha_N} \int_{\mathbb{S}^{N-1}} \int_{r_1 \in \mathbb{R}} \int_{r_2 \in \mathbb{R}} dr_2 e^{-r_1^2 - r_2^2} \int_{V_1 \in \omega^\perp} \int_{V_2 \in \omega^\perp} \left(\int_{V_3 \in \mathcal{D}_{V_1, V_2}} \min \{ \Phi(|V_3 - V_1|), \Phi(|V_2 - V_3|) \} e^{-|V_3|^2} dV_3 \right) \tilde{b}(\theta_{1,2}) k(r_1\omega + V_1, r_2\omega + V_2, r_2\omega + V_1, r_1\omega + V_2) e^{-|V_1|^2 - |V_2|^2} dr_1 dr_2 dV_1 dV_2 d\omega.$$

Under assumption (1.2), an easy computation leads to

$$\left(\int_{V_3 \in \mathcal{D}_{V_1, V_2}} \min \{ \Phi(|V_3 - V_1|), \Phi(|V_2 - V_3|) \} e^{-|V_3|^2} dV_3 \right) \geq c_\Phi \int_{\{V \in \mathbb{R}^{N-1} \mid |V| \geq 2R\}} e^{-|V|^2} dV = c_\Phi \beta_R > 0$$

as soon as $|V_1 - V_2| \leq R$, i.e

$$\left(\int_{V_3 \in \mathcal{D}_{V_1, V_2}} \min \{ \Phi(|V_3 - V_1|), \Phi(|V_2 - V_3|) \} e^{-|V_3|^2} dV_3 \right) \geq c_\Phi \beta_R \mathbf{1}_{|V_1 - V_2| \leq R}.$$

Taking the mean of this estimate and the one obtained by the trivial lower bound $\Phi(r) \geq c_\Phi \mathbf{1}_{\{r \geq R\}}$, we get in the end

$$D_{b,\Phi}^{\mathcal{B}_o}(h) \geq \min \left(\frac{c_\Phi \beta_R}{8\alpha_N}, \frac{c_\Phi}{2} \right) \frac{1}{4} \int_{\mathbb{S}^{N-1}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} b(\theta) M M_* [h'_* + h' - h - h_*]^2 d\sigma dv_* dv,$$

which yields

$$D_{b,\Phi}^{\mathcal{B}o}(h) \geq \left(\frac{c_\Phi \beta_R}{8\alpha_N} \right) D_{b,1}^{\mathcal{B}o}(h)$$

and concludes the proof of the lemma. \square

We now have to take the grazing collisions limit in the entropy dissipation functional to prove inequality (1.6) of Theorem 1.2 (this limit is essentially well-known, see for instance [9]).

Lemma 3.2. *Let us consider b_ε as defined in (3.1) and Φ satisfying assumption (1.2). Then for a given $h \in L^2(M)$,*

$$D_{b_\varepsilon,\Phi}^{\mathcal{B}o}(h) \xrightarrow{\varepsilon \rightarrow 0} c_{N,j} D_\Phi^{\mathcal{L}a}(h)$$

where

$$c_{N,j} = \frac{2^{N-5} |\mathbb{S}^{N-2}|}{N-1} \left(\int_0^\pi j(\chi) \chi^2 d\chi \right)$$

depends only on the dimension N and the mollifier j . $D_{b_\varepsilon,\Phi}^{\mathcal{B}o}$ stands for the Boltzmann entropy dissipation functional with $B = \Phi b_\varepsilon$, and $D_\Phi^{\mathcal{L}a}$ stands for the Landau entropy dissipation functional with collision kernel Φ .

Proof of Lemma 3.2. The idea of the proof is to expand the expression for small ε and is very similar to what is done in [9]. Let us write the angular vector σ

$$\sigma = \frac{v - v_*}{|v - v_*|} \cos(\theta) + \mathbf{n} \sin(\theta),$$

where \mathbf{n} is a unit vector in $(v - v_*)^\perp$. Therefore, we shall write

$$\begin{aligned} D_{b_\varepsilon,\Phi}^{\mathcal{B}o}(h) &= \frac{1}{4} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \Phi(|v - v_*|) M M_* \int_{\mathbb{S}^{N-2}((v-v_*)^\perp)} \int_{\theta=0}^\pi b_\varepsilon(\theta) \\ &\quad \left[h \left(v - \frac{v - v_*}{2} (1 - \cos(\theta)) + \frac{|v - v_*|}{2} \mathbf{n} \sin(\theta) \right) \right. \\ &\quad + h \left(v_* + \frac{v - v_*}{2} (1 - \cos(\theta)) - \frac{|v - v_*|}{2} \mathbf{n} \sin(\theta) \right) \\ &\quad \left. - h(v) - h(v_*) \right]^2 \sin^{N-2} \theta d\theta d\mathbf{n} dv dv_*, \end{aligned}$$

where $\mathbb{S}^{N-2}((v - v_*)^\perp)$ denotes the unit sphere in $(v - v_*)^\perp$. Let us now focus on the integral on θ

$$\begin{aligned} &\int_{\theta=0}^\pi b_\varepsilon(\theta) \left[h \left(v - \frac{v - v_*}{2} (1 - \cos(\theta)) + \frac{|v - v_*|}{2} \mathbf{n} \sin(\theta) \right) \right. \\ &\quad \left. + h \left(v_* + \frac{v - v_*}{2} (1 - \cos(\theta)) - \frac{|v - v_*|}{2} \mathbf{n} \sin(\theta) \right) - h(v) - h(v_*) \right]^2 \sin^{N-2} \theta d\theta, \end{aligned}$$

and make the change of variables $\chi = \theta/\varepsilon$. We get

$$\int_{\chi=0}^{\pi} \frac{\sin^{N-2}(\varepsilon \chi)}{\sin^{N-2}(\frac{\varepsilon \chi}{2})} \frac{j(\chi)}{\varepsilon^2} \left[h \left(v - \frac{v-v_*}{2}(1 - \cos(\varepsilon \chi)) + \frac{|v-v_*|}{2} \mathbf{n} \sin(\varepsilon \chi) \right) \right. \\ \left. + h \left(v_* + \frac{v-v_*}{2}(1 - \cos(\varepsilon \chi)) - \frac{|v-v_*|}{2} \mathbf{n} \sin(\varepsilon \chi) \right) - h(v) - h(v_*) \right]^2 d\chi$$

i.e for small ε ,

$$\int_{\chi=0}^{\pi} (2^{N-2} + O(\varepsilon)) \frac{j(\chi)}{\varepsilon^2} \left(\frac{|v-v_*|}{2} \right)^2 \left[\varepsilon \chi \mathbf{n} \cdot (\nabla_v h(v) - \nabla_{v_*} h(v_*)) + O(\varepsilon^2 \chi^2) \right]^2 d\chi,$$

which writes

$$|v-v_*|^2 \int_{\chi=0}^{\pi} 2^{N-4} j(\chi) \chi^2 \left[\mathbf{n} \cdot (\nabla_v h(v) - \nabla_{v_*} h(v_*)) \right]^2 d\chi + O(\varepsilon) \\ = 2^{N-4} \left(\int_0^{\pi} j(\chi) \chi^2 d\chi \right) |v-v_*|^2 \left[\mathbf{n} \cdot (\nabla_v h(v) - \nabla_{v_*} h(v_*)) \right]^2 + O(\varepsilon).$$

As the unit vector \mathbf{n} is orthogonal to $\frac{v-v_*}{|v-v_*|}$, we can introduce here the orthogonal projection onto $(v-v_*)^\perp$

$$D_{b_\varepsilon, \Phi}^{\mathcal{B}o}(h) = \frac{2^{N-4}}{4} \left(\int_0^{\pi} j(\chi) \chi^2 d\chi \right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \int_{\mathbb{S}^{N-2}((v-v_*)^\perp)} \Phi(|v-v_*|) |v-v_*|^2 \\ \left[\mathbf{n} \cdot \Pi_{(v-v_*)^\perp} (\nabla_v h(v) - \nabla_{v_*} h(v_*)) \right]^2 M M_* d\mathbf{n} dv dv_* + O(\varepsilon).$$

It is straightforward to see that

$$\int_{\mathbb{S}^{N-2}((v-v_*)^\perp)} (\mathbf{n} \cdot u)^2 d\mathbf{n} = \zeta_N \|u\|^2$$

with, for any $\mathbf{u} \in \mathbb{S}^{N-2}$

$$\zeta_N = \int_{\mathbb{S}^{N-2}} (\mathbf{u} \cdot \mathbf{n})^2 d\mathbf{n} = \frac{|\mathbb{S}^{N-2}|}{N-1}.$$

Thus we get in the end

$$D_{b_\varepsilon, \Phi}^{\mathcal{B}o}(h) = \frac{|\mathbb{S}^{N-2}| 2^{N-4}}{4(N-1)} \left(\int_0^{\pi} j(\chi) \chi^2 d\chi \right) \\ \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} |v-v_*|^2 \Phi(|v-v_*|) M M_* \|\nabla_v h(v) - \nabla_{v_*} h(v_*)\|^2 dv_* dv + O(\varepsilon) \\ = \frac{|\mathbb{S}^{N-2}| 2^{N-5}}{N-1} \left(\int_0^{\pi} j(\chi) \chi^2 d\chi \right) D_{\Phi}^{\mathcal{L}^a}(h) + O(\varepsilon).$$

This concludes the proof of Lemma 3.2. □

Coming back to the proof of Theorem 1.2, we first prove (1.6): we write down inequality (3.2) on $D_{\Phi, b_\varepsilon}^{\mathcal{B}^o}$ since \tilde{b}_ε is non-increasing, and we apply Lemma 3.2 on each term, which gives

$$D^{\mathcal{L}^a}(h) \geq C_{\Phi}^{\mathcal{L}^a} D_0^{\mathcal{L}^a}(h),$$

where

$$C_{\Phi}^{\mathcal{L}^a} = \left(\frac{c_{\Phi} \beta_R}{8\alpha_N} \right).$$

Inequality (1.7) follows immediatly.

It remains to prove the lower bound (1.8) on the first non-zero eigenvalue of the Landau linearized operator for Maxwellian molecules in dimension 3. Let us denote by $\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}$ the first non-zero eigenvalue for the Boltzmann linearized operator with $B = b_\varepsilon$: for all $h \in L^2(M)$ orthogonal in $L^2(M)$ to $1, v, |v|^2$,

$$D_{b_\varepsilon}^{\mathcal{B}^o}(h) \geq |\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}| \|h\|_{L^2(M)}^2.$$

We apply Lemma 3.2 to this inequality which leads to

$$D_0^{\mathcal{L}^a}(h) \geq \frac{\lim_{\varepsilon \rightarrow 0} |\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}|}{c_{3, j}} \|h\|_{L^2(M)}^2$$

for all $h \in L^2(M)$ orthogonal in $L^2(M)$ to $1, v, |v|^2$. An explicit formula for $|\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}|$ is given in [1]

$$|\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}| = \pi \int_0^\pi \sin^3(\theta) b_\varepsilon(\theta) d\theta$$

and thus

$$\lim_{\varepsilon \rightarrow 0} |\lambda_{0, b_\varepsilon}^{\mathcal{B}^o}| = 2\pi \left(\int_0^\pi j(\chi) \chi^2 d\chi \right)$$

which concludes the proof.

Acknowledgment : Support by the European network HYKE, funded by the EC as contract HPRN-CT-2002-00282, is acknowledged.

References

- [1] BOBYLĚV, A. V. The theory of the nonlinear spatially uniform Boltzmann equation for Maxwell molecules. In *Mathematical physics reviews, Vol. 7*, vol. 7 of *Soviet Sci. Rev. Sect. C Math. Phys. Rev.* Harwood Academic Publ., Chur, 1988, pp. 111–233.
- [2] CAFLISCH, R. E. The Boltzmann equation with a soft potential. II. Nonlinear, spatially-periodic. *Comm. Math. Phys.* 74, 2 (1980), 97–109.

- [3] CARRILO, J., MCCANN, R., AND VILLANI, C. Kinetic equilibration rates for granular media. Preprint, 2001.
- [4] CERCIGNANI, C. *The Boltzmann equation and its applications*, vol. 67 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1988.
- [5] CERCIGNANI, C., ILLNER, R., AND PULVIRENTI, M. *The Mathematical Theory of Dilute Gases*. Springer, 1994.
- [6] CZECHOWSKY, Z., AND PALCZEWSKY, A. Spectrum of the Boltzmann equation collision operator for radial cut-off potentials. *Bull. Acad. Polon. Sci.* 28 (1980), 387–396.
- [7] DEGOND, P., AND LEMOU, M. Dispersion relations for the linearized Fokker-Planck equation. *Arch. Rational Mech. Anal.* 138, 2 (1997), 137–167.
- [8] DESVILLETES, L. Entropy dissipation rate and convergence in kinetic equations. *Comm. Math. Phys.* 123, 4 (1989), 687–702.
- [9] DESVILLETES, L. On asymptotics of the Boltzmann equation when the collisions become grazing. *Transport Theory Statist. Phys.* 21, 3 (1992), 259–276.
- [10] DESVILLETES, L. Convergence towards the thermodynamical equilibrium. In *Trends in applications of mathematics to mechanics (Nice, 1998)*, vol. 106 of *Chapman & Hall/CRC Monogr. Surv. Pure Appl. Math.* Chapman & Hall/CRC, Boca Raton, FL, 2000, pp. 115–126.
- [11] DESVILLETES, L., AND VILLANI, C. On the spatially homogeneous Landau equation for hard potentials. II. H -theorem and applications. *Comm. Partial Differential Equations* 25, 1-2 (2000), 261–298.
- [12] ELLIS, R. S., AND PINSKY, M. A. The first and second fluid approximations to the linearized Boltzmann equation. *J. Math. Pures Appl. (9)* 54 (1975), 125–156.
- [13] GOLSE, F., AND POUPAUD, F. Un résultat de compacité pour l'équation de Boltzmann avec potentiel mou. Application au problème de demi-espace. *C. R. Acad. Sci. Paris Sér. I Math.* 303, 12 (1986), 583–586.
- [14] KLAUS, M. Boltzmann collision operator without cut-off. *Helv. Phys. Acta* 50, 6 (1977), 893–903.
- [15] PAO, Y. P. Boltzmann collision operator with inverse-power intermolecular potentials. I, II. *Comm. Pure Appl. Math.* 27 (1974), 407–428, 559–581.
- [16] VILLANI, C. A survey of mathematical topics in kinetic theory. *Handbook of fluid mechanics*, S. Friedlander and D. Serre, Eds. Elsevier Publ., 2002.

- [17] VILLANI, C. On the spatially homogeneous Landau equation for Maxwellian molecules. *Math. Models Methods Appl. Sci.* 8, 6 (1998), 957–983.
- [18] VILLANI, C. Contribution à l'étude mathématique des collisions en théorie cinétique. Master's thesis, Univ. Paris-Dauphine France, 2000.
- [19] WANG CHANG, C. S., UHLENBECK, G. E., AND DE BOER, J. In *Studies in Statistical Mechanics, Vol. V*. North-Holland, Amsterdam, 1970.
- [20] WENNBERG, B. On an entropy dissipation inequality for the Boltzmann equation. *C. R. Acad. Sci. Paris Sér. I Math.* 315, 13 (1992), 1441–1446.

C. BARANGER

CMLA, ENS CACHAN
61 AVENUE DU PRÉSIDENT WILSON
94235 CACHAN CEDEX
FRANCE

E-MAIL: Celine.Baranger@cmla.ens-cachan.fr

C. MOUHOT

UMPA, ENS LYON
46 ALLÉE D'ITALIE
69364 LYON CEDEX 07
FRANCE

E-MAIL: cmouhot@umpa.ens-lyon.fr