



HAL
open science

Hydrodynamic turbulence and intermittent random fields

Jean Duchon, Raoul Robert, Vincent Vargas

► **To cite this version:**

Jean Duchon, Raoul Robert, Vincent Vargas. Hydrodynamic turbulence and intermittent random fields. 2006. hal-00088376v1

HAL Id: hal-00088376

<https://hal.science/hal-00088376v1>

Preprint submitted on 1 Aug 2006 (v1), last revised 7 Jul 2008 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HYDRODYNAMIC TURBULENCE AND INTERMITTENT RANDOM FIELDS

Jean Duchon, Raoul Robert
Institut Fourier, universit  Grenoble 1, UMR CNRS 5582,
100, rue des Math matiques, BP 74, 38402 Saint-Martin d'H res cedex, France
e-mail: Jean.Duchon@ujf-grenoble.fr, Raoul.Robert@ujf-grenoble.fr

Vincent Vargas
Universit  Paris 7, Math matiques, case 7012,
2, place Jussieu, 75251 Paris, France
e-mail: vargas@math.jussieu.fr

ABSTRACT. In this article, we construct two families of nonsymmetrical multifractal fields. One of these families is used for the modelization of the velocity field of turbulent flows.

Short Title. Hydrodynamic turbulence and intermittent random fields

Key words and phrases. Gaussian multiplicative chaos, intermittency.

MSC 2000 subject classifications: 60G60, 60G57, 28A80

1. INTRODUCTION

Roughly observed, some random phenomena seem perfectly scale invariant. This is the case for the velocity field of turbulent flows or the (logarithm of) evolution in time of the price of a financial asset. However, a more precise empirical study of these phenomena displays in fact a weakened form of scale invariance commonly called multifractal scale invariance or intermittency (the exponent which governs the power law scaling of the process or field is no longer linear). An important question is therefore to construct intermittent random fields which exhibit the observed characteristics.

Following the work of Kolmogorov and Obukhov ([9], [12]) on the energy dissipation in turbulent flows, Mandelbrot introduced in [10] a "limit-lognormal" model to describe turbulent dissipation or the volatility of a financial asset. This model was rigorously defined and studied in a mathematical framework by Kahane in [8]; more precisely, Kahane constructed a random measure called gaussian multiplicative chaos. A natural extension of this work is to use gaussian multiplicative chaos to construct a field (or a process in the financial case) which describes the whole phenomenon: the velocity field in turbulent flows (the price of an asset on a financial market). This extension was first performed by Mandelbrot himself who proposed to modelize the price of a financial asset with a time changed Brownian motion,

Partially supported by CNRS (UMR 7599 "Probabilit s et Mod les Al atoires").

the time change being random and independent of the Brownian motion. In [2], the authors proposed for the time change to take the primitive of multiplicative chaos: this gives the so called multifractal random walk model (MRW) (Bacry and Muzy later generalized the construction of the MRW model in [3]). The obtained process accounts for many observed properties of financial assets.

The inconvenient of the above construction and of the MRW model is that the laws of the increments are symmetrical. In the case of finance, this is in contradiction with the skewness property observed for certain asset prices. In the case of turbulence, the laws of the increments must be nonsymmetrical: it is a theoretical necessity and stems from the dissipation of the kinetic energy ([7]). In light of these observations, we are naturally led to construct random fields which generalize to any dimension such process and which present multifractal scale invariance as well as nonsymmetrical increments.

We will answer a very natural question: how can one obtain a two parameter family of multifractal fields with nonsymmetrical increments by perturbing a given scale invariant gaussian random field on \mathbb{R}^d ? Finally, in the last part we will mention the difficulties which arise in trying to construct an incompressible multifractal velocity field that verifies the 4/5-law of Kolmogorov with positive dissipation.

2. NOTATIONS AND PRELIMINARY RESULTS

2.1. The underlying gaussian field. Let $dW_0(x)$ denote the gaussian white noise on \mathbb{R}^d and $\varphi : \mathbb{R}^d \rightarrow [0, 1]$ denote a C^∞ , radially symmetric function worth 1 for $|x| \leq 1$ and 0 for $|x| > 2$. We also introduce a fixed correlation scale $R > 0$ and α a number which satisfies

$$d/2 < \alpha < d/2 + 1. \quad (2.1)$$

We define the gaussian field \mathcal{X}^g by the following formula:

$$\mathcal{X}^g(x) = \int_{\mathbb{R}^d} \frac{\varphi_R(x-y)}{|x-y|^{d-\alpha}} dW_0(y), \quad (2.2)$$

where we set the following notation:

$$\varphi_R(x) = R^{d/2-\alpha} \varphi\left(\frac{x}{R}\right).$$

It is easy to show that (2.2) defines a homogeneous, isotropic gaussian field which is almost surely holderian of order $< \alpha - d/2$. Note that condition (2.1) implies that the integrand in (2.2) is square integrable and the $R^{d/2-\alpha}$ factor ensures that the field is dimensionless.

scaling property. Let e be a unitary vector and $\lambda > 0$. we have the following identity in law:

$$\mathcal{X}^g(x + \lambda e) - \mathcal{X}^g(x) \stackrel{(law)}{=} \int_{\mathbb{R}^d} \left(\frac{\varphi_R(y - \lambda e)}{|y - \lambda e|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}} \right) dW_0(y).$$

From the gaussianity of the above law, we deduce that for all $q > 0$, there exists $c_q > 0$ such that:

$$E(|\mathcal{X}^g(x + \lambda e) - \mathcal{X}^g(x)|^q) = \sigma_{\lambda e}^q c_q,$$

with

$$\sigma_{\lambda e}^2 \underset{\lambda \rightarrow 0}{\sim} \left(\frac{\lambda}{R}\right)^{2\alpha-d} \int_{\mathbb{R}^d} \left(\frac{1}{|y-e|^{d-a}} - \frac{1}{|y|^{d-a}}\right)^2 dy.$$

We thus derive the following scaling

$$E(|\mathcal{X}^g(x + \lambda e) - \mathcal{X}^g(x)|^q) \underset{\lambda \rightarrow 0}{\sim} \left(\frac{\lambda}{R}\right)^{q(\alpha-d/2)} C_q,$$

where the constant C_q is independent of e . One says that $(\mathcal{X}^g(x))_{x \in \mathbb{R}^d}$ is at small scales monofractal with scaling exponent $\alpha - d/2$.

A field $(\mathcal{X}(x))_{x \in \mathbb{R}^d}$ is multifractal if there exists a non linear function ζ_q such that:

$$E((\mathcal{X}(x + \lambda e) - \mathcal{X}(x))^q) \underset{\lambda \rightarrow 0}{\sim} \left(\frac{\lambda}{R}\right)^{\zeta_q} C_q.$$

We call ζ_q the structural function of the field $(\mathcal{X}(x))_{x \in \mathbb{R}^d}$.

2.2. Outline of the construction of multifractal fields from the field \mathcal{X}^g .

Our construction is inspired by the work of Kahane in [8]. Let $\epsilon > 0$ and $X^\epsilon(y)$ a regular family of gaussian fields (not necessarily independent of dW_0). We consider a family of fields \mathcal{X}^ϵ defined by:

$$\mathcal{X}^\epsilon(x) = \int_{\mathbb{R}^d} \frac{\varphi_R(x-y)}{|x-y|_\epsilon^{d-\alpha}} e^{X^\epsilon(y)-C_\epsilon} dW_0(y) \quad (2.3)$$

($|x-y|_\epsilon$ is defined in the next subsection and is given by a standard convolution). For an appropriate family X^ϵ , we show that it is possible to find constants C_ϵ such that \mathcal{X}^ϵ tends to a non trivial field \mathcal{X} as ϵ tends to 0. If one chooses X^ϵ independent of dW_0 , we will see that this leads to a field \mathcal{X} that extends the model introduced by Bacry in [2] and that has symmetrical increments. Thus, to obtain nonsymmetrical increments, we must introduce correlation between X^ϵ and dW_0 .

2.3. Notations and construction of the family X^ϵ .

Let k^R be the function

$$k^R(x) = \begin{cases} \frac{1}{|x|^{d/2}} & \text{for } |x| \leq R, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\theta(x)$ be a C^∞ , non negative and radially symmetrical function with compact support in $|x| \leq 1$ such that

$$\int_{\mathbb{R}^d} \theta(x) dx = 1.$$

We define $\theta^\epsilon = \frac{1}{\epsilon^d} \theta(\frac{\cdot}{\epsilon})$ and the corresponding convolutions:

$$k_\epsilon^R = \theta^\epsilon * k^R, \quad |\cdot|_\epsilon = \theta^\epsilon * |\cdot|.$$

Let γ be a strictly positive parameter and dW be a gaussian white noise on \mathbb{R}^d . We consider the following gaussian field:

$$X^\epsilon(y) = \gamma \int_{\mathbb{R}^d} k_\epsilon^R(y-\sigma) dW(\sigma).$$

Its correlation kernel is given by:

$$E(X^\epsilon(x)X^\epsilon(y)) = \gamma^2 \rho_{\epsilon/R}\left(\frac{x-y}{R}\right),$$

where $\rho = k^1 * k^1$ and $\rho_\epsilon = \theta^\epsilon * \theta^\epsilon * \rho$. One can prove the following expansion

$$\rho(x) = \omega_d \ln^+ \frac{1}{|x|} + \phi(x),$$

where ω_d denotes the surface of the unit sphere in \mathbb{R}^d and ϕ is a continuous function that vanishes for $|x| \geq 2$. We will note $|\cdot|_* = 1 \wedge |\cdot|$ and, with this definition, the previous expansion is equivalent to:

$$e^{\rho(x)} = \frac{e^{\phi(x)}}{|x|_*^{\omega_d}}.$$

One can also prove the following expansions with respect to ϵ :

$$k_\epsilon^R(0) = \frac{C_0}{\epsilon^{d/2}} \tag{2.4}$$

with $C_0 = \int_{|u| \leq 1} \frac{\theta(u)}{u^{d/2}} du$ and there exists a constant C_1 such that

$$\rho_{\epsilon/R}(0) = \omega_d \ln \frac{R}{\epsilon} + C_1 + o(\epsilon). \tag{2.5}$$

In the sequel, we will consider the case

$$\gamma dW = \gamma_0(\epsilon) dW_0 + \gamma_1 dW_1,$$

where dW_1 is a white noise independant of dW_0 and $\gamma_0(\epsilon)$ is a function of ϵ that will be defined later. Note that the integral in formula (2.3) has a meaning since dW_0 can be viewed as a random distribution.

2.4. Preliminary technical results. We remind the following integration by parts formula for gaussian vectors (cf. lemma 1.2.1 in [11]):

Lemma 2.1. *Let (g, g_1, \dots, g_n) be a centered gaussian vector and $G : \mathbb{R}^n \rightarrow \mathbb{R}$ a C^1 function such that its partial derivatives have at most exponential growth. Then we have:*

$$E(gG(g_1, \dots, g_n)) = \sum_{i=1}^n E(gg_i) E\left(\frac{\partial G}{\partial x_i}(g_1, \dots, g_n)\right). \tag{2.6}$$

From the above formula, one can easily deduce by induction the following lemma which will be frequently used in the sequel:

Lemma 2.2. *Let $l \in \mathbb{N}^*$ be some positive integer and (g, g_1, \dots, g_{2l}) a centered gaussian vector. Then:*

$$E(g_1 \dots g_{2l} e^g) = \left(\sum_{k=0}^l S_{k,l} \right) e^{\frac{1}{2} E(g^2)},$$

where

$$S_{k,l} = \sum_{\{i_1, \dots, i_{2k}\} \subset \{1, \dots, 2l\}} \sum E(gg_{i_1}) \dots E(gg_{i_{2k}}) E(g_{i_{2k+1}} g_{i_{2k+2}}) \dots E(g_{i_{2l-1}} g_{i_{2l}}),$$

where the second sum is taken over all partitions of $\{1, \dots, 2l\} / \{i_1, \dots, i_{2k}\}$ in subsets of two elements $\{i_{2k+1}, i_{2k+2}\}$.

Similarly, we get the following formula:

$$E(g_1 \dots g_{2l+1} e^g) = \left(\sum_{k=0}^l \tilde{S}_{k,l} \right) e^{\frac{1}{2} E(g^2)},$$

where

$$\tilde{S}_{k,l} = \sum_{\{i_1, \dots, i_{2k+1}\} \subset \{1, \dots, 2l+1\}} \sum E(gg_{i_1}) \dots E(gg_{i_{2k+1}}) E(g_{i_{2k+2}} g_{i_{2k+3}}) \dots E(g_{i_{2l}} g_{i_{2l+1}}),$$

Remark 2.3. In $S_{k,l}$ ($\tilde{S}_{k,l}$), the summation is made of $\frac{2l!}{2k!2^{l-k}(l-k)!}$ ($\frac{(2l+1)!}{(2k+1)!2^{l-k}(l-k)!}$) terms, number we will denote by $\alpha_{k,l}$ ($\tilde{\alpha}_{k,l}$).

We will also use the following lemma essentially due to Kahane ([8]).

Lemma 2.4. Let (T, d) be a metric space and σ a finite positive measure on T equipped with the borelian σ -field induced by d .

Let $q : T \times T \rightarrow \mathbb{R}_+$ a symmetric application and m a positive integer. Then we have the following inequalities:

$$\int_{T^{2m}} e^{\sum_{1 \leq j < k \leq 2m} q(t_j, t_k)} d\sigma(t_1) \dots d\sigma(t_{2m}) \leq \sigma(T) \left(\sup_{s \in T} \int_T e^{mq(t,s)} d\sigma(t) \right)^{2m-1}, \quad (2.7)$$

$$\begin{aligned} & \int_{T^{2m+1}} e^{\sum_{1 \leq j < k \leq 2m+1} q(t_j, t_k)} d\sigma(t_1) \dots d\sigma(t_{2m+1}) \\ & \leq \sigma(T) \sup_{s, \tilde{s}} \left(\int_T e^{q(\tilde{s}, t)} d\sigma(t) \right) \left(\int_T e^{mq(s,t)} e^{q(\tilde{s}, t)} d\sigma(t) \right)^{2m-1}. \end{aligned} \quad (2.8)$$

Proof. The proof of (2.7) can be found in [8]. Thus we just prove how to derive inequality (2.8) from (2.7). By integrating with respect to the first $2m$ variables, we get:

$$\begin{aligned} & \int_{T^{2m+1}} e^{\sum_{1 \leq j < k \leq 2m+1} q(t_j, t_k)} d\sigma(t_1) \dots d\sigma(t_{2m+1}) \\ & = \int_T d\sigma(t_{2m+1}) \int_{T^{2m}} e^{\sum_{1 \leq j < k \leq 2m} q(t_j, t_k)} \prod_{j=1}^{2m} e^{q(t_j, t_{2m+1})} d\sigma(t_1) \dots d\sigma(t_{2m}) \\ & \stackrel{(2.7)}{\leq} \int_T d\sigma(t_{2m+1}) \left(\int_T e^{q(t, t_{2m+1})} d\sigma(t) \right) \left(\sup_s \int_T e^{mq(s,t)} e^{q(t, t_{2m+1})} d\sigma(t) \right)^{2m-1} \\ & \leq \sigma(T) \sup_{s, \tilde{s}} \left(\int_T e^{q(\tilde{s}, t)} d\sigma(t) \right) \left(\int_T e^{mq(s,t)} e^{q(\tilde{s}, t)} d\sigma(t) \right)^{2m-1}. \end{aligned}$$

3. CONSTRUCTION OF MULTIFRACTAL RANDOM FIELDS WITH NONSYMMETRICAL INCREMENTS

In this section, we will suppose that $d/2 < \alpha < (d/2 + 1) \wedge d$ and $\omega_d \gamma_1^2 < d$. We consider the field \mathcal{X}^ϵ defined by formula (2.3) with

$$X^\epsilon(y) = \gamma_0(\epsilon)X_0^\epsilon(y) + \gamma_1 X_1^\epsilon(y),$$

where

$$X_i^\epsilon(y) = \int_{\mathbb{R}^d} k_\epsilon^R(y - \sigma) dW_i(\sigma), \quad i = 0, 1.$$

We set also

$$C_\epsilon = ((\gamma_0(\epsilon))^2 + \gamma_1^2) \rho_{\epsilon/R}(0).$$

and

$$\gamma_0(\epsilon) = \gamma_0^* \left(\frac{\epsilon}{R} \right)^{\frac{d - \omega_d \gamma_1^2}{2}}.$$

Therefore, we introduce a slight correlation between X^ϵ and dW_0 ($\gamma_0(\epsilon)$ tends to 0 as ϵ goes to 0).

3.1. Multiplicative chaos in dimension d . Multiplicative chaos or the "limit-lognormal" model introduced by Mandelbrot is a generalization of the exponential of a gaussian process. As mentioned in the introduction, it was defined rigorously by Kahane in [8]. The construction of Kahane was based on the theory of martingales and thus the generalized correlation kernel (here $\rho(t - s)$) had to verify a condition hard to verify practically (the σ -positivity condition). Our construction is based on L^2 -theory and can be carried out without this condition.

Let γ_1 be some real number such that $\gamma_1^2 \omega_d < d$ and ϵ a positive number. Let $\mathcal{B}(\mathbb{R}^d)$ denote the standard borelian σ -field; we want to consider the limit as ϵ goes to 0 of the random measures Q^{ϵ, γ_1} defined by:

$$\begin{aligned} Q^{\epsilon, \gamma_1}(dy) &= e^{\gamma_1 X_1^\epsilon(y) - \frac{1}{2} E((X_1^\epsilon(y))^2)} dy \\ &= e^{\gamma_1 X_1^\epsilon(y) - \frac{1}{2} \gamma_1^2 \rho_{\epsilon/R}(0)} dy. \end{aligned} \quad (3.1)$$

This leads us to state the following proposition:

Proposition 3.1 (Multiplicative chaos of order γ_1). *There exists a positive random measure $Q^{\gamma_1}(dy)$ independent of the regularizing function θ such that:*

- (1) *for all A bounded in $\mathcal{B}(\mathbb{R}^d)$, $E(Q^{\gamma_1}(A)) = |A|$.*
- (2) *Q^{γ_1} has almost surely no atoms.*
- (3) *Q^{γ_1} is almost surely singular with respect to the Lebesgue measure.*

If q is some positive integer and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a deterministic function that satisfies the following condition:

$$\int_{(\mathbb{R}^d)^{2q}} |f(y_1)| \dots |f(y_{2q})| \prod_{1 \leq i < j \leq 2q} \frac{1}{\left| \frac{y_i - y_j}{R} \right|_*^{\gamma_1^2 \omega_d}} dy_1 \dots dy_{2q} < \infty, \quad (3.2)$$

then we have the following convergence:

$$\int_{\mathbb{R}^d} f(y) Q^{\epsilon, \gamma_1}(dy) \xrightarrow[\epsilon \rightarrow 0]{L^{2q}} \int_{\mathbb{R}^d} f(y) Q^{\gamma_1}(dy).$$

We also have the following expression for the moments of $\int_{\mathbb{R}^d} f(y)Q^{\gamma_1}(dy)$:

$$\forall k \leq 2q, \quad E\left(\left(\int_{\mathbb{R}^d} f(y)Q^{\gamma_1}(dy)\right)^k\right) = \int_{(\mathbb{R}^d)^k} f(y_1) \dots f(y_k) \prod_{1 \leq i < j \leq k} \frac{e^{\gamma_1^2 \phi\left(\frac{y_i - y_j}{R}\right)}}{\left|\frac{y_i - y_j}{R}\right|^{\gamma_1^2 \omega_d}} dy_1 \dots dy_k. \quad (3.3)$$

We will call $Q^{\gamma_1}(dy)$ multiplicative chaos of order γ_1 .

Proof. We first start by considering a positive integer and a function f that satisfies the corresponding integrability condition (3.2). Let ϵ, ϵ' be two positive numbers. By using Fubini, we get for all $j \leq 2q$:

$$\begin{aligned} & E\left(\left(\int_{\mathbb{R}^d} f(y)Q^{\epsilon, \gamma_1}(dy)\right)^j \left(\int_{\mathbb{R}^d} f(y)Q^{\epsilon', \gamma_1}(dy)\right)^{2q-j}\right) \\ &= e^{-\frac{j}{2}\gamma_1^2 \rho_{\epsilon/R}(0) - \frac{2q-j}{2}\gamma_1^2 \rho_{\epsilon'/R}(0)} \int_{(\mathbb{R}^d)^{2q}} f(y_1) \dots f(y_{2q}) \times \\ & \quad e^{\frac{1}{2}\gamma_1^2 E((\sum_{i=1}^j \mathcal{X}_1^\epsilon(y_i) + \sum_{i=j+1}^{2q} \mathcal{X}_1^{\epsilon'}(y_i))^2)} dy_1 \dots dy_{2q} \\ & \xrightarrow{\epsilon, \epsilon' \rightarrow 0} \int_{(\mathbb{R}^d)^{2q}} f(y_1) \dots f(y_{2q}) \prod_{1 \leq i < j \leq 2q} \frac{e^{\gamma_1^2 \phi\left(\frac{y_i - y_j}{R}\right)}}{\left|\frac{y_i - y_j}{R}\right|^{\gamma_1^2 \omega_d}} dy_1 \dots dy_{2q}, \end{aligned}$$

From this, we deduce that:

$$E\left(\left(\int_{\mathbb{R}^d} f(y)Q^{\epsilon, \gamma_1}(dy) - \int_{\mathbb{R}^d} f(y)Q^{\epsilon', \gamma_1}(dy)\right)^{2q}\right) \xrightarrow{\epsilon, \epsilon' \rightarrow 0} 0$$

and therefore that $\int_{\mathbb{R}^d} f(y)Q^{\epsilon, \gamma_1}(dy)$ is a Cauchy sequence in L^{2q} that converges to some random variable $\tilde{Q}^{\gamma_1}(f)$. For $k \leq 2q$, the moment $E((\tilde{Q}^{\gamma_1}(f))^k)$ is the limit as ϵ goes to 0 of $E((Q^{\epsilon, \gamma_1}(f))^k)$; from this one can deduce that the moments of $\tilde{Q}^{\gamma_1}(f)$ are given by formula (3.3).

Since $\gamma_1^2 \omega_d < d$, for any bounded set A in $\mathcal{B}(\mathbb{R}^d)$, we deduce from the proof above and lemma 2.4 that $Q^{\epsilon, \gamma_1}(A)$ converges in L^2 to some random variable $\tilde{Q}^{\gamma_1}(A)$. This defines a family of random variables (indexed by the bounded borelian sets) that satisfies the following properties:

- (1) For all disjoint and bounded sets A_1, A_2 in $\mathcal{B}(\mathbb{R}^d)$,

$$\tilde{Q}^{\gamma_1}(A_1 \cup A_2) = \tilde{Q}^{\gamma_1}(A_1) + \tilde{Q}^{\gamma_1}(A_2) \quad a.s.$$

- (2) For any bounded sequence $(A_n)_{n \geq 1}$ decreasing to \emptyset :

$$\tilde{Q}^{\gamma_1}(A_n) \xrightarrow{n \rightarrow \infty} 0 \quad a.s.$$

By theorem 6.1.VI. in [5], there exists a random measure Q^{γ_1} such that for all bounded A in $\mathcal{B}(\mathbb{R}^d)$ we have:

$$Q^{\gamma_1}(A) = \tilde{Q}^{\gamma_1}(A) \quad a.s.$$

Finally, one can easily show that the limit random variable $\tilde{Q}^{\gamma_1}(f)$ is almost surely equal to $\int_{\mathbb{R}^d} f(y)Q^{\gamma_1}(dy)$. \square

3.2. Convergence towards a field \mathcal{X} . In this subsection, we will prove the following proposition:

Proposition 3.2. *Let α be such that $d/2 < \alpha < (d/2 + 1) \wedge d$ and γ_1 such that $2\gamma_1^2\omega_d < \alpha - d/2$. There exists a field $(\mathcal{X}(x))_{x \in \mathbb{R}^d}$ such that for all k and $x_1, \dots, x_k \in \mathbb{R}^d$ the following convergence in law holds:*

$$(\mathcal{X}^\epsilon(x_1), \dots, \mathcal{X}^\epsilon(x_k)) \xrightarrow{\epsilon \rightarrow 0} (\mathcal{X}(x_1), \dots, \mathcal{X}(x_k)). \quad (3.4)$$

Let l be an integer such that one of the following conditions hold:

- (1) l is even and $l\gamma_1^2\omega_d < \alpha - d/2$.
- (2) l is odd and $(l+1)\gamma_1^2\omega_d < \alpha - d/2$.

Then there exists C such that, for all x in \mathbb{R}^d , the random variable $\mathcal{X}(x)$ has a moment of order $2l$ given by the following expression:

$$\begin{aligned} E((\mathcal{X}(x))^{2l}) &= \sum_{k=0}^l \alpha_{k,l} C^{2k} \int_{(\mathbb{R}^d)^{k+l}} \frac{\varphi_R(y_1)}{|y_1|^{d-\alpha}} \cdots \frac{\varphi_R(y_{2k})}{|y_{2k}|^{d-\alpha}} \frac{\varphi_R(y_{2k+1})^2}{|y_{2k+1}|^{2(d-\alpha)}} \cdots \frac{\varphi_R(y_{k+l})^2}{|y_{k+l}|^{2(d-\alpha)}} \\ &\prod_{1 \leq i < j \leq 2k} \frac{e^{\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{\gamma_1^2 \omega_d}} \prod_{\substack{1 \leq i \leq 2k \\ j > 2k}} \frac{e^{2\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{2\gamma_1^2 \omega_d}} \prod_{2k+1 \leq i < j \leq k+l} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{4\gamma_1^2 \omega_d}} dy_1 \dots dy_{k+l}. \end{aligned} \quad (3.5)$$

We also have:

$$\begin{aligned} E((\mathcal{X}(x+h) - \mathcal{X}(x))^{2l}) &= \sum_{k=0}^l \alpha_{k,l} C^{2k} \int_{(\mathbb{R}^d)^{k+l}} \left(\frac{\varphi_R(y_1 - h)}{|y_1 - h|^{d-\alpha}} - \frac{\varphi_R(y_1)}{|y_1|^{d-\alpha}} \right) \cdots \\ &\left(\frac{\varphi_R(y_{2k} - h)}{|y_{2k} - h|^{d-\alpha}} - \frac{\varphi_R(y_{2k})}{|y_{2k}|^{d-\alpha}} \right) \left(\frac{\varphi_R(y_{2k+1} - h)}{|y_{2k+1} - h|^{d-\alpha}} - \frac{\varphi_R(y_{2k+1})}{|y_{2k+1}|^{d-\alpha}} \right)^2 \cdots \left(\frac{\varphi_R(y_{k+l} - h)}{|y_{k+l} - h|^{d-\alpha}} - \frac{\varphi_R(y_{k+l})}{|y_{k+l}|^{d-\alpha}} \right)^2 \\ &\prod_{1 \leq i < j \leq 2k} \frac{e^{\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{\gamma_1^2 \omega_d}} \prod_{\substack{1 \leq i \leq 2k \\ j > 2k}} \frac{e^{2\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{2\gamma_1^2 \omega_d}} \prod_{2k+1 \leq i < j \leq k+l} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{4\gamma_1^2 \omega_d}} dy_1 \dots dy_{k+l}. \end{aligned} \quad (3.6)$$

Proof. Let γ_1 be such that $2\gamma_1^2\omega_d < \alpha - d/2$. We set

$$C = \frac{\gamma_0^* C_0 e^{-1/2\gamma_1^2 C_1}}{R^{d/2}}.$$

and define two auxiliary fields $\mathcal{Y}^\epsilon, \mathcal{Z}^\epsilon$ by the following expressions:

$$\mathcal{Y}^\epsilon(x) = \int_{\mathbb{R}^d} \frac{\varphi_R(x-y)}{|x-y|^{d-\alpha}} Q^{\epsilon, \gamma_1}(dy) \quad (3.7)$$

and

$$\mathcal{Z}^\epsilon(x) = \int_{\mathbb{R}^d} \frac{\varphi_R(x-y)}{|x-y|^{d-\alpha}} e^{\gamma_1 X_1^\epsilon(y) - C_\epsilon} dW_0(y). \quad (3.8)$$

Note that $\mathcal{Z}^\epsilon(x)$ exists since X_1^ϵ and dW_0 are independent with:

$$E\left(\int_{\mathbb{R}^d} \frac{\varphi_R(x-y)^2}{|x-y|^{2(d-\alpha)}} e^{2\gamma_1 X_1^\epsilon(y) - 2C_\epsilon} dy\right) < \infty. \quad (3.9)$$

We can compute, for all x in \mathbb{R}^d , $E((\mathcal{X}^\epsilon(x) - C\mathcal{Y}^\epsilon(x) - \mathcal{Z}^\epsilon(x))^2)$ (cf. the more complicated computations in the proof of proposition 3.7) and derive the following limit:

$$\mathcal{X}^\epsilon(x) - (C\mathcal{Y}^\epsilon(x) + \mathcal{Z}^\epsilon(x)) \xrightarrow[\epsilon \rightarrow 0]{L^2} 0.$$

Thus, we must show that the finite dimensional distributions of the field $C\mathcal{Y}^\epsilon + \mathcal{Z}^\epsilon$ converge in law. Let k be some positive integer and x_1, \dots, x_k points in \mathbb{R}^d . For all $\xi = (\xi_1, \dots, \xi_k)$ in \mathbb{R}^k , by conditioning on the field generated by the white noise dW_1 and using proposition 3.1, we get:

$$\begin{aligned} & E(e^{i\xi \cdot (C\mathcal{Y}^\epsilon(x_1) + \mathcal{Z}^\epsilon(x_1), \dots, C\mathcal{Y}^\epsilon(x_k) + \mathcal{Z}^\epsilon(x_k))}) \\ &= E(e^{iC \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}}) Q^{\epsilon, \gamma_1}(dy) - \frac{1}{2} e^{2\gamma_0(\epsilon)^2 \rho_{\epsilon/R}(0)} \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}})^2 Q^{\epsilon, 2\gamma_1}(dy)}) \\ &\xrightarrow[\epsilon \rightarrow 0]{} E(e^{iC \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}}) Q^{\gamma_1}(dy) - \frac{1}{2} \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}})^2 Q^{2\gamma_1}(dy)}). \end{aligned}$$

Thus, by applying Levy's theorem, we conclude that the finite dimensional distributions of the field $C\mathcal{Y}^\epsilon + \mathcal{Z}^\epsilon$ converge in law to those of a field \mathcal{X} . We also get from the proof above the characteristic function of $(\mathcal{X}(x_1), \dots, \mathcal{X}(x_k))$:

$$E(e^{i\xi \cdot (\mathcal{X}(x_1), \dots, \mathcal{X}(x_k))}) = E(e^{iC \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}}) Q^{\gamma_1}(dy) - \frac{1}{2} \int_{\mathbb{R}^d} (\sum_{i=1}^k \xi_i \frac{\varphi_R(x_i - y)}{|x_i - y|^{d-\alpha}})^2 Q^{2\gamma_1}(dy)}). \quad (3.10)$$

Suppose l is a positive integer that satisfies the condition of the proposition. By applying (3.10), we get for all ξ in \mathbb{R} :

$$E(e^{i\xi \cdot \mathcal{X}(x)}) = E(e^{iC\xi \int_{\mathbb{R}^d} \frac{\varphi_R(x-y)}{|x-y|^{d-\alpha}} Q^{\gamma_1}(dy) - \frac{1}{2} \xi^2 \int_{\mathbb{R}^d} (\frac{\varphi_R(x-y)}{|x-y|^{d-\alpha}})^2 Q^{2\gamma_1}(dy)}).$$

We derive expression (3.5) by computing $\frac{\partial^{2l} E(e^{i\xi \cdot \mathcal{X}(x)})}{\partial \xi^{2l}} \Big|_{\xi=0}$ thanks to the above formula and proposition 3.1. We derive expression (3.6) similarly.

3.3. Scaling of \mathcal{X} and tightness of \mathcal{X}^ϵ . The purpose of this subsection is to show that the field $(\mathcal{X}(x))_{x \in \mathbb{R}^d}$ satisfies the multifractal scaling relation (this is what propositions 3.5 and 3.6 below assert) and to study the tightness of the family \mathcal{X}^ϵ .

We first state two preliminary lemmas we will use in the rest of the paper.

Lemma 3.3. *Let δ be some real number such that $0 \leq \delta < \alpha$ and $\delta \neq \alpha - 1$. There exists $C = C(\delta)$ such that we have the following inequality for $|h| \leq R$:*

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} \left| \frac{\varphi_R(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}} \right| \frac{1}{|\frac{x-y}{R}|_*^\delta} dy \leq R^{d/2} C \left| \frac{h}{R} \right|^{(\alpha-\delta) \wedge 1}. \quad (3.11)$$

Proof. By homogeneity, we suppose that $R = 1$ and for simplicity, we suppose $d \geq 2$. Since $\frac{1}{|x|_*} \leq \frac{1}{|x|} + 1$, we have to show that for $\delta \in [0, \alpha[$:

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} \left| \frac{\varphi(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi(y)}{|y|^{d-\alpha}} \right| \frac{1}{|x-y|^\delta} dy \leq C |h|^{(\alpha-\delta) \wedge 1}.$$

There exists C such that for all y and h , we have:

$$|\varphi(y-h) - \varphi(y)| \leq C|h| \text{ and } \varphi(y) \leq C1_{|y| \leq 2}. \quad (3.12)$$

We set

$$I(x) = \int_{\mathbb{R}^d} \left| \frac{\varphi(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi(y)}{|y|^{d-\alpha}} \right| \frac{1}{|x-y|^\delta} dy.$$

Therefore we get

$$I(x) \leq C|h| \int_{|y| \leq 3} \frac{1}{|y|^{d-\alpha}} \frac{1}{|x-y|^\delta} dy \quad (3.13)$$

$$\begin{aligned} &+ C \int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right| \frac{1}{|x-y|^\delta} dy \\ &\leq C|h| + C \int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right| \frac{1}{|x-y|^\delta} dy. \end{aligned} \quad (3.14)$$

First case: $\delta < \alpha - 1$.

Plugging inequality

$$\left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right| \leq \frac{(d-\alpha)|h|}{|y-h|^{d-\alpha+1} \wedge |y|^{d-\alpha+1}}$$

in (3.14), we get

$$I(x) \leq C|h| \int_{|y| \leq 3} \frac{1}{|y-h|^{d-\alpha+1} \wedge |y|^{d-\alpha+1}} \frac{1}{|x-y|^\delta} dy,$$

which concludes the proof.

Second case: $\delta > \alpha - 1$.

By the change of variable $y = |h|u$ and setting $h = |h|e$ with $|e| = 1$, we get:

$$\begin{aligned} &\int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right| \frac{1}{|x-y|^\delta} dy \\ &= |h|^{\alpha-\delta} \int_{|u| \leq \frac{3}{|h|}} \left| \frac{1}{|u-e|^{d-\alpha}} - \frac{1}{|u|^{d-\alpha}} \right| \frac{1}{|x/|h| - u|^\delta} du \\ &\leq |h|^{\alpha-\delta} \sup_{a \in \mathbb{R}^d} \int_{\mathbb{R}^d} \left| \frac{1}{|u-e|^{d-\alpha}} - \frac{1}{|u|^{d-\alpha}} \right| \frac{1}{|a-u|^\delta} du. \end{aligned}$$

□

Lemma 3.4. *Let δ be some real number such that $0 \leq \delta < 2\alpha - d$ or, if $d = 1$ and $\alpha > 1$, $\delta < 1$. There exists $C = C(\delta)$ such that we have the following inequality for $|h| \leq R$:*

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} \left| \frac{\varphi_R(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}} \right|^2 \frac{1}{|x-y|_*^\delta} dy \leq C \left| \frac{h}{R} \right|^{2\alpha-d-\delta}. \quad (3.15)$$

Proof. As in the proof above, we can replace $|\cdot|_*$ by $|\cdot|$ and suppose that $R = 1$; thus we have to show inequality (3.15) with $J(x)$ where we set:

$$J(x) = \int_{\mathbb{R}^d} \left| \frac{\varphi(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi(y)}{|y|^{d-\alpha}} \right|^2 \frac{1}{|x-y|^\delta} dy.$$

Using inequality (3.12), we get

$$\begin{aligned}
J(x) &\leq C|h|^2 \int_{|y| \leq 3} \frac{1}{|y|^{2(d-\alpha)}} \frac{1}{|x-y|^\delta} dy \\
&\quad + C \int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right|^2 \frac{1}{|x-y|^\delta} dy \\
&\leq C|h|^2 + C \int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right|^2 \frac{1}{|x-y|^\delta} dy. \tag{3.16}
\end{aligned}$$

Since $2 > 2\alpha - d - \delta$, we only have to consider the second term in inequality (3.16). By the change of variable $y = |h|u$ and setting $h = |h|e$ with $|e| = 1$, we get:

$$\begin{aligned}
&\int_{|y| \leq 3} \left| \frac{1}{|y-h|^{d-\alpha}} - \frac{1}{|y|^{d-\alpha}} \right|^2 \frac{1}{|x-y|^\delta} dy \\
&= |h|^{2\alpha-d-\delta} \int_{|u| \leq \frac{3}{|h|}} \left| \frac{1}{|u-e|^{d-\alpha}} - \frac{1}{|u|^{d-\alpha}} \right|^2 \frac{1}{|x/|h| - u|^\delta} du \\
&\leq |h|^{2\alpha-d-\delta} \sup_{a \in \mathbb{R}^d} \int_{\mathbb{R}^d} \left| \frac{1}{|u-e|^{d-\alpha}} - \frac{1}{|u|^{d-\alpha}} \right|^2 \frac{1}{|a-u|^\delta} du.
\end{aligned}$$

□

Proposition 3.5. (*Scaling along the even integers*)

Let l be an integer such that one of the following conditions hold:

- (1) l is even and $l\gamma_1^2\omega_d < \alpha - d/2$
- (2) l is odd and $(l+1)\gamma_1^2\omega_d < \alpha - d/2$.

Let e be a unit vector ($|e| = 1$). Then there exists $C_l \neq 0$ independent of e such that the following scaling relation holds:

$$E((\mathcal{X}(x + \lambda e) - \mathcal{X}(x))^{2l}) \underset{\lambda \rightarrow 0}{\sim} C_l \left(\frac{\lambda}{R}\right)^{\zeta_{2l}}, \tag{3.17}$$

where we have

$$\zeta_{2l} = l(2\alpha - d) - 2\gamma_1^2\omega_d l(l-1). \tag{3.18}$$

Proof. For simplicity, we will suppose that l is even and that $l\gamma_1^2\omega_d < \alpha - d/2$. We introduce the following notation:

$$f_h(y) = \frac{\varphi_R(y-h)}{|y-h|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}}.$$

We shall see that the scaling at small scale of the sum (3.6) is given by the term $k = 0$. Indeed for all $k \geq 1$ let us consider the integral

$$\begin{aligned}
& \int_{(\mathbb{R}^d)^{k+l}} f_h(y_1) \dots f_h(y_{2k}) (f_h(y_{2k+1}))^2 \dots (f_h(y_{k+l}))^2 \prod_{1 \leq i < j \leq 2k} \frac{e^{\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{\gamma_1^2 \omega_d}} \times \\
& \prod_{\substack{1 \leq i \leq 2k \\ j > 2k}} \frac{e^{2\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{2\gamma_1^2 \omega_d}} \prod_{2k+1 \leq i < j \leq k+l} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{4\gamma_1^2 \omega_d}} dy_1 \dots dy_{k+l}. \\
& \leq I_k J_{k,l}, \tag{3.19}
\end{aligned}$$

where we set

$$\begin{aligned}
I_k &= \sup_{y_{2k+1}, \dots, y_{k+l}} \int_{(\mathbb{R}^d)^{2k}} f_h(y_1) \dots f_h(y_{2k}) \prod_{\substack{1 \leq i \leq 2k \\ j > 2k}} \frac{e^{2\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{2\gamma_1^2 \omega_d}} \times \\
& \prod_{1 \leq i < j \leq 2k} \frac{e^{\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{\gamma_1^2 \omega_d}} dy_1 \dots dy_{2k}.
\end{aligned}$$

and

$$J_{k,l} = \int_{(\mathbb{R}^d)^{l-k}} (f_h(y_{2k+1}))^2 \dots (f_h(y_{k+l}))^2 \prod_{2k+1 \leq i < j \leq k+l} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{4\gamma_1^2 \omega_d}} dy_{2k+1} \dots dy_{k+l}.$$

By using the estimates (3.11), (3.15) and the inequalities (2.7), (2.8), one can show that for all $k \geq 1$, we have

$$I_k J_{k,l} \leq CR^{dk} \left| \frac{h}{R} \right|^{c_{k,l}},$$

with

$$c_{k,l} = (\alpha - 2(l-k)\gamma_1^2 \omega_d) \wedge 1 + ((\alpha - (2l-k)\gamma_1^2 \omega_d) \wedge 1)(2k-1) + (2\alpha - d)(l-k) - 2\gamma_1^2 \omega_d (l-k)(l-k-1).$$

If $\alpha - 2(l-k)\gamma_1^2 \omega_d < 1$, then $c_{k,l} = \zeta_{2l} + k(d - \gamma_1^2 \omega_d)$; If $\alpha - 2(l-k)\gamma_1^2 \omega_d \geq 1$ and $\alpha - (2l-k)\gamma_1^2 \omega_d < 1$, then $c_{k,l} = \zeta_{2l} + 1 - \alpha + dk + (2l-3k)\gamma_1^2 \omega_d$; otherwise $c_{k,l} = 2k + (2\alpha - d)(l-k) - 2\gamma_1^2 \omega_d (l-k)(l-k-1)$. In all cases, it is easy to show that $c_{k,l} > \zeta_{2l}$ under the conditions of the proposition..

Finally, we study the term where $k = 0$. We get for $h = \lambda e$ with $|e| = 1$:

$$\begin{aligned}
& \int_{(\mathbb{R}^d)^l} (f_h(y_1))^2 \dots (f_h(y_l))^2 \prod_{1 \leq i < j \leq l} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|y_i - y_j|_*^{4\gamma_1^2 \omega_d}} dy_1 \dots dy_l. \\
& = \left(\frac{\lambda}{R}\right)^{l(2\alpha - d)} \int_{(\mathbb{R}^d)^l} \left(\frac{\varphi_R(\lambda(u_1 - e))}{|u_1 - e|^{d-\alpha}} - \frac{\varphi_R(\lambda u_1)}{|u_1|^{d-\alpha}}\right)^2 \dots \left(\frac{\varphi_R(\lambda(u_l - e))}{|u_l - e|^{d-\alpha}} - \frac{\varphi_R(\lambda u_l)}{|u_l|^{d-\alpha}}\right)^2 \times \\
& \quad \prod_{1 \leq i < j \leq l} \frac{e^{4\gamma_1^2 \phi(\frac{\lambda(u_i - u_j)}{R})}}{|\frac{\lambda(u_i - u_j)}{R}|_*^{4\gamma_1^2 \omega_d}} du_1 \dots du_l. \\
& \stackrel{\lambda \rightarrow 0}{\sim} e^{2l(l-1)\gamma_1^2 \phi(0)} \left(\frac{\lambda}{R}\right)^{\zeta_{2l}} \int_{(\mathbb{R}^d)^l} \left(\frac{1}{|u_1 - e|^{d-\alpha}} - \frac{1}{|u_1|^{d-\alpha}}\right)^2 \dots \left(\frac{1}{|u_l - e|^{d-\alpha}} - \frac{1}{|u_l|^{d-\alpha}}\right)^2 \times \\
& \quad \prod_{1 \leq i < j \leq l} \frac{1}{|u_i - u_j|^{4\gamma_1^2 \omega_d}} du_1 \dots du_l,
\end{aligned}$$

and inequality (2.7) shows that this integral is finite when $l\gamma_1^2 \omega_d < \alpha - d/2$. \square

In the next proposition, we state the scaling relations of \mathcal{X} along the odd integers.

Proposition 3.6. *(Scaling along the odd integers) We suppose in this proposition that $\alpha > 2$ and thus $d \geq 3$. Let l be an integer such that $l + 1$ satisfies one of the conditions in proposition 3.5.*

Let e be a unit vector ($|e| = 1$). Then there exists C_l (with $C_3 \neq 0$) independent of e such that the following scaling relation holds:

$$E((\mathcal{X}(x + \lambda e) - \mathcal{X}(x))^{2l+1}) \stackrel{\lambda \rightarrow 0}{\sim} \gamma_0^* R^{d/2} C_l \left(\frac{\lambda}{R}\right)^{\tilde{\zeta}_{2l+1}}, \quad (3.20)$$

where we have

$$\tilde{\zeta}_{2l+1} = l(2\alpha - d) - 2\gamma_1^2 \omega_d l(l - 1) + 2. \quad (3.21)$$

Proof. As in proposition 3.2, setting $C = \frac{\gamma_0^* C_0 e^{-1/2\gamma_1^2 C_1}}{R^{d/2}}$, it is possible to show that:

$$\begin{aligned}
& E((\mathcal{X}(x + h) - \mathcal{X}(x))^{2l+1}) \\
& = \sum_{k=0}^l \tilde{\alpha}_{k,l} C^{2k+1} \int_{(\mathbb{R}^d)^{k+l+1}} f_h(y_1) \dots f_h(y_{2k+1}) (f_h(y_{2k+2}))^2 \dots (f_h(y_{k+l+1}))^2 \times \\
& \quad \prod_{1 \leq i < j \leq 2k+1} \frac{e^{\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{\gamma_1^2 \omega_d}} \prod_{\substack{1 \leq i \leq 2k+1 \\ j > 2k+1}} \frac{e^{2\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{2\gamma_1^2 \omega_d}} \prod_{2k+2 \leq i < j \leq k+l+1} \frac{e^{4\gamma_1^2 \phi(\frac{y_i - y_j}{R})}}{|\frac{y_i - y_j}{R}|_*^{4\gamma_1^2 \omega_d}} dy_1 \dots dy_{k+l+1},
\end{aligned}$$

where, as usual, we set

$$f_h(y) = \frac{\varphi_R(y - h)}{|y - h|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}}.$$

Similarly to proposition 3.5, to get the main contribution as $|h|$ goes to 0, we examine the term $k = 0$. We set:

$$\psi(y) = \frac{\varphi(y)}{|y|^{d-\alpha}}.$$

Note that the condition $2l\gamma_1^2\omega_d < \alpha - 2$ ensures that:

$$\int_{\mathbb{R}^d} \left| \frac{\partial^2 \psi}{\partial y^i \partial y^j} \right| e^{2l\gamma_1^2\rho(y)} dy < \infty.$$

We get for $h = \lambda e$ with $|e| = 1$ the following equivalent:

$$\begin{aligned} & \int_{(\mathbb{R}^d)^{l+1}} f_h(y_1)(f_h(y_2))^2 \dots (f_h(y_{l+1}))^2 \prod_{j \geq 2} \frac{e^{2\gamma_1^2\phi(\frac{y_1-y_j}{R})}}{\left|\frac{y_1-y_j}{R}\right|^{2\gamma_1^2\omega_d}} \times \\ & \prod_{2 \leq i < j \leq l+1} \frac{e^{4\gamma_1^2\phi(\frac{y_i-y_j}{R})}}{\left|\frac{y_i-y_j}{R}\right|^{4\gamma_1^2\omega_d}} dy_1 \dots dy_{l+1}. \\ & = \int_{\substack{y_i = \lambda u_i, i \geq 2 \\ y_1 = R y}} R^{d/2} \left(\frac{\lambda}{R}\right)^{l(2\alpha-d)} \int_{(\mathbb{R}^d)^{l+1}} (\psi(y - \frac{\lambda}{R}e) - \psi(y)) \prod_{j \geq 2} \left(\frac{\varphi_R(\lambda(u_j - e))}{|u_j - e|^{d-\alpha}} - \frac{\varphi_R(\lambda u_j)}{|u_j|^{d-\alpha}}\right)^2 \times \\ & \prod_{j \geq 2} \frac{e^{2\gamma_1^2\phi(y - \frac{\lambda u_j}{R})}}{\left|y - \frac{\lambda u_j}{R}\right|^{2\gamma_1^2\omega_d}} \prod_{2 \leq i < j \leq l+1} \frac{e^{4\gamma_1^2\phi(\frac{\lambda(u_i - u_j)}{R})}}{\left|\frac{\lambda(u_i - u_j)}{R}\right|^{4\gamma_1^2\omega_d}} dy du_2 \dots du_{l+1}. \\ & \underset{\lambda \rightarrow 0}{\sim} R^{d/2} \tilde{C}_l e^{2l(l-1)\gamma_1^2\phi(0)} \left(\frac{\lambda}{R}\right)^{\tilde{c}_{2l+1}}, \end{aligned}$$

with $\tilde{C}_l = A_l B_l$ where

$$A_l = \sum_{1 \leq i, j \leq d} e_i e_j \int_{\mathbb{R}^d} \frac{\partial^2 \psi}{\partial y^i \partial y^j} e^{2l\gamma_1^2\rho(y)} dy$$

and

$$\begin{aligned} B_l &= \int_{(\mathbb{R}^d)^l} \left(\frac{1}{|u_1 - e|^{d-\alpha}} - \frac{1}{|u_1|^{d-\alpha}}\right)^2 \dots \left(\frac{1}{|u_l - e|^{d-\alpha}} - \frac{1}{|u_l|^{d-\alpha}}\right)^2 \times \\ & \prod_{2 \leq i < j \leq l+1} \frac{1}{|u_i - u_j|^{4\gamma_1^2\omega_d}} du_2 \dots du_{l+1}. \end{aligned}$$

A direct computation shows that $\tilde{C}_3 \neq 0$ from which we deduce that for $\gamma_0^* \neq 0$ the distribution of $\mathcal{X}(x + \lambda e) - \mathcal{X}(x)$ is nonsymmetrical. \square

Proposition 3.7. (*Tightness*) *Let l be some positive integer that satisfies the condition of proposition 3.5 and γ a positive parameter such that $\gamma_1^2 < \gamma^2$. Then there exists $\epsilon_0 > 0$ and C independent of ϵ such that for $\epsilon < \epsilon_0$ and $|h| \leq R$:*

$$\forall x, \quad E((\mathcal{X}^\epsilon(x+h) - \mathcal{X}^\epsilon(x))^{2l}) \leq C|h|^{l(2\alpha-d)-2\gamma^2\omega_d l(l-1)}, \quad (3.22)$$

and

$$E((\mathcal{X}^\epsilon(0))^{2l}) \leq C. \quad (3.23)$$

Proof. We only prove (3.22) (the proof of 3.23 is similar). We are going to compute the moment

$$E\left(\left(\int_{\mathbb{R}^d} f_{\epsilon,h}(y) e^{X^\epsilon(y)-C_\epsilon} dW_0(y)\right)^{2l}\right)$$

where we set

$$f_{\epsilon,h}(y) = \frac{\varphi_R(y-h)}{|y-h|_\epsilon^{d-\alpha}} - \frac{\varphi_R(y)}{|y|_\epsilon^{d-\alpha}}.$$

We get

$$E\left(\left(\int_{\mathbb{R}^d} f_{\epsilon,h}(y) e^{X^\epsilon(y)-C_\epsilon} dW_0(y)\right)^{2l}\right) = e^{-2lC_\epsilon} \int f_{\epsilon,h}(y_1) \dots f_{\epsilon,h}(y_{2l}) E(e^{\hat{X}^\epsilon} dW_0(y_1) \dots dW_0(y_{2l})), \quad (3.24)$$

where

$$\hat{X}^\epsilon = X^\epsilon(y_1) + \dots + X^\epsilon(y_{2l}).$$

The rest of the computation can be performed rigorously by regularizing the white noise dW_0 , using lemma 2.2 and going to the limit. It is easy to see that we obtain the same result by introducing the following formal rules:

$$E(dW_0(y)dW_0(y')) = \delta_{y-y'} dy \quad (3.25)$$

and

$$E(dW_0(y)X^\epsilon(y')) = \gamma_0(\epsilon)k_\epsilon^R(y'-y)dy \quad (3.26)$$

As a consequence of lemma 2.2, $E(e^{\hat{X}^\epsilon} dW_0(y_1) \dots dW_0(y_q))$ is the sum of terms of the form

$$E(dW_0(y_1)\hat{X}^\epsilon) \dots E(dW_0(y_k)\hat{X}^\epsilon) E(dW_0(y_{k+1})dW_0(y_{k+2})) \dots E(dW_0(y_{q-1})dW_0(y_q)) e^{\frac{1}{2}E((\hat{X}^\epsilon)^2)}. \quad (3.27)$$

We will compute the limit of each one of these terms. By using (3.26), we get

$$\begin{aligned} E(dW_0(y)X^\epsilon(y_l)) &= \gamma_0(\epsilon) \left(\sum_{i=1}^q k_\epsilon^R(y_i - y_l)\right) dy_l \\ &= \gamma_0(\epsilon) k_\epsilon^R(0) (1 + Q_l^\epsilon) dy_l \end{aligned}$$

where

$$Q_l^\epsilon = \frac{1}{k_\epsilon^R(0)} \left(\sum_{i \neq l} k_\epsilon^R(y_i - y_l)\right).$$

We also have:

$$e^{\frac{1}{2}E((\hat{X}^\epsilon)^2)} = e^{\left(\frac{q}{2}\rho_{\epsilon/R}(0) + \sum_{i < j} \rho_{\epsilon/R}\left(\frac{y_i - y_j}{R}\right)\right)(\gamma_0(\epsilon)^2 + \gamma_1^2)}.$$

By using lemma 2.2, expression (3.24) and the rules above, we get:

$$\begin{aligned} E\left(\left(\int_{\mathbb{R}^d} f_{\epsilon,h}(y) e^{X^\epsilon(y)-C_\epsilon} dW_0(y)\right)^{2l}\right) &= \sum_{k=0}^l \alpha_{k,l} (\gamma_0(\epsilon))^{2k} (k_\epsilon^R(0))^{2k} e^{(2l-k)((\gamma_0(\epsilon))^2 + \gamma_1^2)\rho_{\epsilon/R}(0) - 2lC_\epsilon} \\ &\int_{(\mathbb{R}^d)^{k+l}} f_{\epsilon,h}(y_1) \dots f_{\epsilon,h}(y_{2k}) (f_{\epsilon,h}(y_{2k+1}))^2 \dots (f_{\epsilon,h}(y_{k+l}))^2 \prod_{i=1}^{2k} (1 + Q_{i,k,l}^\epsilon) e^{S_{k,l}^\epsilon} dy_1 \dots dy_{k+l} \end{aligned}$$

where

$$Q_{i,k,l}^\epsilon = \frac{1}{k_\epsilon^R(0)} \left(\sum_{\substack{1 \leq j \leq 2k \\ j \neq i}} k_\epsilon^R(y_i - y_j) + 2 \sum_{j > 2k} k_\epsilon^R(y_i - y_j) \right)$$

and

$$\begin{aligned} S_{k,l}^\epsilon &= ((\gamma_0(\epsilon))^2 + \gamma_1^2) \left(\sum_{1 \leq i < j \leq 2k} \rho_{\epsilon/R} \left(\frac{y_i - y_j}{R} \right) + 2 \sum_{1 \leq i \leq 2k} \sum_{j > 2k} \rho_{\epsilon/R} \left(\frac{y_i - y_j}{R} \right) \right) \\ &+ 4 \sum_{2k+1 \leq i < j \leq k+l} \rho_{\epsilon/R} \left(\frac{y_i - y_j}{R} \right). \end{aligned}$$

We first take care of the normalizing constant outside each integral:

$$(\gamma_0(\epsilon) k_\epsilon^R(0) e^{-1/2((\gamma_0(\epsilon))^2 + \gamma_1^2) \rho_{\epsilon/R}(0)})^{2k} e^{2l((\gamma_0(\epsilon))^2 + \gamma_1^2) \rho_{\epsilon/R}(0) - 2lC_\epsilon}.$$

By the choice of C_ϵ , we have $e^{2l((\gamma_0(\epsilon))^2 + \gamma_1^2) \rho_{\epsilon/R}(0) - 2lC_\epsilon} = 1$. Using expansions (2.4) and (2.5), we derive the following equivalent:

$$\gamma_0(\epsilon) k_\epsilon^R(0) e^{-1/2((\gamma_0(\epsilon))^2 + \gamma_1^2) \rho_{\epsilon/R}(0)} \underset{\epsilon \rightarrow 0}{\sim} \frac{\gamma_0^* C_0 e^{-1/2\gamma_1^2 C_1}}{R^{d/2}}.$$

In conclusion, the constant outside the integral of term k in the above sum is $\alpha_{k,l} \left(\frac{\gamma_0^* C_0 e^{-1/2\gamma_1^2 C_1}}{R^{d/2}} \right)^{2k}$.

Let γ be such that $\gamma_1^2 < \gamma^2$. One can choose $\epsilon_0 > 0$ such that $\gamma_0(\epsilon_0)^2 + \gamma_1^2 < \gamma^2$. Using the fact that, for all y , $\rho_{\epsilon/R}(y/R) \leq \omega_d \ln^+ \frac{R}{y} + C$ with C independent of ϵ , we get:

$$\begin{aligned} e^{S_{k,l}^\epsilon} &\leq C \prod_{1 \leq i < j \leq 2k} \frac{1}{\left| \frac{y_i - y_j}{R} \right|_*^{\gamma^2 \omega_d}} \prod_{\substack{1 \leq i \leq 2k \\ j > 2k}} \frac{1}{\left| \frac{y_i - y_j}{R} \right|_*^{2\gamma^2 \omega_d}} \times \\ &\prod_{2k+1 \leq i < j \leq k+l} \frac{1}{\left| \frac{y_i - y_j}{R} \right|_*^{4\gamma^2 \omega_d}}. \end{aligned} \quad (3.28)$$

Finally, we conclude by using inequality (3.11) and (3.15) similarly as in the proof of proposition 3.5. □

Remark 3.8. *One can easily deduce from this that for γ_1^2 sufficiently small, by Kolmogorov's compactness theorem, \mathcal{X}^ϵ tends to \mathcal{X} in the functional sense and that \mathcal{X} is locally holderian.*

Comment 3.9. *Starting with a two parameter (R, α) monofractal gaussian field, we constructed a four parameter $(R, \alpha, \gamma_1, \gamma_0^*)$ multifractal field with nonsymmetrical increments. In dimension $d = 1$, this family can be used for financial modeling and, in any case, has it's own interest. Unfortunately, this family is inappropriate to modelize the velocity of turbulent flows where, as we shall see, the 4/5-law of Kolmogorov imposes the condition $\tilde{\zeta}_3 = 1$: indeed, a look at expression (3.21) shows that, for this family, $\tilde{\zeta}_3 > 2$.*

In the case where $\gamma_0^* = 0$, we obtain symmetrical random fields which extend to higher dimensions the model introduced in [2].

In the next section, we will study a multifractal field which is not in this family but that can be seen as a limit case where $\gamma_1 = 0$ and γ_0 is constant (independent of ϵ). As we will see, this family will be compatible with the 4/5-law.

4. THE FIELD \mathcal{X}_0

4.1. Construction of the field \mathcal{X}_0 . In this section, we only outline the main steps of the construction of \mathcal{X}_0 . The field \mathcal{X}_0^ϵ is given by formula (2.3) where X^ϵ is now defined by:

$$X^\epsilon(y) = \gamma_0 \int_{\mathbb{R}^d} k_\epsilon^R(y - \sigma) dW_0(\sigma).$$

We suppose that α is in the interval $]0, 1[$. We choose the normalizing constant C_ϵ such that:

$$\gamma_0 k_\epsilon^R(0) e^{-C_\epsilon + \frac{1}{2} \gamma_0^2 \rho_{\epsilon/R}(0)} = 1.$$

We start by stating a lemma we will use in the proof of the proposition below:

Lemma 4.1. *let δ be some real number different from d . Then there exists $C = C(\delta) > 0$ with:*

$$\int_{|u| \leq R} \frac{du}{|u|_\epsilon^\delta} \leq C \epsilon^{(d-\delta) \wedge 0}. \quad (4.1)$$

Proof. We suppose $\delta > d$, the other case being obvious. We have:

$$\begin{aligned} \int_{|u| \leq R} \frac{du}{|u|_\epsilon^\delta} & \stackrel{u = \epsilon \tilde{u}}{=} \epsilon^{d-\delta} \int_{|u| \leq R/\epsilon} \frac{d\tilde{u}}{(\int_{|v| \leq 1} \theta(v) |v + \tilde{u}| dv)^\delta} \\ & \leq \epsilon^{d-\delta} \int_{\mathbb{R}^d} \frac{d\tilde{u}}{(\int_{|v| \leq 1} \theta(v) |v + \tilde{u}| dv)^\delta}. \end{aligned}$$

□

We can now state the following proposition:

Proposition 4.2. *Let q be some positive integer satisfying:*

- (1) $q = 1$ or $q = 2$ with $\gamma_0^2 \omega_d < \alpha$.
- (2) q is even, greater or equal to 4 with $(q - 3/2) \gamma_0^2 \omega_d < \alpha \wedge \frac{d}{2}$.
- (3) q is odd, greater or equal to 3 with $(q - \frac{1}{2}) \gamma_0^2 \omega_d < \alpha \wedge \frac{d}{2}$.

Under the above condition, for all x , $\mathcal{X}_0^\epsilon(x)$ converges in L^q to a random variable $\mathcal{X}_0(x)$ such that, if e is a unit vector, we get the following scaling:

$$E((\mathcal{X}_0(x + \lambda e) - \mathcal{X}_0(x))^q) \underset{\lambda \rightarrow 0}{\sim} C_q \left(\frac{\lambda}{R}\right)^{\zeta_q}, \quad (4.2)$$

where

$$\zeta_q = q\alpha - \frac{1}{2} q(q-1) \gamma_0^2 \omega_d$$

and

$$C_q = e^{\frac{q(q-1)}{2}\phi(0)} \int_{(\mathbb{R}^d)^q} \prod_{1 \leq i < j \leq q} \frac{1}{|u_i - u_j|^{\gamma_0^2 \omega_d}} \prod_{1 \leq i \leq q} \left(\frac{1}{|u_i - e|^{\alpha}} - \frac{1}{|u_i|^{\alpha}} \right) du_1 \dots du_q. \quad (4.3)$$

Proof. In the proof, we suppose that $q = 2l$ with $l \geq 1$; we will first prove that:

$$E((\mathcal{X}_0^\epsilon(x))^q) \xrightarrow{\epsilon \rightarrow 0} \int_{(\mathbb{R}^d)^q} \prod_{1 \leq i < j \leq q} \frac{e^{\gamma_0^2 \phi(\frac{y_i - y_j}{R})}}{\left| \frac{y_i - y_j}{R} \right|^{\gamma_0^2 \omega_d}} \prod_{1 \leq i \leq q} \frac{\varphi_R(y_i)}{|y_i|^{\alpha}} dy_1 \dots dy_q. \quad (4.4)$$

We remind that the right hand side of the above limit exists by lemma 2.4. In order to prove the above relation, we develop $E((\mathcal{X}_0^\epsilon(x))^q)$ in $l + 1$ terms similarly as in the proof of proposition 3.7; then, using formula (2.4) and the fact that, for all y , $\rho_{\epsilon/R}(y) \leq \omega_d \ln \frac{1}{\epsilon} + C$, we are led to show that, for all $k \leq l - 1$, we have the following convergence:

$$\begin{aligned} & \epsilon^{(l-k)(d-\gamma_0^2 \omega_d)} \epsilon^{-2(l-k)(l-k-1)\gamma_0^2 \omega_d} \epsilon^{-4k(l-k)\gamma_0^2 \omega_d} \int_{(\mathbb{R}^d)^{k+l}} \frac{\varphi_R(y_1)}{|y_1|_\epsilon^{d-\alpha}} \dots \frac{\varphi_R(y_{2k})}{|y_{2k}|_\epsilon^{d-\alpha}} \times \\ & \frac{\varphi_R(y_{2k+1})^2}{|y_{2k+1}|_\epsilon^{2(d-\alpha)}} \dots \frac{\varphi_R(y_{k+l})^2}{|y_{k+l}|_\epsilon^{2(d-\alpha)}} \prod_{1 \leq i < j \leq 2k} \frac{e^{\gamma_0^2 \phi(\frac{y_i - y_j}{R})}}{\left| \frac{y_i - y_j}{R} \right|^{\gamma_0^2 \omega_d}} dy_1 \dots dy_{k+l} \xrightarrow{\epsilon \rightarrow 0} 0. \end{aligned}$$

We apply inequality (4.1) and obtain (if $\alpha = d/2$, one can work with $\alpha - \eta$ for $\eta > 0$ sufficiently small) :

$$\int_{\mathbb{R}^d} \frac{\varphi_R(y)^2}{|y|_\epsilon^{2(d-\alpha)}} dy \leq \epsilon^{(2\alpha-d) \wedge 0}$$

Therefore the above convergence to 0 amounts to showing that, for all $k \leq l - 1$, we have the following inequality:

$$d + (2\alpha - d) \wedge 0 - \gamma_0^2 \omega_d > 2(l - k - 1)\gamma_0^2 \omega_d + 4k\gamma_0^2 \omega_d.$$

This is equivalent to $(2l - \frac{3}{2})\gamma_0^2 \omega_d < \alpha \wedge \frac{d}{2}$. One can show, for all x , that $(\mathcal{X}_0^\epsilon(x))_{\epsilon > 0}$ is a Cauchy sequence in L^q by computing $E((\mathcal{X}_0^\epsilon(x) - \mathcal{X}_0^{\epsilon'}(x))^q)$ and letting ϵ, ϵ' go to 0. Thus, $E((\mathcal{X}_0(x))^q)$ is given by the left hand side of (4.4).

To show the scaling (4.2), observe that we can prove the following analogue to (4.4):

$$E((\mathcal{X}_0(x + \lambda e) - \mathcal{X}_0(x))^q) = \int_{(\mathbb{R}^d)^q} \prod_{1 \leq i < j \leq q} \frac{e^{\gamma_0^2 \phi(\frac{y_i - y_j}{R})}}{\left| \frac{y_i - y_j}{R} \right|^{\gamma_0^2 \omega_d}} \prod_{1 \leq i \leq q} f_{\lambda e}(y_i) dy_1 \dots dy_q, \quad (4.5)$$

where

$$f_{\lambda e}(y) = \frac{\varphi_R(y - \lambda e)}{|y - \lambda e|^{d-\alpha}} - \frac{\varphi_R(y)}{|y|^{d-\alpha}}. \quad (4.6)$$

By setting $y_i = \lambda \tilde{y}_i$ in the integral of (4.5), we deduce easily (4.2). \square

Remark 4.3. Similarly as in the previous section, for γ_0 sufficiently small, \mathcal{X}_0^ϵ converges in law to \mathcal{X}_0 in the space of continuous fields.

4.2. **Nonsymmetry of the increments of \mathcal{X}_0 .** Let q be some odd integer and consider C_q given by formula (4.3). It is clear that C_q does not depend on the unitary vector e . By making the change of variable $\tilde{u}_i = u_i - e$ in (4.3), we get:

$$C_q(e) = -C_q(-e) = -C_q(e).$$

Thus, we get $C_q = 0$. Therefore, it is not obvious to see if the law of $\mathcal{X}_0(x+h) - \mathcal{X}_0(x)$ is nonsymmetrical or not. Nevertheless, one can repeat the same construction of \mathcal{X}_0 as above replacing the kernel $\frac{\varphi_R(x-y)}{|x-y|_\epsilon^{d-\alpha}}$ by the kernel:

$$\varphi_R(x-y) \frac{x_i - y_i}{|x-y|_\epsilon^{d-\alpha+1}}, \quad i = 1 \dots d. \quad (4.7)$$

In this case, $C_1 = 0$ but, by Monte Carlo simulation, one can verify that $C_3 \neq 0$. In particular, with the kernel (4.7), the law of $\mathcal{X}_0(x+h) - \mathcal{X}_0(x)$ is nonsymmetrical.

5. A STEP TOWARDS A MODEL OF THE VELOCITY FIELD OF TURBULENT FLOWS

An acceptable solution to the problem of hydrodynamical turbulence in dimension 3 would be to construct a random velocity field U solution to the dynamics (Euler or Navier Stokes typically) that is stationary, incompressible, space-homogeneous, isotropic and that satisfies the main statistical properties of the velocity field of turbulent flows. These properties are:

- (1) The 4/5-law of Kolmogorov that links the energy dissipation of the turbulent flow to the statistics of the increments of the velocity. This law is widely accepted since it is the only one that can be proven with the dynamics ([6], [7], [13]). More precisely, this law states:

$$E\left(\left(U(x+\xi) - U(x) \cdot \frac{\xi}{|\xi|}\right)^3\right) = -\frac{4}{5}D|\xi|. \quad (5.1)$$

In the above formula, D denotes the average dissipation of the kinetic energy per unit mass in the fluid.

Remark 5.1. *To obtain this law, it is sufficient to suppose that the field U is space homogeneous and isotropic.*

- (2) The intermittency of the field U :

$$E\left(|U(x+\xi) - U(x) \cdot \frac{\xi}{|\xi|}|^q\right) \underset{|\xi| \rightarrow 0}{\sim} C_q |\xi|^{\zeta_q}, \quad (5.2)$$

where q is a positive real number and the ζ_q are called the structural exponents.

It is a very challenging task to construct a field with all the aforementioned properties, especially because this field must be invariant by the Euler or Navier-Stokes equation.

Nevertheless, one can in the first place forget the invariance condition and simply try to construct a field that satisfies all the other properties. The 4/5-law shows that the nonsymmetry of the increments is an essential feature: this is one of the

main difficulties in trying to extend the previous construction of scalar fields to 3-dimensional incompressible fields.

Quite naturally, we consider the incompressible family U^ϵ defined by:

$$U^\epsilon(x) = \int_{\mathbb{R}^3} \varphi_R(x-y) \frac{x-y}{|x-y|_\epsilon^{d-\alpha+1}} \wedge e^{X^\epsilon(y)-C_\epsilon} dW(y),$$

where $dW(y) = (dW_1(y), dW_2(y), dW_3(y))$ denotes a three dimensional white noise and X^ϵ is defined by the following formula:

$$X^\epsilon(y) = \gamma \int_{\mathbb{R}^3} K_\epsilon^R(y-\sigma) \cdot dW(\sigma),$$

with $K^R(x) = \frac{x}{|x|^{1+d/2}} 1_{|x| \leq R}$. As in the previous sections, we choose the constant C_ϵ such that U^ϵ converges to a non trivial field U as ϵ goes to 0. The vector field U we obtain is incompressible, homogeneous, isotropic and intermittent with structural exponents ζ_q defined by:

$$\zeta_q = q\alpha - 2\pi\gamma^2 q(q-1). \quad (5.3)$$

We can derive the energy dissipation D of the field U by computing the following limit:

$$-\frac{5}{4} \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} E(((U(x+\lambda e) - U(x)) \cdot e)^3),$$

where e is some unitary vector. In order to get D finite and $D \neq 0$, we must have $\zeta_3 = 1$ or equivalently:

$$\alpha = 1/3 + 4\pi\gamma^2. \quad (5.4)$$

Unfortunately, since the field $e^{X^\epsilon(y)} dW(y)$ is isotropic with respect to all unitary transformations (and not just the rotations), we get $D = 0$. Thus, the construction of an intermittent incompressible field with positive dissipation remains an open question.

Comment 5.2. *If one plugs relation (5.4) into (5.3), we get the following expression for ζ_q :*

$$\zeta_q = (1/3 + 6\pi\gamma^2)q - 2\pi\gamma^2 q^2.$$

The small scale behavior of the field depends only on the intermittency parameter γ^2 . One can easily identify it using the experimental curve obtained in [1] (cf. fig. 8.8 p. 132 in [7]): with their data, we find $4\pi\gamma^2 = 0.023$. Thus, the intermittency parameter is small, a situation that seems to be similar in finance for certain assets; indeed, if one denotes by γ the intermittency parameter of the MRW model introduced by Bacry, Delour and Muzy ([2]), the authors of [4] estimate $\gamma^2 \approx 0.03$ for the S&P500 future index using intraday data over the period 1988-1999.

REFERENCES

- [1] Anselmet, F., Antonia, R.A., Gagne, Y., Hopfinger, E.J.: High-order velocity structure functions in turbulent shear flow, *J.Fluid Mech.*, **140** (1984), 63-89.
- [2] Bacry E., Delour J., Muzy J.-F.: Modelling financial time series using multifractal random walks, *Physica A*, **299** (2001), 84-92.

- [3] Bacry E., Muzy J.-F.: Log-infinitely divisible multifractal processes, *Commun. Math. Phys.*, **236** (2003), 449-475.
- [4] Bacry E., Kozhemyak, A., Muzy J.-F.: Continuous cascade models for asset returns, available at www.cmap.polytechnique.fr/~bacry/biblio.html, to appear in *Journal of Economic Dynamics and Control*.
- [5] Daley D.J., Vere-Jones D.: An introduction to the theory of point processes, Springer-Verlag, (1988).
- [6] Duchon, J., Robert, R.: Inertial energy dissipation for weak solutions of incompressible Euler and Navier-Stokes equations, *Nonlinearity*, **13** (2000), no.1, 249-255.
- [7] Frisch, U.: Turbulence, Cambridge University Press, (1995).
- [8] Kahane, J.-P.: Sur le chaos multiplicatif, *Ann. Sci. Math. Québec*, **9** (1985), no.2, 105-150.
- [9] Kolmogorov A.N.: A refinement of previous hypotheses concerning the local structure of turbulence, *J. Fluid. Mech.*, **13** (1962), 83-85.
- [10] Mandelbrot B.B.: A possible refinement of the lognormal hypothesis concerning the distribution of energy in intermittent turbulence, *Statistical Models and Turbulence*, La Jolla, CA, Lecture Notes in Phys. no. 12, Springer, (1972), 333-335.
- [11] Nualart D.: The Malliavin Calculus and Related Topics, Springer Verlag, (1995).
- [12] Obukhov A.M.: Some specific features of atmospheric turbulence, *J. Fluid. Mech.*, **13** (1962), 77-81.
- [13] Robert, R.: Mathématiques et turbulence, *Images des mathématiques, CNRS*, (2004), 91-100.