



HAL
open science

Integers with a large friable component

Gérald Tenenbaum

► **To cite this version:**

| Gérald Tenenbaum. Integers with a large friable component. 2006. hal-00091306v1

HAL Id: hal-00091306

<https://hal.science/hal-00091306v1>

Preprint submitted on 5 Sep 2006 (v1), last revised 18 Jan 2007 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Integers with a large friable component

Gérald Tenenbaum

1. Introduction and statement

It is well-known (see, e.g., [‘Te95’], chap. III.3) that the logarithms of the prime factors of an integer normally have exponential growth. Therefore, it is expected that the product of the small prime factors of a typical integer remains small—a device which has been employed by Erdős in many different contexts and for which various effective versions appear in the literature. We return here to the problem of finding a quantitative estimate for the number of exceptional integers. Some similar results have been obtained concomitantly, through a more elementary approach, by Banks and Shparlinski [‘BS06’].

Given an integer n and a real parameter $y \geq 1$, we define

$$n_y := \prod_{p^\nu \parallel n, p \leq y} p^\nu$$

as the y -friable component of n and we put

$$\Theta(x, y, z) := \sum_{\substack{n \leq x \\ n_y > z}} 1 \quad (x \geq 1, y \geq 1, z \geq 1).$$

We also write, for complex s with positive real part,

$$\zeta(s, y) := \sum_{P(n) \leq y} 1/n^s = \prod_{p \leq y} (1 - p^{-s})^{-1},$$

where $P(n)$ denotes the largest prime factor of n with the convention that $P(1) := 1$, we let ϱ designate Dickman’s function and we set

$$S(y, z) := \sum_{\substack{P(m) \leq y \\ m > z}} \frac{1}{m}.$$

It has been shown in [‘Te99’], Corollary 2, that, writing $u := (\log x)/\log y$, we have

$$(1.1) \quad \Theta(x, y, z) = \frac{xS(y, z)}{\zeta(1, y)} + O\left(x\varrho(u)2^{(1+\varepsilon)u} + x^\varepsilon\right)$$

uniformly for $x \geq 2$, $y \geq 2$, $z \geq 2$, and also that, denoting Euler’s constant by γ , and writing

$$\tau(w) := \int_w^\infty \varrho(t) dt \quad (w \geq 0),$$

we have, with $v := (\log z)/\log y$,

$$(1.2) \quad S(y, z) = \left\{1 + O\left(\frac{\log(v+2)}{\log y}\right)\right\} \tau(v) \log y$$

for all $\varepsilon > 0$ and uniformly for

$$(1.3) \quad y \geq 2, \quad 1 \leq z \leq \exp \exp \{(\log y)^{3/5-\varepsilon}\}.$$

Thus, for all $x \geq 2$ and y, z satisfying (1.3), we have

$$(1.4) \quad \Theta(x, y, z) = \left\{e^{-\gamma} + O\left(\frac{\log(v+2)}{\log y}\right)\right\} x\tau(v) + O\left(x\varrho(u)2^{(1+\varepsilon)u} + x^\varepsilon\right).$$

Note that $\Theta(x, y, z) = 0$ unless $z \leq x$ and that, when the latter holds, (1.3) is implied by Hildebrand’s condition

$$(H_\eta) \quad x \geq 2, \quad \exp \{(\log_2 x)^{5/3+\eta}\} \leq y \leq x$$

with $\eta = 3\varepsilon$.

In particular, if condition (1.3) holds, $u \rightarrow \infty$, $y \rightarrow \infty$, and, say, $z \leq x^{1-\varepsilon}$, then

$$(1.5) \quad \Theta(x, y, z) \sim e^{-\gamma} \tau(v)x.$$

Formula (1.1) has been derived in [‘Te99’] as a by-product of a general result on the Kubilius model of probabilistic number theory. However, the estimates established in [‘Te99’] easily yield an asymptotic formula that is valid also when u is bounded. Let ω denote Buchstab’s function. We obtain the following result in which we put

$$\begin{aligned}\sigma(u, v) &:= \int_v^\infty \varrho(t)\omega(u-t) dt \\ \vartheta(u, v) &:= \varrho(u) + \sigma(u, v) \\ \kappa(u, v) &:= (1-\gamma)\varrho(u-1) + \gamma\varrho(v)\omega(u-v)\end{aligned}\quad (u \geq 1, v \geq 0).$$

Theorem 1.1. *Let $\varepsilon > 0$. Under conditions (1.3) and $1 \leq z \leq x/y$, we have uniformly*

$$(1.6) \quad \Theta(x, y, z) = x \left\{ \vartheta(u, v) - \frac{\kappa(u, v)}{\log y} + O\left(\frac{\tau(v)}{\log y} + \frac{\varrho(v) \log(v+2)}{(\log y)^2} + \frac{1}{z}\right) \right\}.$$

In particular, if $(\log y)^2 \leq z \leq \min\left(x/y^{1+\varepsilon/\log(u+1)}, e^{\sqrt{y}}\right)$, we have

$$(1.7) \quad \Theta(x, y, z) = \left\{1 + O\left(\frac{\log(v+2)}{\log y}\right)\right\} \vartheta(u, v)x.$$

Remarks. (i) The condition $z \leq x/y$ is not restrictive since $\Theta(x, y, z) = \Psi(x, y) - \Psi(z, y)$ otherwise.

(ii) It will be clear from the proof that, under suitable assumptions, more precise estimates may be derived by the same method.

(iii) It follows from classical estimates on Dickman’s function (see e.g. [‘Te95’], chap. III.5) that

$$(1.8) \quad \tau(w) = \frac{\varrho(w)}{\log(w+1)} \left\{1 + O\left(\frac{1}{\log(w+1)}\right)\right\} \quad (w \geq 1).$$

Since $\frac{1}{2} \leq \omega(t) \leq 1$ ($t \geq 1$) and $\varrho(v+h) \gg \varrho(v)\{v \log(v+1)\}^{-h}$ ($0 \leq h \leq 1 \leq v$) (see e.g. [‘Te95’], chaps. III.5 and III.6), this implies, for instance, that $\vartheta(u, v) \asymp \tau(v)$ whenever $u-1-v \gg 1/\log(v+1)$.

2. Proof of ‘evalT’

We start with an improvement of (1.2) established by the same method. As usual, we define the derivatives of the Dickman function by right continuity at integer points. It is known (see e.g. [‘Te95’], cor. III.5.8.3) that $\varrho^{(k)}(w) \sim (-1)^k (\log w)^k \varrho(w)$ as $w \rightarrow \infty$. We also denote by $\{a_j\}_{j=0}^\infty$ the Taylor coefficients of $s\zeta(s+1)/(s+1)$ at $s=0$.

Lemma 2.1. *Let $\varepsilon > 0$ and $k \in \mathbb{N}$ be given. Then, uniformly under condition (1.3), we have*

$$(2.1) \quad S(y, z) = \tau(v) \log y - \sum_{0 \leq j \leq k} a_{j+1} \frac{\varrho^{(j)}(v)}{(\log y)^j} - \frac{\Psi(z, y)}{z} + O\left(\frac{\varrho(v)\{\log(v+2)\}^{k+1}}{(\log y)^{k+1}}\right).$$

In particular, we have

$$(2.2) \quad S(y, z) = \tau(v) \log y - \gamma\varrho(v) + O\left(\frac{\varrho(v) \log(v+2)}{\log y} + \frac{1}{z}\right).$$

Proof. We have

$$S(y, z) = (\log y) \int_v^\infty \frac{\Psi(y^w, y)}{y^w} dw - \frac{\Psi(z, y)}{z}.$$

Inserting Saias’ estimate for $\Psi(y^w, y)$ (see [‘Sa89’] or [‘Te95’], th. III.5.9) in its range of validity and estimating the contribution of large w as in [‘Te99’], we obtain

$$\int_v^\infty \frac{\Psi(y^w, y)}{y^w} dw = \left\{1 + O\left(e^{-(\log y)^{3/5-\varepsilon}}\right)\right\} \int_{0-}^\infty \tau(v-t) d\left(\frac{[y^t]}{y^t}\right).$$

Formula (2.1) follows by integrating by parts and inserting the generalized Taylor expansion for the Dickman function established in [‘FT91’], lemma 4.2. We omit the details, which are very similar to those in [‘FT91’]. Then, we derive (2.2) by appealing to Hildebrand’s formula [‘Hi86a’]

$$(2.3) \quad \Psi(z, y) = z\varrho(v) \left\{1 + O\left(\frac{\log(v+1)}{\log y}\right)\right\} + O(1) \quad (z \geq 2, y \geq e^{(\log_2 z)^{5/3+\varepsilon}}).$$

□

For $y \geq 2$, $t \geq 1$, we write $u_t := (\log t)/\log y$.

Lemma 2.2. Under conditions (1.3) and $1 \leq z \leq x/y$, we have uniformly

$$(2.4) \quad \sum_{\substack{P(m) \leq y \\ m > z}} \frac{\omega(u - u_m)}{m} = \sigma(u, v) \log y - \kappa(u, v) + \varrho(u - 1) + O\left(\tau(v) + \frac{\varrho(v) \log(v + 2)}{\log y} + \frac{1}{z}\right).$$

Proof. Since $\omega(s)$ is continuous for $s \geq 1$ and differentiable for $s > 1$, we have

$$\begin{aligned} \sum_{\substack{P(m) \leq y \\ m > z}} \frac{\omega(u - u_m)}{m} &= \sum_{\substack{P(m) \leq y \\ z < m \leq x/y}} \frac{1}{m} \left\{ 1 + \int_{u_m}^{u-1} \omega'(u - t) dt \right\} \\ &= S(y, z) - S(y, x/y) + \int_v^{u-1} \omega'(u - t) \{S(y, z) - S(y, y^t)\} dt \\ &= \omega(u - v)S(y, z) - S(y, x/y) - \int_v^{u-1} \omega'(u - t)S(y, y^t) dt. \end{aligned}$$

The required formula then follows by inserting (2.2), using the estimate (1.8). We omit the details which only involve standard partial integration and the fact that $\omega' \in L^1(\mathbb{R})$. \square

We are now in a position to complete the proof of ‘evalT’.

We first consider the case when (x, y) lies outside the region (H_ε) . Appealing to the bound $\omega(t) - e^{-\gamma} \ll \varrho(t)$ ($t \geq 0$) established, in a more precise form, in lemma 4 of [‘Te99’] and noting that, say, $u > 3v$ provided y is large enough, we see that

$$\begin{aligned} \sigma(u, v) &= e^{-\gamma} \tau(v) \{1 + O(1/(\log y)^2)\}, \\ \kappa(u, v) &= \gamma e^{-\gamma} \varrho(v) \{1 + O(1/(\log y)^2)\}. \end{aligned}$$

Thus (1.6) is in this case an immediate consequence of (1.1) and (2.2).

When $(x, y) \in H_\varepsilon$, $z \leq x/y$, we apply the formula

$$\Theta(x, y, z) = \sum_{\substack{P(m) \leq y \\ z < m \leq x/y}} \Phi\left(\frac{x}{m}, y\right) + \Psi(x, y) - \Psi(x/y, y),$$

where $\Phi(t, y)$ denotes the number of positive integers not exceeding t and all of whose prime factors exceed y . The last two terms may be evaluated by (2.3). Note that $\Psi(x/y, y)$ may be regarded as an error term since this is $\ll x\varrho(v)/y$. Thus, we may restrict our attention to evaluating the m -sum. To this end, we apply Corollary 3 of [‘Te99’] in the form

$$\Phi\left(\frac{x}{m}, y\right) = \frac{e^\gamma x \omega(u - u_m)}{m \zeta(1, y)} - \frac{e^\gamma y}{\zeta(1, y)} + O\left(\frac{x \varrho(u - u_m)}{m (\log y)^2}\right) \quad (z < m \leq x/y).$$

By (1.2), the contribution of remainder term of the left-hand side is dominated by that of (1.6) and that of the second term equals $-x\varrho(u - 1)/\log y$ to within an acceptable error. Since the remaining sum depends on (2.4), this completes the proof of (1.6). Formula (1.7) then follows from the bound $\tau(v) \gg v^{-2v}$.

References

- [‘BS06’] W.D. Banks & I. Shparlinski, Integers with a large smooth divisor, preprint.
- [‘FT91’] E. Fouvry & G. Tenenbaum, Entiers sans grand facteur premier en progressions arithmétiques, *Proc. London Math. Soc.* (3) **63** (1991), 449–494.
- [‘Hi86a’] A. Hildebrand, On the number of positive integers $\leq x$ and free of prime factors $> y$, *J. Number Theory* **22** (1986), 265–290.
- [‘Sa89’] E. Saias, Sur le nombre des entiers sans grand facteur premier, *J. Number Theory* **32** (1989), 78–99.
- [‘Te95’] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge studies in advanced mathematics, no. 46, Cambridge University Press (1995).

- [Te99] G. Tenenbaum, Crible d'Ératosthène et modèle de Kubilius, in: K. Györy , H. Iwaniec, J. Urbanowicz(eds.), *Number Theory in Progress*, Proceedings of the conference in honor of Andrzej Schinzel, Zakopane, Poland 1997, 1099–1129, Walter de Gruyter, Berlin, New York, 1999.

Gérald Tenenbaum
Institut Élie Cartan
Université Henri Poincaré–Nancy 1
BP 239
54506 Vandœuvre Cedex
France
internet: gerald.tenenbaum@iecn.u-nancy.fr