



# The brachiopod Lingula in the Middle Miocene of the Central parathethys

Christian Emig, Maria Aleksandra Bitner

## ► To cite this version:

Christian Emig, Maria Aleksandra Bitner. The brachiopod Lingula in the Middle Miocene of the Central parathethys. Acta Palaeontologica Polonica, 2005, 50, pp.181-184. hal-00093666

HAL Id: hal-00093666

<https://hal.science/hal-00093666>

Submitted on 13 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## The brachiopod *Lingula* in the Middle Miocene of the Central Paratethys

Christian C. Emig and Maria Aleksandra Bitner

*Acta Palaeontologica Polonica* 50 (1), 2005: 181-184

In the Middle Miocene deposits of the Central Paratethys, lingulides, because of their low taphonomic potential, are

very rare and difficult to identify. Two species of *Lingula*, *L. dumortieri* Nyst, 1843 and *L. dregeri* Andreæ, 1893, have been recorded from the Badenian (Middle Miocene) of Poland.

Re-examination of the specimens has shown that none of them can be referred to *Lingula dumortieri* (now *Glottidia dumortieri*). The rounded outline of the scar of the posterior adductor muscle and the lack of the septa indicate that all the specimens must be referred to the genus *Lingula*. Their valves appear to be slightly more convex than in other species of *Lingula* and closely resemble the Recent *L. tumidula* Reeve, 1841. We assign all the specimens of *Lingula* found in the Miocene of Poland to *L. dregeri*. Because our specimens are juveniles and some are broken we cannot adequately redescribe this species. *Lingula dregeri* was distributed in the Middle Miocene of the Central Paratethys (Austria, Poland, Ukraine, Romania), while *Glottidia dumortieri* occurs in the Pliocene of Northern Europe (Atlantic Province).

Christian C. Emig [[christian.emig@com.univ-mrs.fr](mailto:christian.emig@com.univ-mrs.fr)], CNRS–UMR 6540, Centre d’Océanologie, Chemin de la Batterie-des-Lions, F 13007 Marseille, France;  
Maria Aleksandra Bitner [[bitner@twarda.pan.pl](mailto:bitner@twarda.pan.pl)], Instytut Paleobiologii, Polska Akademia Nauk, ul. Twarda 51/55, PL 00–818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

