



HAL
open science

Non-diffusive phase spreading of a Bose-Einstein condensate at finite temperature

Alice Sinatra, Yvan Castin, Emilia Witkowska

► **To cite this version:**

Alice Sinatra, Yvan Castin, Emilia Witkowska. Non-diffusive phase spreading of a Bose-Einstein condensate at finite temperature. 2006. hal-00109021v1

HAL Id: hal-00109021

<https://hal.science/hal-00109021v1>

Preprint submitted on 23 Oct 2006 (v1), last revised 18 Jan 2007 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non-diffusive phase spreading of a Bose-Einstein condensate at finite temperature

A. Sinatra and Y. Castin

*Laboratoire Kastler Brossel, Ecole Normale Supérieure,
24 Rue Lhomond, 75231 Paris Cedex 05, France*

E. Witkowska

*Institute of Physics, Polish Academy of Sciences,
Aleja Lotników 32/46, 02-668 Warszawa, Poland*

We show that the phase of a condensate in a finite temperature gas spreads linearly in time at long times rather than in a diffusive way. This result is supported by classical field simulations, and analytical calculations which are generalized to the quantum case. We show that this super-diffusive behavior is intimately related to conservation of energy during the free evolution of the system and to fluctuations of energy in the prepared initial state.

PACS numbers: 03.75.Kk, 03.75.Pp

I. INTRODUCTION

Phase coherence is one of the fundamental properties of Bose-Einstein condensates. It is also a key feature in the present developments of the research on condensates which, ten years after the first experimental realization, go in the direction of integrating this powerful tool into other branches of physics, of which metrology and quantum information are two promising examples [1].

The problem of the condensate phase dynamics due to atomic interactions at zero temperature has been analyzed by different authors in theory [2] and in experiment [3, 4, 5]. It is now well understood that an initially prepared relative phase between two condensates will spread in time due to the corresponding uncertainty in the relative particle number as the relative phase and the relative particle number are conjugate variables. The phase dynamics of a two component condensate in realistic situations including harmonic traps, non stationarity and fluctuations in the total number of particles was analyzed in [6], where a comparison to the experiments of [4] is also performed. An important conclusion was that the zero temperature theory could not account for the coherence times observed in experiment, which raises the question of the role of the non-condensed fraction.

In this paper we address the fundamental problem of phase spreading of a Bose-Einstein condensate in a finite temperature atomic gas. In order to obtain simple and general results, we consider the ideal case of a spatially uniform condensate at thermodynamical equilibrium, and we assume that one has access to the first order temporal correlation function $\langle a_0^\dagger(t)a_0 \rangle$ of the component a_0 of the atomic field in the condensate mode. In real life, the situation is more complex: the atoms are trapped in harmonic potentials, and the measurement of phase coherence is a delicate procedure, usually relying on the interference between two condensates [4]. It was predicted that at finite temperature the condensate phase would show a diffusive behavior (variance growing linearly in time) [7, 8, 9, 10]. We study this problem first

with a classical field model [11, 12, 13, 14, 15, 16], where exact numerical simulations can be performed. We then explain the numerics analytically, and extend the analytical approach to the quantum case.

The important result that we obtain is that the variance of the phase increases quadratically in time. This is at variance with the prediction of phase diffusion from the “quantum optics” open system approaches of [7, 8, 9, 10] assuming the condensate to evolve under the influence of Langevin short memory fluctuating forces. Our prediction results from two ingredients, (i) the system is prepared in an initial state with an energy fluctuating from one experimental realization to the other, here sampling the canonical ensemble, and (ii) the system is isolated in its further evolution and therefore keeps a constant energy. As we shall see, the combination of these two ingredients prevents some temporal correlation functions to vanish at long times.

In section II we present the classical field model; numerical predictions for this model are presented in section III, and analytical results reproducing the numerics at short or long times are given in section IV. These analytical results are extended to the case of the quantum field in section V. We conclude in section VI.

II. THE CLASSICAL FIELD MODEL

In this section we develop a classical field model that has the advantage that it can be exactly simulated numerically. This will allow us to understand the physics governing the spreading of the condensate phase and to test the validity of various approximations, paving the way to the quantum treatment.

We consider a lattice model for a classical field $\psi(\mathbf{r})$ in three dimensions. The lattice spacings are l_1, l_2, l_3 along the three directions of space and $dV = l_1 l_2 l_3$ is the volume of the unit cell in the lattice. We enclose the atomic field in a spatial box of sizes L_1, L_2, L_3 and volume $V = L_1 L_2 L_3$, with periodic boundary conditions.

The discretized field has the following Poisson brackets

$$i\hbar\{\psi(\mathbf{r}_1), \psi^*(\mathbf{r}_2)\} = \frac{\delta_{\mathbf{r}_1, \mathbf{r}_2}}{dV} \quad (1)$$

where the Poisson brackets are such that $df/dt = \{f, H\}$ for a time-independent functional f of the field ψ . The field ψ may be expanded over the plane waves

$$\psi(\mathbf{r}) = \sum_{\mathbf{k}} a_{\mathbf{k}} \frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{V}}, \quad (2)$$

where \mathbf{k} is restricted to the first Brillouin zone, $k_\alpha \in [-\pi/l_\alpha, \pi/l_\alpha[$ where α labels the directions of space.

We assume that, in the real physical system, the total number of atoms is fixed, equal to N . In the classical field model, this fixes the norm squared of the field:

$$dV \sum_{\mathbf{r}} |\psi(\mathbf{r})|^2 = N. \quad (3)$$

Equivalently the density of the system

$$\rho = \frac{N}{V} \quad (4)$$

is fixed for each realization of the field. The evolution of the field is governed by the Hamiltonian

$$H = \sum_{\mathbf{k}} \tilde{E}_k a_{\mathbf{k}}^* a_{\mathbf{k}} + \frac{g}{2} \sum_{\mathbf{r}} dV \psi^*(\mathbf{r}) \psi^*(\mathbf{r}) \psi(\mathbf{r}) \psi(\mathbf{r}), \quad (5)$$

where \tilde{E}_k is the dispersion relation of the non-interacting waves, and the binary interaction between particles in the real gas is reflected in the classical field model by a field self-interaction with a coupling constant $g = 4\pi\hbar^2 a/m$, where a is the s -wave scattering length of two atoms.

In general, we expect the predictions of a classical field model to be cut-off dependent, i.e. the predictions of our model may depend on the lattice spacings l_α . We use here a refinement to the usual classical field model, which makes it cut-off independent for *some* observables like the condensate fraction, a quantity expected to play an important role here. An obvious example of a quantity which will remain cut-off dependent is the mean value of the Hamiltonian H in thermal equilibrium.

Let us consider first the non-interacting case ($g = 0$) in presence of a condensate. For a thermalized classical field the occupation numbers of the excited plane wave modes are given by the equipartition formula

$$\langle a_{\mathbf{k}}^* a_{\mathbf{k}} \rangle = \frac{k_B T}{\tilde{E}_k}. \quad (6)$$

We adjust the dispersion relation \tilde{E}_k in order to reproduce the Bose law for the occupation numbers of the quantum field in the Bose-condensed regime:

$$\frac{1}{e^{\beta\hbar^2 k^2/2m} - 1} = \frac{k_B T}{\tilde{E}_k}. \quad (7)$$

For all modes with large occupation number $\tilde{E}_k \simeq \hbar^2 k^2/2m$, while the occupation of modes with $\hbar^2 k^2/2m \gg k_B T$, whose quantum dynamics is not well approximated by the classical field model anyway, is exponentially suppressed as in the quantum theory.

In the interacting case, one could adapt the same trick of a modified dispersion relation, by including the fact that the relevant spectrum is not $\hbar^2 k^2/2m$ but the Bogoliubov spectrum [17]. The resulting \tilde{E}_k would now start growing exponentially with k when the Bogoliubov energy $[(\hbar^2 k^2/2m)(2\rho g + \hbar^2 k^2/2m)]^{1/2}$ reaches $k_B T$.

In the classical field model we restrict our analysis to the regime $k_B T \gg \rho g$ so that at energies of the order of $k_B T$, the Bogoliubov energy is dominated by the kinetic term $\hbar^2 k^2/2m$. One can then simply use in the Hamiltonian the modified dispersion relation \tilde{E}_k as given by Eq.(7). This is what we did in the simulations of this paper, so that the classical field ψ evolves according to the non-linear equation [18]:

$$i\hbar \partial_t \psi = \left\{ k_B T \left[\exp\left(-\beta \frac{\hbar^2}{2m} \Delta\right) - 1 \right] + g|\psi(\mathbf{r}, t)|^2 \right\} \psi. \quad (8)$$

In practice this equation is integrated numerically with the FFT splitting technique.

We then introduce the density and the phase of the condensate mode

$$a_0 = e^{i\theta} \sqrt{n_0}. \quad (9)$$

In what follows, we will concentrate on three physical quantities: the condensate amplitude correlation function

$$\langle a_0^*(t) a_0(0) \rangle, \quad (10)$$

the condensate atom number correlation function

$$\langle \delta n_0(t) \delta n_0(0) \rangle \quad \text{where} \quad \delta n_0 = n_0 - \langle n_0 \rangle, \quad (11)$$

and the variance of the condensate phase change during t :

$$\text{Var } \varphi(t) = \langle \varphi(t)^2 \rangle - \langle \varphi(t) \rangle^2 \quad \text{where} \quad \varphi(t) = \theta(t) - \theta(0). \quad (12)$$

The averages are taken over stochastic realizations of the classical field, as the initial field samples a thermal probability distribution.

III. CLASSICAL FIELD: NUMERICAL RESULTS

We consider a gas of $N = 4 \times 10^5$ atoms with $\rho g = 700 \hbar^2/mV^{2/3}$ in a box of non commensurable square lengths to guarantee efficient ergodicity in the system, in the ratio $L_1^2 : L_2^2 : L_3^2 = \sqrt{2} : (1 + \sqrt{5})/2 : \sqrt{3}$. We choose the number of the lattice points in a temperature dependent way, such that the maximal Bogoliubov

energy $[(\hbar^2 k^2/2m)(2\rho g + \hbar^2 k^2/2m)]^{1/2}$ on the lattice is equal to $3k_B T$.

To generate the stochastic initial values of the classical field we proceed as follows. (i) For each realization, we generate a non condensed field $\psi_{\perp}(\mathbf{r})$ at temperature T in the Bogoliubov approximation as explained in [19]. In practice we generate complex numbers $\{b_{\mathbf{k}}\}$ for each vector \mathbf{k} on the grid according to the probability distribution

$$P(b_{\mathbf{k}}) = \frac{1}{\pi} \frac{\tilde{\epsilon}_k}{k_B T} e^{-(|b_{\mathbf{k}}|^2 \tilde{\epsilon}_k/k_B T)} \quad (13)$$

where $\tilde{\epsilon}_k = [\tilde{E}_k(\tilde{E}_k + 2\rho g)]^{1/2}$. With a set of $\{b_{\mathbf{k}}\}$ for a given realization we build the non condensed field

$$\psi_{\perp}(\mathbf{r}) = e^{i\theta} \sum_{\mathbf{k} \neq 0} \left(b_{\mathbf{k}} \tilde{U}_k \frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{V}} + b_{\mathbf{k}}^* \tilde{V}_k \frac{e^{-i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{V}} \right) \quad (14)$$

where the initial value of the condensate phase θ is randomly chosen with the uniform law in $[0, 2\pi[$, and where the real amplitudes \tilde{U}_k, \tilde{V}_k , normalized as $\tilde{U}_k^2 - \tilde{V}_k^2 = 1$, are given by the usual Bogoliubov theory, here with the modified dispersion relation, so that

$$\tilde{U}_k + \tilde{V}_k = \left(\frac{\tilde{E}_k}{\tilde{E}_k + 2\rho g} \right)^{1/4}. \quad (15)$$

(ii) We create the classical field with the constraint that the total number of atoms N is fixed:

$$\psi(\mathbf{r}) = \frac{a_0}{\sqrt{V}} + \psi_{\perp}(\mathbf{r}) \quad (16)$$

where $a_0 = \sqrt{N - N_{\perp}} e^{i\theta}$, N_{\perp} is the number of non condensed atoms,

$$N_{\perp} = \sum_{\mathbf{r}} dV |\psi_{\perp}(\mathbf{r})|^2. \quad (17)$$

(iii) We let the field evolve for some time interval with the Eq.(8) to eliminate transients due to the fact that the Bogoliubov approximation used in the sampling does not produce an exactly stationary distribution. After this ‘thermalization’ period we start calculating the relevant observables, as ψ evolves with the same Eq.(8).

First we investigate the mean condensate phase change $\langle \varphi \rangle(t)$. We find a linear dependence with time, with a slope slightly different from the value $-\rho g$ naively expected, e.g. from the zero temperature Gross-Pitaevskii equation. The slope difference is temperature dependent and is expected physically to correspond to the discrepancy between the zero temperature chemical potential ρg and the actual finite temperature one $\mu(T)$. This we shall confirm using Bogoliubov theory in Sec. IV (see also [20]).

In figure 1, we show the real part of the amplitude correlation function of the condensate $\langle a_0^*(t) a_0(0) \rangle$ as a function of time, for a temperature $T = 0.17T_c$, where

T_c is the critical temperature of the ideal gas. The zero-temperature evolution $e^{-i\rho g t/\hbar}$ is subtracted so that the oscillations in the figure are due to the above mentioned effect $\mu(T) \neq \rho g$. Due to the finite temperature in the system, the correlation function of the condensate amplitude is smeared out at long times.




FIG. 1: (Color online) Real part of the condensate amplitude correlation function (10) normalized to its $t = 0$ value, and divided by the zero temperature evolution $e^{-i\rho g t/\hbar}$, as a function of time: in the vertical axis label, $\tilde{a}_0(t)$ stands for $a_0(t)e^{i\rho g t/\hbar}$. (a) Short times behavior and (b) long times behavior. In solid line from an average over 500 solutions of Eq.(8), in dashed line (red) the Bogoliubov approximation (39). Here the temperature is $k_B T = 3077.3 \hbar^2/mV^{2/3} = 0.1711T_c$, where T_c is the critical temperature $k_B T_c = (2\pi\hbar^2/m)(\rho/\zeta(3/2))^{2/3}$ of the ideal gas, the number of particles is $N = 4 \times 10^5$ and the coupling constant is such that the zero-temperature chemical potential is $\rho g = gN/V = 700\hbar^2/mV^{2/3}$.

Correspondingly the standard deviation of the condensate phase change increases with time, as we show in figure 2 for five different values of the temperature, up to $T = 0.65T_c$. In all cases, at long times, we observe a quadratic growth of $\text{Var } \varphi$ contrarily to the phase diffusion behavior $\propto t$ predicted in the literature [7, 8, 9, 10].

To complete the physical picture, we show in figure 3 the correlation function of the condensate atom number (11). At very short times, see the beginning of the curves in Fig.3a, the simulation (square symbols) confirms the




FIG. 2: (Color online) Standard deviation of the condensate phase change $\varphi(t)$ (12) as a function of time for (a) $T = 0.08245T_c$ (lower curve) and $T = 0.1711T_c$ (upper curve), (b) $T = 0.29467T_c$, (c) $T = 0.453T_c$, (d) $T = 0.6473T_c$. Thick solid line (black): numerical solution from the classical field model Eq.(8) averaged over 500 realizations. Thin solid line (red): a linear fit. The parameters N and ρg have the same values as in Fig.1.

Bogoliubov prediction (dashed line); at long times, see Fig.3b, the correlation function drops to a value significantly smaller than the Bogoliubov prediction (fast oscillations are not shown in the figure); a key point is that this long time value of the correlation function of the condensate atom number is not zero.

One may fear at this stage that the classical field model is missing some source of damping in the dynamics of the system. However it is a well established fact that the classical field model is able to simulate damping processes, including the finite temperature Beliaev-Landau processes [21, 22, 23], since the interaction among the Bogoliubov modes is included in this model [11, 12, 14, 15, 19, 20, 24, 25, 26, 27]. More quantitatively we now check that the damping times due to the Beliaev-Landau processes in the simulation are much shorter than the evolution times considered here. To this end, we extract from the simulations the temporal correlation functions $\langle b_{\mathbf{k}}^*(t)b_{\mathbf{k}}(0) \rangle$ obtained by projecting the classical field over the corresponding Bogoliubov mode and averaging over many realizations. We show an example of such correlation function for the lowest energy Bogoliubov mode in Fig.4.

We come then into a paradox. On one side, the various Bogoliubov oscillators $b_{\mathbf{k}}$ decorrelate at long times. On the other side, the variance of the phase change φ of




FIG. 3: (Color online) Correlation function of the condensate atom number (11). (a) Short times, (b) long times. The classical field results are obtained from an average over 500 solutions of Eq.(8); they are represented by symbols in (a) and a solid line in (b). The dashed lines (red) are the Bogoliubov approximations (21) and (22) in (a), and only (22) in (b). The dashed-dotted line (purple) is the Gaussian model. For clarity in (b) we washed out fast oscillations in the simulation result and in the Gaussian model, by averaging over consecutive points over a time width $0.45mV^{2/3}/\hbar$. The horizontal dashed-dotted-dotted line (blue) in (a) and (b) is the ergodic long time limit prediction, described in section IV. The parameters N and ρg have the same values as in Fig.1.

the condensate varies quadratically at long times, which implies, as we shall see in Sec. IV, that the derivative of the phase $\dot{\varphi}$ does not decorrelate at long times, even if it is a function of the $b_{\mathbf{k}}$'s; similarly, the fluctuations of the number of condensate atoms δn_0 , which are functions of the $b_{\mathbf{k}}$'s, do not decorrelate at long times.

This paradox will be explained in Sec. IV, and quantitative predictions for long times behavior of the condensate atom number correlation function and of the variance of the condensate phase change will be derived. Anticipating these analytical results, we show in Fig.5a the long time limit of $(\text{Var } \varphi)^{1/2}/t$ as a function of T/T_c , from




FIG. 4: (Color online) Correlation function $\langle b_{\mathbf{k}}^*(t)b_{\mathbf{k}}(0) \rangle$ for the mode $\mathbf{k} = (0, 0, 2\pi/L_3)$ as a function of time. The solid line results from an average over 8000 solutions of Eq.(8). Here the squares of the box sizes are in the ratio 1.082712 : 0.779954 : 1.18417. The dashed-dotted line (purple) is an exponential function of t whose time constant has been calculated by time-dependent perturbation theory including the discrete nature of the spectrum as in [19]. The agreement is very good, and we have checked that the agreement is good for all vectors \mathbf{k} lying in an arbitrarily chosen plane in k -space. Here the temperature is $T = 0.0845T_c$. The parameters N and ρg have the same values as in Fig.1.

the results of the classical field simulations, but also from the predictions of the Bogoliubov approximation Eq.(37), and of the ergodic theory of Sec. IV. In figure 5b we show the same results and predictions for the asymptotic value of the condensate atom number correlation function.

IV. CLASSICAL FIELD: ANALYTICAL RESULTS

The general procedure used here to obtain analytical results is the following. First one expresses the quantity of interest (the number of condensate atoms or the time derivative of the condensate phase) in terms on the amplitudes $b_{\mathbf{k}}$ of the field ψ over the Bogoliubov modes,

$$b_{\mathbf{k}}(t) = dV \sum_{\mathbf{r}} \frac{\tilde{U}_{\mathbf{k}}}{\sqrt{V}} \frac{e^{-i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{V}} e^{-i\theta(t)} \psi_{\perp}(\mathbf{r}, t) + \tilde{V}_{\mathbf{k}} \frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{\sqrt{V}} e^{i\theta(t)} \psi_{\perp}^*(\mathbf{r}, t) \quad (18)$$

where ψ_{\perp} is the component of ψ orthogonal to the condensate mode. Second one evaluates the correlation functions of products of $b_{\mathbf{k}}$ in various physical limits.




FIG. 5: (Color online) (a) Long time limit of $(\text{Var } \varphi)^{1/2}/t$, in units of $\hbar/mV^{2/3}$, as a function of the temperature over the ideal gas critical temperature. (b) Long time limit of the condensate atom number correlation function, as a function of T/T_c . In square symbols the numerical results of the classical field simulation (averaged over 500 realizations), in dashed line (red) the Bogoliubov prediction and in dashed-dotted (blue) the ergodic prediction. The parameters N and ρg have the same values as in Fig.1.

A. Correlation function of the condensate atom number

As the total number of particles is fixed, it is equivalent to calculate the correlation function of δn_0 in Eq.(11) and of the number of non-condensed particles N_{\perp} . Injecting the expansion Eq.(14) for the time dependent non-condensed field ψ_{\perp} over the Bogoliubov modes into Eq.(17) we obtain

$$N_{\perp}(t) = \sum_{\mathbf{k} \neq 0} |\tilde{U}_{\mathbf{k}} b_{\mathbf{k}}(t) + \tilde{V}_{\mathbf{k}} b_{-\mathbf{k}}^*(t)|^2. \quad (19)$$

Bogoliubov theory: In Bogoliubov theory interaction among the Bogoliubov modes is neglected so that at all times

$$b_{\mathbf{k}}(t) = b_{\mathbf{k}}(0) e^{-i\omega_{\mathbf{k}} t} \quad \text{with} \quad \omega_{\mathbf{k}} = \tilde{\epsilon}_{\mathbf{k}}/\hbar. \quad (20)$$

As Wick's theorem applies for the initial thermal distribution we obtain for the correlation function of the condensate atom number:

$$\langle \delta n_0(t) \delta n_0(0) \rangle_{\text{Bog}} = \sum_{\mathbf{k} \neq 0} \tilde{n}_k^2 \left[|\tilde{U}_k^2 e^{i\omega_k t} + \tilde{V}_k^2 e^{-i\omega_k t}|^2 + (\tilde{U}_k \tilde{V}_k)^2 |e^{i\omega_k t} + e^{-i\omega_k t}|^2 \right] \quad (21)$$

where $\tilde{n}_k = k_B T / \tilde{\epsilon}_k$ is the Bogoliubov mean occupation number of a mode for the classical field. At very short times, a good agreement of the Bogoliubov prediction with the simulation is observed in Fig.3a. Smearing out the terms oscillating rapidly at Bohr frequencies $2\omega_k$, we obtain a prediction directly comparable to the coarse grained numerical result of Fig.3b:

$$\langle \delta n_0(t) \delta n_0(0) \rangle_{\text{Bog non osc}} = \sum_{\mathbf{k} \neq 0} (\tilde{U}_k^2 + \tilde{V}_k^2)^2 \tilde{n}_k^2. \quad (22)$$

As expected, Bogoliubov theory fails at long times. Note that in the thermodynamical limit, where the above sum is dominated by the low k terms, one may approximate $\tilde{V}_k \sim -\tilde{U}_k$, so that Eq.(22) is roughly half of the $t = 0$ value of Eq.(21); in other words, it is approximately half of the variance of the condensate number. In the numerical result of Fig.3, the correlation function drops by much more than a factor 2.

Gaussian theory: A heuristic approach to improve Bogoliubov theory consists in assuming that the $b_{\mathbf{k}}$ are Gaussian variables with a finite time correlation due to the Beliaev-Landau mechanism:

$$|\langle b_{\mathbf{k}}^*(t) b_{\mathbf{k}}(0) \rangle|^2 = \tilde{n}_k^2 e^{-2\Gamma_k |t|} \quad (23)$$

where Γ_k is calculated with time dependent perturbation theory as in [19], which amounts to weighting each term of Eq.(22) by $\exp(-2\Gamma_k |t|)$. The resulting prediction, while looking promising at short times, see Fig.3a, is in clear disagreement with the simulation at long times, see Fig.3b. Since the assumption of a long time decorrelation of $b_{\mathbf{k}}^*(t)$ with $b_{\mathbf{k}}(0)$ is physically reasonable, one may suspect that the Gaussian hypothesis is not strictly correct. This is indeed the case, as we now show.

Ergodic theory: A systematic way to calculate the long time limit of the correlation function is to assume that the non-linear dynamics generated by Eq.(8) is ergodic: at long times, the $b_{\mathbf{k}}(t)$'s for a given realization of the field explore uniformly a fixed energy surface in phase space [28]. In the Bogoliubov approximation for the energy, this means that the $b_{\mathbf{k}}(t)$'s sample the unnormalized probability distribution

$$P_{\infty}(\{b_{\mathbf{k}}\}) = \delta \left(E - \sum_{\mathbf{k} \neq 0} \tilde{\epsilon}_k b_{\mathbf{k}}^* b_{\mathbf{k}} \right) \quad (24)$$

where the Bogoliubov energy E is fixed by the initial value of the field:

$$E = \sum_{\mathbf{k} \neq 0} \tilde{\epsilon}_k b_{\mathbf{k}}^*(0) b_{\mathbf{k}}(0). \quad (25)$$

First, for a given initial condition of the field, we calculate the expectation value of $N_{\perp}(t)$ as given by Eq.(19) over the ergodic distribution Eq.(24), which is equivalent to the temporal average of $N_{\perp}(t)$ over an infinite time interval. The terms of the form bb or b^*b^* have a zero mean, since the phases of the $b_{\mathbf{k}}$'s are uniformly distributed over 2π , according to Eq.(24). To calculate the expectation value of the b^*b terms, it is convenient to introduce rescaled variables

$$B_{\mathbf{k}} = \left(\frac{\tilde{\epsilon}_k}{E} \right)^{1/2} b_{\mathbf{k}}. \quad (26)$$

According to Eq.(24) the real parts and the imaginary parts of all the $B_{\mathbf{k}}$ are uniformly distributed over the unit hypersphere in a space of dimension $2(\mathcal{N} - 1)$, where $\mathcal{N} = V/dV$ is the number of modes of the field. Assuming $1 \ll \mathcal{N}$ we obtain $\overline{|B_{\mathbf{k}}|^2} = 1/\mathcal{N}$ where the overline stands for the average over the ergodic distribution (24). As a consequence the ergodic average of N_{\perp} is

$$\overline{N_{\perp}} = \frac{1}{\mathcal{N}} \sum_{\mathbf{k} \neq 0} (\tilde{U}_k^2 + \tilde{V}_k^2) \frac{E}{\tilde{\epsilon}_k}. \quad (27)$$

Note that this ergodic average depends on the $t = 0$ value of the $b_{\mathbf{k}}$'s *via* (25).

Second, we average the product $\overline{N_{\perp}} N_{\perp}(0)$ over the thermal canonical distribution for the initial values $b_{\mathbf{k}}(0)$. This gives the long time limit of the correlation function of the number of condensate atoms:

$$\langle \delta n_0(t \rightarrow +\infty) \delta n_0(0) \rangle_{\text{ergo}} = \frac{1}{\mathcal{N}} \left(\sum_{\mathbf{k} \neq 0} (\tilde{U}_k^2 + \tilde{V}_k^2) \tilde{n}_k \right)^2. \quad (28)$$

This prediction is in good agreement with the simulations at long times, see Fig.3b for a fixed value of the temperature, and Fig.5b as a function of temperature. Note that, according to Schwartz inequality, the ergodic value is lower than the coarse grained Bogoliubov prediction Eq.(22), as was expected physically.

This clearly shows that the existence of infinite time correlations in the number of condensate atoms is a consequence of the conservation of energy during the free evolution of the system.

B. Variance of the condensate phase change

To reproduce the approach of the previous subsection for the phase, one should express the phase change $\varphi(t)$ of the condensate amplitude a_0 as a function of the $b_{\mathbf{k}}$'s. It turns out that the quantity easily expressed in terms of the $b_{\mathbf{k}}$'s is the time derivative $\dot{\varphi}$. The variance of φ is then related to the correlation function \mathcal{C} of $\dot{\varphi}$:

$$\text{Var } \varphi = \int_0^t d\tau \int_0^t d\tau' \mathcal{C}(|\tau - \tau'|) \quad (29)$$

where time translational invariance in steady state imposes for a classical field that \mathcal{C} depends only on $|\tau - \tau'|$:

$$\mathcal{C}(|\tau - \tau'|) = \langle \dot{\varphi}(\tau)\dot{\varphi}(\tau') \rangle - \langle \dot{\varphi}(\tau) \rangle \langle \dot{\varphi}(\tau') \rangle. \quad (30)$$

If $\mathcal{C}(\tau) \rightarrow 0$ fast enough when $\tau \rightarrow \infty$ then $\text{Var } \varphi$ grows linearly in time. On the other hand, if $\mathcal{C}(\tau)$ has a non-zero limit at long times, then $\text{Var } \varphi$ grows quadratically in time [29].

To express $\dot{\varphi}$ in terms of the $b_{\mathbf{k}}$'s, we write the equation of motion for a_0 :

$$\begin{aligned} i\hbar\dot{a}_0 &= i\hbar\{a_0, H\} = \partial_{a_0^*} H \\ &= \frac{g}{\sqrt{V}} \sum_{\mathbf{r}} \psi^*(\mathbf{r})\psi^2(\mathbf{r}) \end{aligned} \quad (31)$$

where we used $\partial_{a_0^*} \psi^*(\mathbf{r}) = 1/\sqrt{V}$ obtained from Eq.(2). We split ψ as in Eq.(16); we eliminate the condensate amplitude in the resulting expression for \dot{a}_0/a_0 (i) by using $|a_0|^2 = N - N_{\perp}$, where N_{\perp} is a function of the $b_{\mathbf{k}}$'s, see Eq.(19), and (ii) by introducing the field [30]

$$\Lambda(\mathbf{r}) = e^{-i\theta} \psi_{\perp}(\mathbf{r}) \quad (32)$$

which is a function of the $b_{\mathbf{k}}$'s only according to Eq.(14). This leads to

$$i\hbar \frac{\dot{a}_0}{a_0} = \rho g + \frac{g}{V} \sum_{\mathbf{r}} dV [\Lambda(\mathbf{r})^2 + |\Lambda(\mathbf{r})|^2] \quad (33)$$

$$+ \frac{g}{\sqrt{V}} \sum_{\mathbf{r}} dV \frac{\Lambda^*(\mathbf{r})\Lambda^2(\mathbf{r})}{\sqrt{N - N_{\perp}}}. \quad (34)$$

The real part of the above equation gives $-\hbar\dot{\theta}$, which is also $-\hbar\dot{\varphi}$.

Restricting to a weak non-condensed fraction, we drop the cubic terms in Eq.(34), to obtain

$$\begin{aligned} \hbar\dot{\varphi} &\simeq -\rho g - \frac{1}{2} \frac{g}{V} \sum_{\mathbf{r}} dV [\Lambda(\mathbf{r}) + \Lambda^*(\mathbf{r})]^2 \\ &= -\rho g - \frac{1}{2} \frac{g}{V} \sum_{\mathbf{k} \neq 0} (\tilde{U}_{\mathbf{k}} + \tilde{V}_{\mathbf{k}})^2 |b_{\mathbf{k}} + b_{-\mathbf{k}}^*|^2. \end{aligned} \quad (35)$$

It turns out that the products $b_{\mathbf{k}}b_{-\mathbf{k}}$ generate oscillating terms which do not contribute to a coarse grained time average. It is thus useful to define

$$\hbar\dot{\varphi}_{\text{non osc}} = -\rho g - \frac{g}{V} \sum_{\mathbf{k} \neq 0} (\tilde{U}_{\mathbf{k}} + \tilde{V}_{\mathbf{k}})^2 |b_{\mathbf{k}}|^2. \quad (36)$$

Bogoliubov theory: By using (20) and Wick's theorem we calculate the correlation function of Eq.(35). By temporal integration we obtain the variance of the condensate phase change

$$(\text{Var } \varphi)_{\text{Bog}} = \left(\frac{g}{\hbar V} \right)^2 \sum_{\mathbf{k} \neq 0} (\tilde{U}_{\mathbf{k}} + \tilde{V}_{\mathbf{k}})^4 \tilde{n}_{\mathbf{k}}^2 \left[t^2 + \frac{\sin^2 \omega_{\mathbf{k}} t}{(2\omega_{\mathbf{k}})^2} \right]. \quad (37)$$

Qualitatively Bogoliubov theory correctly predicts a linear growth of the standard deviation of φ at long times. As we show in Fig.5a, however, it is not fully quantitative: it does not reproduce the value of the dephasing rate obtained from the simulations. This is not surprising as in the full non linear theory the $b_{\mathbf{k}}$'s interact and do not follow Eq.(20).

To be complete, we also give the Bogoliubov approximation for the correlation function of the condensate amplitude a_0 . Neglecting the fluctuations of the modulus of a_0 , one can set

$$\langle a_0^*(t) a_0(0) \rangle \simeq \langle N_0 \rangle \langle e^{-i\varphi(t)} \rangle. \quad (38)$$

Dropping the oscillating terms in $b_{\mathbf{k}}b_{-\mathbf{k}}$ and $b_{\mathbf{k}}^*b_{-\mathbf{k}}^*$ in $\dot{\varphi}(t)$, which give a small contribution, we get

$$\langle a_0^*(t) \hat{a}_0(0) \rangle_{\text{Bog}} \simeq \langle N_0 \rangle \prod_{\mathbf{k} \neq 0} \frac{1}{1 + i \frac{g}{\hbar V} (\tilde{V}_{\mathbf{k}} + \tilde{U}_{\mathbf{k}})^2 \tilde{n}_{\mathbf{k}} t}. \quad (39)$$

The resulting expression is plotted as a dashed line in Fig.1 against the result of the simulation.

Gaussian theory: If we add by hand a decorrelation of the $b_{\mathbf{k}}$'s and assume Gaussian statistics, we get for the variance a prediction growing linearly at long times:

$$(\text{Var } \varphi)_{\text{Gauss}} = \left(\frac{g}{\hbar V} \right)^2 \sum_{\mathbf{k} \neq 0} (\tilde{U}_{\mathbf{k}} + \tilde{V}_{\mathbf{k}})^4 \tilde{n}_{\mathbf{k}}^2 \left[\frac{e^{-2\Gamma_{\mathbf{k}} t} - 1}{2\Gamma_{\mathbf{k}}^2} + \frac{t}{\Gamma_{\mathbf{k}}} \right], \quad (40)$$

in clear contradiction with the numerical simulations. This prediction corresponds to a correlation function \mathcal{C} vanishing at long times, whereas the correct correlation function has a finite limit, see Fig.6.

Ergodic theory: as in subsection IV A we calculate the long time value of the correlation function for $\dot{\varphi}$ using the ergodic assumption. The various steps of the calculation are rigorously the same as in Sec. IV A and lead to

$$\mathcal{C}(\tau \rightarrow +\infty)_{\text{ergo}} = \left(\frac{g}{\hbar V} \right)^2 \frac{1}{\mathcal{N}} \left(\sum_{\mathbf{k} \neq 0} (\tilde{U}_{\mathbf{k}} + \tilde{V}_{\mathbf{k}})^2 \tilde{n}_{\mathbf{k}} \right)^2. \quad (41)$$

This prediction is in excellent agreement with the simulations: it gives the correct asymptotic value of \mathcal{C} , see Fig.6, and from the asymptotic expression $\text{Var } \varphi \simeq \mathcal{C}(+\infty)t^2$ it gives the correct values of the long time limit of $(\text{Var } \varphi)^{1/2}/t$, see Fig.5a, as a function of the temperature.

V. QUANTUM TREATMENT: ANALYTICAL RESULTS

So far the classical field model was very useful in revealing the physical processes governing the long time behavior of the phase and atom number fluctuations in the condensate. However it is not a fully quantitative theory, as the long time limits of the correlation functions considered here depend on the precise choice of the




FIG. 6: (Color online) Correlation function $C_{\text{non osc}}$ of the quantity $\hat{\varphi}_{\text{non osc}}$ defined in Eq.(36) calculated from the simulation (square symbols; the solid line is a guide to the eye), or using the Gaussian theory (purple dashed-dotted line going to zero at long times). In dashed line (red) the Bogoliubov prediction. In dashed-dotted-dotted line (blue) the long time prediction (41) of the ergodic theory. The parameters T , N and ρg have the same values as in Fig.1.

energy cut-off, that is on the number of modes \mathcal{N} in the simulation, as is apparent on Eqs.(28,41). In this section, we therefore adapt the previous physical reasonings to the quantum field case.

A. The quantum model

We use a straightforward generalization of the classical field lattice model, taking here for simplicity a cubic lattice, as discussed in [19, 31, 32]. The bosonic field $\hat{\psi}$ evolves according to the Hamiltonian

$$H = \sum_{\mathbf{k}} \frac{\hbar^2 k^2}{2m} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \frac{g_0}{2} \sum_{\mathbf{r}} dV \hat{\psi}^\dagger \hat{\psi}^\dagger \hat{\psi} \hat{\psi}, \quad (42)$$

where $\hat{a}_{\mathbf{k}}$ annihilates a particle of wavevector \mathbf{k} in the first Brillouin zone. The dispersion relation of the wave is now the usual one. The total number of atoms is fixed, equal to N . The coupling constant g_0 depends on the lattice spacing l in order to ensure a l -independent scattering length for the discrete delta interaction potential among the particles [31, 32]:

$$V(\mathbf{r}_1 - \mathbf{r}_2) = \frac{g_0}{dV} \delta_{\mathbf{r}_1, \mathbf{r}_2}. \quad (43)$$

Since we consider here the weakly interacting regime $\sqrt{\rho a^3} \ll 1$, where $\rho = N/V$ is the gas density, we can restrict to a lattice spacing much larger than the scattering length a so that g_0 is actually very close to $g = 4\pi\hbar^2 a/m$.

To be able to use Bogoliubov theory as we did in the classical field reasoning, we restrict to the low temperature regime $T \ll T_c$ with a macroscopic occupation of the condensate mode. We thus neglect the possibility

that the condensate is empty, which allows us to use the modulus-phase representation of the condensate mode:

$$\hat{a}_0 \simeq e^{i\hat{\theta}} \sqrt{\hat{n}_0} \quad (44)$$

where $\hat{n}_0 = \hat{a}_0^\dagger \hat{a}_0$ and where $\hat{\theta}$ is a Hermitian ‘phase’ operator obeying the commutation relation

$$[\hat{n}_0, \hat{\theta}] = i. \quad (45)$$

This allows to consider the correlation of the condensate atom number fluctuation $\delta\hat{n}_0 \equiv \hat{n}_0 - \langle \hat{n}_0 \rangle$ but also the variance of the condensate phase change $\hat{\varphi}(t) \equiv \hat{\theta}(t) - \hat{\theta}(0)$, as we did for the classical field.

B. Correlation function of the condensate atom number

To predict the correlation function of $\delta\hat{n}_0$, we use Bogoliubov theory at short times and the quantum analog of the ergodic theory at long times.

In the number conserving Bogoliubov theory [30, 33], written here for a spatially homogeneous system, one introduces the field conserving the total number of particles

$$\hat{\Lambda}(\mathbf{r}) \equiv e^{-i\hat{\theta}} \hat{\psi}_\perp(\mathbf{r}) \quad (46)$$

where the non-condensed field $\hat{\psi}_\perp$ is obtained by projecting out the component of the field $\hat{\psi}$ on the condensate mode. The field $\hat{\Lambda}$ then admits the modal expansion on the Bogoliubov modes

$$\hat{\Lambda}(\mathbf{r}) = \sum_{\mathbf{k} \neq 0} \hat{b}_{\mathbf{k}} U_{\mathbf{k}} \frac{e^{i\mathbf{k} \cdot \mathbf{r}}}{\sqrt{V}} + \hat{b}_{\mathbf{k}}^\dagger V_{\mathbf{k}} \frac{e^{-i\mathbf{k} \cdot \mathbf{r}}}{\sqrt{V}} \quad (47)$$

where the real amplitudes $U_{\mathbf{k}}$, $V_{\mathbf{k}}$, normalized as $U_{\mathbf{k}}^2 - V_{\mathbf{k}}^2 = 1$, are given by the usual Bogoliubov theory,

$$U_{\mathbf{k}} + V_{\mathbf{k}} = \left(\frac{\hbar^2 k^2 / 2m}{2\rho g_0 + \hbar^2 k^2 / 2m} \right)^{1/4}. \quad (48)$$

Since the total number of particles is fixed to N , it is equivalent to consider the fluctuations of \hat{n}_0 or of the number of non-condensed atoms

$$\hat{N}_\perp = \sum_{\mathbf{r}} dV \hat{\Lambda}^\dagger(\mathbf{r}) \hat{\Lambda}(\mathbf{r}). \quad (49)$$

This, together with the expansion (47), expresses \hat{N}_\perp as a function of the $\hat{b}_{\mathbf{k}}$'s.

The equilibrium state of the system is approximated in the canonical ensemble by the Bogoliubov thermal density operator

$$\hat{\rho}_{\text{Bog}}(T) = \frac{1}{Z} e^{-\sum_{\mathbf{k} \neq 0} \epsilon_{\mathbf{k}} \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} / k_B T} \quad (50)$$

where the normalization factor Z is the Bogoliubov approximation for the partition function, and where we have introduced the Bogoliubov spectrum

$$\epsilon_k = \left[\frac{\hbar^2 k^2}{2m} \left(\frac{\hbar^2 k^2}{2m} + 2\rho g_0 \right) \right]^{1/2}. \quad (51)$$

Bogoliubov theory: In the Bogoliubov approximation for the time evolution, the $\hat{b}_{\mathbf{k}}$ merely accumulate a phase, at the frequency $\omega_k = \epsilon_k/\hbar$, similarly to the classical field case. From Wick's theorem one then obtains

$$\frac{1}{2} \langle \{ \delta \hat{n}_0(t), \delta \hat{n}_0(0) \} \rangle_{\text{Bog}} = \sum_{\mathbf{k} \neq 0} \bar{n}_{\mathbf{k}} (\bar{n}_{\mathbf{k}} + 1) (U_{\mathbf{k}}^2 + V_{\mathbf{k}}^2)^2 + 2U_{\mathbf{k}}^2 V_{\mathbf{k}}^2 \cos(2\omega_{\mathbf{k}} t) [\bar{n}_{\mathbf{k}}^2 + (\bar{n}_{\mathbf{k}} + 1)^2] \quad (52)$$

where

$$\bar{n}_{\mathbf{k}}(T) = \frac{1}{\exp(\epsilon_k/k_B T) - 1} \quad (53)$$

is the mean occupation number of the Bogoliubov mode \mathbf{k} . Note that we have considered here the so-called symmetric correlation function (as $\{X, Y\}$ stands for the anticommutator $XY + YX$ of two operators) which is a real quantity, equal to the real part of the non-symmetrized correlation function. The time coarse grained version of the prediction (52) is obtained by averaging out the oscillating terms, which amounts to considering the correlation function of the temporally smoothed operator number of non-condensed particles

$$\hat{N}_{\perp}^{\text{non osc}} \equiv \sum_{\mathbf{k} \neq 0} \left[(U_{\mathbf{k}}^2 + V_{\mathbf{k}}^2) \hat{b}_{\mathbf{k}}^{\dagger} \hat{b}_{\mathbf{k}} + V_{\mathbf{k}}^2 \right]. \quad (54)$$

Quantum Ergodic theory: Discarding from the start the oscillating terms in \hat{N}_{\perp} , as in (54), we face here the problem of calculating the long time limit of $\langle A(t)A(0) \rangle$, where A is a linear function of the Bogoliubov mode occupancies,

$$A = \sum_{\mathbf{k} \neq 0} \gamma_{\mathbf{k}} \hat{b}_{\mathbf{k}}^{\dagger} \hat{b}_{\mathbf{k}}. \quad (55)$$

As the quantum state of the system is given by the Bogoliubov approximation Eq.(50), we may inject a closure relation in the Bogoliubov Fock eigenbasis:

$$\langle A(t)A(0) \rangle = \frac{1}{Z} \sum_{\{n_{\mathbf{k}}\}} e^{-\beta \sum_{\mathbf{k} \neq 0} \epsilon_k n_{\mathbf{k}}} \left(\sum_{\mathbf{k} \neq 0} \gamma_{\mathbf{k}} n_{\mathbf{k}} \right) \times \langle \{n_{\mathbf{k}}\} | A(t) | \{n_{\mathbf{k}}\} \rangle, \quad (56)$$

where the sum is taken over all possible integer values of the occupation numbers, not to be confused with the mean occupation numbers (53).

The non-explicit piece of this expression is the matrix element of $A(t)$, which may be reinterpreted as follows:

$$\langle \{n_{\mathbf{k}}\} | A(t) | \{n_{\mathbf{k}}\} \rangle = \text{Tr} [A\sigma(t)] \quad (57)$$

where the density operator σ , initially a pure state in the Bogoliubov Fock basis,

$$\sigma(0) = |\{n_{\mathbf{k}}\}\rangle \langle \{n_{\mathbf{k}}\}| \quad (58)$$

evolves during t with the full Hamiltonian H . We know that this evolution involves Beliaev-Landau processes that will spread σ over the various Fock states $|\{n'_{\mathbf{k}}\}\rangle$. This evolution is complex. But we need here the long time limit only, in which we may assume that an equilibrium statistical description is possible. Since the system is isolated during its evolution, we take for $\sigma(t \rightarrow +\infty)$ the equilibrium density operator in the *microcanonical* ensemble [34], and we calculate the expectation value of A with $\sigma(t \rightarrow +\infty)$ as we did for the classical field model. The calculation can be done in the thermodynamical limit. As shown in the Appendix A, one can calculate to leading order in this limit the difference between canonical and microcanonical averages.

Here the microcanonical ensemble has an energy $E = E_0^{\text{Bog}} + \sum_{\mathbf{k} \neq 0} \epsilon_k n_{\mathbf{k}}$, where E_0^{Bog} is the ground state Bogoliubov energy. We introduce the effective temperature T_{eff} such that the mean energy in the canonical ensemble at temperature T_{eff} is equal to E ,

$$0 = \langle H_{\text{Bog}} \rangle(T_{\text{eff}}) - E = \sum_{\mathbf{k} \neq 0} \epsilon_k [\bar{n}_{\mathbf{k}}(T_{\text{eff}}) - n_{\mathbf{k}}] \quad (59)$$

where $\langle \dots \rangle$ stands for an average in the canonical ensemble and H_{Bog} is the Bogoliubov Hamiltonian. Using the results of Appendix A one gets

$$\bar{A}(E) - \langle A \rangle(T_{\text{eff}}) = -\frac{1}{2} k_B T_{\text{eff}}^2 \left(\frac{\langle A \rangle'}{\langle H_{\text{Bog}} \rangle'} \right)' (T_{\text{eff}}) \quad (60)$$

where $\bar{A}(E)$ is the microcanonical average of A at energy E and where the appex $'$ stands for derivation with respect to temperature. We further use the fact that, in the thermodynamical limit, for typical values of the occupation numbers $n_{\mathbf{k}}$, T_{eff} weakly deviates from the physical temperature T . We calculate T_{eff} by expanding (59) up to second order in $T - T_{\text{eff}}$ [35]. Evaluating (60) with this value of T_{eff} , keeping terms up to the relevant order [35], gives the desired result

$$\begin{aligned} & \langle \{n_{\mathbf{k}}\} | A(t \rightarrow +\infty) | \{n_{\mathbf{k}}\} \rangle = \langle A \rangle \\ & + \left[\sum_{\mathbf{k} \neq 0} \epsilon_k (n_{\mathbf{k}} - \bar{n}_{\mathbf{k}}) \right] \frac{\langle A \rangle'}{\langle H_{\text{Bog}} \rangle'} + \frac{1}{2} \left(\frac{\langle A \rangle'}{\langle H_{\text{Bog}} \rangle'} \right)' \\ & \times \left\{ \frac{\left[\sum_{\mathbf{k} \neq 0} \epsilon_k (n_{\mathbf{k}} - \bar{n}_{\mathbf{k}}) \right]^2}{\langle H_{\text{Bog}} \rangle'} - k_B T^2 \right\} \end{aligned} \quad (61)$$

where all the canonical averages are now evaluated at the physical temperature T [36].

It remains to inject this expression into Eq.(56). The resulting average over $n_{\mathbf{k}}$ leads to the long time value of

the correlation function:

$$\langle A(t \rightarrow +\infty)A(0) \rangle - \langle A \rangle^2 = \frac{\left[\sum_{\mathbf{k} \neq 0} \gamma_k \epsilon_k \bar{n}_k (\bar{n}_k + 1) \right]^2}{\sum_{\mathbf{k} \neq 0} \epsilon_k^2 \bar{n}_k (\bar{n}_k + 1)}. \quad (62)$$

Using Schwartz inequality, one can show that this long time value of the correlation function is less than its zero time value $\sum \gamma_k^2 \bar{n}_k (\bar{n}_k + 1)$. Replacing in Eq.(62) the coefficients γ_k by their expression from Eq.(54), $\gamma_k = U_k^2 + V_k^2$, we obtain the long time value of the condensate atom number correlation function predicted by the quantum version of ergodic theory. Note that, in the thermodynamical limit, this long time value scales as the volume V , whereas the $t = 0$ value scales as $V^{4/3}$.

C. Correlation function of the time derivative of the condensate phase

As in the classical field case, we first look for an expression of the first order time derivative of the condensate phase operator $\hat{\theta}$ in terms of the amplitudes of the field $\hat{\Lambda}$ on the Bogoliubov modes. Taking as a starting point in Heisenberg picture

$$i\hbar \frac{d}{dt} \hat{\theta} = [\hat{\theta}, H], \quad (63)$$

we split the quantum field in a condensate part and a non-condensed part,

$$\hat{\psi}(\mathbf{r}) = \frac{\hat{a}_0}{\sqrt{V}} + \hat{\psi}_\perp(\mathbf{r}), \quad (64)$$

and we insert this splitting in the expression of H . Using the modulus-phase representation of \hat{a}_0 and the commutation relation Eq.(45), we obtain, after using $\hat{a}_0^\dagger \hat{a}_0 = \hat{N} - \hat{N}_\perp$,

$$\begin{aligned} -\hbar \frac{d}{dt} \hat{\theta} &= \frac{g_0}{V} \left[\hat{N} - \frac{1}{2} + \sum_{\mathbf{r}} dV \hat{\Lambda}^\dagger \hat{\Lambda} \right] \\ &+ \frac{g_0}{2V} \sum_{\mathbf{r}} dV \left[\hat{\Lambda} \frac{n_0 + 1/2}{\sqrt{n_0(n_0 + 1)}} \hat{\Lambda} + \text{h.c.} \right] \\ &+ \frac{g_0}{2\sqrt{V}} \sum_{\mathbf{r}} dV \left[\frac{1}{\sqrt{n_0}} \hat{\Lambda}^\dagger \hat{\Lambda}^2 + \text{h.c.} \right]. \quad (65) \end{aligned}$$

The quantity $(n_0 + 1/2)/\sqrt{n_0(n_0 + 1)}$ is actually $1 + O(1/n_0^2)$ so it can to a high accuracy be replaced by unity. Furthermore, as we did in the classical field model, we now keep the leading terms in $\hat{\Lambda}$, under the assumption of a weak non-condensed fraction. We can also replace $\hat{\theta}$ by $\hat{\varphi}$ under the temporal derivative, since $\hat{\theta}(0)$ is time independent. We obtain

$$-\hbar \frac{d}{dt} \hat{\varphi} \simeq \frac{g_0}{V} \left[\hat{N} - \frac{1}{2} + \sum_{\mathbf{r}} dV \left(\hat{\Lambda}^\dagger \hat{\Lambda} + \frac{1}{2} \hat{\Lambda}^2 + \frac{1}{2} \hat{\Lambda}^{\dagger 2} \right) \right]. \quad (66)$$

Taking the expectation value of this expression over the thermal state in the Bogoliubov approximation leads to an expression coinciding with the value of the chemical potential predicted by Eq.(103) of [32], which includes in a systematic way the first correction to the pure condensate prediction ρg_0 [37]:

$$\mu(T) = \frac{g_0}{V} \left[N - \frac{1}{2} + \sum_{\mathbf{k} \neq 0} (U_k + V_k)^2 \bar{n}_k + V_k (U_k + V_k) \right]. \quad (67)$$

At this order of the expansion, this analytically shows that $-\hbar \langle d\hat{\theta}/dt \rangle$ is the chemical potential of the system. We now turn to various predictions for the symmetrized correlation function of $d\hat{\varphi}/dt$,

$$C_S(\tau) = \frac{1}{2} \left\langle \left\{ \left(\frac{d}{dt} \hat{\varphi} \right) (\tau), \left(\frac{d}{dt} \hat{\varphi} \right) (0) \right\} \right\rangle - \left\langle \frac{d}{dt} \hat{\varphi} \right\rangle^2. \quad (68)$$

Bogoliubov theory: At a time short enough for the interactions between the Bogoliubov modes to remain negligible, one can apply Bogoliubov theory to get

$$\begin{aligned} C_S^{\text{Bog}}(\tau) &= \left(\frac{g_0}{2V\hbar} \right)^2 \sum_{\mathbf{k} \neq 0} (U_k + V_k)^4 \{ 2\bar{n}_k (\bar{n}_k + 1) \\ &+ \cos(2\omega_k \tau) [\bar{n}_k^2 + (\bar{n}_k + 1)^2] \}. \quad (69) \end{aligned}$$

The temporal coarse grained version of this correlation function is obtained by averaging out the cosine terms, which amounts to considering a temporal derivative of $\hat{\varphi}$ freed from the oscillating terms $\hat{b}\hat{b}$ and $\hat{b}^\dagger\hat{b}^\dagger$:

$$\begin{aligned} \left(\frac{d}{dt} \hat{\varphi} \right)_{\text{non osc}} &= -\frac{g_0}{\hbar V} \left[\hat{N} - \frac{1}{2} + \sum_{\mathbf{k} \neq 0} V_k (U_k + V_k) \right] \\ &- \frac{g_0}{\hbar V} \sum_{\mathbf{k} \neq 0} (U_k + V_k)^2 \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}. \quad (70) \end{aligned}$$

Quantum ergodic theory: We directly apply to the smoothed temporal derivative (70) the reasoning performed in the previous subsection. Up to an additive constant, Eq.(70) is indeed of the form (55), with $\gamma_k = -(g_0/\hbar V)(U_k + V_k)^2$. From (62) we therefore obtain the long time behavior of the phase derivative correlation function

$$C_S^{\text{ergo}}(\tau \rightarrow +\infty) = \left(\frac{g_0}{\hbar V} \right)^2 \frac{\left[\sum_{\mathbf{k} \neq 0} (U_k + V_k)^2 \epsilon_k \bar{n}_k (\bar{n}_k + 1) \right]^2}{\sum_{\mathbf{k} \neq 0} \epsilon_k^2 \bar{n}_k (\bar{n}_k + 1)}. \quad (71)$$

In the thermodynamical limit, analytical expressions can be obtained for this ergodic prediction in the low temperature limit $k_B T \ll \rho g$,

$$C_S^{\text{ergo}}(+\infty) \sim \frac{8\pi^4}{15} \frac{a^2 \xi}{V} \left(\frac{k_B T}{\hbar} \right)^2 \left(\frac{k_B T}{\rho g} \right)^3, \quad (72)$$

where ξ is the healing length such that $\hbar^2/m\xi^2 = \rho g$, and in the high temperature limit $k_B T \gg \rho g$,

$$C_S^{\text{ergo}}(+\infty) \sim \frac{12\zeta(3/2)^2}{5\zeta(5/2)} \frac{a^2 \lambda}{V} \left(\frac{k_B T}{\hbar} \right)^2 \quad (73)$$




FIG. 7: (Color online) In the regime $T \ll T_c$ for a weakly interacting Bose gas, quantum ergodic prediction (71) for the long time limit of $(\text{Var } \hat{\varphi})^{1/2}/t$, in the thermodynamical limit. When expressed in units of $(a^2 \xi/V)^{1/2} \rho g/\hbar$, $(\text{Var } \hat{\varphi})^{1/2}/t$ is a function of $k_B T/\rho g$ only, that is readily calculated numerically (solid line) or that may be approximated by asymptotic equivalents (72,73) in the low temperature or high temperature limit (green dashed line). Note that the dimensionless quantity $a^2 \xi/V$ may also be written as $\sqrt{\rho a^3}/(N\sqrt{4\pi})$.

where the thermal de Broglie wavelength obeys $\lambda^2 = 2\pi\hbar^2/mk_B T$ and where ζ is the Riemann Zeta function. Here we have identified g_0 to g . In Fig.7 we give the quantum ergodic prediction for $\lim_{t \rightarrow \infty} (\text{Var } \hat{\varphi})^{1/2}/t$ calculated from numerical integration, which is a universal function of $k_B T/\rho g$ when expressed in the right units.

VI. CONCLUSION

We have investigated theoretically the phase spreading of a finite temperature weakly interacting condensate. The gas is assumed to be prepared at thermal equilibrium in the canonical ensemble, and then to freely evolve as an isolated system. After average over many realizations of the system, we find in classical field simulations that the variance of the phase change of the condensate $\text{Var } \varphi$ grows quadratically in time. This non-diffusive behavior is quantitatively explained by an ergodic theory for the Bogoliubov modes, the key point being that conservation of energy during the free evolution prevents some correlation functions of the field from vanishing at long times. We have extended the analytical treatment to the quantum field case and we have determined the coefficient of the t^2 term in the long time behavior of $\text{Var } \hat{\varphi}$, see Eq.(71).

A more physical insight is obtained from the following rewriting of the result,

$$\text{Var } \hat{\varphi}(t) \sim \frac{t^2}{\hbar^2} \left(\frac{\partial_T \mu}{\partial_T \langle H \rangle} \right)^2 \text{Var } H \quad (74)$$

where $\text{Var } H$ is the variance of the energy of the gas, here in the Bogoliubov approximation and in the canonical

ensemble, $\mu(T)$ is the chemical potential of the system as given by Eq.(67) and $\langle H \rangle(T)$ is its mean energy in the Bogoliubov approximation.

This formula may also be obtained from the following reasoning. For a given realization of the system, of energy E , the long time limit of the condensate phase change $\varphi(t)$ can be shown to behave as $-\mu_{\text{micro}}(E)t/\hbar$, where μ_{micro} is the chemical potential calculated in the microcanonical ensemble [38]. For a large system, energy fluctuations around $\bar{E} = \langle H \rangle(T)$ are weak so that one may expand $\mu_{\text{micro}}(E)$ to first order in $E - \bar{E}$. Taking the variance of $\varphi(t)$ leads to (74), since $\partial_T \mu/\partial_T \langle H \rangle \simeq \partial_E \mu_{\text{micro}}(\bar{E})$ for a large system.

This reasoning shows that a necessary condition for the observation of an intrinsic diffusive spreading of the condensate phase change is a strong suppression of the energy fluctuations of the gas. To this end one may try to prepare the system in a clever way, starting with a pure condensate and giving to the system a well defined amount of energy, e.g. by a reproducible change of the trapping potential [39]. Alternatively one may try to follow a given experimental realization of the system, measuring the phase of the condensate in a non-destructive way and replacing the ensemble average by a time average.

We acknowledge discussions with Fabrice Gerbier and thank Mikhail Kolobov for interest in the problem. One of us (E. W.) acknowledges financial support from QU-FAR. Laboratoire Kastler Brossel is a research unit of University Paris 6 and Ecole normale supérieure, associated to CNRS. Our group is a member of IFRAF.

APPENDIX A: DEVIATION OF MICROCANONICAL AND CANONICAL AVERAGES

We wish to calculate the thermal expectation value of an observable A in the microcanonical ensemble rather than in the canonical one. For convenience, we shall parametrize the problem by the temperature T of the canonical ensemble. Restricting to the thermodynamical limit, where $k_B T$ is much larger than the typical level spacing of the system, we calculate the first order deviation of the two ensembles.

We start with the usual integral representation of the canonical ensemble in terms of the microcanonical one:

$$\langle A \rangle(T) = \frac{\int dE \bar{A}(E) e^{S(E)/k_B} e^{-\beta E}}{\int dE e^{S(E)/k_B} e^{-\beta E}} \quad (A1)$$

where the density of states is written in terms of the exponential of the microcanonical entropy $S(E)$, $\bar{A}(E)$ stands for the expectation value of A in the microcanonical ensemble of energy E , $\langle A \rangle(T)$ is the expectation value of A in the canonical ensemble of temperature T , and $\beta = 1/k_B T$.

In the thermodynamical limit we expect the integrand

to be strongly peaked around the value $E_0(T)$ such that

$$\frac{d}{dE} \left[\frac{S(E)}{k_B} - \beta E \right]_{E=E_0(T)} = S'[E_0(T)]/k_B - \beta = 0, \quad (\text{A2})$$

where $f'(x)$ stands for the derivative of a function f with respect to its argument x . We then expand $u(E) \equiv S(E)/(k_B) - \beta E$ up to third order in $E - E_0$ and we approximate the integrand as

$$e^{u(E)} = e^{u(E_0)} e^{S''(E_0)(E-E_0)^2/2k_B} \times \left(1 + \frac{1}{6}(E-E_0)^3 S^{(3)}(E_0)/k_B + \dots \right). \quad (\text{A3})$$

We also expand $\bar{A}(E)$ up to second order in $E - E_0$. Performing the resulting Gaussian integrals leads to

$$\langle A \rangle(T) - \bar{A}[E_0(T)] = \frac{k_B}{2|S''(E_0)|} \times \left[\bar{A}'(E_0) \frac{S^{(3)}(E_0)}{|S''(E_0)|} + \bar{A}''(E_0) \right] + \dots \quad (\text{A4})$$

This relation can be inverted to first order, to give the microcanonical average as a function of the canonical one; to this order, we can indeed assume that $\bar{A}[E_0(T)] = \langle A \rangle(T)$ in the right hand side of (A4). Furthermore, using the implicit equation (A2) one is able to express the

derivatives with respect to E_0 in terms of derivatives with respect to T , e.g. $S''[E_0(T)] = -1/[T^2 E_0'(T)]$. This leads to

$$\bar{A}[E_0(T)] - \langle A \rangle(T) = -k_B T \left[\frac{\langle A \rangle'(T)}{E_0'(T)} + \frac{T \langle A \rangle''(T)}{2E_0'(T)} \right] + \dots \quad (\text{A5})$$

It is actually more convenient to parametrize the result in terms of the mean canonical energy $\langle H \rangle(T)$ rather than in terms of $E_0(T)$. Applying (A5) to $A = H$ allows to calculate $E_0(T) - \langle H \rangle(T)$ to first order. One then uses the first order expansion

$$\bar{A}[\langle H \rangle(T)] = \bar{A}[E_0(T)] + [\langle H \rangle(T) - E_0(T)] \times \frac{1}{E_0'(T)} \frac{d}{dT} \{ \bar{A}[E_0(T)] \} + \dots \quad (\text{A6})$$

In the first order term of this expression, we can replace $\bar{A}[E_0(T)]$ by the canonical average $\langle A \rangle(T)$, and we can identify $E_0(T)$ with $\langle H \rangle(T)$; we can do the same identification in the right hand side of (A5). We finally obtain [40]

$$\bar{A}[\langle H \rangle(T)] - \langle A \rangle(T) = -\frac{1}{2} k_B T^2 \frac{d}{dT} \left(\frac{d\langle A \rangle/dT}{d\langle H \rangle/dT} \right) + \dots \quad (\text{A7})$$

-
- [1] K. Bongs, K. Sengstock, Reports on Progress in Physics **67**, 907-963 (2004); T. Schumm, S. Hofferberth, L.M. Anderson, S. Wildermuth, S. Groth, I. Bar-Joseph, J. Schmiedmayer, P. Krüger, Nature Physics **1**, 57 (2005); P. Treutlein, P. Hommelhoff, T. Steinmetz, T. W. Hänsch, J. Reichel, Phys. Rev. Lett. **92**, 203005 (2004); O. Mandel, M. Greiner, A. Widera, T. Rom, T.W. Hänsch, I. Bloch, Nature **425**, 937 (2003); A. Micheli, D. Jaksch, I. Cirac, P. Zoller, Phys. Rev. A **67**, 013607 (2003).
- [2] E.M. Wright, D.F. Walls, J.C. Garrison, Phys. Rev. Lett. **77**, 2158 (1996); J. Javanainen, M. Wilkens, Phys. Rev. Lett. **78**, 4675 (1997) [see also the comment by A. Leggett, F. Sols, Phys. Rev. Lett. **81**, 1344 (1998), and the related answer by J. Javanainen, M. Wilkens, Phys. Rev. Lett. **81**, 1345 (1998)]; M. Lewenstein, Li You, Phys. Rev. Lett. **77**, 3489 (1997); Y. Castin, J. Dalibard, Phys. Rev. A **55**, 4330 (1997); P. Villain, M. Lewenstein, R. Dum, Y. Castin, Li You, A. Imamoglu, T.A.B. Kennedy, Journal of Modern Optics, **44** 1775-1799 (1997); A. Sinatra, Y. Castin, Eur. Phys. J. D **4**, 247-260 (1998).
- [3] M.R. Andrews, C.G. Townsend, H.J. Miesner, D.S. Durfee, D.M. Kurn, W. Ketterle, Science **275**, 637 (1997).
- [4] D.S. Hall, M.R. Matthews, C.E. Wieman, and E.A. Cornell, Phys. Rev. Lett. **81**, 1543 (1998).
- [5] C. Orzel, A.K. Tuchman, M.L. Fenselau, M. Yasuda, M. Kasevich, Science **291**, 2386 (2001); M. Greiner, O. Mandel, T.W. Hänsch, I. Bloch, Nature **419**, 51 (2002); Y. Shin, M. Saba, T.A. Pasquini, W. Ketterle, D.E. Pritchard, A.E. Leanhardt, Phys. Rev. Lett. **92**, 050405 (2004); G.-B. Jo, Y. Shin, S. Will, T. A. Pasquini, M. Saba, W. Ketterle, D. E. Pritchard, M. Vengalattore, M. Prentis, cond-mat/0608585 (2006).
- [6] A. Sinatra, Y. Castin, Eur. Phys. J. D **8**, 319-332 (2000).
- [7] D. Jaksch, C. W. Gardiner, K. M. Gheri, P. Zoller, Phys. Rev. A **58**, 1450 (1998).
- [8] R. Graham, Phys. Rev. Lett. **81**, 5262 (1998).
- [9] R. Graham, Phys. Rev. A **62**, 023609 (2000).
- [10] R. Graham, Journal of Mod. Opt. **47**, 2615 (2000).
- [11] Yu. Kagan, B. V. Svistunov, and G. V. Shlyapnikov, Sov. Phys. JETP **75**, 387 (1992); Yu. Kagan and B. Svistunov, Phys. Rev. Lett. **79** 3331 (1997).
- [12] K. Damle, S. N. Majumdar and S. Sachdev, Phys. Rev. A **54**, 5037 (1996).
- [13] M. J. Steel, M. K. Olsen, L. I. Plimak, P. D. Drummond, S. M. Tan, M. J. Collett, D. F. Walls, R. Graham, Phys. Rev. A **58**, 4824 (1998).
- [14] K. Góral, M. Gajda, K. Rzążewski, Opt. Express **8**, 92 (2001); D. Kadio, M. Gajda and K. Rzążewski, Phys. Rev. A **72**, 013607 (2005).
- [15] M.J. Davis, S.A. Morgan and K. Burnett, Phys. Rev. Lett. **87**, 160402 (2001).
- [16] A. Sinatra, C. Lobo, Y. Castin, Phys. Rev. Lett. **87**, 210404 (2001).
- [17] To get the equation defining \tilde{E}_k , one would then replace, in Eq.(7), $\hbar^2 k^2/2m$ by $[(\hbar^2 k^2/2m)(2\rho g + \hbar^2 k^2/2m)]^{1/2}$ in the left hand side and \tilde{E}_k by $[\tilde{E}_k(\tilde{E}_k + 2\rho g)]^{1/2}$ in the right hand side.

- [18] Δ stands here for the operator acting on functions on the lattice, such that its eigenvectors are the plane waves $e^{i\mathbf{k}\cdot\mathbf{r}}$ with eigenvalues $-k^2$.
- [19] A. Sinatra, C. Lobo, Y. Castin, J. Phys. B **35**, 3599 (2002).
- [20] M. Brewczyk, P. Borowski, M. Gajda, K. Rzążewski, J. Phys. B **37**, 2725 (2004).
- [21] Vincent Liu, Phys. Rev. Lett. **79**, 4056 (1997).
- [22] L.P. Pitaevskii, and S. Stringari, Phys. Lett. A **235**, 398 (1997).
- [23] P.O. Fedichev and G.V. Shlyapnikov, Phys. Rev. A **58**, 3146 (1998).
- [24] A. Sinatra, P. Fedichev, Y. Castin, J. Dalibard, G. Shlyapnikov, Phys. Rev. Lett. **82** 251-254 (1998).
- [25] H. Schmidt, K. Góral, F. Floegel, M. Gajda, K. Rzążewski, J. Opt. B **5**, S96 (2003).
- [26] C. Lobo, A. Sinatra and Y. Castin, Phys. Rev. Lett. **92**, 020403 (2004); N.G. Parker, C.S. Adams, Phys. Rev. Lett. **95**, 145301 (2005).
- [27] A. A. Norrie, R. J. Ballagh, C. W. Gardiner, Phys. Rev. Lett. **94**, 040401 (2005).
- [28] One may argue that the total momentum of the system provides three additional constants of motion. Actually on the lattice model considered here, the momentum along the direction α is conserved modulo $2\pi\hbar/l_\alpha$ only by the interaction term. Since the energy cut-off in the simulations is of order of $k_B T$, $2\pi\hbar/l_\alpha$ is also of the order of the typical momentum of the populated modes of the field, so that we do not expect conservation of momentum. The simulations indeed confirm this expectation.
- [29] This results e.g. from the identity:
- $$\int_0^t d\tau \int_0^t d\tau' \mathcal{C}(|\tau - \tau'|) = 2 \int_0^t (t - \tau) \mathcal{C}(\tau) d\tau.$$
- [30] Y. Castin, R. Dum, Phys. Rev. A **57**, 3008 (1998).
- [31] Y. Castin, Lecture notes of the 2003 Les Houches Spring School, *Quantum Gases in Low Dimensions*, M. Olshanii, H. Perrin, L. Pricoupenko, Eds., J. Phys. IV France **116**, p.89-132 (2004) and cond-mat/0407118.
- [32] C. Mora, Y. Castin, Phys. Rev. A **67**, 053615 (2003).
- [33] C. Gardiner, Phys. Rev. A **56** 1414-1423 (1997).
- [34] J.M. Deutsch, Phys. Rev. A **43**, 2046 (1991) and references therein.
- [35] Expanding (59) to first order in $T_{\text{eff}} - T$ gives
- $$T_{\text{eff}} - T \simeq \frac{\sum_{\mathbf{k} \neq \mathbf{0}} \epsilon_k [n_{\mathbf{k}} - \bar{n}_k(T)]}{\langle H_{\text{Bog}} \rangle'(T)}.$$
- As the $n_{\mathbf{k}}$ are randomly distributed according to the canonical distribution of temperature T , this expression has a vanishing average, and a variance scaling as the inverse of the volume V . When $V \rightarrow \infty$, $\langle A \rangle$ is of order V , the right hand side of (60) is of order V^0 , and $T_{\text{eff}} - T$ is of order $V^{-1/2}$, so that, to be consistent, one actually has to expand the left hand side of (59) up to *second* order in $T_{\text{eff}} - T$ to obtain the next order correction to $T_{\text{eff}} - T$. In evaluating (60) for T replaced by T_{eff} , we expand $\langle A \rangle(T_{\text{eff}})$ up to second order in $T_{\text{eff}} - T$, keeping only terms up to $O(V^0)$; in the right hand side of (60) one can to this order replace T_{eff} by T . One then obtains the microcanonical expectation value \bar{A} up to the desired $O(V^0)$ order, as given by Eq.(61).
- [36] As a consistency check one calculates the average of \bar{A} , that is of Eq.(61), over the canonical distribution for the occupation numbers $n_{\mathbf{k}}$: using $d\bar{n}_k/dT = \epsilon_k \bar{n}_k (\bar{n}_k + 1)/k_B T^2$, one indeed finds $\langle \bar{A} \rangle(T) = \langle A \rangle(T)$.
- [37] More precisely it exactly coincides if one replaces the value of the chemical potential μ by its lowest order value ρg_0 in the Bogoliubov expectation values of products of \hat{A} in the right hand side of Eq.(103) of [32]. This is allowed at the order of accuracy of Eq.(66).
- [38] In the Bogoliubov frame, we use, from the Hellman-Feynman theorem, $\mu_{\text{micro}} = \partial_N E_0 + \sum_{\mathbf{k} \neq \mathbf{0}} \langle \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} \rangle_{\text{micro}} \partial_N \epsilon_k$, where $E_0 = g_0 N(N-1)/2 - \sum_{\mathbf{k} \neq \mathbf{0}} \epsilon_k V_k^2$ is the Bogoliubov ground state energy. We then calculate all the derivatives with respect to N . E.g., $\partial_N \epsilon_k = (g_0/V)(U_k + V_k)^2$. On the other hand, in the spirit of the ergodic method we calculate the expectation value of $(d\hat{\varphi}/dt)_{\text{non osc}}$ in the microcanonical ensemble. The expressions of the chemical potential obtained in these two ways coincide.
- [39] J. Kinast, A. Turlapov, J.E. Thomas, Q. Chen, J. Stajic, K. Levin, Science **307**, 1296 (2005); J. E. Thomas, J. Kinast, and A. Turlapov, Phys. Rev. Lett. **95**, 120402 (2005).
- [40] If we apply this formula to the number of non-condensed particles in the 1D harmonically trapped ideal Bose gas, we recover equation (64) of [41].
- [41] C. Weiss, M. Block, M. Holthaus, G. Schmieder, J. Phys. A **36**, 1827 (2003).