

A simple proof of a recurrence theorem for random walks in $^{\rm 2}$

Jean-Marc Derrien

▶ To cite this version:

Jean-Marc Derrien. A simple proof of a recurrence theorem for random walks in 2 . 2006. hal-00109869

HAL Id: hal-00109869 https://hal.science/hal-00109869

Preprint submitted on 25 Oct 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. A simple proof of a recurrence theorem for random walks in \mathbb{Z}^2

Jean-Marc Derrien

October, 2006

Abstract In this note, we prove without using Fourier analysis that the symmetric square integrable random walks in \mathbb{Z}^2 are recurrent.

George Pòlya related in [3, pp. 582-583] an incident that enables him to formulate the question of recurrence for random walks : during a stroll through the woods, he felt embarrassed because he met "certainly much too often" a student with his girlfriend.

Let x be an element of \mathbb{Z}^2 . A random walk in \mathbb{Z}^2 starting at x is a sequence $(S_n)_{n \in \mathbb{N}}$ of random variables such that

$$S_0 = x$$
 and $S_n = x + X_1 + X_2 + \dots + X_n$, $n \in \mathbb{N}$,

where $(X_n)_{n \in \mathbb{N}}$ is a sequence of independent and identically distributed (i.i.d.), \mathbb{Z}^2 -valued random variables. $(S_n)_{n \in \mathbb{N}}$ is characterized in law by x and the law of X_1 .

In this note, we give an elementary proof of the following result.

Theorem If $(S_n^{(1)})_{n \in \mathbb{N}}$ and $(S_n^{(2)})_{n \in \mathbb{N}}$ are two square integrable, independent and identically distributed random walks in \mathbb{Z}^2 , then

$$S_n^{(1)} = S_n^{(2)}$$
 infinitely often

with probability one.

This theorem is a straightforward consequence of the following proposition (first proved in [2] in a more general context) applied to $(S_n := S_n^{(1)} - S_n^{(2)})_{n \in \mathbb{N}}$.

Proposition If $(S_n)_{n \in \mathbb{N}}$ is a symmetric, square integrable random walk starting at 0 in \mathbb{Z}^2 , then

 $S_n = 0$ infinitely often

with probability one.

Proof It is classical that, in order to prove this proposition, we have only to establish that

$$\sum_{n=0}^{+\infty} \mathbb{P}[S_n = 0] = +\infty$$

(see, for instance, [1]).

One can write

$$S_n = X_1 + X_2 + \dots + X_n \; ,$$

where $(X_k)_{k\geq 1}$ is a sequence of i.i.d., square integrable, \mathbb{Z}^2 -valued random variables with $X_1 \equiv -X_1$ in distribution.

Since the square integrable random walk $(S_n)_{n \in \mathbb{N}}$ is symmetric, it is centered and we have

$$\mathbb{E}(\|S_n\|_2^2) = \sum_{k=1}^n \mathbb{E}(\|X_k\|_2^2) + \sum_{1 \le k < l \le n} \mathbb{E}(X_k) \cdot \mathbb{E}(X_l) = n \ \mathbb{E}(\|X_1\|_2^2)$$

 $(|| \cdot ||_2 \text{ denotes the euclidean norm}).$

The symmetry also gives

$$\mathbb{P}[S_{2n} = 0] = \sum_{x \in \mathbb{Z}^2} \mathbb{P}[S_{2n} = 0 \mid S_n = x] \mathbb{P}[S_n = x]$$

=
$$\sum_{x \in \mathbb{Z}^2} \mathbb{P}[-X_{n+1} - X_{n+2} - \dots - X_{2n} = x] \mathbb{P}[S_n = x] = \sum_{x \in \mathbb{Z}^2} \mathbb{P}[S_n = x]^2.$$

Hence if we introduce

$$B_n := \left\{ x \in \mathbb{Z}^2 : \|x\|_2^2 < 2n \mathbb{E}\left(\|X_1\|_2^2\right) \right\}$$

we deduce from Cauchy-Schwarz's and Markov's inequalities that, if n is large enough,

$$\mathbb{P}[S_{2n} = 0] \geq \frac{1}{|B_n|} \left(\sum_{x \in B_n} \mathbb{P}[S_n = x] \right)^2 \\
\geq \frac{C}{n} \left(1 - \mathbb{P}[\|S_n\|_2^2 \ge 2n \mathbb{E}(\|X_1\|_2^2)] \right)^2 \\
\geq \frac{C}{n} \left(1 - \frac{\mathbb{E}(\|S_n\|_2^2)}{2n \mathbb{E}(\|X_1\|_2^2)} \right)^2 = \frac{C}{4n},$$

where C > 0 depends only on $\mathbb{E}(||X_1||_2^2)$. The proposition follows.

Acknowledgement I am grateful to Emmanuel Lesigne for useful remarks.

REFERENCES

[1] L. Breiman, *Probability*, Addison-Wesley, Reading, 1968.

[2] K. L. Chung and W. H. J. Fuchs, On the distribution of values of sums of random variables, *Mem. Amer. Math. Soc.* **6**, 1951.

[3] G. Pòlya, Collected papers, volume IV, The MIT Press, Cambridge, Massachusetts, 1984.

Département de Mathématiques. Université de Brest. France. Jean-Marc.Derrien@univ-brest.fr