



**HAL**  
open science

# On strongly Petrovskii's parabolic SPDEs in arbitrary dimension and the stochastic Cahn-Hilliard equation

Caroline Cardon-Weber, Annie Millet

► **To cite this version:**

Caroline Cardon-Weber, Annie Millet. On strongly Petrovskii's parabolic SPDEs in arbitrary dimension and the stochastic Cahn-Hilliard equation. *Journal of Theoretical Probability*, 2004, 17, pp.1-49. 10.1023/B:JOTP.0000020474.79479.fa . hal-00111189

**HAL Id: hal-00111189**

**<https://hal.science/hal-00111189>**

Submitted on 3 Nov 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On strongly Petrovskii's parabolic SPDEs in arbitrary dimension

C. Cardon-Weber & A. Millet

*Laboratoire de Probabilités et Modèles Aléatoires, CNRS-UMR 7599,  
Université Paris 6, 4 place Jussieu, Tour 56, F-75252 Paris Cedex 05  
and MODAL'X, Université Paris 10  
email: weber@ccr.jussieu.fr and amil@ccr.jussieu.fr*

## Abstract

In this paper we show that the Cahn-Hilliard stochastic SPDE has a function valued solution in dimension 4 and 5 when the perturbation is driven by a space-correlated Gaussian noise. This is done proving general results on SPDEs with globally Lipschitz coefficients associated with operators on smooth domains of  $\mathbb{R}^d$  which are parabolic in the sense of Petrovskii, and do not necessarily define a semi-group of operators. We study the regularity of the trajectories of the solutions and the absolute continuity of the law at some given time and position.

*Keywords:* Parabolic operators, Cahn-Hilliard equation, Green function, SPDEs, Malliavin calculus.

*AMS Classification:* 60H07, 60H15, 35R60.

## 1 Introduction - Weak solution

Let  $Q$  be a compact subset of  $\mathbb{R}^d$ ,  $\sigma$  and  $b_i$   $1 \leq i \leq N$  be real-valued functions defined on  $[0, T] \times Q \times \mathbb{R}$  and  $(k_i, 1 \leq i \leq N)$  be multi-indices. Let  $F$  denote a one-dimensional  $(d+1)$ -parameter Gaussian noise (either a space-time white noise, or a space correlated noise),  $A(t, x, D_x)$  denote a differential operator of order  $2n$ . Consider the following stochastic partial differential equation

$$\frac{\partial}{\partial t} u(t, x) = A(t, x, D_x) u(t, x) + \sigma(t, x, u(t, x)) \dot{F}(t, x) + \sum_{i=1}^N D_x^{k_i} (b_i(t, x, u(t, x))), \quad (1.1)$$

with some homogeneous boundary conditions, denoted by **(BC)**, defined for  $1 \leq q \leq n$  by

$$B_q(t, x, D_x) u(t, x) = \sum_{|k| \leq r_q} B_{q,k}(t, x) D_x^k u(t, x) = 0 \text{ on } [0, T] \times \partial Q, \quad r_q \leq 2n - 1,$$

and the initial condition  $u(0, \cdot) = u_0$ . We suppose that the operator  $L = \frac{\partial}{\partial t} - A(t, x, D_x)$  is uniformly parabolic in the sense of Petrovskii and that the boundary conditions are complementary and satisfy the normality assumption (for the complete definition see e.g. S.D. Eidelman and N.V. Zhitarashu [13], p. 2-17). The most important class of uniformly strongly parabolic operator in the sense of Petrovskii is defined by:

$$A(t, x, D_x) = \sum_{|k| \leq 2n} a_k(x, t) D_x^k,$$

where there exists a positive constant  $\delta_0$  such that for any  $(x, t) \in Q \times [0, T]$ ,  $\xi \in \mathbb{R}^d$ ,

$$(-1)^n \left( \sum_{|k|=2n} a_k(x, t) \xi^k \right) \leq -\delta_0 |\xi|^{2n}.$$

In this particular case, the definitions can be found in [18], p.113-121. A simple example is provided by  $A(t, x, D_x) = a_1(t, x)D_n(a_2(t, x)D_n)$ , where  $D_n$  is a differential operator of order  $n$ , and  $(-1)^n \sup\{(a_1 a_2)(t, x) : (t, x) \in [0, T] \times Q\} < 0$ . When  $n = 2$  and  $D_2 = \Delta$ , the Dirichlet boundary conditions ( $u = \Delta u = 0$  on  $[0, T] \times \partial Q$ ) or Neumann boundary conditions ( $\frac{\partial}{\partial \nu} u = \frac{\partial}{\partial \nu} \Delta u = 0$  on  $[0, T] \times \partial Q$ , where  $\nu$  denotes the outer normal) are normal and complementary.

A similar equation has been studied by Z. Brzezniak and S. Peszat [3] when  $Q = \mathbb{R}^d$  for smooth bounded coefficients  $a_k$  which do not depend on the time parameter  $t$ . Using time-homogeneous semi-group techniques, these authors prove the existence and uniqueness of a mild solution to (1.1) in some weighted  $L^p$  spaces (or the set of continuous functions having some decay property at infinity), depending on the hypothesis on the initial condition  $u_0$ . Unlike this paper, we allow the case where the differential operator does not yield a time-homogeneous semi-group and work with martingale-measures as in J.B. Walsh [28].

Let  $l \geq 0$  be an integer and  $\lambda \in ]0, 1[$ . According to S.D. Eidelman and N.V. Ivasisen [12], if  $\partial Q$  is of class  $\mathcal{C}^{2n+l+\lambda}$ , the coefficients  $a_k(t, x)$  are of class  $\mathcal{C}^{2n(l+\lambda), l+\lambda}([0, T] \times Q)$  for  $|k| \leq 2n$ , the coefficients  $B_{q,k}(t, x)$  are of class  $\mathcal{C}^{(2n-r_q+l+\lambda)2n, 2n-r_q+l+\lambda}([0, T] \times \partial Q)$ , then if  $G$  denotes the Green function associated with  $L$  and the boundary conditions **(BC)**, for  $|a| + 2nb \leq 2n + l$ , one has for every  $t > s \geq 0$  and  $x, y \in Q$

$$\left| D_x^a \frac{\partial^b}{\partial t^b} G(t, x; s, y) \right| \leq C(t-s)^{-(\alpha+a\delta+b\eta)} \exp\left(-c \frac{|x-y|^\beta}{(t-s)^\gamma}\right), \quad (1.2)$$

$$\text{with } \alpha = \frac{d}{2n}, \beta = \frac{2n}{2n-1}, \gamma = \frac{1}{2n-1}, \delta = \frac{1}{2n} \text{ and } \eta = 1. \quad (1.3)$$

Note that in some cases, it is possible to extend this upper estimate on  $G$  to the case the subset  $Q$  is not smooth (see e.g. [4] for the case  $A = -\Delta^2$  on  $Q = [0, \pi]^d$  and homogeneous Neumann's boundary conditions). Therefore, for  $d < 2n$ , the integral  $\int_0^t \int_Q G^2(t, x; s, y) dy ds$  converges, so that the stochastic integrals of  $G(t, x; s, y)$  with respect to the space-time white noise  $F(ds, dy)$  are well-defined. Usual arguments show that in the particular case of selfadjoint operators  $A(t, x, D_x)$ , such as  $A = a_1(t)D_n(a_2(t, x)D_n)$  with appropriate normal and complementary Dirichlet boundary conditions, the Green function  $G(t, \cdot; s, \cdot)$  is symmetrical in  $x$  and  $y$ , so that  $D_y^a G(t, x; s, y) = D_y^{\tilde{a}} G(t, x; s, y)$  with  $|a| = |\tilde{a}|$ ; then (1.2) holds for  $D_y^a$  instead of  $D_x^a$ .

We now generalize the setting of [28], in order to define a "weak" solution to (1.1), which is an alternative to mild solutions. Return time, and consider the adjoint operator  $L^* = -\frac{\partial}{\partial t} - A^*(t, x, D_x)$  and the adjoint boundary conditions  $B_q^* = 0$ ,  $1 \leq q \leq n$ , on  $[0, t] \times \partial Q$ ; then for fixed  $t > 0$ , exchanging the role of  $(t, x)$  and  $(s, y)$ ,  $G(t, x; s, y)$  is the fundamental solution to the adjoint problem on the time interval  $[0, t]$ . Thus, for any smooth function  $\phi$  on  $Q$ , the function

$$v(s, y) = \int_Q G(t, x; s, y) \phi(x) dx \quad (1.4)$$

is the solution to the equation  $L^*v = 0$  on  $[0, t] \times Q$ , with adjoint boundary conditions ( $B_q^*v = 0$ ,  $1 \leq q \leq n$ ) on  $[0, t] \times \partial Q$ , and such that  $v(t, \cdot) = \phi$ . Then for Dirichlet's systems ( $r_q = q - 1$ ,  $1 \leq q \leq n$ ) or in particular cases (see e.g. example 1.1), for  $v$  defined by (1.4) and "regular"  $u$ , the following Green formula holds (see e.g. [13], p. 231 or [18], p. 133):

$$\int_0^t \int_Q (Lu)(s, y) v(s, y) dy ds + \int_Q u(0, y) v(0, y) dy = \int_Q u(t, x) v(t, x) dx.$$

Furthermore, if integration by parts yields

$$\int_0^t \int_Q D_y^{k_i} (b_i(s, y, u(s, y)) v(s, y)) dy ds = (-1)^{|k_i|} \int_0^t \int_Q b_i(s, y, v(s, y)) D_y^{k_i} v(s, y) dy ds$$

and if  $\int_0^t \int_Q \sigma(s, y, u(s, y)) v(s, y) F(ds, dy)$  is defined as the stochastic integral with respect to a worthy martingale measure, then even if  $u$  is not "regular", we can define the weak solution to (1.1), by requiring that the following form of equation (1.5) holds for every function  $v$  in  $\mathcal{C}^{2n,1}([0, T] \times Q)$ :

$$\int_0^t \int_Q \left[ \sigma(s, y, u(s, y)) v(s, y) F(ds, dy) + \sum_{i=1}^N (-1)^{|k_i|} b_i(s, y, u(s, y)) D_y^{k_i} v(s, y) dy ds \right] + \int_Q u(0, y) v(0, y) dy = \int_Q u(t, x) v(t, x) dx . \quad (1.5)$$

Using (stochastic) Fubini's theorem we obtain the following evolution equation, which is equivalent to (1.5):

$$u(t, x) = \int_Q G(t, x; 0, y) u_0(y) dy + \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u(s, y)) F(ds, dy) + \sum_{i=1}^N \int_0^t \int_Q H_i(t, x; s, y) b_i(s, y, u(s, y)) dy ds , \quad (1.6)$$

where  $H_i(t, x; s, y) = (-1)^{|k_i|} D_y^{k_i} G(t, x; s, y)$ ; if  $G(t, x; s, y)$  is symmetric in  $x$  and  $y$ , the upper estimate (1.2) implies that  $|H_i(t, x; s, y)| \leq C (t - s)^{-(\alpha + |k_i| \delta)} \exp\left(-c \frac{|x - y|^\beta}{(t - s)^\gamma}\right)$ . We give an example where all the requirements (except that on the existence of the stochastic integral) are fulfilled.

**Example 1.1** *The boundary of the set  $Q$  is of class  $\mathcal{C}^{4+l+\lambda}$ , the functions  $a_1(t) \in \mathcal{C}^{4(l+\lambda)}([0, T])$ ,  $a_2(t, x) \in \mathcal{C}^{4(l+\lambda), l+\lambda}([0, T] \times Q)$ ,  $A(t, x, D_x) = a_1(t) \Delta(a_2(t, x) \Delta)$  and  $\sup_{(t, x) \in [0, T] \times Q} a_1(t) a_2(t, x) < 0$ .*

**Case 1** *Dirichlet's boundary conditions:  $u = \Delta u = 0$  on  $[0, T] \times \partial Q$ , either  $0 \leq |k_i| \leq 1$ , or  $|k_i| = 2$  and  $b_i(s, y, z) = \tilde{b}_i(z)$  for some function  $\tilde{b}_i$  of class  $\mathcal{C}_2$  such that  $\tilde{b}_i(0) = 0$ .*

**Case 2** *Neumann's boundary conditions: let  $\nu$  denote the outer normal,  $\frac{\partial}{\partial \nu} u = \frac{\partial}{\partial \nu} \Delta u = 0$ ,  $\frac{\partial}{\partial \nu} a_2(t, x) = 0$  on  $[0, T] \times \partial Q$ ,  $k_i = 0$  or  $|k_i| = 2$ ,  $k_i$  has even components and  $b_i(s, y, z) = \tilde{b}_i(z)$  for some function  $\tilde{b}_i$  of class  $\mathcal{C}_2$ .*

For  $d \geq 2n$ , the function  $G^2(s, x; s, y)$  need not be in  $L^2([0, T] \times Q, ds dy)$ , so that the Gaussian noise  $F$  need not be the space-time white noise; we require the noise  $F$  to be a Gaussian process which is white in time, but has a space correlation defined in terms of a function  $f$  depending on the difference of two vectors of  $\mathbb{R}^d$  (or such that when  $|x - y| \rightarrow 0$ , the product  $f(x, y) |x - y|^a$  remains bounded for some  $a > 0$ ). We just mention a few previous papers on this subject, stressing the type of noise which is used. A particular case of this noise (where the function  $f$  only depends of the norm  $|x - y|$ , such as  $f(x - y) = |x - y|^{-a}$  for  $0 < a < d$ ) has been used in C. Mueller [22], R. Dalang and N. Frangos [7], A. Millet and M. Sanz-Solé [21] in the case  $Q = \mathbb{R}^2$  for the wave operator. In these papers, the existence and uniqueness of a continuous solution is proved by precise estimates of integrals involving the corresponding Green function. A more general covariance structure (depending on the radon measure  $\mu$  with Fourier transform  $f$ , or more generally a tempered distribution  $\Gamma = \hat{\mu}$ ) has been used in [3], [16], [17], [26],[24] for the wave and heat operators on  $\mathbb{R}^d$ ; in the last references, the existence of a solution is proved in some weighted  $L^p$ -space, or in the space of continuous functions with some decay at infinity, and the method uses the Fourier transform of the Green function  $G$  (see e.g. S. Peszat and J. Zabczyk [26] for a detailed account of existence and uniqueness results to parabolic SPDEs with a semi-group structure in any dimension). This general covariance was also used by R. Dalang [6], who proves the existence of continuous processes solutions to the heat and wave stochastic

SPDEs by means of an extension of stochastic integrals with respect to martingale measures for distribution-valued integrands. In these references, the coefficients of the differential operator  $A$  do not depend on  $(t, x)$ .

On the other hand, several attempts have been made to find function-valued solutions to "highly non-linear" stochastic SPDEs, namely PDEs with a polynomial forcing term  $b_i$  (such as the Burgers PDE ( $d = 1$ ,  $A = \Delta$ ,  $N = 1$ ,  $a_1 = 1$  and  $b_1(t, x, y) = y^2$ ), or the Cahn-Hilliard's PDE ( $d \leq 3$ ,  $A = -\Delta^2$ , and  $\sum_i D_x^{k_i} b_i(t, x, u(t, x)) = \Delta R(u(t, x))$ ), where  $R$  is a polynomial of odd degree with positive dominant coefficient) and with a stochastic perturbation driven by the space-time white noise. Thus, G. Da Prato A. Debussche and Temam [9] and then I. Gyöngy [15] have proved the existence of a function-valued solution to the stochastic Burgers equation in dimension 1. G. Da Prato and A. Debussche [8] have proved the existence of a function-valued solution to the stochastic Cahn-Hilliard equation in dimension 1 (up to 3) when the perturbation is driven by a space-time white noise (a Gaussian noise with some spatial correlation). C. Cardon-Weber [4] and [5] has proved the existence of a function-valued solution to the stochastic Cahn-Hilliard equation in dimension  $d \leq 3$  when  $R$  is a polynomial of degree 3 and when the stochastic perturbation is driven by the space-time white noise. The method used in these papers is the following: using a truncation procedure and the existence and uniqueness results proved in the case of globally Lipschitz coefficients, one proves the existence and uniqueness of a solution to the SPDE where the polynomial coefficients have been changed. Then the uniqueness property of the solution allows to use concatenation to obtain the existence of a solution up to some stopping time. Finally, a priori estimates for a deterministic PDE obtained by isolating the stochastic integral (whose behavior is controlled by means of the Garsia lemma), prove that this stopping time is the terminal time  $T$ . These last estimates use methods of analysis which heavily depend on the specific form of the PDE, and no general scheme can be given. Let us finally mention that, using semi-group techniques, Z. Brzezniak and S. Peszat [3] have proved the existence of solutions to some SPDE with a polynomial drift term (when  $N = 1$ ,  $a_1 = 0$ , and when the operator  $A$  is of order 2 and yields a semi-group of operators). Also in [20], the existence of the solution to a stochastic wave equation in dimension 2 with a non-uniformly Lipschitz drift has been proved, while the existence to the stochastic KDV equation has been shown by A. Debussche and A. de Bouard [11].

The aim of this paper is two-fold. On one hand, we prove that in this general context with time and space dependent coefficients, the upper estimates (1.2) of the Green function  $G$  and its time and space derivatives are sufficient to ensure the existence and uniqueness of the solution  $u$  to (1.6), provided that some integrability condition of the covariance function  $f$  on a neighborhood of 0 is required. We prove that, when the Green function  $G$  has a lower estimate by  $t^{-\alpha}$  on the diagonal (which can be the case when it admits an explicit eigenvectors-eigenvalues expansion), this condition is necessary to be able to consider stochastic integrals of  $G$ . As in [28] and [6], we use stochastic integrals with respect to martingale measures. We give sufficient conditions on the covariance function  $f$  for the trajectories of  $u$  to be Hölder-continuous. We then use these results to extend the existence and uniqueness of a function-valued solution to the stochastic Cahn-Hilliard equation when  $Q = [0, \pi]^d$  or a bounded "smooth" subset of  $\mathbb{R}^d$ ,  $d = 4, 5$ . We give necessary and sufficient conditions on the covariance function  $f$  to ensure that the stochastic integral of the corresponding Green function  $G$  is well-defined. We study regularity properties of the trajectories of  $u$  and prove that, if  $Q = [0, \pi]^d$  and the diffusion coefficient  $\sigma$  is strictly elliptic, the law of  $u(t, x)$  has a density for  $t > 0$  and  $x \in Q$ . This extends the results proved in [8] and [4] to higher dimensions. For the sake of simplicity, we mostly restrict ourselves to the case  $F$  is a space-correlated noise; this could be avoided in "small" dimension for arbitrary Petrovskii's parabolic SPDEs. Also note that the proof of the existence of a solution to the stochastic Cahn-Hilliard equation extends directly to the more general

situation described in Example 1.1, when  $R$  is a polynomial of degree 3 with positive dominant coefficient (which has no constant term in the case of the Dirichlet boundary conditions).

The paper is organized as follows. Section 2 gives necessary and sufficient conditions to ensure that stochastic integrals of a function  $G$  which satisfies (1.2) are well defined, and sufficient conditions to ensure Hölder properties of stochastic integrals appearing in (1.6), provided that the process  $u$  has bounded moments. In section 3, we prove both the existence of solutions to (1.6) either in  $\mathcal{C}([0, T], L^q(Q))$  or  $\mathcal{C}([0, T] \times Q)$  when the coefficients are globally Lipschitz functions. We then concentrate on the proof of the a priori estimates which allow to deduce the existence and uniqueness of a solution to the stochastic Cahn-Hilliard equation in dimension 4 and 5. Section 4 establishes Hölder regularity of the trajectories, while section 5 shows the absolute continuity of the law of  $u(t, x)$  for  $t > 0$  and  $x \in Q$  and the solution  $u$  to the Cahn-Hilliard equation.

All the constants  $C$  appearing in the statements can change from one line to the next one. When we want to stress the fact that  $C$  depends on some parameter  $k$ , we denote it by  $C_k$ .

## 2 Stochastic integrals with respect to a space correlated noise

Let  $Q$  be a compact subset of  $\mathbb{R}^d$ ,  $\mathcal{D}(\mathbb{R}_+ \times Q)$  denote the space of functions  $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}_+ \times Q)$  with compact support, endowed with the topology defined by the following convergence:  $\varphi_n \rightarrow \varphi$  if:

- (i) There exists a compact subset  $K$  of  $\mathbb{R}^+ \times Q$  such that  $\text{support}(\varphi_n - \varphi) \subset K$  for all  $n$ .
- (ii)  $\lim_{n \rightarrow +\infty} D^a \varphi_n = D^a \varphi$  uniformly on  $K$  for every multi-index  $a$ .

Let  $F = (F(\varphi), \varphi \in \mathcal{D}(\mathbb{R}_+ \times Q))$  be an  $L^2(P)$ -valued centered Gaussian process, which is white in space but has a space correlation defined as follows: given  $\varphi$  and  $\psi$  in  $\mathcal{D}(\mathbb{R}_+ \times Q)$ , the covariance functional of  $F(\varphi)$  and  $F(\psi)$  is

$$J(\varphi, \psi) = E(F(\varphi) F(\psi)) = \int_0^{+\infty} dt \int_Q dy \int_Q \varphi(t, y) f(y - z) \psi(t, z) dz \quad (2.1)$$

where  $(Q - Q)^* = \{y - z : y, z \in Q, y \neq z\}$  and  $f : (Q - Q)^* \rightarrow [0, +\infty[$  is a continuous function. According to [27], the bilinear form  $J$  defined by (2.1) is non-negative definite if and only if  $f$  is the Fourier transform of a non-negative tempered distribution  $\mu$  on  $Q$ . Then  $F$  defines a martingale-measure (still denoted by  $F$ ), which allows to define stochastic integrals (see [28]).

In this section, we consider fairly general functions  $H : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$ , including Green functions associated with parabolic operators in the sense of Petrovskii: more precisely, we suppose that  $H$  satisfies the following upper estimate for some  $\beta \geq 1$ , some strictly positive parameters  $\alpha$  and  $\gamma$  and some positive constants  $c$  and  $C$ : for any  $t > 0, x, y \in Q$ :

$$|H(t, x; s, y)| \leq C (t - s)^{-\alpha} \exp\left(-c \frac{|x - y|^\beta}{(t - s)^\gamma}\right). \quad (2.2)$$

Then the change of variables  $u = (y - x)(t - s)^{-\frac{\gamma}{\beta}}$  yields

$$\int_Q |H(t, x; s, y)| dy \leq C (t - s)^{-\alpha + \frac{\gamma}{\beta} d}. \quad (2.3)$$

We now give a sufficient integrability condition on the space-correlation function  $f$  of the Gaussian noise  $F$  to ensure that the stochastic integral of a bounded adapted process multiplied by a kernel satisfying (2.2) is a well-defined stochastic process. Under some additional assumptions on  $H$  we prove that this condition is necessary.

**Lemma 2.1** . (i) Let  $H : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$  satisfy (2.2) and suppose that either

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < \infty, \quad \text{if } d = \frac{\beta}{\gamma}(2\alpha - 1), \quad (2.4)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-[\frac{\beta}{\gamma}(2\alpha-1)-d]^+} dv < \infty, \quad \text{if } d \neq \frac{\beta}{\gamma}(2\alpha - 1). \quad (2.5)$$

Then for any  $t \in [0, T]$ ,

$$I(t) = \int_0^T \int_Q \int_Q |H(t, x; s, y)| f(y - z) |H(t, x; s, z)| dy dz dt < +\infty. \quad (2.6)$$

(ii) Let  $H : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$  satisfy

$$\inf\{H(t, x; s, x) : x \in Q\} \geq C_0(t - s)^{-\alpha} \quad (2.7)$$

and for every multi-index  $k$  with  $|k| = 1$ :

$$\sup\{|D_y^k H(t, x; s, y)| : x, y \in Q\} \leq C_1(t - s)^{-(\alpha+\delta)}, \quad (2.8)$$

for  $\alpha, \delta > 0$ , some constant  $C > 0$  and any  $t > s > 0$ . Then if (2.6) holds for every  $x \in Q$  and  $t \in ]0, T]$ , one has either (2.9) or (2.10) depending on  $d$ , where:

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < \infty, \quad \text{if } d = \frac{1}{\delta}(2\alpha - 1), \quad (2.9)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-[\frac{1}{\delta}(2\alpha-1)-d]^+} dv < \infty, \quad \text{if } d \neq \frac{1}{\delta}(2\alpha - 1). \quad (2.10)$$

**Remark 2.2** . When  $H$  is the Green function of the operator  $\frac{\delta}{\delta t} + \Delta^2$  on  $Q = [0, \pi]^d$  with homogeneous Neumann's or Dirichlet's boundary conditions, condition (2.7) holds with  $\alpha = \frac{d}{4}$ . We only sketch the proof for Neumann's boundary conditions; in that case,  $G(t, x; s, x) \geq \sum_{k \in \mathbb{N}^{*d}} \exp(-|k|^4(t - s)) \left( \prod_{1 \leq i \leq d} \cos^2(k_i x_i) \right)$ . Set  $\mathcal{I} = ]\frac{\pi}{2} - \theta, \frac{\pi}{2} + \theta[$  for some  $\theta > 0$  small enough; then for  $k \in \mathbb{N}^*$  and  $x \in ]0, \pi[$ ,  $kx \in \mathcal{I}$  implies that  $(k + 1)x \notin \mathcal{I}$ . Skipping at most every other term and using the monotonicity of  $(\exp(-k^4(t - s)); k \geq 1)$ , we deduce that  $G(t, x; s, x) \geq C \sum_{k \in \mathbb{N}^{*d}} \exp(-c|k|^4(t - s)) = C(t - s)^{-\frac{d}{4}}$  for some positive constants  $c$  and  $C$  (see Remark 3.5 in [4] for a similar argument). Since in the case of Green functions of parabolic operators,  $\delta = \frac{2}{\beta}$ , the conditions (2.4) and (2.9) (respectively (2.5) and (2.10)) are identical.

*Proof of Lemma 2.1:* (i) Since  $\beta \geq 1$ ,  $|y - z|^\beta \leq 2^{\beta-1}(|x - y|^\beta + |x - z|^\beta)$ , so that for  $\lambda \in ]0, 1[$  and  $t_0 \in [0, T]$ ,

$$\begin{aligned} I(t_0) &\leq C \int_0^T t^{-2\alpha} dt \int_Q \exp\left(-c(1 - \lambda) \frac{|x - y|^\beta}{t^\gamma}\right) dy \\ &\quad \times \int_Q \exp\left(-c2^{1-\beta} \lambda \frac{|y - z|^\beta}{t^\gamma}\right) f(y - z) dz \\ &\leq C \int_0^T t^{-2\alpha} dt \int_Q \exp\left(-\bar{c} \frac{|x - y|^\beta}{t^\gamma}\right) dy \int_Q \exp\left(-\bar{c} \frac{|y - z|^\beta}{t^\gamma}\right) f(y - z) dz. \end{aligned} \quad (2.11)$$

Set  $t^{\frac{\gamma}{\beta}}\eta = x - y$ ,  $v = y - z$  and then  $u = |\eta|$ ; there exist positive constants  $C$ ,  $c$  and  $R$  such that

$$\begin{aligned} I(t_0) &\leq C \int_0^T t^{\frac{\gamma}{\beta}d-2\alpha} dt \int_0^{+\infty} \exp(-cu^\beta) u^{d-1} du \int_{B_d(0,R)} \exp\left(-c\frac{|v|^\beta}{t^\gamma}\right) f(v) dv \\ &\leq C \int_0^T \psi(t) dt, \end{aligned} \quad (2.12)$$

where for any  $0 < t \leq T$  one sets

$$\psi(t) = t^{\frac{\gamma}{\beta}d-2\alpha} \int_{B_d(0,R)} \exp\left(-c\frac{|v|^\beta}{t^\gamma}\right) f(v) dv. \quad (2.13)$$

For fixed  $v \neq 0$ , set  $r = |v|^\beta t^{-\gamma}$ ; then Fubini's theorem yields

$$\int_0^T \psi(t) dt \leq C \int_{B_d(0,R)} f(v) |v|^{d+\frac{\beta}{\gamma}(1-2\alpha)} dv \int_{|v|^\beta T^{-\gamma}}^{+\infty} r^{-1+\frac{1}{\gamma}(2\alpha-1-\frac{\gamma d}{\beta})} \exp(-cr) dr.$$

We now distinguish three cases:

*Case 1* If  $2\alpha > 1 + \frac{\gamma d}{\beta}$ , then the second integral is bounded by a constant independent of  $|v|$  and  $\int_0^T \psi(t) dt \leq C \int_{B_d(0,R)} f(v) |v|^{d+\frac{\beta}{\gamma}(1-2\alpha)} dv$ , which yields (2.5).

*Case 2* If  $2\alpha = 1 + \frac{\gamma d}{\beta}$ , since for  $0 < |v| \leq R$ ,

$$\int_{|v|^\beta T^{-\gamma}}^{+\infty} r^{-1} \exp(-cr) dr \leq C |1 - \ln(|v|) 1_{\{|v| \leq 1\}}|,$$

we have  $\int_0^T \psi(t) dt \leq C \int_{B_d(0,R)} f(v) [1 + \ln(|v|^{-1}) 1_{\{|v| \leq 1\}}] dv$ , which yields (2.4)

*Case 3* Finally, if  $2\alpha < 1 + \frac{\gamma d}{\beta}$ , then for  $0 < |v| \leq R$ ,

$$\int_{|v|^\beta T^{-\gamma}}^{+\infty} r^{-1+\frac{1}{\gamma}(2\alpha-1-\frac{\gamma d}{\beta})} \exp(-cr) dr \leq C \left(1 + |v|^{\frac{\beta}{\gamma}(2\alpha-1)-d}\right),$$

so that  $\int_0^T \psi(t) dt \leq C \int_{B_d(0,R)} f(v) dv$ , which yields (2.5).

Note that for small  $T$ , the following computation gives a more precise upper estimate of  $\int_0^T \psi(t) dt$ , which will be used in the sequel. Indeed, for  $\nu \in ]0, \gamma[$ , the decomposition of the integral over  $B_d(0, R)$  into  $\{|v|^\beta T^{-\gamma} \geq T^{-\nu}\}$  and its complement yields

$$\begin{aligned} \int_0^T \psi(t) dt &\leq C \int_{B_d\left(0, T^{\frac{\gamma-\nu}{\beta}}\right)} f(v) |v|^{d+\frac{\beta}{\gamma}(1-2\alpha)} dv \int_{|v|^\beta T^{-\gamma}}^{+\infty} r^{-1+\frac{1}{\gamma}(2\alpha-1-\frac{\gamma d}{\beta})} \exp(-cr) dr \\ &\quad + C I \exp(-\bar{c}T^{-\nu}). \end{aligned}$$

Thus for  $0 < \nu < \gamma$  and  $2\alpha \neq 1 + \frac{\gamma d}{\beta}$  one has

$$\int_0^T \psi(t) dt \leq C \left[ \exp(-\bar{c}T^{-\nu}) + \int_{B_d\left(0, T^{\frac{\gamma-\nu}{\beta}}\right)} f(v) |v|^{\left[-\frac{\beta}{\gamma}(2\alpha-1)-d\right]^+} dv \right], \quad (2.14)$$

while for  $0 < \nu < \gamma$  and  $2\alpha = 1 + \frac{\gamma d}{\beta}$  one has

$$\int_0^T \psi(t) dt \leq C \left[ \exp(-\bar{c}T^{-\nu}) + \int_{B_d\left(0, T^{\frac{\gamma-\nu}{\beta}}\right)} f(v) \ln(|v|^{-1}) dv \right]. \quad (2.15)$$



(ii) The assumptions (2.7) and (2.8) imply that for  $|x - y| \leq 2C_2 t^\delta$  with  $C_2 < \frac{C_0}{4C_1}$  small enough, one has

$$\begin{aligned} H(t, x; s, y) &\geq H(t, x; s, x) - |H(t, x; s, x) - H(t, x; s, y)| \\ &\geq (C_0 - 2C_1 C_2) (t - s)^{-\alpha} \geq \frac{C_0}{2} (t - s)^{-\alpha}. \end{aligned}$$

Let  $a > 0$  be such that  $Q_a = \{x \in Q : d(x, \partial Q) > a\} \neq \emptyset$ ; then for  $0 < 2C_2 t_0^\delta \leq a$ ,  $0 < s \leq t_0 \leq T$ ,  $x \in Q_{2C_2 t_0^\delta}$ ,  $y \in B_d(x, C_2 s^\delta)$  and  $z \in B_d(y, C_2 s^\delta)$ , one has  $y, z \in Q$ . Thus Fubini's theorem implies for  $x \in Q_{2C_2 t_0^\delta} \neq \emptyset$ :

$$\begin{aligned} I(t_0) &\geq \int_0^{t_0} ds \int_{Q \cap B_d(x, C_2 s^\delta)} dy \int_{Q \cap B_d(y, C_2 s^\delta)} |H(t_0, x; s, y)| f(y - z) |H(t_0, x; s, z)| dz \\ &\geq C \int_0^{t_0} s^{-2\alpha + d\delta} ds \int_{B_d(0, C_2 s^\delta)} f(v) dv \geq C \int_{B_d(0, R)} f(v) dv \int_{\left(\frac{|v|}{C_2}\right)^\frac{1}{\delta}}^{t_0} s^{-2\alpha + d\delta} ds \end{aligned}$$

for  $R = C_2 t_0^\delta$ . Again we have to study three cases depending on the power of  $s$ .

*Case 1* If  $d\delta + 1 < 2\alpha$ , let  $\bar{R} = \frac{R}{2}$ ; then one has

$$I(t_0) \geq C \int_{B_d(0, \bar{R})} f(v) dv \int_{\left(\frac{|v|}{C_2}\right)^\frac{1}{\delta}}^{\left(\frac{2|v|}{C_2}\right)^\frac{1}{\delta}} s^{-2\alpha + d\delta} ds \geq C \int_{B_d(0, \bar{R})} f(v) |v|^{\frac{1-2\alpha}{\delta} + d} dv, \quad (2.16)$$

which yields (2.10).

*Case 2* If  $d\delta + 1 = 2\alpha$ , let  $\nu > 0$  and let  $\bar{R} = R \wedge 1 \wedge (C_2^{-\nu} R^{1+\nu})$ ; then  $|v| \leq \bar{R}$  and  $t_0 \leq C_2^\frac{1}{\delta}$  imply  $\left(\frac{|v|}{C_2}\right)^{\frac{1}{(1+\nu)\delta}} \leq t_0$  and  $|v|^{\frac{1}{2}} C_2^{-1} \leq 1$ ; hence  $C_2 |v|^{-1} \geq |v|^{-\frac{1}{2}}$  and

$$I(t_0) \geq C \int_{B_d(0, \bar{R})} f(v) dv \int_{\left(\frac{|v|}{C_2}\right)^\frac{1}{\delta}}^{\left(\frac{|v|}{C_2}\right)^{\frac{1}{(1+\nu)\delta}}} s^{-1} ds \geq C \int_{B_d(0, \bar{R})} f(v) \ln(|v|^{-1}) dv, \quad (2.17)$$

which yields (2.9).

*Case 3* Finally, if  $d\delta + 1 > 2\alpha$ , one has  $I(t_0) \geq C \int_{B_d(0, R)} f(v) dv$ , which yields (2.10).  $\square$

The following lemma gives sufficient conditions on the covariance function  $f$  to obtain moment estimates of stochastic integrals which yield Hölder regularity of the corresponding process. For this, we impose an upper estimate of the space and time partial derivatives of the kernel  $H$ : there exist positive constants  $\delta, \eta, c, C$  such that for any  $t > 0$ ,  $x, y \in Q$  and  $k \in \mathbb{N}^d$  with  $|k| = 1$ ,

$$\left| D_x^k H(t, x; s, y) \right| \leq C(t - s)^{-(\alpha + \delta)}, \quad \left| \frac{\partial}{\partial t} H(t, x; s, y) \right| \leq C(t - s)^{-(\alpha + \eta)} \exp\left(-c \frac{|x - y|^\beta}{(t - s)^\gamma}\right). \quad (2.18)$$

In order to deal with time increments, we impose also that  $f$  satisfies the following "monotonicity" condition:

**(C1)** There exist strictly positive constants  $C_1$  and  $c_1$  such that

$$f(u) \leq C_1 f(v) \quad \text{for } |v| \leq c_1 |u|. \quad (2.19)$$

Note that (C1) holds if  $f(u) = |u|^{-a}$  for some  $a > 0$ .

**Lemma 2.3** . Suppose that  $Q$  is convex and let  $H : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$  satisfy the condition (2.2). Let  $F$  be a Gaussian noise with spatial covariance defined by (2.1) such that the correlation function  $f$  satisfies (C1) and either (2.4) or (2.5). Fix  $p \in [1, +\infty[$ , let  $u : \Omega \rightarrow \mathbb{R}$  be an adapted process such that  $\sup_{(t,x) \in [0,T] \times Q} E(|u(t,x)|^{2p}) < +\infty$  and for  $t \in [0, T]$  and  $x \in Q$ , let

$$I(t, x) = \int_0^t \int_Q H(t, x; s, y) u(s, y) F(ds, dy). \quad (2.20)$$

(i) Suppose that  $H$  satisfies (2.18) and let  $a \in ]0, 1[$ ; if either

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < +\infty \quad \text{for } d = \frac{\beta}{\gamma} (2\alpha + a\delta - 1), \quad (2.21)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-[\frac{\beta}{\gamma}(2\alpha+a\delta-1)-d]^+} dv < +\infty \quad \text{for } d \neq \frac{\beta}{\gamma} (2\alpha + a\delta - 1); \quad (2.22)$$

then there exists  $C_p > 0$  such that for every  $x, x' \in Q$ ,

$$A(x, x') = \sup_{t \in [0, T]} E(|I(t, x) - I(t, x')|^{2p}) \leq C_p |x - x'|^{ap}. \quad (2.23)$$

(ii) Suppose that  $H$  satisfies (1.2) and (1.3) and let  $b \in ]0, 1[$ ; if either

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < +\infty \quad \text{for } d = \frac{\beta}{\gamma} (2\alpha + b\eta - 1), \quad (2.24)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-[\frac{\beta}{\gamma}(2\alpha+b\eta-1)-d]^+} dv < +\infty \quad \text{for } d \neq \frac{\beta}{\gamma} (2\alpha + b\eta - 1); \quad (2.25)$$

then there exists  $C_p > 0$  such that for every  $0 \leq t < t' \leq T$ ,

$$B(t, t') = \sup_{x \in Q} E \left( \left| \int_0^t \int_Q [H(t, x; s, y) - H(t', x; s, y)] u(s, y) F(ds, dy) \right|^{2p} \right) \leq C_p |t - t'|^{bp},$$

and

$$C(t, t') = \sup_{x \in Q} E \left( \left| \int_t^{t'} \int_Q H(t', x; s, y) u(s, y) F(ds, dy) \right|^{2p} \right) \leq C_p |t - t'|^{bp}. \quad (2.26)$$

*Proof:* (i) Burkholder's inequality implies that for every  $p \in [1, +\infty[$ ,

$$\begin{aligned} A(x, x') &\leq C_p \sup_{t \in [0, T]} E \left| \int_0^t ds \int_Q dy \int_Q |H(t, x; s, y) - H(t, x'; s, y)| |u(s, y)| \right. \\ &\quad \left. \times f(y - z) |H(t, x; s, z) - H(t, x'; s, z)| |u(s, z)| dz \right|^p. \end{aligned}$$

We prove that for  $\Delta(x, x') = \sup_{t \in [0, T]} \int_0^t ds \int_Q dy \int_Q dz |H(t, x; s, y) - H(t, x'; s, y)| f(y - z) \times |H(t, x; s, z) - H(t, x'; s, z)|$ , one has

$$\Delta(x, x') \leq C_p |x - x'|^a, \quad (2.27)$$

provided that  $H$  satisfies either (2.21) or (2.22). Assuming that (2.27) holds, Hölder's and Schwarz's inequalities yield

$$\begin{aligned} A(x, x') &\leq C_p \Delta(x, x')^{p-1} \sup_{t \in [0, T]} \int_0^t ds \int_Q dy |H(t, x; s, y) - H(t, x'; s, y)| \\ &\quad \times \int_Q f(y - z) |H(t, x; s, z) - H(t, x'; s, z)| E(|u(s, y)|^p |u(s, z)|^p) dz \\ &\leq C_p \Delta(x, x')^p \sup_{(s, y) \in [0, T] \times Q} E(|u(s, y)|^{2p}) \leq C_p |x - x'|^{ap}. \end{aligned}$$

In order to prove (2.27), we use Taylor's formula, the convexity of  $Q$  and the inequalities (2.2) and (2.18); thus for any  $a \in ]0, 1[$ ,  $\Delta(x, x') \leq |x - x'|^a (T_1 + T_2)$ , where

$$\begin{aligned} T_1 &= \sup_{x \in Q} \int_0^T s^{-(2\alpha + a\delta)} ds \int_Q dy \int_Q \exp\left(-c \frac{|x - y|^\beta}{s^\gamma}\right) f(y - z) \exp\left(-c \frac{|x - z|^\beta}{s^\gamma}\right) dz, \\ T_2 &= \int_0^T s^{-(2\alpha + a\delta)} ds \int_Q dy \int_Q \exp\left(-c \frac{|x - y|^\beta}{s^\gamma}\right) f(y - z) \exp\left(-c \frac{|x' - z|^\beta}{s^\gamma}\right) dz. \end{aligned}$$

Replacing  $2\alpha$  by  $2\alpha + a\delta$ , the arguments used to prove part (i) of Lemma 2.1 show that if either (2.21) or (2.22) holds, then  $T_1 < +\infty$ . To study  $T_2$ , we have to distinguish several cases. Let  $0 < c_2 < 1$ ,  $k \geq 1$  be such that  $c_2^k < \frac{1}{3}$  and set  $\varepsilon = c_2^{k+1}$ ,  $\bar{\varepsilon} = \frac{c_2^k}{1 - c_2^{k+1}}$ ; then  $\frac{1}{1 + \varepsilon} > \bar{\varepsilon}$  and  $\varepsilon < 1$ .

We study three cases:

*Case 1* If  $|x - y| \geq \varepsilon |x' - y|$  or  $|x' - z| \geq \varepsilon |x - z|$ , we have (changing the constant in the exponential functions)  $T_2 \leq C(\varepsilon) T_1$ , and the proof is complete.

*Case 2* If  $|x - y| < \varepsilon |x' - y|$ ,  $|x' - z| < \varepsilon |x - z|$  and  $|y - z| \leq \bar{\varepsilon} |x - x'|$ , then  $|x - z| \geq \frac{|x - x'|}{1 + \varepsilon}$  and we have

$$|x - y| \geq ||x - z| - |y - z|| \geq \left(\frac{1}{1 + \varepsilon} - \bar{\varepsilon}\right) |x - x'| \geq \left(\frac{1}{\bar{\varepsilon}(1 + \varepsilon)} - 1\right) |y - z|.$$

This implies for  $\bar{c} = c \min\left(1, \frac{1}{\bar{\varepsilon}(1 + \varepsilon)} - 1\right)$  which is positive by the choice of  $k$ :

$$T_2 \leq \int_0^T s^{-(2\alpha + a\delta)} ds \int_Q dy \int_Q \exp\left(-\bar{c} \frac{|y - z|^\beta}{s^\gamma}\right) f(y - z) \exp\left(-\bar{c} \frac{|x' - z|^\beta}{s^\gamma}\right) dz,$$

and since this is the upper estimate of (2.11), the proof is again concluded by an argument similar to that in Lemma 2.1 (i), with  $2\alpha + a\delta$  instead of  $2\alpha$ .

*Case 3* Suppose finally that  $|x - y| < \varepsilon |x' - y|$ ,  $|x' - z| < \varepsilon |x - z|$  and  $|y - z| > \bar{\varepsilon} |x - x'|$ . Then  $|x' - y| \leq \frac{|x - x'|}{1 - \varepsilon}$  and  $|x - y| \leq \frac{\varepsilon}{1 - \varepsilon} |x - x'| \leq \frac{\varepsilon}{(1 - \varepsilon)\bar{\varepsilon}} |y - z| = c_2 |y - z|$ , so that  $\frac{c_1}{c_2} |x - y| \leq c_1 |y - z|$ . Since  $f$  satisfies (2.19), we have  $f(y - z) \leq C_1 f\left(\frac{c_1}{c_2} (y - x)\right)$ . Set  $\bar{y} = x + \frac{c_1}{c_2} (y - x)$  and  $\tilde{c} = c \min\left(\frac{c_1}{c_2}, 1\right)$ ; then

$$T_2 \leq C \int_0^T s^{-(2\alpha + a\delta)} ds \int_Q d\bar{y} \int_Q \exp\left(-\tilde{c} \frac{|x - \bar{y}|^\beta}{s^\gamma}\right) f(\bar{y} - x) \exp\left(-\tilde{c} \frac{|x' - z|^\beta}{s^\gamma}\right) dz,$$

and again the proof is complete, since the right hand-side is similar to (2.11).

(ii) For  $0 \leq t < t' \leq T$ , set

$$\bar{\Delta}(t, t') = \sup_{x \in Q} \left| \int_0^t ds \int_Q dy |H(t, x; s, y) - H(t', x; s, y)| \right|$$

$$\times \int_Q f(y-z) |H(t, x; s, z) - H(t', x; s, z)| dz \Big|^p.$$

Again we prove that under either condition (2.24) or (2.25) we have

$$\overline{\Delta}(t, t') \leq C|t - t'|^b. \quad (2.28)$$

If (2.28) holds, using again Burkholder's, Hölder's and Schwarz's inequalities, we deduce that

$$B(t, t') \leq C_p \overline{\Delta}(t, t')^p \sup_{(t, y) \in [0, T] \times Q} E(|u(s, y)|^{2p}) \leq C_p |t - t'|^{bp}.$$

We now prove (2.28). Using (2.2), (2.18) and Taylor's formula, we obtain for  $h = t' - t$ : for any  $b \in ]0, 1[$ ,  $\overline{\Delta}(t, t') \leq |t - t'|^b (T'_1 + T'_2)$ , where

$$\begin{aligned} T'_1 &= \sup_{x \in Q} \int_0^t s^{-(2\alpha + b\eta)} ds \int_Q dy \int_Q \exp\left(-c \frac{|x-y|^\beta}{s^\gamma}\right) f(y-z) \exp\left(-c \frac{|x-z|^\beta}{s^\gamma}\right) dz, \\ T'_2 &= \sup_{x \in Q} \int_0^t s^{-(\alpha + b\eta)} (s+h)^{-\alpha} ds \int_Q dy \exp\left(-c \frac{|x-y|^\beta}{s^\gamma}\right) \\ &\quad \times \int_Q f(y-z) \exp\left(-c \frac{|x-z|^\beta}{(s+h)^\gamma}\right) dz. \end{aligned}$$

Clearly,  $T'_1$  is similar to  $T_1$  with  $b\eta$  instead of  $a\delta$ ; thus the proof of (i) yields (2.26) if either (2.24) or (2.25) holds. To estimate  $T'_2$ , we distinguish two cases.

*Case 1* If  $|x - y| \leq c_1 |y - z|$ , condition (2.19) on  $f$  yields

$$\begin{aligned} T'_2 &\leq C_1 \int_0^t s^{-(\alpha + b\eta)} (s+h)^{-\alpha} ds \int_Q \exp\left(-c \frac{|x-y|^\beta}{s^\gamma}\right) f(x-y) dy \\ &\quad \times \int_Q \exp\left(-c \frac{|x-z|^\beta}{(s+h)^\gamma}\right) dz. \end{aligned}$$

Set  $v = x - y$ ; then since  $\alpha \geq \frac{\gamma}{\beta}d$ , we have for some  $R > 0$ ,

$$\begin{aligned} T'_2 &\leq C_1 \int_0^t s^{-(\alpha + b\eta)} (s+h)^{-\alpha + \frac{\gamma}{\beta}d} ds \int_{B_d(0, R)} \exp\left(-c \frac{|v|^\beta}{s^\gamma}\right) f(v) dv \\ &\leq C_1 \int_0^t s^{-(2\alpha + b\eta) + \frac{\gamma}{\beta}d} ds \int_{B_d(0, R)} \exp\left(-c \frac{|v|^\beta}{s^\gamma}\right) f(v) dv. \end{aligned}$$

This last upper estimate is similar to the right hand side of (2.12) with  $2\alpha + b$  instead of  $2\alpha$ ; thus the end of the proof of Lemma 2.1 (i) concludes the proof.

*Case 2* If  $|x - y| > c_1 |y - z|$ , then for  $\bar{c} = \min(c, c c_1^\beta)$ , we have

$$\begin{aligned} T'_2 &\leq \sup_{x \in Q} \int_0^t s^{-(\alpha + b\eta)} (s+h)^{-\alpha} ds \int_Q dy \\ &\quad \times \int_Q \exp\left(-\bar{c} \frac{|y-z|^\beta}{s^\gamma}\right) f(y-z) \exp\left(-\bar{c} \frac{|x-z|^\beta}{(s+h)^\gamma}\right) dz. \end{aligned}$$

Since  $(s+h)^{-\alpha + \frac{\gamma}{\beta}d} \leq s^{-\alpha + \frac{\gamma}{\beta}d}$ , the change of variables  $\zeta = x - z$  and  $v = y - z$  shows that  $T'_2$  is dominated by the right hand side of (2.12) with  $2\alpha + b$  instead of  $2\alpha$  and  $\bar{c}$  instead of  $c$ ; this concludes the proof of (2.26)

Finally, using again Burkholder's inequality and (2.2), we have  $C(t, t') \leq C_p T_3^p \sup_{(s,y) \in [0,T] \times Q} E(|u(s,y)|^{2p})$ , where

$$T_3 = \sup_{x \in Q} \int_0^{t'-t} s^{-2\alpha} ds \int_Q dy \int_Q \exp\left(-c \frac{|x-y|^\beta}{s^\gamma}\right) f(y-z) \exp\left(-c \frac{|x-z|^\beta}{s^\gamma}\right) dz.$$

Computations similar to those in the proof of Lemma 2.1 (i) imply that for some  $R > 0$ ,  $T_3 \leq \int_0^{t'-t} \psi(s) ds$ , where  $\psi$  is defined by (2.13). Fubini's theorem and Hölder's inequality with respect to  $ds$  with the conjugate exponents  $\lambda = (b\eta)^{-1}$  and  $\mu$  imply

$$T_3 \leq |t' - t|^b \int_{B_d(0,R)} f(v) dv \left( \int_0^T \exp\left(-c \frac{|v|^\beta}{s^\gamma}\right) s^{\mu(\frac{\gamma}{\beta}d - 2\alpha)} ds \right)^{\frac{1}{\mu}}.$$

For  $v \neq 0$  set  $r = |v|^\beta s^{-\gamma}$ ; then since  $\frac{1}{\mu} = 1 - b$ , we obtain

$$T_3 \leq C |t' - t|^b \int_{B_d(0,R)} f(v) |v|^{d + \frac{\beta}{\gamma}(1 - 2\alpha - b)} dv \left( \int_{|v|^\beta T^{-\gamma}}^{+\infty} r^{-1 + \frac{\mu}{\gamma}[2\alpha - \frac{\gamma}{\beta}d - 1 + b]} \exp(-cr) dr \right)^{\frac{1}{\mu}}.$$

As in the proof of Lemma 2.1, we distinguish three cases, according to the power of  $r$  in the last integral, with  $2\alpha + b$  instead of  $2\alpha$ ; this concludes the proof of (2.26).  $\square$

### 3 Existence of solutions

#### 3.1 The case of Lipschitz coefficients

Let  $\sigma : [0, T] \times Q \times \mathbb{R} \rightarrow \mathbb{R}$  and  $b_i : [0, T] \times Q \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $1 \leq i \leq N$  be continuous functions such that the following boundness and Lipschitz conditions hold:

**(L1)** Uniform linear growth with respect to the last variable: for every  $y \in Q$

$$\sup_{(t,x) \in [0,T] \times Q} \left( |\sigma(t, x, y)| + \sum_{i=1}^N |b_i(t, x, y)| \right) \leq C(1 + |y|). \quad (3.1)$$

**(L2)** Uniform Lipschitz condition with respect to the last variable: for any  $y, z \in Q$

$$\sup_{(t,x) \in [0,T] \times Q} \left( |\sigma(t, x, y) - \sigma(t, x, z)| + \sum_{i=1}^N |b_i(t, x, y) - b_i(t, x, z)| \right) \leq C|y - z|. \quad (3.2)$$

We then consider the following non-linear evolution equation for  $t \in [0, T]$  and  $x \in Q$ :

$$\begin{aligned} u(t, x) &= \int_Q G(t, x; 0, y) u_0(y) dy + \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u(s, y)) F(ds, dy) \\ &\quad + \sum_{i=1}^N \int_0^t \int_Q H_i(t, x; s, y) b_i(s, y, u(s, y)) dy ds, \end{aligned} \quad (3.3)$$

for a function  $u_0 : Q \rightarrow \mathbb{R} \in L^2(Q)$ .

In this section we will make the following assumptions (restricting ourselves to the case  $G$  is the Green function of an operator which is parabolic in the sense of Petrovskiĭ, and the functions  $H_i$  are partial derivatives of  $G$  with respect to the space variable  $y$ ):

**(C2)** The continuous function  $G : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$  (respectively for  $i = 1, \dots, N$  each continuous functions  $H_i : ([0, T] \times Q)^2 \rightarrow \mathbb{R}$ ) satisfies (2.2) with constants  $\beta, \gamma$  and  $\alpha = \frac{\gamma}{\beta}d$  (respectively  $\alpha_i, \beta$  and  $\gamma$ ).

**(C3)** The covariance of the Gaussian process  $F$  is defined in terms of  $f$  by (2.1), and the constants  $\alpha, \beta, \gamma$  in (C2) satisfy either

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < \infty, \quad \text{if } d = \frac{\beta}{\gamma}, \quad (3.4)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-[d-\frac{\beta}{\gamma}]^+} dv < \infty, \quad \text{if } d \neq \frac{\beta}{\gamma}. \quad (3.5)$$

**(C4)** The constants  $\alpha_i, \beta, \gamma$  in (C2) satisfy  $\alpha_i < \alpha + 1$  for every  $i \in \{1, \dots, N\}$ .

Condition (C3) will allow to define stochastic integrals of  $G(t, x; \cdot)$  with respect to the noise  $F$ , while (C4) will allow to define deterministic integrals involving  $H_i(t, x; \cdot)$ .

We at first study moment estimates of deterministic integrals. Let  $\lambda, \rho \in [1, +\infty]$ ; for  $v \in L^\lambda([0, T], L^\rho(Q))$ , and  $0 \leq t_0 \leq t \leq T$ ,  $x \in Q$  set

$$J(v)(t_0, t, x) = \int_{t_0}^t \int_Q H(t, x; s, y) v(s, y) dy ds. \quad (3.6)$$

The following lemma provides  $L^q$  estimates of  $J(v)(t_0, t, \cdot)$  in terms of  $L^\rho$  estimates of  $v(s, \cdot)$ . It extends similar results proved in I. Gyöngy [15] and C. Cardon-Weber [4].

**Lemma 3.1** . Fix  $\rho \in [1, +\infty]$ ,  $q \in [\rho, +\infty]$ , and let  $r$  be defined by  $\frac{1}{r} = \frac{1}{q} - \frac{1}{\rho} + 1$ . Then for  $0 \leq t_0 \leq t \leq T$

$$\|J(v)(t_0, t, \cdot)\|_q \leq C \int_{t_0}^t (t-s)^{-\alpha + \frac{\gamma d}{\beta r}} \|v(s, \cdot)\|_\rho ds. \quad (3.7)$$

Hence given any  $\lambda \in [0, +\infty]$ , if  $\alpha + \frac{1}{\lambda} < \frac{\gamma d}{\beta r} + 1$ , then  $J(0, \cdot, \cdot)$  is a bounded operator from  $L^\lambda([0, T], L^\rho(Q))$  into  $L^\infty([0, T], L^q(Q))$ .

*Proof:* Using Minkowski's inequality, (2.2), then Young's inequality with  $\frac{1}{q} = \frac{1}{r} + \frac{1}{\rho} - 1$  and (2.3), we obtain

$$\begin{aligned} \|J(v)(t_0, t, \cdot)\|_q &\leq C \int_{t_0}^t \left\| \int_Q (t-s)^{-\alpha} \exp\left(-c \frac{|\cdot - y|^\beta}{(t-s)^\gamma}\right) |v(s, y)| dy \right\|_q ds \\ &\leq C \int_{t_0}^t (t-s)^{-\alpha} \|v(s, \cdot)\|_\rho \left\| \exp\left(-c \frac{|\cdot|^\beta}{(t-s)^\gamma}\right) \right\|_r ds \\ &\leq C \int_{t_0}^t (t-s)^{-\alpha + \frac{\gamma d}{\beta r}} \|v(s, \cdot)\|_\rho ds. \end{aligned}$$

Finally, Hölder's inequality applied with  $\lambda$  and  $\mu = \frac{\lambda}{\lambda-1}$  yields that

$$\|J(v)(t_0, t, \cdot)\|_q \leq C \left( \int_{t_0}^t (t-s)^{\mu(-\alpha + \frac{\gamma d}{\beta r})} ds \right)^{\frac{1}{\mu}} \left( \int_{t_0}^t \|v(s, \cdot)\|_\rho^\lambda ds \right)^{\frac{1}{\lambda}}.$$

Hence for  $t_0 = 0$ , the right hand side is finite and bounded with respect to  $t \in [0, T]$  if and only if  $\mu(-\alpha + \frac{\gamma d}{\beta r}) > -1$ ; this completes the proof.  $\square$

The following result proves that the evolution equation (3.3) has a unique solution with moments of all finite order. However, in order to prove that, when  $u_0 \in L^q(Q)$  for  $2 \leq q < +\infty$ , the  $\|\cdot\|_q$ -norm of the solution has bounded  $L^p$  moments for  $q < p < +\infty$ , we have to reinforce condition (C3) as follows (clearly when  $p = q$ , the conditions (C3) and (C'3)(q,p) coincide, while if  $p < q$ , (C'3)(q,p) implies (C3)):

**(C'3)(q,p)** Let  $f$  define the covariance of the Gaussian noise according to (2.1),  $2 \leq q \leq p < +\infty$ ; the constants  $\alpha, \beta, \gamma$  in (C2) satisfy one of the following conditions:

$$\int_{B_d(0,1)} f(v) \ln(|v|^{-1}) dv < \infty, \quad \text{if } \frac{\beta}{\gamma}(2\alpha - 1) = \frac{q}{p}d, \quad (3.8)$$

or

$$\int_{B_d(0,1)} f(v) |v|^{-\left[\frac{\beta}{\gamma}(2\alpha-1)-d\frac{q}{p}\right]^+} dv < \infty, \quad \text{if } \frac{\beta}{\gamma}(2\alpha - 1) \neq \frac{q}{p}d. \quad (3.9)$$

**Theorem 3.2** . Suppose that the functions  $G$  and  $H_i$ ,  $1 \leq i \leq N$  satisfy the conditions (C2) and (C4) and that the functions  $\sigma$  and  $b_i$ ,  $1 \leq i \leq N$  satisfy the assumptions (L1) and (L2).

(i) Let  $u_0 \in L^\infty(Q)$ , and let  $F$  denote either the space-time white noise if  $\alpha < 1$ , or a Gaussian process with covariance defined by (2.1) such that (C3) holds. Then the evolution equation (3.3) has a unique solution  $u \in L^\infty([0, T], L^\infty(Q))$ , such that for any  $p \in [1, \infty[$ ,

$$\sup_{(t,x) \in [0,T] \times Q} E(|u(t,x)|^p) < +\infty. \quad (3.10)$$

(ii) Let  $u_0 \in L^q(Q)$  for  $2 \leq q < +\infty$ , let  $p \in [q, +\infty[$ . Suppose that the following assumptions holds:

(a)  $F$  is the space-time white noise,  $\alpha < 1$  and  $p < \frac{2\alpha}{\left[\frac{2\alpha}{q}-1+\alpha\right]^+}$ .

(b)  $F$  is a Gaussian process with covariance defined by (2.1) such that (C'3)(q,p) holds.

Then the evolution equation (3.3) has a unique solution  $u \in L^\infty([0, T], L^q(Q))$ , such that

$$\sup_{t \in [0,T]} E(\|u(t, \cdot)\|_q^p) < +\infty. \quad (3.11)$$

*Proof:* In the case of the space-time white noise, the proof which is easier and more classical is omitted, except that of (3.11) in case (ii). Unless specified otherwise, we assume that  $F$  is Gaussian with a space-correlation function  $f$ . We use the following Picard iteration scheme;  $u_0(t, x) = G_t u_0(x) = \int_Q G(t, x; 0, y) u_0(y) dy$  and for  $n > 0$  let

$$\begin{aligned} u_{n+1}(t, x) &= u_0(t, x) + \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u_n(s, y)) F(ds, dy) \\ &\quad + \sum_{i=1}^N \int_0^t \int_Q H_i(t, x; s, y) b_i(s, y, u_n(s, y)) dy ds. \end{aligned} \quad (3.12)$$

Case (ii) Let  $2 \leq q \leq p < +\infty$  and suppose that condition (b) holds; set

$$M_n(t) = E\left(\|u_n(t, \cdot)\|_q^p\right),$$

and let  $\psi_p$  be the function defined by  $\psi_p(t) = t^{\alpha(-2+\frac{q}{p})} \int_{B_d(0,R)} \exp\left(-c\frac{|v|^\beta}{t^\gamma}\right) f(v) dv$ . Using (C'3)(q,p), computations similar to that proving (2.6) from (2.12) using (2.13) show that  $\psi_p$  is integrable; set  $I_p = \int_0^T \psi_p(s) ds < \infty$  and let

$$\varphi_p(s) = C_p \left( \psi_p(s) + \sum_{i=1}^N s^{-\alpha_i + \alpha} \right);$$

the assumptions (C'3)(q,p), (C4), (2.3) and the proof of lemma 2.1 imply that  $\varphi_p \in L_+^1([0, T])$ . Let  $\psi$  and  $I$  be defined as in the proof of lemma 2.1; then for  $q \leq p$ ,  $\psi_p \geq \psi_q = \psi$  and  $I_q \geq I_p = I$ . We prove that

$$\sup_{t \in [0, T]} M_0(t) < +\infty, \quad (3.13)$$

and for any  $n \geq 0$ ,  $t \in [0, T]$ ,

$$M_{n+1}(t) \leq \int_0^t \varphi_p(t-s) [1 + M_n(s)] ds; \quad (3.14)$$

then Lemma 15 in [6] shows that

$$\sup_n \sup_{0 \leq t \leq T} E(\|u_n(t, \cdot)\|_q^p) < +\infty. \quad (3.15)$$

Fubini's theorem, (2.2), Hölder's inequality and (2.3) imply that

$$\begin{aligned} \sup_{0 \leq t \leq T} M_0(t) &\leq C \sup_{0 \leq t \leq T} \left[ \int_Q \left| \int_Q u_0(y) t^{-\alpha} \exp\left(-c \frac{|x-y|^\beta}{t^\gamma}\right) dy \right|^q dx \right]^{\frac{p}{q}} \\ &\leq C \left[ \int_Q |u_0(y)|^q \left( t^{-\alpha} \int_Q \exp\left(-\bar{c} \frac{|x-y|^\beta}{t^\gamma}\right) dx \right) dy \right]^{\frac{p}{q}} \leq C \|u_0\|_q^p. \end{aligned}$$

We now prove (3.14); for  $n \geq 0$ ,  $M_{n+1}(t) \leq C_p [M_0(t) + T_n^1(t, p) + \sum_{i=1}^N T_n^{2,i}(t, p)]$ , where for  $q \leq p < +\infty$ ,

$$\begin{aligned} T_n^1(t, p) &= E \left( \left\| \int_0^t \int_Q G(t, \cdot; s, y) \sigma(s, y, u_n(s, y)) F(ds, dy) \right\|_q^p \right), \\ T_n^{2,i}(t, p) &= E \left( \left\| \int_0^t \int_Q H_i(t, \cdot; s, y) b_i(s, y, u_n(s, y)) dy ds \right\|_q^p \right). \end{aligned}$$

Since  $\beta \geq 1$ ,  $|y-z|^\beta \leq 2^{\beta-1} [|x-y|^\beta + |x-z|^\beta]$ . Therefore, Fubini's theorem, Burkholder's and Hölder's inequalities yield the existence of a constant  $\bar{c}$  such that

$$\begin{aligned} T_n^1(t, p) &\leq C_p \int_Q dx E \left( \left\| \int_0^t ds (t-s)^{-2\alpha} \int_Q dy \exp\left(-\bar{c} \frac{|x-y|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, y)|) \right. \right. \\ &\quad \left. \left. \times \int_Q f(y-z) \exp\left(-\bar{c} \frac{|x-y|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, z)|) dz \right\|_{\frac{p}{2}} \right) \\ &\leq C_p I_q^{\frac{p}{2} - \frac{p}{q}} \int_Q dx E \left( \left[ \int_0^t (t-s)^{-2\alpha} \int_Q dy \exp\left(-\bar{c} \frac{|x-y|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, y)|^{\frac{q}{2}}) \right. \right. \\ &\quad \left. \left. \times \int_Q f(y-z) \exp\left(-\bar{c} \frac{|y-z|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, z)|^{\frac{q}{2}}) dz ds \right]^{\frac{p}{q}} \right). \end{aligned}$$

Fubini's theorem and Jensen's inequality imply that

$$\begin{aligned} T_n^1(t, p) &\leq C_p I_q^{\frac{p}{2} - \frac{p}{q}} E \left( \left[ \int_0^t (t-s)^{-2\alpha} \left\| \int_Q dy \exp\left(-\bar{c} \frac{|\cdot - y|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, y)|^{\frac{q}{2}}) \right. \right. \right. \\ &\quad \left. \left. \times \int_Q f(y-z) \exp\left(-\bar{c} \frac{|y-z|^\beta}{(t-s)^\gamma}\right) (1 + |u_n(s, z)|^{\frac{q}{2}}) dz \right\|_{\frac{p}{q}} ds \right]^{\frac{p}{q}} \right). \end{aligned}$$



Then Young's, Schwarz's inequalities, (2.3), Hölder's inequality with respect to  $\psi_p(t-s) ds$  and  $I_q \leq I_p$  yield

$$\begin{aligned}
T_n^1(t, p) &\leq C_p I_q^{\frac{p}{2}-\frac{p}{q}} E \left( \left| \int_0^t (t-s)^{-2\alpha} \left\| \exp \left( -\bar{c} \frac{|\cdot|^\beta}{(t-s)^\gamma} \right) * \left[ \left( 1 + |u_n(s, \cdot)|^{\frac{q}{2}} \right) \right. \right. \right. \right. \\
&\quad \times \left. \left. \left. \left. \left( f(\cdot) \exp \left( -\bar{c} \frac{|\cdot|^\beta}{(t-s)^\gamma} \right) \right) * \left( 1 + |u_n(s, \cdot)|^{\frac{q}{2}} \right) \right] \right\|_{L^{\frac{p}{q}}(Q, dx)} ds \right|^{\frac{p}{q}} \right) \\
&\leq C_p I_q^{\frac{p}{2}-\frac{p}{q}} E \left( \left| \int_0^t (t-s)^{-2\alpha} \left\| \exp \left( -\bar{c} \frac{|\cdot|^\beta}{(t-s)^\gamma} \right) \right\|_{L^{\frac{p}{q}}(Q, dx)} \left( 1 + \|u_n(s, \cdot)\|_{\frac{q}{2}}^{\frac{q}{2}} \right) \right. \right. \\
&\quad \times \left. \left. \left\| f(\cdot) \exp \left( -\bar{c} \frac{|\cdot|^\beta}{(t-s)^\gamma} \right) \right\|_{L^1(Q, dx)} \left( 1 + \|u_n(s, \cdot)\|_{\frac{q}{2}}^{\frac{q}{2}} \right) ds \right|^{\frac{p}{q}} \right) \\
&\leq C_p I_q^{\frac{p}{2}-\frac{p}{q}} E \left( \left| \int_0^t (t-s)^{\alpha(-2+\frac{q}{p})} [1 + (\|u_n(s, \cdot)\|_q^q)] \right. \right. \\
&\quad \times \left. \left. \int_Q f(v) \exp \left( -\bar{c} \frac{|v|^\beta}{(t-s)^\gamma} \right) dv ds \right|^{\frac{p}{q}} \right) \\
&\leq C_q I_p^{\frac{p}{2}-1} \int_0^t \psi_p(t-s) [1 + M_n(s)] ds. \tag{3.16}
\end{aligned}$$

For every  $1 \leq i \leq N$ , using (C4), (L1), (3.7) with  $\rho = q$ ,  $r = 1$ , and Hölder's inequality (since  $\alpha_i < \alpha + 1$ ) and Fubini's theorem, we deduce that for  $q \leq p < +\infty$ ,

$$\begin{aligned}
T_{n,i}^2(t, p) &\leq C E \left( \left| \int_0^t (t-s)^{-\alpha_i+\alpha} (1 + \|u_n(s, \cdot)\|_q) ds \right|^p \right) \\
&\leq C \int_0^t (t-s)^{-\alpha_i+\alpha} [1 + E(\|u_n(s, \cdot)\|_q^p)] ds.
\end{aligned}$$

This concludes the proof of (3.14). Let  $\Delta_n(t) = E \left( \|u_{n+1}(t, \cdot) - u_n(t, \cdot)\|_q^p \right)$ ; a similar computation using the global Lipschitz property (L2) of the coefficients with respect to the last variable shows that

$$\Delta_{n+1}(t) \leq C_p \int_0^t \varphi_p(t-s) \Delta_n(s) ds \tag{3.17}$$

where the function  $\varphi_p$  is the previous one. Using again Lemma 15 in [6], we conclude that  $\sum_{n \geq 0} \Delta_n(t)$  converges uniformly on  $[0, T]$ . Therefore, usual arguments show that the solution  $u$  to (3.3) exists in  $L^\infty([0, T], L^q(Q))$  and satisfies (3.11).

We now suppose that condition (a) holds. Set  $M_n(t) = E(\|u_n(t, \cdot)\|_q^p)$ . According to the results proved above, it suffices to check that (using the previous notations),  $T_n^1(t, p) \leq C_p \int_0^t (t-s)^{-a} (1 + M_n(s, p)) ds$  for some  $a < 1$ . Using Hölder's and Burkholder's inequalities, Fubini's theorem, then (3.7) with  $\frac{p}{2}$  instead of  $q$ ,  $2\alpha$  instead of  $\alpha$  and  $\rho = \frac{q}{2}$ , we deduce that for  $1 + \frac{2}{p} = \frac{2}{q} + \frac{1}{r}$ , and  $a = (2 - \frac{1}{r})\alpha < 1$  by the choice of  $p$ , we have

$$\begin{aligned}
T_n^1(t, p) &\leq C_p E \left( \left| \int_Q \left| \int_0^t \int_Q (t-s)^{-2\alpha} \exp \left( -c \frac{|x-y|^\beta}{(t-s)^\gamma} \right) (1 + |u_n(s, y)|^2) dy ds \right|^{\frac{p}{2}} dx \right. \right. \\
&\quad \left. \left. \leq \int_0^t (t-s)^{-a} (1 + E(\|u_n(s, \cdot)\|_q^p)) ds. \right. \right)
\end{aligned}$$

The rest of the proof, similar to that of the case (ii)(b), is omitted.

*Case (i)* Let  $u_0 \in L^\infty(Q)$ ,  $p \in [1, +\infty[$  and suppose that the covariance function  $f$  satisfies (C3); set

$$M_n(t) = \sup_{x \in Q} E(|u_n(t, x)|^{2p}).$$

We again prove (3.13) and (3.14). Since  $u_0 \in L^\infty(Q)$ , the inequality (2.3) proves (3.13). Let  $\psi$  be defined by (2.13) and let  $\varphi(t) = \psi(t) + \sum_{i=1}^N t^{-\alpha_i + \alpha} \in L^1([0, T])$ ; then Burkholder's, Hölder's inequalities and (L1) yield

$$\begin{aligned} M_{n+1}(t, x) &\leq C_p \left[ M_0(t) + E \left( \left| \int_0^t ds \int_Q dy \int_Q |G(t, x; s, y)| |\sigma(s, y, u_n(s, y))| f(y - z) \right. \right. \right. \\ &\quad \left. \left. \left. \times |G(t, x; s, z)| |\sigma(s, y, u_n(s, z))| dz \right|^p \right) \right. \\ &\quad \left. + \int_0^t \int_Q \sum_{i=1}^N |H_i(t, x; s, y)| (1 + M_n(s)) dy ds \right] \\ &\leq C_p \left( M_0(t) + \int_0^t \varphi(t - s) [1 + M_n(s)] ds \right). \end{aligned}$$

This implies (3.14) and again Lemma 15 in [6] shows that

$$\sup_n \sup_{(t, x) \in [0, T] \times Q} E(|u_n(t, x)|^{2p}) < +\infty. \quad (3.18)$$

A similar computation for  $\Delta_n(t) = \sup_{x \in Q} E(|u_{n+1}(t, x) - u_n(t, x)|^{2p})$  and the global Lipschitz property (L2) of the coefficients with respect to the last variable show that (3.17) holds. As in case 2, usual arguments prove that the solution  $u$  to (3.3) exists and satisfies (3.10).  $\square$

### 3.2 Cahn-Hilliard equation in dimension $d = 4, 5$

The following stochastic Cahn-Hilliard equation has been studied in dimension 1 up to 3 by C. Cardon-Weber [4] and [5]; see also G. Da Prato and A. Debussche [8]. Let  $Q = [0, \pi]^d$  or a compact convex subset of  $\mathbb{R}^d$ ,  $(t, x) \in [0, T] \times Q$  and multi-indices  $(k_i, 1 \leq i \leq N)$  with  $|k_i| \leq 3$  which satisfy the conditions in Remark 1.1, the following equation is defined in a weak sense:

$$\begin{aligned} \frac{\partial u}{\partial t}(t, x) + (\Delta^2 u(t, x) - \Delta R(u)(t, x)) &= \sigma(t, x, u(t, x)) \dot{F} + g(t, x, u(t, x)) \\ &\quad + \sum_{i=1}^N D_x^{k_i} (b_i(t, x, u(t, x))), \end{aligned} \quad (3.19)$$

with the initial condition  $u(0, \cdot) = u_0$  and the homogeneous Neumann boundary conditions:

$$\frac{\partial u}{\partial n} = \frac{\partial \Delta u}{\partial n} = 0 \text{ on } \partial Q. \quad (3.20)$$

We will also consider the homogeneous Dirichlet boundary conditions:

$$u = \Delta u = 0 \text{ on } \partial Q. \quad (3.21)$$

In this section, we will suppose that  $d = 4, 5$  and make the following assumptions:

**(H.1)**  $R$  is a polynomial of degree 3 with positive dominant coefficient, and such that  $R(0) = 0$  if (3.21) holds.

**(H.2)**  $\sigma : [0, T] \times Q \times \mathbb{R} \mapsto \mathbb{R}$  is bounded, and the functions  $\sigma$  and  $b_i : [0, T] \times Q \times \mathbb{R} \mapsto \mathbb{R}$  are globally Lipschitz with respect to the last variable, while the function  $g : [0, T] \times Q \times \mathbb{R} \mapsto \mathbb{R}$  has quadratic growth with respect to the last variable uniformly with respect to the first ones, and satisfies for  $y, z \in Q$ :

$$\sup_{(t,x) \in [0,T] \times Q} |g(t, x, y) - g(t, x, z)| \leq C(1 + |y| + |z|) |y - z|. \quad (3.22)$$

**(H.3)**  $u_0$  belongs to  $L^q(Q)$  for some  $q > d$ .

**(H.4)** Set  $H_i(t, x; s, y) = D_x^{k_i} G(t, x; s, y)$  for  $|k_i| \leq 3$ , where  $G$  is the Green function associated with the operator  $\frac{\partial}{\partial t} + \Delta^2$  on  $Q$  with the homogeneous boundary conditions (3.20) or (3.21), and the multi-indices  $k_i$  and the functions  $b_i$  satisfy the assumptions in Example 1.1.

The upper estimates of  $G$  stated in condition (C2) are given in the introduction:  $\alpha = \frac{d}{4}$ ,  $\beta = \frac{4}{3}$ ,  $\gamma = \frac{1}{3}$ ,  $\delta = \frac{1}{4}$  and  $\eta = 1$ . Clearly for  $|k_i| \leq 3$ ,  $\alpha_i \leq \alpha + \frac{3}{4}$  and (C4) holds. As explained in the introduction, the Green formula shows that the weak formulation of (3.19) is equivalent with the evolution formulation: for  $x \in Q, t \in [0, T]$ :

$$\begin{aligned} u(t, x) &= \int_Q G(t, x; 0, y) u_0(y) dy + \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u(s, y)) F(ds, dy) \\ &+ \int_0^t \int_Q \Delta G(t, x; s, y) R(u(s, y)) dy ds \\ &+ \int_0^t \int_Q \left[ \sum_{i=1}^N H_i(t, x; s, y) b_i(s, y, u(s, y)) + G(t, x; s, y) g(s, y, u(s, y)) \right] dy ds. \end{aligned} \quad (3.23)$$

The following theorem completes the existence and uniqueness of the solution to (3.19) in dimension 4 and 5.

**Theorem 3.3** . Let  $Q$  denote either  $[0, \pi]^d$  or a compact subset of  $\mathbb{R}^d$  with boundary of class  $\mathcal{C}^{4+\lambda}$  for  $\lambda > 0$ ,  $d = 4, 5$  and assume that (H.1)- (H.4) hold. Let  $f$  be the covariance function of the Gaussian noise  $F$  defined by (2.1), which satisfies (C1) and such that for  $\varepsilon \in ]0, 1[$ ,

$$\int_{B_d(0,1)} f(v) |v|^{-d(1+\varepsilon)+4} dv < \infty. \quad (3.24)$$

There exists a unique adapted process  $u$  in  $L^\infty([0, T], L^q(Q))$  that satisfies equation (3.23).

**Remark 3.4** Under assumptions (H.1), (H.3) and (H.4), the existence result proved in [4] in dimension 1-3 (respectively Theorem 3.3) extends to a compact  $Q$  with boundary of class  $\mathcal{C}^{4+\lambda}$  for  $\lambda > 0$ , for the differential operator  $a(t)\Delta^2$  where the function  $a$  is such that  $\sup_{0 \leq t \leq T} a(t) < 0$ , and when  $F$  is the space-time white noise (respectively when (H.2) holds).

*Proof:* To prove this theorem we at first prove the existence of a solution when the coefficients are truncated. Let  $K_n : \mathbb{R}^+ \rightarrow \mathbb{R}$  be a  $\mathcal{C}^1$  function such that

$$K_n(x) = 1 \text{ if } x < n, \quad K_n(x) = 0 \text{ if } x \leq n + 1, \quad |K_n| \leq 1 \text{ and } |K_n'| \leq 2. \quad (3.25)$$

We denote by  $u_n$  the solution to the following evolution equation with truncated coefficients:

$$u_n(t, x) = \int_Q G(t, x; 0, y) u_0(y) dy + \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u_n(s, y)) F(ds, dy) \quad (3.26)$$

$$\begin{aligned}
& + \int_0^t \int_Q K_n(\|u_n(s, \cdot)\|_q) [\Delta G(t, x; s, y) R(u_n(s, y)) + G(t, x; s, y) g(s, y, u_n(s, y))] dy ds \\
& + \sum_{i=1}^N \int_0^t \int_Q H_i(t, x; s, y) b_i(s, y, u_n(s, y)) dy ds.
\end{aligned}$$

Given an adapted process  $u$ , let  $L(u)$  be defined by

$$L(u)(t, x) = \int_0^t \int_Q G(t, x; s, y) \sigma(s, y, u(s, y)) F(ds, dy).$$

The arguments used in the proof of Theorem 3.2 show that  $u_n$  exists, is unique and that for any  $p \in [q, \frac{q}{1-\varepsilon}]$ ,

$$\sup_{t \in [0, T]} E(\|u_n(t, \cdot)\|_q^p) < +\infty.$$

Indeed, let  $\mathcal{K}$  denote the set of adapted  $L^q(Q)$ -valued processes such that for  $q \leq p \leq \frac{q}{1-\varepsilon}$ ,  $\|u\|_{\mathcal{K}}^p = \sup_{0 \leq t \leq T} E(\|u(t, \cdot)\|_q^p) < +\infty$ . For any  $u \in \mathcal{K}$ ,  $(t, x) \in [0, T] \times Q$ , set

$$\begin{aligned}
H_n(u)(t, x) &= \int_0^t \int_Q \Delta G(t, x; s, y) K_n(\|u(s, \cdot)\|_q) R(u(s, y)) dy ds, \\
J_n(u)(t, x) &= \int_0^t \int_Q G(t, x; s, y) K_n(\|u(s, \cdot)\|_q) g(s, y, (u(s, y))) dy ds, \\
B(u)(t, x) &= \sum_{i=1}^N \int_0^t \int_Q H_i(t, x; s, y) b_i(s, y, u(s, y)) dy ds.
\end{aligned}$$

Since  $\sigma$  is bounded and (3.24) implies (2.5), Burkholder's inequality and Lemma 2.1 yield that for any adapted process  $u$  and  $2 \leq p < +\infty$ ,  $\sup\{E(\|L(u)(t, x)\|^p) : (t, x) \in [0, T] \times Q\} < +\infty$ , so that  $\|L(u)\|_{\mathcal{K}} < \infty$ . Furthermore, given  $u, v \in \mathcal{K}$ , (H.2), the fact that (3.24) implies (C'3)(q,p) for  $q \leq p \leq \frac{q}{1-\varepsilon}$  and the argument used to show (3.16) in the proof of Theorem 3.2 yield

$$E(\|L(u)(s, \cdot) - L(v)(s, \cdot)\|_q^p) \leq C \left( \int_0^T \psi_p(s) ds \right)^{\frac{p}{2}} \sup_{0 \leq s \leq T} E(\|u(s, \cdot) - v(s, \cdot)\|_q^p).$$

Since  $\psi_p$  is integrable,  $L$  is a contraction of  $\mathcal{K}$  for small enough  $T$ . For the polynomial term  $H_n$ , we just need to notice that if  $u$  and  $v$  belong to  $L^\infty([0, T], L^q(Q))$

$$\left\| K_n(\|u(s, \cdot)\|_q) R(u(s, \cdot)) - K_n(\|v(s, \cdot)\|_q) R(v(s, \cdot)) \right\|_{\frac{q}{3}} \leq C_n \|u(s, \cdot) - v(s, \cdot)\|_q. \quad (3.27)$$

Using (3.7), (3.27) and Hölder's inequality, we obtain that for  $d < q \leq p < +\infty$ ,

$$\begin{aligned}
\sup_{0 \leq t \leq T} \|H_n(u)(t, \cdot) - H_n(v)(t, \cdot)\|_{\mathcal{K}}^p &\leq C_n T^{(p-1)\left(\frac{1}{2} - \frac{d}{2q}\right)} \int_0^T (t-s)^{-\frac{d+2}{4} + \frac{d}{4}\left(1 - \frac{2}{q}\right)} \\
&\quad \times E(\|u(s, \cdot) - v(s, \cdot)\|_q^p) ds \\
&\leq C_n T^{p\left(\frac{1}{2} - \frac{d}{2q}\right)} \|u - v\|_{\mathcal{K}}^p.
\end{aligned}$$

A similar computation based on the quadratic growth and increments property of  $g$  with respect to the third variable shows that for  $\frac{d}{4} < q \leq p < +\infty$ ,

$$\sup_{0 \leq t \leq T} \|J_n(u)(t, \cdot) - J_n(v)(t, \cdot)\|_{\mathcal{K}} \leq C_n T^{1 - \frac{d}{4q}} \|u - v\|_{\mathcal{K}}.$$

Finally, the estimation of  $T_n^2$  made in the proof of Theorem 3.2 shows that

$$\sup_{0 \leq t \leq T} \|B(u)(t, \cdot) - B(v)(t, \cdot)\|_{\mathcal{K}} \leq C \sup_{1 \leq i \leq N} T^{1+\alpha-\alpha_i} \|u - v\|_{\mathcal{K}}.$$

Hence, there exists  $T_0 > 0$ , independent of the initial condition  $u_0$ , such that for  $0 < T \leq T_0$ ,  $L + H_n + J_n + B$  is a contraction of  $\mathcal{K}$ , and hence admits a unique fixed point such that  $u(0, \cdot) = u_0$ . A concatenation argument implies that (3.26) has a unique solution on  $[0, T]$  for an arbitrary terminal time  $T$ .

To prove the existence and uniqueness of  $u$  we follow the proof in C. Cardon-Weber [4]. Let  $\tau_n$  be the stopping time defined by:  $\tau_n = \inf\{t \geq 0, \|u_n(t, \cdot)\|_q \geq n\}$ . By uniqueness of the solution to (3.26), the local property of the stochastic integrals yields for  $m > n$ ,  $u_m(t, \cdot) = u_n(t, \cdot)$  if  $t \leq \tau_n$ , so that we can define a process  $u$  by setting  $u(t, \cdot) = u_n(t, \cdot)$  on  $t \leq \tau_n$ . Set  $\tau_\infty = \lim_n \tau_n$ . Then  $u$  is the unique solution of (3.19) on the interval  $[0, \tau_\infty)$ . We just need to prove that  $\tau_\infty = +\infty$  a.s.

Set  $v_n = u_n - L(u_n)$ ; then for every  $T > 0$ ,  $v_n$  is the weak solution on  $[0, T]$  to the SPDE (with the same boundary conditions as (3.19)):

$$\left\{ \begin{array}{l} \frac{\partial v_n}{\partial t}(t, x) + \Delta^2 v_n(t, x) - \Delta \left[ K_n(\|v_n(t, \cdot) + L(u_n)(t, \cdot)\|_q) R(v_n(t, x) + L(u_n)(t, x)) \right] = \\ \quad K_n(\|v_n(t, \cdot) + L(u_n)(t, \cdot)\|_q) g(t, x, v_n(t, x) + L(u_n)(t, x)) \\ \quad + \sum_{i=1}^N D_x^{k_i} b_i(t, x, v_n(t, x) + L(u_n)(t, x)), \\ v_n(0, \cdot) = u_0(\cdot), \\ \frac{\partial v_n}{\partial n} = \frac{\partial \Delta v_n}{\partial n} = 0 \text{ (resp. } v_n = \Delta v_n = 0) \text{ on } \partial Q. \end{array} \right. \quad (3.28)$$

Since  $\sigma$  is bounded, the Garsia-Rodemich-Ramsay Lemma (cf. eg. [14]), (3.24) and lemma 2.3 yield that for any  $p \in [2, +\infty[$ ,

$$\sup_n E(\|L(u_n)\|_\infty^p) < \infty. \quad (3.29)$$

Since  $u_0$  belongs to  $L^q(Q)$ ,

$$\sup_{t \leq T} \|G_t u_0\|_q \leq \|u_0\|_q. \quad (3.30)$$

We need to prove a uniform upper estimate for the drift terms  $H_n(u_n)$  and  $J_n(u_n)$  (the estimation of the other drift term  $B(u_n)$ , which is easier, will be omitted and to lighten the notations, we will assume that  $b_i = 0$ ,  $1 \leq i \leq N$ ). Since the function  $\Delta G$  has a regularizing effect, we first show that  $u_n$  belongs to the sets  $L^a([0, T], L^q(Q))$  for some well-chosen  $a$ .

Let us introduce some notations: set  $\mathcal{A} = -\Delta$ , let  $\langle \cdot, \cdot \rangle$  denote the usual scalar product in  $L^2(Q)$ , let  $(e_n, n \in \mathbb{N}^d)$  be a basis of  $L^2(Q)$  made of eigenfunctions of  $\mathcal{A}$  (namely  $e_n(x) = \prod_{i=1}^d e_{n_i}(x_i)$  and either  $e_0 = \frac{1}{\sqrt{\pi}}$ ,  $e_n(x) = \sqrt{\frac{2}{\pi}} \cos(nx)$  for  $n > 0$  in the case of the Neumann boundary conditions (3.20), or  $e_n(x) = \sqrt{\frac{2}{\pi}} \sin(nx)$  for  $n > 0$  in the case of the Dirichlet boundary conditions (3.21) if  $Q = [0, \pi]^d$ ). The corresponding eigenvalues are  $\lambda_n^2$ , where  $\lambda_n = \sum_{i=1}^d n_i^2$ . For  $\mu \neq 0$  and  $u \in \text{Dom}(\mathcal{A}^\mu)$ , let

$$\mathcal{A}^\mu u = \sum_{k \in \mathbb{N}^{d^*}} \lambda_k^\mu \langle \varepsilon_k, u \rangle \varepsilon_k;$$

$\mathcal{A}^\mu u$  exists for every  $u$  such that  $\sum_{k \in \mathbb{N}^{d^*}} \lambda_k^{2\mu} \langle \varepsilon_k, u \rangle^2 < \infty$ . In the sequel, for a function  $u : [0, T] \times Q \rightarrow \mathbb{R}$ , we will set (if  $e_0$  is a constant eigenfunction):

$$m(u)(t) = \langle \varepsilon_0, u(t, \cdot) \rangle = \pi^{-\frac{d}{2}} \int_Q u(t, x) dx \text{ and } \tilde{u}(t, y) = u(t, y) - m(u)(t).$$

Apply  $\mathcal{A}^{-1}$  to the equation (3.33) and take its scalar product in  $L^2(Q)$  with  $\tilde{v}_n(t, \cdot)$ ; this leads to

$$\begin{aligned} & \|\mathcal{A}^{-1/2}\tilde{v}_n(t, \cdot)\|_2^2 - \|\mathcal{A}^{-1/2}\tilde{v}_n(0, \cdot)\|_q^2 + \int_0^t \|\mathcal{A}^{1/2}v_n(s, \cdot)\|_2^2 ds \\ & + \int_0^t K_n(\|v_n(s, \cdot) + L(u_n)(s, \cdot)\|_q) \int_Q \left[ R(v_n(s, x) + L(u_n)(s, x))\tilde{v}_n(s, x) \right. \\ & \quad \left. + g(s, x, v_n(s, x) + L(u_n)(s, x)) \mathcal{A}^{-1}\tilde{v}_n(s, x) \right] dx ds = 0. \end{aligned} \quad (3.31)$$

This equation is justified because  $v_n$  belongs to  $L^\infty([0, T] \times Q)$ . Using the properties of the polynomial  $R$ , computations made to obtain (2.19)-(2.21) in [4], we obtain that for some  $b > 0$ ,

$$\begin{aligned} & \|\mathcal{A}^{-1/2}\tilde{v}_n(t, \cdot)\|_2^2 + \int_0^t \|\mathcal{A}^{1/2}v_n(s, \cdot)\|_2^2 ds \\ & + \frac{b}{4} \int_0^t K_n(\|v_n(s, \cdot) + L(u_n)(s, \cdot)\|_q) \|v_n(s, \cdot) + L(u_n)(s, \cdot)\|_4^4 ds \\ & \leq \int_0^t C(1 + m(u_0)(s))^4 + \|L(v_n)(s, \cdot)\|_4^4 ds + \|\mathcal{A}^{-1/2}\tilde{u}_0(\cdot)\|_2^2. \end{aligned} \quad (3.32)$$

Let us find a second ‘‘a priori’’ estimate. Denote by  $v_n^m$  the Galerkin approximation of  $v_n$  and let  $P_m$  be the orthogonal projector on  $\text{Span}\{e_0, \dots, e_m\}$ . For every  $\omega$ ,  $v_n^m$  is the ‘‘strong’’ solution to the following PDE:

$$\begin{cases} \frac{\partial v_n^m}{\partial t}(t, x) + \Delta^2 v_n^m(t, x) \\ \quad - \Delta \left[ K_n(\|v_n^m(t, \cdot) + L(u_n)(t, \cdot)\|_q) P_m(R(v_n^m(t, x) + L(u_n)(t, x))) \right] \\ \quad - K_n(\|v_n^m(t, \cdot) + L(u_n)(t, \cdot)\|_q) P_m(g(t, x, v_n^m(t, x) + L(u_n)(t, x))) = 0. \end{cases} \quad (3.33)$$

The solution  $v_n^m$  to (3.33) is unique on some random time interval  $[0, t_n^m[$  and we prove that  $t_n^m = +\infty$ . The boundary conditions satisfied by  $v_n^m$  and the Green Formula yield

$$\int_Q \Delta^2 v_n^m(t, x) \times v_n^m(t, x) dx = \|\Delta v_n^m(t, x)\|_2^2.$$

We now take the scalar product in  $L^2(Q)$  of (3.33) with  $v_n^m$ ; using once more the Green formula, we obtain

$$\begin{aligned} & \frac{1}{2} \frac{\partial}{\partial t} \|v_n^m(t, \cdot)\|_2^2 + \int_Q \Delta^2 v_n^m(t, x) \times v_n^m(t, x) dx = K_n(\|v_n^m(t, \cdot) + L(u_n)(t, \cdot)\|_q) \\ & \times \int_Q \left[ R(v_n^m(t, x) + L(u_n)(t, x)) \Delta v_n^m(t, x) + g(t, x, v_n^m(t, x) + L(u_n)(t, x)) v_n^m(t, x) \right] dx. \end{aligned}$$

Using the local Lipschitz property of  $R$  and  $g(t, x, \cdot)$  and the fact that  $\int_Q (v_n^m(t, x))^3 \Delta v_n^m(t, x) dx$  is negative, that the leading coefficient of  $R$  is positive and that  $\|K_n\|_\infty \leq 1$ , we obtain:

$$\begin{aligned} & \|v_n^m(t, \cdot)\|_2^2 + \int_0^t \left[ \|\Delta v_n^m(s, \cdot)\|_2^2 + m(v_n^m(s, \cdot))^2 \right] ds \leq \|u_0\|_2^2 + C_T (1 + \|L(u_n)\|_\infty^6 + m(u_0)^4) \\ & + C(1 + \|L(u_n)\|_\infty^2) \int_0^t \|v_n^m(s, \cdot)\|_4^4 K_n(\|v_n^m(s, \cdot) + L(u_n)(s, \cdot)\|_q) ds. \end{aligned} \quad (3.34)$$

The norm  $(\|\Delta \bullet\|_{L^2(Q)}^2 + m(\bullet)^2)^{\frac{1}{2}}$  is equivalent with the Sobolev norm of  $W^{2,2}(Q)$  (cf. eg. Da Prato and Debussche (1996) p. 245). The sequence  $(v_n^m)_m$  is bounded in  $L^2([0, T], W^{2,2}(Q))$ .

Thus,  $t_n^m = \infty$  and this sequence converges as  $m \rightarrow +\infty$  in the weak\* topology of  $L^2([0, T], W^{2,2}(Q))$ . Its weak limit is the weak solution to (3.28) and hence is equal to  $v_n$ . Therefore  $v_n$  belongs to  $L^2([0, T], W^{2,2}(Q))$ , and we can repeat the above computation with  $v_n$  instead of  $v_n^m$ , which yields

$$\begin{aligned} \|v_n(t, \cdot)\|_2^2 + \int_0^t \left[ \|\Delta v_n(s, \cdot)\|_2^2 + m(v_n(s, \cdot))^2 \right] ds &\leq \|u_0\|_2^2 + C_T(1 + \|L(u_n)\|_\infty^6 + m(u_0)^4) \\ &+ C(1 + \|L(u_n)\|_\infty^2) \int_0^t \|v_n(s, \cdot)\|_4^4 K_n(\|v_n(s, \cdot) + L(u_n)(s, \cdot)\|_q) ds. \end{aligned}$$

Thus, (3.32) and Schwarz's inequality imply that

$$\begin{aligned} \|v_n(t, \cdot)\|_2^2 + \int_0^t \left[ \|\Delta v_n(s, \cdot)\|_2^2 + m(v_n(s, \cdot))^2 \right] ds &\leq \|u_0\|_2^2 \\ &+ C_T(1 + \|L(u_n)\|_\infty^6) + C_T(1 + \|L(u_n)\|_\infty^2) \left[ \|\mathcal{A}^{-\frac{1}{2}} u_0\|_2^2 + m(u_0)^4 \right]. \end{aligned}$$

Inequality (3.29) yields that for  $\beta \in ]1, +\infty[$ ,

$$\sup_n E\left(\sup_{t \in [0, T]} \|v_n(t, \cdot)\|_2^{2\beta}\right) < \infty, \quad (3.35)$$

$$\sup_n E\left(\left[\int_0^T \left\{ \|\Delta v_n(t, \cdot)\|_2^2 + m(v_n(t, \cdot))^2 \right\} dt\right]^\beta\right) < \infty. \quad (3.36)$$

Moreover, by Sobolev's embedding theorem (Adams 1975, Corollary 5.16) there exists  $C > 0$  such that for  $d \geq 4$  and  $2 \leq r \leq \frac{2d}{d-4}$ , if  $u \in W^{2,2}(Q)$ ,  $\|u\|_{L^r(Q)} \leq C\|u\|_{W^{2,2}(Q)}$ . Thus, (3.36) becomes for  $2 \leq r \leq \frac{2d}{d-4}$ ,  $\beta \in [1, +\infty[$ ,

$$\sup_n E\left(\left[\int_0^T \|v_n(t, \cdot)\|_r^2 dt\right]^\beta\right) < \infty. \quad (3.37)$$

The inequalities (3.29), (3.35) and (3.37) imply for  $2 \leq r < \frac{2d}{d-4}$ ,  $\beta \in [1, +\infty[$ ,

$$\sup_n E\left(\sup_{t \in [0, T]} \|u_n(t, \cdot)\|_2^{2\beta}\right) < \infty, \quad (3.38)$$

$$\sup_n E\left(\left[\int_0^T \|u_n(t, \cdot)\|_r^2 dt\right]^\beta\right) < \infty. \quad (3.39)$$

Let us use the interpolation method to prove that  $u_n$  belongs a.s. to  $L^a([0, T], L^{\bar{R}}(Q))$  for  $\frac{2d}{d-4} > r \geq \bar{R} \geq 2$ ,  $a \geq 1 \vee \frac{2\bar{R}}{r}$ . Set  $\bar{R} = (1 - \lambda)2 + \lambda r$  for  $\lambda \in [0, 1]$ ; Hölder's inequality implies that

$$\int_0^T \|u_n(t, \cdot)\|_{\bar{R}}^a dt \leq \int_0^T \|u_n(t, \cdot)\|_2^{\frac{2a(1-\lambda)}{\bar{R}}} \|u_n(t, \cdot)\|_r^{\frac{a\lambda}{\bar{R}}} dt.$$

Let  $\lambda = \frac{2\bar{R}}{ar}$ , we obtain

$$\int_0^T \|u_n(t, \cdot)\|_{\bar{R}}^a dt \leq \sup_{t \in [0, T]} \|u_n(t, \cdot)\|_2^{\frac{2}{\bar{R}}a(1-\lambda)} \times \int_0^T \|u_n(t, \cdot)\|_r^2 dt;$$

(3.38) and (3.39) imply that for  $\bar{R} \in [2, \frac{2d}{d-4}[$  and  $a \geq 2$ ,

$$\sup_n E\left(\left[\int_0^T \|u_n(t, \cdot)\|_{\bar{R}}^a dt\right]^\beta\right) < \infty. \quad (3.40)$$

Using lemma 3.1 with  $\rho = \frac{\bar{R}}{3}$ , so that  $\frac{1}{r'} = 1 + \frac{1}{q} - \frac{3}{\bar{R}}$ , we obtain

$$\|H_n(u_n)(t, \cdot)\|_q \leq C \int_0^t (t-s)^{-\frac{d+2}{4} + \frac{d}{4r'}} (\|u_n(s, \cdot)\|_{\frac{3}{\bar{R}}} + 1) ds.$$

Let  $\gamma, \gamma' \in ]1, +\infty[$  be conjugate exponents, with  $\gamma$  close enough to one to ensure  $\gamma(-\frac{d+2}{4} + \frac{d}{4r'}) > -1$ ; this is possible (choosing  $\bar{R}$  close enough to  $\frac{2d}{d-4}$ ) if  $\frac{1}{2} + \frac{d}{4q} > \frac{3}{8}(d-4)$ , i.e.,  $(\frac{3}{8} - \frac{1}{4q})d < 2$ . Since  $q > d$ , this yields  $d \leq 5$  for any  $q \in ]d, +\infty[$ . Then Hölder's inequality implies

$$\|H_n(u_n)(t, \cdot)\|_q \leq C \left[ \int_0^t (t-s)^{(-\frac{d+2}{4} + \frac{d}{4r'})\gamma} ds \right]^{\frac{1}{\gamma}} \left[ \int_0^t (\|u_n(s, \cdot)\|_{\frac{3}{\bar{R}}} + 1)^{\gamma'} ds \right]^{\frac{1}{\gamma'}}.$$

Using (3.40), we obtain

$$\sup_n E \left( \sup_{t \in [0, T]} \|H_n(u_n)(t, \cdot)\|_q^\beta \right) < \infty; \quad (3.41)$$

A similar computation (using the quadratic growth of  $g$ ) yields for  $\rho = \frac{\bar{R}}{2}$ ,  $\frac{1}{r''} = 1 - \frac{2}{\bar{R}} + \frac{1}{q}$  and  $\gamma$  close enough to one to ensure that  $\gamma d(-\frac{1}{4} + \frac{1}{4r''}) > -1$  (i.e., for  $d < 8$  and  $\bar{R}$  close to  $\frac{2d}{d-4}$ ) yields

$$\sup_n E \left( \sup_{t \in [0, T]} \|J_n(u_n)(t, \cdot)\|_q^\beta \right) < +\infty. \quad (3.42)$$

The equations (3.29)-(3.30), (3.41) and (3.42) imply that for  $\beta \in [q, +\infty[$ ,  $d < q$ :

$$\sup_n E \left( \sup_{t \in [0, T]} \|u_n(t, \cdot)\|_q^\beta \right) < \infty.$$

We can now conclude that  $\tau_\infty = +\infty$  a.s.; indeed, for every  $T > 0$ ,

$$P(\tau_n \leq T) = P \left( \sup_{t \leq T} \|u_n(t, \cdot)\|_q \geq n \right) \leq n^{-\beta} E \left( \sup_{t \leq T} \|u_n(t, \cdot)\|_q^\beta \right),$$

so that  $\lim_{n \rightarrow \infty} P(\tau_n \leq T) = 0$ . Therefore, we can construct the solution to the SPDE (3.23) on any interval  $[0, T]$ .  $\square$

## 4 Regularity of the solution

The following lemma studies the Hölder regularity of the term involving the initial condition. There are many possible situations, depending on the boundary conditions, whether  $\int_Q G(t, x; 0, y) dy = 1$  or not, which requires two different arguments. For  $(t, x) \in ]0, T] \times Q$ , set  $G_t u_0(x) = \int_Q G(t, x; 0, y) u_0(y) dy$  and set  $G_0 u_0 = u_0$ .

**Lemma 4.1** . *Suppose that  $Q$  is convex and that  $G$  satisfies (1.2) with  $a, b \in \{0, 1\}$ .*

- 1) (i) *Let  $u_0 \in L^q(Q)$  for some  $q \in [1, +\infty[$ ; then  $G u_0 \in \mathcal{C}([0, T], L^q(Q))$ .*  
(ii) *Let  $u_0$  be bounded; then for  $0 < \lambda < 1$  and  $0 < t_0 < T$ ,  $G u_0 \in \mathcal{C}^\lambda([t_0, T] \times Q)$ .*
- 2) *Assume furthermore that  $\int_Q G(t, x; s, y) dy = 1$  for all  $(s, t, x) \in ]0, T]^2 \times Q$  with  $s < t$ .*

(i) *Let  $u_0$  be continuous; then  $G u_0 \in \mathcal{C}([0, T] \times Q)$ .*

(ii) *Let  $u_0 \in L^q(Q)$ ; then  $G u_0 \in \mathcal{C}([0, T], L^q(Q))$ .*

(iii) *Let  $u_0 \in \mathcal{C}^\lambda(Q)$  for some  $\lambda \in ]0, 1[$ ; then for  $0 \leq s < t \leq T$ ,  $\sup_{x \in Q} |G_t u_0(x) - G_s u_0(x)|$*

$$\leq C(t-s)^{\frac{\lambda}{4}}.$$



3) Let  $Q = [0, M]^d$ ,  $u_0 \in C^\lambda(Q)$  for some  $\lambda \in ]0, 1[$  and suppose that for every  $1 \leq i \leq d$ , if  $\hat{x}_i = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d)$ , there exists a function  $\phi_i : [0, T] \times \mathbb{R} \times [0, M]^{d-1} \times [0, M]^{d-1}$  such that

$$G(t, x; 0, y) = \phi_i(t, x_i + y_i, \hat{x}_i, \hat{y}_i) + \varepsilon_i \phi_i(t, x_i - y_i, \hat{x}_i, \hat{y}_i),$$

with  $\varepsilon_i \in \{-1, 1\}$ , with  $\sup_{(t,x) \in [0,T] \times Q} \int_Q |\phi_i(t, x_i + \varepsilon y_i, \hat{x}_i, \hat{y}_i)| dy < +\infty$  for  $\varepsilon \in \{-1, +1\}$ , and suppose that either one of the conditions (a) or (b) holds:

(a)  $\varepsilon_i = +1$  and  $\phi_i(t, r + 2M, \hat{x}_i, \hat{y}_i) = \phi_i(t, r, \hat{x}_i, \hat{y}_i)$  for every  $r \in \mathbb{R}$ .

(b)  $u_0(x) = 0$  for  $x \in \partial Q$ .

Then for any  $x, x' \in Q$  one has  $\sup_{t \in [0, T]} |G_t u_0(x) - G_t u_0(x')| \leq C |x - x'|^\lambda$ .

**Remark 4.2** . This proposition can be used for any convex compact subset  $Q$ ; in (1),  $G$  need not to be a semi-group. If  $Q = [0, \pi]^d$  and  $G$  is the Green function of the operator  $\frac{\partial}{\partial t} + \Delta^2 = 0$ , for  $u_0 \in C^\lambda(Q)$ ,  $0 < \lambda < 1$ , the function  $G \cdot u_0(\cdot) \in C^{\frac{\lambda}{4}, \lambda}([0, T] \times Q)$  under the homogeneous Neumann boundary conditions (3.20), while under the homogeneous Dirichlet boundary conditions (3.21), one has  $\sup_{0 \leq t \leq T} |G_t u_0(x) - G_t u_0(x')| \leq C |x - x'|^\lambda$ , and for  $0 < t_0 \leq s < t \leq T$ , one has for any  $0 < \mu < 1$ ,  $\sup_{x \in Q} |G_t u_0(x) - G_s u_0(x)| \leq C(t_0) |t - s|^\mu$ .

*Proof of Lemma 4.1:* 1) (i) Given  $0 < t_0 \leq s < t \leq T$ ,  $0 < \lambda < 1$ ,  $\|G_t u_0 - G_s u_0\|_q^q \leq C |t - s|^{\lambda q} C(M_1 + M_2)$ , where

$$M_1 = \int_Q dx \left| \int_Q \left| \int_0^1 (\theta t + (1 - \theta)s)^{-(\alpha + \eta)} \exp\left(-c \frac{|x - y|^\beta}{(\theta t + (1 - \theta)s)^\gamma}\right) d\theta \right|^\lambda t^{-\alpha(1 - \lambda)} u_0(y) dy \right|^q$$

$$M_2 = \int_Q dx \left| \int_Q \left| \int_0^1 (\theta t + (1 - \theta)s)^{-(\alpha + \eta)} d\theta \right|^\lambda s^{-\alpha(1 - \lambda)} \exp\left(-c \frac{|x - y|^\beta}{s^\gamma}\right) u_0(y) dy \right|^q.$$

Clearly,  $M_1 + M_2 \leq t_0^{-c}$  for some  $c > 0$ .

(ii) A similar argument for  $q = +\infty$  shows that for  $0 < t_0 \leq t < t' \leq T$ ,  $x, x' \in Q$ ,  $0 < \lambda < 1$  and some  $c > 0$ ,

$$|G_t u_0(x) - G_{t'} u_0(x')| \leq C t_0^{-c} \left( |t' - t|^\lambda + |x - x'|^\lambda \right)$$

2) Since  $\int_Q G(t, x; s, y) dy = 1$  for every  $s < t$ ,  $G_t u_0(x) - u_0(x) = \int_Q G(t, x; 0, y) [u_0(y) - u_0(x)] dy$  and since  $G$  is a semi-group, for  $t < t'$ ,  $G_{t'} u_0(x) - G_t u_0(x) = \int_Q G(t, x; 0, y) dy \int_Q G(t', y; t, z) [u_0(z) - u_0(y)] dz$ , so that the study of the time-regularity is completed by that at 0 .

(i) One has to check that for any  $x \in Q$ ,  $G_t u_0(x) - u_0(x)$  converges to 0 as  $t \rightarrow 0$ . The argument, based on the continuity of  $u_0$  at  $x$ , is similar to the previous one (see e.g. [4], Lemma 2.1).

(ii) Let  $u_0 \in L^q(Q)$ , let  $(u_0^n)_{n \geq 1}$  be a sequence of continuous function converging to  $u_0$  in  $L^q(Q)$ . According to (i),  $(G u_0^n)$  belong  $C([0, T] \times Q)$  and it suffices to check that  $\sup_t \|G_t u_0^n\|_q \leq C \|u_0^n\|_q$ . This follows from Hölder's inequality and (2.3).

(iii) Using the Hölder continuity of  $u_0$ , one has

$$\begin{aligned} \sup_{x \in Q} |G_t u_0(x) - u_0(x)| &\leq \sup_{x \in Q} C \int_Q t^{-\alpha} \exp\left(-c \frac{|x - y|^\beta}{t^\gamma}\right) |u_0(y) - u_0(x)| dy \\ &\leq \sup_{x \in Q} C \int_Q t^{-\alpha} \exp\left(-c \frac{|x - y|^\beta}{t^\gamma}\right) \left(\frac{|x - y|}{t^{\frac{\gamma}{\beta}}}\right)^\lambda t^{\lambda \frac{\gamma}{\beta}} dy \leq C t^{\lambda \frac{\gamma}{\beta}}. \end{aligned}$$

(3) The proof of the space regularity under condition (a), which is a straightforward extension of that of [2] Lemma A.2 and [4], Lemma 2.2, is omitted.

We suppose that (b) holds and compare the function  $Gu_0$  at points  $x = (x_1, \hat{x}_1)$  and  $x' = (x'_1, \hat{x}_1)$  with  $x_1 < x'_1$ ; increments of other components are similarly dealt with, and provide the required regularity. Obvious changes of variables yield  $G_t u_0(x) - G_t u_0(x') = \sum_{i=1}^4 D_i(t, x, x')$ , where if we set  $\tilde{Q} = [0, M]^{d-1}$ ,

$$\begin{aligned} D_1(t, x, x') &= \int_0^{M-(x'_1-x_1)} \int_{\tilde{Q}} \phi_1(t, x'_1 + y_1, \hat{x}_1, \hat{y}_1) [u_0(y) - u_0(y_1 + (x'_1 - x_1), \hat{y}_1)] d\hat{y}_1 dy_1, \\ D_2(t, x, x') &= \varepsilon_1 \int_{x'_1-x_1}^M \int_{\tilde{Q}} \phi_1(t, x'_1 - y_1, \hat{x}_1, \hat{y}_1) [u_0(y) - u_0(y_1 - (x'_1 - x_1), \hat{y}_1)] d\hat{y}_1 dy_1, \\ D_3(t, x, x') &= \int_{M-(x'_1-x_1)}^M \int_{\tilde{Q}} \phi_1(t, x'_1 + y_1, \hat{x}_1, \hat{y}_1) [u_0(y) - u_0(M, \hat{y}_1)] d\hat{y}_1 dy_1 \\ &\quad - \int_{-(x'_1-x_1)}^0 \int_{\tilde{Q}} \phi_1(t, x'_1 + y_1, \hat{x}_1, \hat{y}_1) [u_0(y_1 + (x'_1 - x_1), \hat{y}_1) - u_0(0, \hat{y}_1)] d\hat{y}_1 dy_1, \\ D_4(t, x, x') &= \varepsilon_1 \int_0^{x'_1-x_1} \int_{\tilde{Q}} \phi_1(t, x'_1 - y_1, \hat{x}_1, \hat{y}_1) [u_0(y) - u_0(0, \hat{y}_1)] d\hat{y}_1 dy_1 \\ &\quad - \varepsilon_1 \int_M^{M+(x'_1-x_1)} \int_{\tilde{Q}} \phi_1(t, x'_1 - y_1, \hat{x}_1, \hat{y}_1) [u_0(y_1 - (x'_1 - x_1), \hat{y}_1) - u_0(M, \hat{y}_1)] d\hat{y}_1 dy_1. \end{aligned}$$

The Hölder regularity of  $u_0$  and the integrability property of  $\phi_1(t, \cdot, \hat{x}_1, \cdot)$ , uniformly with respect to  $(t, \hat{x}_1)$ , conclude the proof.  $\square$

We suppose that  $u_0 \in C^a(Q)$  for some  $a \in ]0, 1[$ ; then  $u_0 \in L^q(Q)$  for any  $q > d$ , so that by Theorem 3.3, the solution  $u$  to (3.23) belongs to  $L^\infty([0, T], L^q(Q))$  for every  $q \in ]d, +\infty[$ . Remark 4.2 gives the regularity of  $Gu_0$  depending on the boundary conditions, while Lemma 2.3 gives the regularity of the stochastic integral in (3.23). Thus it suffices to study the regularity of the drift terms of (3.23) with coefficients which may have polynomial growth.

**Lemma 4.3** . *Let  $G$  be a (non-necessarily time-homogeneous) semi-group satisfying (1.2) with  $\alpha = \frac{\gamma}{\beta}d$ , let  $a$  be a multi-index such that  $|a|\delta < 1$ ,  $H(t, x; s, y) = D_x^a G(t, x; s, y)$ . Let  $b : [0, T] \times Q \times \mathbb{R} \rightarrow \mathbb{R}$  be measurable such that  $\sup\{|b(t, x, y)|, (t, x) \in [0, T] \times Q\} \leq C|y|^m$  for some  $m \geq 1$ , let  $0 < \lambda < \frac{1-|a|\delta}{\eta\sqrt{1}}$  and  $0 < \mu < \left[ (1 - |a|\delta) \left( \frac{\beta}{\gamma} \wedge \frac{1}{\delta} \right) \right] \wedge 1$ . Then if  $u : \Omega \times [0, T] \times Q \rightarrow \mathbb{R}$  is a process in  $L^\infty([0, T], L^q(Q))$  for  $q$  large enough,*

$$B(u)(t, x) = \int_0^t \int_Q H(t, x; s, y) b(s, y, u(s, y)) dy ds.$$

the map  $B(u)$  belongs to  $\mathcal{C}^{\lambda, \mu}([0, T] \times Q)$ .

*Proof:* The argument, based on the factorization method (see e.g. G. Da Prato and J. Zabczyk [10]) is similar to that in the proof of section 2.3 in [4]; it is briefly sketched.

Let  $\bar{\delta} = |a|\delta$ ,  $\varepsilon \in ]0, 1[$  and set

$$\begin{aligned} \mathcal{J}(v)(t, x) &= \int_0^t \int_Q G(t, x; s, y) (t-s)^{-\varepsilon} v(s, y) dy ds, \\ \mathcal{K}(v)(t, x) &= \int_0^t \int_Q H(t, x; s, y) (t-s)^{\varepsilon-1} b(s, y, v(s, y)) dy ds. \end{aligned}$$

The semi-group property of  $G$  implies that for every  $(t, x) \in [0, T] \times Q$ ,  $B(u)(t, x) = \frac{\sin(\varepsilon\pi)}{\pi} \times \mathcal{J}(\mathcal{K}(u))(t, x)$ . We prove that for  $\varepsilon > \bar{\delta} + \alpha \frac{m-1}{q}$ ,  $\mathcal{K}$  maps  $L^\infty([0, T], L^q(Q))$  into itself. Indeed,

it suffices to use Lemma 3.1 with  $q$  and  $\frac{q}{m}$ , so that  $\frac{1}{r} = 1 - \frac{m-1}{q}$ , and (2.3). We then prove that for  $v \in L^\infty([0, T], L^q(Q))$ , the trajectories of  $\mathcal{J}(v)$  have the required Hölder regularity. Let  $x, x' \in Q$ ; then  $|\mathcal{J}(v)(t, x) - \mathcal{J}(v)(t, x')| \leq A_1(t, x, x') + A_2(t, x, x')$ , where

$$\begin{aligned} A_1(t, x, x') &= \int_0^t \int_Q 1_{\{|y-x| \leq |x'-x|\}} (t-s)^{-\varepsilon} \left( |G(t, x; s, y)| + |G(t, x'; s, y)| \right) |v(s, y)| dy ds, \\ A_2(t, x, x') &= \int_0^t \int_Q 1_{\{|y-x| > |x'-x|\}} (t-s)^{-\varepsilon} |G(t, x; s, y) - G(t, x'; s, y)| |v(s, y)| dy ds. \end{aligned}$$

Hölder's inequality and a change of variables yield that for  $0 < \mu < \frac{q-1}{q}$ ,

$$\begin{aligned} A_1(t, x, x') &\leq C \int_0^t (t-s)^{-(\varepsilon+\alpha)} \|v(s, \cdot)\|_q \left( \int_{|z| \leq |x-x'| t^{-\frac{\gamma}{\beta}}} \exp(-c|z|^\beta) t^\alpha dz \right)^{\frac{q-1}{q}} ds \\ &\leq |x-x'|^\mu \int_0^t (t-s)^{-\varepsilon - \frac{\alpha}{q} - \frac{\gamma\mu}{\beta}} ds. \end{aligned} \quad (4.1)$$

The convergence of this last integral requires  $\mu < \frac{\beta}{\gamma} \left(1 - \varepsilon - \frac{\alpha}{q}\right)$  and  $1 - \varepsilon - \frac{\alpha}{q} > 0$ . Furthermore, if  $|x-y| > |x'-x|$ , and  $\tilde{x}$  is a convex combination of  $x$  and  $x'$ , then  $|\tilde{x}-y| \geq \frac{1}{\sqrt{2}}(|2x-x'-y| \wedge |x'-y|)$ . Therefore, Taylor's formula and Hölder's inequality imply that for  $0 < \mu < 1$ ,

$$A_2(t, x, x') \leq C |x-x'|^\mu \int_0^t (t-s)^{-\varepsilon - \mu\delta - \frac{\alpha}{q}} \|v(s, \cdot)\|_q ds, \quad (4.2)$$

and the last integral converges if  $\mu < \frac{1-\varepsilon-\frac{\alpha}{q}}{\delta}$ .

Similarly, for  $0 < t < t' \leq T$  and  $x \in Q$ ,  $|\mathcal{J}(v)(t, x) - \mathcal{J}(v)(t', x)| \leq B_1(t, t', x) + B_2(t, t', x)$ , where

$$\begin{aligned} B_1(t, t', x) &= \int_0^t \int_Q |(t'-s)^{-\varepsilon} G(t', x; s, y) - (t-s)^{-\varepsilon} G(t, x; s, y)| |v(s, y)| dy ds, \\ B_2(t, t', x) &= \int_t^{t'} \int_Q (t'-s)^{-\varepsilon} |G(t', x; s, y)| |v(s, y)| dy ds. \end{aligned}$$

Computations similar to the previous ones yield for  $\lambda \in ]0, 1[$

$$B_1(t, t', x) \leq C |t-t'|^\lambda \int_0^t (t-s)^{-\varepsilon - \eta\lambda - \frac{\alpha}{q}} \|v(s, \cdot)\|_q ds, \quad (4.3)$$

$$B_2(t, t', x) \leq C \int_t^{t'} (t'-s)^{-\varepsilon - \frac{\alpha}{q}} \|v(s, \cdot)\|_q ds \leq C |t-t'|^{1-\varepsilon - \frac{\alpha}{q}}. \quad (4.4)$$

The integral in the right hand-side of (4.3) converges if  $\lambda < \frac{1-\varepsilon-\frac{\alpha}{q}}{\eta}$ , while that in the right hand-side of (4.4) converges for  $\varepsilon < 1 - \frac{\alpha}{q}$ . Thus for  $q$  arbitrary large and  $\varepsilon$  close enough to  $\bar{\delta}$ , we see that the inequalities (4.1)-(4.4) conclude the proof.  $\square$

The following theorem summarizes the results proved in Lemmas 2.3, 4.3 and Remark 4.2.

**Theorem 4.4** . Assume (H.1), (H.2) and (H.4), let  $d = 4, 5$ ,  $u_0 \in \mathcal{C}^a(Q)$  for some  $a \in ]0, 1[$ ,  $F$  be a Gaussian process with covariance define in terms of the function  $f$  by (2.1), such that (C1) holds and  $\int_{B_d(0,1)} f(v) |v|^{-d(1+\varepsilon)+4} dv < +\infty$  for some  $\varepsilon > 0$ . Then the solution  $u$  to (3.23) belongs to  $\mathcal{C}^{\lambda, \bar{\mu}}([0, T] \times Q)$  under the Neumann boundary conditions (3.20) (resp. the map

$x \rightarrow u(t, x) \in C^{\bar{\mu}}(Q)$  uniformly in  $t \in [0, T]$  while the map  $t \rightarrow u(t, x) \in C^\lambda([t_0, T])$  uniformly in  $x \in Q$  for  $0 < t_0 < T$  under the Dirichlet boundary conditions (3.21)), where

$$0 < \lambda < \left(1 - \frac{\max_i k_i \vee 2}{4}\right) \wedge \frac{\varepsilon d}{8}, \quad \bar{\lambda} = \lambda \wedge \frac{a}{4} \quad \text{and} \quad 0 < \bar{\mu} < a \wedge \frac{\varepsilon d}{2}.$$

Finally, a straightforward extension of the preceding computations, using Lemmas 2.3 and 4.3 (with  $\alpha_i$  instead of  $\alpha + |a|\delta$ ), provides Hölder regularity for the solution to (3.3).

**Theorem 4.5** *Let  $Q$  be convex, suppose that  $G$  satisfies (1.2) and that the assumptions of Theorem 3.2 are fulfilled. Let  $u_0$  be a continuous function on  $Q$ , let  $u$  be the solution to (3.3) and set  $v(t, x) = u(t, x) - G_t u_0(x)$ . Then*

(i) *If  $\alpha < 1$  and  $F$  is the space-time white noise, then  $v \in C^{\lambda, \mu}([0, T] \times Q)$  for  $0 < \lambda < \inf_i (1 + \alpha - \alpha_i) \wedge \frac{1-\alpha}{2}$  and  $0 < \mu < \frac{\inf_i (1 + \alpha - \alpha_i)}{\delta} \wedge \frac{1-\alpha}{2\delta} \wedge 1$ .*

(ii) *If  $\alpha \geq 1$  and  $F$  is a Gaussian process with covariance function  $f$  defined by (2.1) such that (C1) holds and  $\int_{B_d(0,1)} f(v) |v|^{-d(1+\epsilon)+\frac{1}{\delta}} dv < +\infty$  for some  $\epsilon > 0$ , then  $v \in C^{\lambda, \mu}([0, T] \times Q)$  for  $0 < \lambda < \inf_i (1 + \alpha - \alpha_i) \wedge (\epsilon d \delta) \wedge 1$  and  $0 < \mu < \frac{\inf_i (1 + \alpha - \alpha_i)}{\delta} \wedge (\epsilon d) \wedge 1$ .*

## 5 Density of the solution to the stochastic Cahn-Hilliard PDE

In this section, we concentrate on the solution to (3.23) in dimension 4 and 5 under either the homogeneous Neumann or Dirichlet boundary conditions on  $Q = [0, \pi]^d$ . Thus, we prove that under proper non-degeneracy conditions on the "diffusion" coefficient  $\sigma$ , the law of  $u(t, x)$  has a density for  $t > 0$  and  $x \in Q$ . This extends results proved in [4] and [5] to higher dimension. Since the noise  $F$  has a space correlation, the setting of the corresponding stochastic calculus of variations is that used in [21]. Let  $Q = [0, \pi]^d$ ,  $\mathcal{E}$  denote the inner product space of measurable functions  $\varphi : Q \rightarrow \mathbb{R}$  such that  $\int_Q dx \int_Q dy |\varphi(x)| f(x-y) |\varphi(y)| < +\infty$ , endowed with the inner product

$$\langle \varphi, \psi \rangle_{\mathcal{E}} = \int_Q dx \int_Q dy \varphi(x) f(x-y) \psi(y).$$

Let  $\mathcal{H}$  denote the completion of  $\mathcal{E}$  and set  $\mathcal{E}_T = L^2([0, T], \mathcal{E})$  and  $\mathcal{H}_T = L^2([0, T], \mathcal{H})$ . Note that  $\mathcal{H}$  and  $\mathcal{H}_T$  need not be spaces of functions, and that  $\mathcal{H}_T$  is a Hilbert space which is isomorphic to the reproducing kernel Hilbert space of the Gaussian noise  $(F(\varphi); \varphi \in \mathcal{D}([0, T] \times Q))$ . This noise can be identified with a Gaussian process  $(W(h), h \in \mathcal{H}_T)$  defined as follows. Let  $(e_j, j \geq 0) \subset \mathcal{E}$  be a CONS of  $\mathcal{H}$ ; then  $(W_j(t) = \int_0^t \int_Q e_j(x) F(ds, dx), j \geq 0)$  is a sequence of independent standard Brownian motions such that

$$F(\varphi) = \sum_{j \geq 0} \int_0^t \langle \varphi(s, *), e_j \rangle_{\mathcal{H}} dW_j(s), \varphi \in \mathcal{D}([0, T] \times Q).$$

For  $h \in \mathcal{H}_T$ , we set  $W(h) = \sum_j \int_0^T \langle h(s), e_j \rangle_{\mathcal{H}} dW_j(s)$ , and use the framework of the Malliavin calculus described in [23] to define the Malliavin derivative  $DX$  of a random variable  $X$  and the corresponding Sobolev spaces  $\mathbb{D}^{N,p}$ . Given  $X \in \mathbb{D}^{1,2}$ ,  $h \in \mathcal{H}_T$ , set  $D_h X = \langle DX, h \rangle_{\mathcal{H}_T} = \int_0^T \langle D_{r,*} X, h(r) \rangle_{\mathcal{H}} dr$ , where  $D_{r,*} X \in \mathcal{H}$  for every  $r \in [0, T]$ . Finally for  $r \in [0, T]$  and  $\varphi \in \mathcal{H}$ , set  $D_{r,\varphi} X = \langle D_{r,*} X, \varphi \rangle_{\mathcal{H}}$ .

Since the coefficients  $R$  and  $g(t, x, \cdot)$  are locally Lipschitz, we need to localize the Sobolev spaces as follows. A random variable  $X$  belongs to  $\mathbb{D}_{loc}^{1,p}$  if there exists an increasing sequence  $\Omega_n \subset \Omega$  such that  $\lim_n P(\Omega_n) = 1$  and for every  $n$ , there exists a random variable  $X_n \in \mathbb{D}^{1,p}$  and  $X = X_n$  on  $\Omega_n$ . Let  $u_0 \in \mathcal{C}(Q)$  and suppose that the conditions (H1) and (H'2) hold, where

**(H'2)** The function  $\sigma : \mathbb{R} \rightarrow \mathbb{R}$  is bounded, globally Lipschitz, the map  $g(t, x, \cdot)$  is of class  $\mathcal{C}^1$  with quadratic growth and satisfies (3.22), and the maps  $b_i(t, x, \cdot), 1 \leq i \leq N$  are of class  $\mathcal{C}^1$  with derivatives bounded uniformly in  $(t, x)$ .

Let  $u$  denote the solution to (3.23) with either the homogeneous Neumann or Dirichlet boundary conditions. Lemmas 2.3 and 4.3 imply that the trajectories of  $u - Gu_0$  are almost surely Hölder continuous on  $[0, T] \times Q$ , while the function  $Gu_0$  is clearly bounded by  $\|u_0\|_\infty$ . Therefore,  $\lim_n P(\Omega_n) = 1$  if for every  $n \geq 1$  one sets

$$\Omega_n = \left\{ \omega \in \Omega : \sup\{|u(t, x)|, (t, x) \in Q\} \leq n \right\}.$$

We now construct a sequence of processes  $u(n) \in \mathbb{D}^{1,p}$  for every  $p \in [2, +\infty[$  such that  $u = u(n)$  on  $\Omega_n$ . Let  $K_n$  be the sequence defined at the beginning of the proof of Theorem 3.3, which satisfies (3.25); set  $\bar{R}_n(x) = K_n(|x|)R(x)$  and  $\bar{g}_n(t, x, y) = K_n(|y|)g(t, x, y)$ . The functions  $\bar{R}_n$  and  $y \rightarrow \bar{g}_n(t, x, y)$  are of class  $\mathcal{C}^1$  with bounded derivatives. Hence Theorem 3.2 yields the existence and uniqueness of the process  $u(n)$  solution to the evolution equation:

$$\begin{aligned} u(n)(t, x) &= G_t u_0(x) + \int_0^t \int_Q G(t, x; s, y) \sigma(u(n)(s, y)) F(ds, dy) \\ &+ \int_0^t \int_Q \left[ \Delta G(t, x; s, y) \bar{R}_n(u(n)(s, y)) + G(t, x; s, y) \bar{g}_n(s, y, u(n)(s, y)) \right. \\ &\quad \left. + \sum_{i=1}^N H_i(t, x; s, y) b_i(s, y, u(n)(s, y)) \right] dy ds. \end{aligned} \quad (5.1)$$

The local property of stochastic integrals implies that  $u(n) = u$  on  $\Omega_n$ . The following proposition shows that  $u(n) \in \mathbb{D}^{1,p}$  for every  $p \in [2, +\infty[$ .

**Proposition 5.1** . *Let  $Q$  be a compact subset of  $\mathbb{R}^d$ ,  $u_0 \in \mathcal{C}(Q)$ , let  $\sigma : \mathbb{R} \rightarrow \mathbb{R}$  be globally Lipschitz and for  $i \in \{1, \dots, N\}$ , let  $b_i : [0, T] \times Q \times \mathbb{R}$  satisfy the conditions (L1) and (L2), and suppose that the maps  $y \rightarrow b_i(t, x, y)$  are of class  $\mathcal{C}^1$  with bounded derivatives, let  $G$  and  $(H_i, 1 \leq i \leq N)$  satisfy the conditions (C2) and (C4). Let  $F$  be a Gaussian process with space covariance defined by (2.1) in terms of a function  $f$  which satisfies the condition (C3). Then for every  $p \in [2, +\infty[$  and  $(t, x) \in [0, T] \times Q$ , the solution  $v(t, x)$  to (3.3) belongs to  $\mathbb{D}^{1,p}$ . Furthermore, for every  $r \in [0, T]$  and  $\varphi \in \mathcal{H}$ ,  $D_{r,\varphi} v(t, x) = 0$  if  $r > t$ , while there exists a bounded, adapted family of random variables  $(S(s, y), (s, y) \in [0, T] \times Q)$  such that for  $0 \leq r \leq t$ :*

$$\begin{aligned} D_{r,\varphi} v(t, x) &= \langle G(t, x; r, *) \sigma(v(r, *)), \varphi \rangle_{\mathcal{H}} + \int_r^t \int_Q \left[ G(t, x; s, y) S(s, y) \right. \\ &\quad \left. \times D_{r,\varphi} v(s, y) F(ds, dy) + \sum_{i=1}^N H_i(t, x; s, y) \partial_3 b_i(s, y, v(s, y)) D_{r,\varphi} v(s, y) dy ds \right], \end{aligned} \quad (5.2)$$

and for every  $p \in [1, +\infty[$ ,

$$\sup_{(t,x) \in [0,T] \times Q} E \left( \left| \int_0^t \|D_{r,*} v(t, x)\|_{\mathcal{H}}^2 dr \right|^p \right) = C(p) < +\infty. \quad (5.3)$$

Furthermore, given  $0 \leq s < t \leq T$ , if  $\psi$  denotes the function defined by (2.13),

$$\sup_{x \in Q} E \left( \int_s^t \|D_{r,\varphi} v(t, x)\|_{\mathcal{H}}^2 dr \right) \leq C \left[ \int_0^{t-s} \psi(\tau) d\tau + \sum_{i=1}^N (t-s)^{2(1+\alpha-\alpha_i)} \right]. \quad (5.4)$$

*Proof:* Let  $(v_k, k \geq 0)$  be the Picard approximation scheme of  $v$  defined by (3.12); then by the proof of Theorem 3.2, the sequence  $(v_k(t, x), k \geq 0)$  is bounded in  $L^p(\Omega)$  uniformly in  $(t, x)$  and converges in  $L^p(\Omega)$  to  $v(t, x)$ . Following a classical argument, we prove by induction on  $k$  that  $v_k(t, x) \in \mathbb{D}^{1,p}$  and that

$$\sup_k \sup_{(t,x) \in [0,T] \times Q} E \left( \|Dv_k(t, x)\|_{\mathcal{H}_T}^p \right) < +\infty, \quad (5.5)$$

$$\sum_k \sup_{(t,x) \in [0,T] \times Q} E \left( \|Dv_{k+1}(t, x) - Dv_k(t, x)\|_{\mathcal{H}_T}^p \right) < +\infty. \quad (5.6)$$

Then using [23], Lemma 1.2.3, we conclude that  $Dv_k(t, x)$  converges to  $Dv(t, x)$  in the weak topology of  $L^p(\Omega, \mathcal{H}_T)$ ; furthermore, this yields (5.3). Since  $v_0$  is deterministic, it belongs to  $\mathbb{D}^{1,p}$ ; suppose that  $v_k \in \mathbb{D}^{1,p}$ ; since  $\sigma$  is globally Lipschitz, Proposition 1.2.3 in [23] implies that  $\sigma(v_k(s, y)) \in \mathbb{D}^{1,p}$  and that  $D_{r,\varphi}(\sigma(v_k(s, y))) = S_k(s, y) D_{r,\varphi}v_k(s, y)$ , where  $S_k(s, y)$  is a bounded adapted process. Furthermore, for every  $r \in [0, T]$  and  $\varphi \in \mathcal{H}$ ,  $D_{r,\varphi}v_{k+1}(t, x) = 0$  if  $r > t$  and for  $r \leq t$ :

$$\begin{aligned} D_{r,\varphi}v_{k+1}(t, x) = & \langle G(t, x; r, *) \sigma(v_k(r, *)), \varphi \rangle_{\mathcal{H}} + \int_r^t \int_Q \left[ G(t, x; s, y) S_k(s, y) \right. \\ & \left. \times D_{r,\varphi}v_k(s, y) F(ds, dy) + \sum_{i=1}^N H_i(t, x; s, y) \partial_3 b_i(s, y, v_k(s, y)) D_{r,\varphi}v(s, y) dy ds \right]. \end{aligned}$$

Let  $\psi$  be the  $L^1([0, T])$  function defined by (2.13) and set  $I = \int_0^T \psi(r) dr$ . The linear growth condition on  $\sigma$  and equations (3.18) and (2.12) imply that for any  $p \in [2, +\infty[$ , there exists a constant  $C_p$  (which does not depend on  $k$ ) such that for every  $k$

$$\sup_{x \in Q} E \left( \|G(t, x; \cdot, *) \sigma(v_k(\cdot, *))\|_{\mathcal{H}_T}^{2p} \right) \leq C I^{p-1} \int_0^t \psi(t-r) \sup_{(r,x)} E(|v_k(r, x)|^{2p}) dr \leq C_p. \quad (5.7)$$

For  $t \in [0, T]$ ,  $x \in Q$ , let  $(Y_\tau(x), 0 \leq \tau \leq T)$  be the  $\mathcal{H}_T$ -valued martingale defined by

$$Y_\tau(x) = \int_0^{\tau \wedge t} \int_Q G(t, x; s, y) S_k(s, y) Dv_k(s, y) F(ds, dy).$$

Let  $(\epsilon_j, j \geq 0)$  be a CONS of  $\mathcal{H}_T$ ; then Burkholder's inequality for Hilbert-valued martingales (see e.g. [19], p. 212) and Parseval's identity yield

$$\sup_{x \in Q} \|Y_t(x)\|_{L^{2p}(\Omega, \mathcal{H}_T)}^{2p} \leq C_p I^{p-1} \int_0^t \psi(t-s) \sup_y E(\|Dv_k(s, y)\|_{\mathcal{H}_T}^{2p}) ds. \quad (5.8)$$

Finally, Lemma 3.1 and the conditions (C2) - (C4) imply that the function  $\bar{\psi}(s) = \psi(s) + \sum_{i=1}^N s^{-\alpha_i + \alpha} \in L^1([0, T])$ , and together with the inequalities (5.7) and (5.8) this yields that there exists  $C_p > 0$  such that for every  $k \geq 0$ ,

$$\sup_{x \in Q} \|Dv_k(t, x)\|_{L^{2p}(\Omega, \mathcal{H}_T)}^{2p} \leq C_p + C_p \int_0^t \bar{\psi}(t-s) \sup_{y \in Q} \|Dv_k(s, y)\|_{L^{2p}(\Omega, \mathcal{H}_T)}^{2p} ds.$$

Thus Lemma 15 in [6] concludes the proof of (5.5). A similar argument shows (5.6). We conclude that each  $v_k(t, x) \in \mathbb{D}^{1,p}$ , and that (5.2) and (5.3) hold. To prove (5.4), we use (3.10), (5.2) and arguments similar to the previous ones; then for  $0 \leq s < t \leq T$ ,

$$E \left( \int_s^t \|G(t, x; r, *) \sigma(v(r, *))\|_{\mathcal{H}}^2 dr \right) \leq C \left( \int_s^t \|G(t, x; r, *)\|_{\mathcal{H}}^2 dr \right) \left[ 1 + \sup_{(r,y)} E(|v(r, y)|^{2p}) \right]$$

$$\leq C \int_0^{t-s} \psi(\tau) d\tau.$$

Furthermore, the isometry of Hilbert-spaced valued martingales, Fubini's theorem and (5.3) imply

$$\begin{aligned} & E \left( \int_s^t \left\| \int_r^t \int_Q G(t, x; \tau, y) S(\tau, y) D_{r,*} v(\tau, y) F(d\tau, dy) \right\|_{\mathcal{H}}^2 dr \right) \\ & \leq C \int_s^t d\tau \int_Q dy \int_Q dz G(t, x; \tau, y) f(y-z) G(t, x; \tau, z) \\ & \quad \times \int_s^\tau E(\langle D_{r,*} v(s, y), D_{r,*} v(s, z) \rangle_{\mathcal{H}}) dr \leq C \int_0^{t-s} \psi(\tau) d\tau. \end{aligned}$$

Finally, conditions (C2) and (C4), Minkowski's and Schwarz's inequalities and Fubini's theorem imply that for every  $i \leq N$ ,

$$\begin{aligned} & E \left( \int_s^t \left\| \int_r^t \int_Q H_i(t, x; \tau, y) \partial_3 b_i(\tau, y, v(s, y)) D_{r,*} v(\tau, y) dy d\tau \right\|_{\mathcal{H}}^2 dr \right) \\ & \leq C (t-s)^{\alpha+1-\alpha_i} \int_s^t d\tau \int_Q dy |H_i(t, x; \tau, y)| \sup_{(\tau, y) \in [0, T] \times Q} \int_s^\tau E(\|D_{r,*} v(\tau, y)\|_{\mathcal{H}}^2) dr \\ & \leq (t-r)^{2(\alpha+1-\alpha_i)}. \end{aligned}$$

This completes the proof of (5.4).  $\square$

The following theorem, which establishes the absolute continuity of the law to the stochastic Cahn-Hilliard PDE, is the main result of this section.

**Theorem 5.2** . Let  $Q = [0, \pi]^d$ , suppose that the conditions (H.1), (H'2) and (H.4) hold, let  $F$  be a Gaussian noise with covariance defined by (2.1) in terms of a function  $f$  which satisfies the conditions (C1) and (3.24). Let  $u_0 \in \mathcal{C}(Q)$  and let  $u$  be the solution to (3.23) with initial conditions  $u_0$  and the homogeneous Neumann or Dirichlet boundary conditions. Let  $t_0 \in ]0, T]$ ,  $x_1, \dots, x_l$  be pairwise distinct points in  $]0, \pi[^d$ , and set  $u(t_0, \underline{x}) = (u(t_0, x_1), \dots, u(t_0, x_l))$ . For any  $\tau > 0$ , let

$$I(\tau) = \int_{B_d(0, \tau)} f(v) |v|^{-d+4} \ln(|v|^{-1})^{(5-d)^+} dv. \quad (5.9)$$

(i) Suppose that  $|\sigma| \geq C > 0$ ; then if for some  $0 < \nu < \frac{1}{4}$

$$\lim_{\tau \rightarrow 0} I(\tau^{\frac{1}{4}+\nu})^{-1} \left[ \tau + \tau^{\frac{1}{2}} I(\tau^{\frac{1}{4}-\nu}) \right] = 0,$$

the law of  $u(t_0, \underline{x})$  is absolutely continuous with respect to Lebesgue's measure.

(ii) Let  $u_0 \in \mathcal{C}^a(Q)$  for some  $a > 0$  and suppose that for some  $\nu > 0$ :

$$\lim_{\tau \rightarrow 0} I(\tau^{\frac{1}{4}+\nu})^{-1} \left[ \tau^A + \tau^{\frac{1}{2}} I(\tau^{\frac{1}{4}-\nu}) \right] = 0 \text{ for } A \in ]0, \frac{d\varepsilon}{4} + \frac{\mu}{8}[ \text{ and } \mu \in ]0, 1 \wedge \frac{d\varepsilon}{2} \wedge a[. \quad (5.10)$$

Then the law of  $u(t_0, \underline{x})$  is absolutely continuous on  $\{\sigma \neq 0\}^l$ .

**Remark 5.3** . Let  $f(v) = |v|^{-B}$  for some  $B > 0$ ; then the condition (C1) holds for any  $B > 0$ , while (3.24) holds if and only if  $d\varepsilon + B < 4$ . Since for small  $\tau > 0$ ,  $\int_0^\tau \rho^{d-1-B-d+4} d\rho = C \tau^{4-B} \leq I(\tau) \leq \int_0^\tau \rho^{d-1-B-d+4-\xi(d-4)^+} d\rho = C \tau^{4-B-\xi(d-4)^+}$  for any small  $\xi > 0$ , one has  $\lim_{\tau \rightarrow 0} I(\tau^{\frac{1}{4}+\nu})^{-1} [\tau + \tau^{\frac{1}{2}} I(\tau^{\frac{1}{4}-\nu})] = 0$  for every  $B > 0$  and  $0 < \nu < \frac{B \wedge 1}{16}$ , while (5.10) holds if and only if  $B + d\varepsilon > \frac{7}{2} \vee \left( 4 - \left( \frac{d\varepsilon}{4} \wedge \frac{a}{2} \right) \right)$ .

**Remark 5.4** *The proofs of Theorems 1.5 in [4] and Theorems 1.2-1.4 in [5] extend to the case of the space-time white noise  $F$  in dimension  $d \leq 3$  under the homogeneous Dirichlet boundary conditions on  $[0, \pi]^d$ .*

*Proof of Theorem 5.2:* The proof, which is similar to that of Theorem 1.2 in [5] and Theorem 3.1 in [21] is only sketched in case (ii). According to Theorem 4.4, the trajectories of the solution  $u(n)$  to (5.1) almost surely belong to  $\mathcal{C}^{\lambda, \mu}([\frac{t_0}{2}, T] \times Q)$  for  $0 < \lambda < \frac{1}{2} \wedge \frac{d\varepsilon}{8}$  and  $0 < \mu < 1 \wedge \frac{d\varepsilon}{2} \wedge a$ . (Note that according to Theorem 4.4, the trajectories of  $u$  have the same Hölder regularity.) Using Theorem 2.1.2 and the following remark in [23], it suffices to show that for every  $n \geq 1$  and  $M \geq 1$ , the  $l \times l$  Malliavin covariance matrix  $\Gamma(n)$  defined by

$$\Gamma(n)(i, j) = \langle Du(n)(t_0, x_i), Du(n)(t_0, x_j) \rangle_{\mathcal{H}_T}$$

is almost surely invertible on the set  $\tilde{\Omega}(M) = \cap_{i=1}^l \{|\sigma(u(n)(t_0, x_i))| \geq \frac{1}{M}\}$ . As usual, this reduces to proving that for any vector  $v \in \mathbb{R}^l$ , with  $|v| = 1$ ,  $\langle \Gamma(n)v, v \rangle_{\mathbb{R}^l} > 0$  a.s. on  $\tilde{\Omega}(M)$ .

For  $1 \leq i \leq l$ ,  $r \leq t$ , (5.2) in Proposition 5.1 shows that  $D_{r,*}u(n)(t_0, x_i) = G(t_0, x_i; r, *) \times \sigma(u(n)(r, *)) + U(t_0, r, x_i)$ ; for a fixed unit vector  $v$  of  $\mathbb{R}^d$ , a usual argument shows that given  $\tau \in ]0, \frac{t_0}{2}]$ ,

$$\langle \Gamma(n)v, v \rangle \geq \frac{I_1}{4} + \frac{1}{4} \sum_{i=1}^l \sum_{j \neq i; j=1}^l v_i v_j I_2(i, j) - \frac{l}{2} \sum_{i=1}^l v_i^2 I_3(i) - l \sum_{i=1}^l v_i^2 I_4(i), \quad (5.11)$$

where

$$\begin{aligned} I_1 &= \sum_{i=1}^l v_i^2 \int_{t_0-\tau}^{t_0} \|G(t_0, x_i; r, y) \sigma(u(n)(r, x_i))\|_{\mathcal{H}}^2 dr, \\ I_2(i, j) &= \int_{t_0-\tau}^{t_0} \langle G(t_0, x_i; r, *) \sigma(u(n)(r, x_i)), G(t_0, x_j; r, *) \sigma(u(n)(r, x_j)) \rangle_{\mathcal{H}} dr, \\ I_3(i) &= \int_{t_0-\tau}^{t_0} \|G(t_0, x_i; r, y) [\sigma(u(n)(r, y)) - \sigma(u(n)(r, x_i))]\|_{\mathcal{H}}^2 dr, \\ I_4(i) &= \int_{t_0-\tau}^{t_0} \|U(t_0, r, x_i)\|_{\mathcal{H}}^2 dr. \end{aligned}$$

Remark 2.2 shows that condition (2.7) is satisfied. Let  $\bar{c} = \inf\{d(x_i, \partial Q), 1 \leq i \leq l\}$  and suppose that for the constant  $C_2$  defined in the proof of Lemma 2.1 (ii),  $2C_2\tau^{\frac{1}{4}} \leq \bar{c}$ . Then (2.17) and (2.16) imply that on  $\tilde{\Omega}(M)$ , for  $\nu > 0$ ,  $\tau$  small enough and  $d = 4, 5$ ,

$$I_1 \geq C \left( \sum_{i=1}^l |v_i|^2 \right) \frac{1}{M^2} \int_{B_d(0, C_2\tau^{\frac{1}{4}+\nu})} f(v) |v|^{-d+4} \ln(|v|^{-1})^{(5-d)^+} dv. \quad (5.12)$$

We now prove upper estimates of  $I_2(i, j)$  up to  $I_4(i)$ . Let  $m = \inf\{|x_i - x_j|, 1 \leq i < j \leq l\}$ , let  $c_1$  and  $C_1$  denote the constants appearing in condition (C1), and let  $k \in ]0, \frac{1}{3}[$  be such that  $k(1 + c_1) < \frac{1}{2}$ . Fix  $1 \leq i < j \leq l$ ; then  $I_2(i, j) \leq C \|\sigma\|_{\infty}^2 J_2(i, j)$ , where

$$J_2(i, j) = \int_0^{\tau} r^{-\frac{d}{2}} dr \int_Q dy \int_Q dz \exp\left(-c \frac{|x_i - y|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) f(y - z) \exp\left(-c \frac{|x_j - z|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right).$$

We split the integral on  $Q \times Q$  in several parts. Indeed, if  $|y - z| \geq km$ , the continuity of  $f$  implies that  $f(y - z) \leq C < +\infty$ . Suppose now that,  $|y - z| \leq km$ . Then if  $|y - x_i| \leq c_1 |y - z|$ ,



(C1) implies that  $f(y-z) \leq C_1 f(y-x_i)$ , while  $|z-x_j| \geq \left| |x_i-x_j| - (|x_i-y| + |y-z|) \right| \geq m \left(1 - k(1+c_1)\right) \geq \frac{m}{2}$ . Similarly, if  $|z-x_j| \leq c_1|y-z|$ , then  $f(y-z) \leq C_1 f(z-x_j)$  and  $|y-x_i| \geq \frac{m}{2}$ . Finally, suppose that  $|y-z| \leq km$ ,  $c_1|y-z| \leq |y-x_i| \wedge |z-x_j|$ ; then since  $k < \frac{1}{3}$ , one of the norms  $|y-x_i|$  or  $|z-x_j|$  (say  $|z-x_j|$ ) is larger than  $\frac{m}{3}$ . Thus,  $J_2(i, j) \leq J_{2,1}(i, j) + 2J_{2,2}(i, j) + 2J_{2,3}(i, j)$ , where

$$\begin{aligned}
J_{2,1}(i, j) &= \int_0^\tau r^{-\frac{d}{2}} \int \int_{|y-z| \geq km} \exp\left(-c \frac{|y-x_i|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) f(y-z) \exp\left(-c \frac{|z-x_j|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) dy dz \leq C\tau, \\
J_{2,2}(i, j) &\leq C_1 \int_0^\tau r^{-\frac{d}{4}} dr \int_{B_d(0, R)} f(v) \exp\left(-c \frac{|v|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) dv \int_{|z| \geq \frac{m}{2} r^{-\frac{1}{4}}} \exp\left(-c|z|^{\frac{4}{3}}\right) dz \\
&\leq C_1 \int_{B_d(0, R)} f(v) dv \int_0^\tau r^{-\frac{d}{4}} \exp\left(-c \frac{|v|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) \exp\left(-\tilde{c}r^{-\frac{1}{3}}\right) dr \\
&\leq C \exp(-\tilde{c}\tau^{-\frac{1}{3}}) \int_{B_d(0, R)} f(v) |v|^{-d+4} \ln(|v|^{-1})^{(5-d)^+} dv \leq C \exp(-\tilde{c}\tau^{-\frac{1}{3}}), \\
J_{2,3}(i, j) &\leq C_1 \int_0^\tau r^{-\frac{d}{4}} dr \int_Q dy \int_{|z-x_j| \geq \frac{m}{3}} f(y-z) \exp\left(-\tilde{c} \frac{|y-z|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) \exp\left(-\tilde{c} \frac{|z-x_j|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) dz \\
&\leq C \exp(-\tilde{c}\tau^{-\frac{1}{3}}).
\end{aligned}$$

Hence, for  $\tau$  small enough,

$$\left| \sum_{i=1}^l \sum_{j \neq i; j=1}^l v_i v_j I_2(i, j) \right| \leq C\tau. \quad (5.13)$$

Fubini's theorem, the Lipschitz property of  $\sigma$ , the Hölder regularity of  $x \rightarrow u(n)(t, x)$  uniformly for  $\frac{t_0}{2} \leq t \leq t_0$ , Schwarz's inequality and  $|y-z|^{\frac{4}{3}} \leq 2^{\frac{1}{3}} \left(|y-x_i|^{\frac{4}{3}} + |z-x_j|^{\frac{4}{3}}\right)$  yield for any  $i \leq l$ ,  $0 < \tau \leq \frac{t_0}{2}$ :

$$\begin{aligned}
E(|I_3(i)|) &\leq C \int_{t_0-\tau}^{t_0} \int_Q \int_Q dy dz |G(t, x_i; r, y)| f(y-z) |G(t, x_i; r, z)| \\
&\quad \times E\left(|u(n)(r, y) - u(n)(r, x_i)| |u(n)(r, z) - u(n)(r, x_i)|\right) dr \\
&\leq C \int_0^\tau r^{-\frac{d}{2}} dr \int_Q \int_Q dy dz \exp\left(-\tilde{c} \frac{|y-z|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) f(y-z) |z-x_i|^{\frac{\mu}{2}} \exp\left(-\tilde{c} \frac{|z-x_i|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) \\
&\leq C \int_{B_d(0, R)} f(v) dv \int_0^\tau r^{-\frac{d}{4} + \frac{\mu}{8}} \exp\left(-\tilde{c} \frac{|v|^{\frac{4}{3}}}{r^{\frac{1}{3}}}\right) dr \int_{B_d(0, R)} |z|^{\frac{\mu}{2}} \exp(-\tilde{c}|z|^{\frac{4}{3}}) dz \\
&\leq C \int_{B_d(0, R)} f(v) |v|^{-d+4+\frac{\mu}{2}} dv \int_{|v|^{\frac{4}{3}} \tau^{-\frac{1}{3}}}^{+\infty} s^{-4+\frac{3d}{4}-\frac{3\mu}{8}} \exp(-\tilde{c}s) ds,
\end{aligned}$$

where to obtain the last integral, we have set  $s = |v|^{\frac{4}{3}} r^{-\frac{1}{3}}$ . We split the last integral on  $\{|v| \leq \tau^{\frac{1}{4}-\nu}\}$  and its complement for some  $\nu > 0$ . Then using (3.24), a straightforward computation yields for  $d = 4, 5$  and  $\tau$  small enough:

$$E(|I_3(i)|) \leq C \int_{B_d(0, \tau^{\frac{1}{4}-\nu})} f(v) |v|^{4-d+\frac{\mu}{2}} \left(1 + |v|^{-\frac{\mu}{2}} \tau^{\frac{\mu}{8}}\right) dv$$

$$\begin{aligned}
& +C \int_{B_d(0,R)} 1_{\{|v| \geq \tau^{\frac{1}{4}-\nu}\}} f(v) |v|^{4-d+\frac{\mu}{2}} \exp(-\tilde{c}\tau^{-\frac{4\nu}{3}}) dv \\
& \leq C \tau^{\frac{d\varepsilon}{4} + \frac{\mu}{8} - \nu(d\varepsilon + \frac{\mu}{2})}.
\end{aligned}$$

Thus, choosing  $\nu$  close enough to 0 and  $\tau_0$  small enough, we deduce that for  $0 < \tau \leq \tau_0$ :

$$E(|I_3(i)|) \leq C\tau^A, \quad 0 < A < \frac{d\varepsilon}{4} + \frac{\mu}{8}. \quad (5.14)$$

Finally, using the decomposition of  $U(t_0, r, x_i)$ , and Fubini's theorem, we obtain that  $E(|I_4(i)|) \leq \sum_{i=1}^4 T_j$ , where

$$\begin{aligned}
T_1 &= \int_{t_0-\tau}^{t_0} dr E \left( \left\| \int_r^{t_0} \int_Q G(s, x_i; r, y) S_i(n)(s, y) D_{r,*} u(n)(s, y) F(ds, dy) \right\|_{\mathcal{H}}^2 \right), \\
T_2 &= \int_{t_0-\tau}^{t_0} dr E \left( \left\| \int_r^{t_0} \int_Q \Delta G(s, x_i; r, y) R'_n(u(n)(s, y)) D_{r,*} u(n)(s, y) dy ds \right\|_{\mathcal{H}}^2 \right), \\
T_3 &= \int_{t_0-\tau}^{t_0} dr E \left( \left\| \int_r^{t_0} \int_Q G(s, x_i; r, y) \partial_3 g(s, y, (u(n)(s, y))) D_{r,*} u(n)(s, y) dy ds \right\|_{\mathcal{H}}^2 \right), \\
T_4 &= \sum_{j=1}^N \int_{t_0-\tau}^{t_0} dr E \left( \left\| \int_r^{t_0} \int_Q H_j(s, x_i; r, y) \partial_3 b_j(s, y, (u(n)(s, y))) D_{r,*} u(n)(s, y) dy ds \right\|_{\mathcal{H}}^2 \right).
\end{aligned}$$

The isometry property for Hilbert-space valued martingales, (5.3) with  $p = 1$ , Schwarz's inequality and (2.14) or (2.15) imply for  $\nu$  small enough

$$\begin{aligned}
T_1 &\leq C \int_{t_0-\tau}^{t_0} ds \int_Q \int_Q dy dz |G(s, x_i; r, y)| f(y-z) |G(s, x_i; r, z)| \\
&\quad \times \sup_{(s,y) \in [t_0-\tau, t_0] \times Q} E \left( \int_{t_0-\tau}^s \|D_{r,*} u(n)(s, y)\|_{\mathcal{H}}^2 dr \right) \\
&\leq C \tau \left[ I(\tau^{\frac{1}{4}-\nu}) + \exp\left(-\tilde{c}\tau^{\frac{3\nu}{4}}\right) \right]. \quad (5.15)
\end{aligned}$$

Minkowski's and Schwarz's inequalities, then Fubini's theorem and (5.4) imply

$$\begin{aligned}
T_2 &\leq C \int_{t_0-\tau}^{t_0} (t_0 - r)^{\frac{1}{2}} dr \int_r^{t_0} \int_Q \Delta G(t_0, x_i; s, y) R'_n(u(n)(s, y)) E(\|D_{r,*} u(n)(s, y)\|_{\mathcal{H}}^2) dy ds \\
&\leq C_n \tau^{\frac{1}{2}} \int_{t_0-\tau}^{t_0} ds \int_Q dy \Delta G(t_0, x_i; s, y) \sup_{(s,y) \in [t_0-\tau, t_0] \times Q} E \left( \int_{t_0-\tau}^s \|D_{r,*} u(n)(s, y)\|_{\mathcal{H}}^2 dr \right) \\
&\leq C(n) \tau \left[ I(\tau) + \tau^{\frac{1}{2}} \right]. \quad (5.16)
\end{aligned}$$

A similar computation, using Minkowski's inequality, Fubini's theorem, Lemma 3.1 with  $\rho = \infty$  and  $q = 1$  and (5.4) yields

$$\begin{aligned}
T_3 &\leq C_n \tau \int_{t_0-\tau}^{t_0} ds \int_Q dy G(t_0, x_i; s, y) \sup_{(s,y) \in [t_0-\tau, t_0] \times Q} E \left( \int_{t_0-\tau}^s \|D_{r,*} u(n)(s, y)\|_{\mathcal{H}}^2 dr \right) \\
&\leq C_n \tau^2 \left[ I(\tau) + \tau^{\frac{1}{2}} \right], \quad (5.17) \\
T_4 &\leq C_n \tau^{\frac{1}{4}} \sum_{j=1}^N \int_{t_0-\tau}^{t_0} ds \int_Q dy H_j(t_0, x_i; s, y) \sup_{(s,y) \in [t_0-\tau, t_0] \times Q} E \left( \int_{t_0-\tau}^s \|D_{r,*} u(n)(s, y)\|_{\mathcal{H}}^2 dr \right)
\end{aligned}$$

$$\leq C_n \tau^{\frac{1}{2}} \left[ I(\tau) + \tau^{\frac{1}{2}} \right], \quad (5.18)$$

The inequalities (5.15)-(5.18) yield that

$$E(|I_4(i)|) \leq C(n) \left[ \tau^{\frac{1}{2}} I(\tau^{\frac{1}{4}-\nu}) + \tau \right]. \quad (5.19)$$

Finally, the inequalities (5.11)-(5.14) and (5.19) imply that for  $\rho > 0$  such that  $\tau + \tau^\rho \leq \frac{1}{2}I(\tau^{\frac{1}{4}+\nu})$  (which exists because of (5.10)), we have for small enough  $\tau$ :

$$\begin{aligned} P(\langle v, \Gamma(n)v \rangle_{\mathbb{R}^l} < \tau^\rho) &\leq P\left(I_3 + I_4 \geq C [I(\tau^{\frac{1}{4}+\nu}) - \tau - \tau^\rho]\right) \\ &\leq P\left(I_3 + I_4 \geq \frac{1}{2} I(\tau^{\frac{1}{4}+\nu})\right) \leq C I(\tau^{\frac{1}{4}+\nu})^{-1} \left[ \tau^A + \tau^{\frac{1}{2}} I(\tau^{\frac{1}{4}-\nu}) \right] \end{aligned}$$

for some  $A < \frac{\mu}{8} + \frac{d\varepsilon}{2}$ ; then the condition (5.10) concludes the proof of (ii).  $\square$

## References

- [1] Adams, R.A. (1975), Sobolev Spaces, Academic Press, New York.
- [2] Bally, V., Millet, A. and Sanz-Solé, M. (1995), Approximation and support theorem in Hölder norm for parabolic stochastic partial differential equations, Ann. Probab. 23, 178-222.
- [3] Brzezniak, Z. and Peszat, S. (1999), Space-time continuous solutions to SPDEs driven by a homogeneous Wiener process, Studia Mathematica 137, 262-299.
- [4] Cardon-Weber, C., Cahn-Hilliard equation: existence of the solution and of its density, (to appear in Bernoulli), Prépublication 543 du Laboratoire de Probabilités et Modèles Aléatoires (1999).
- [5] Cardon-Weber, C., Cahn-Hilliard stochastic equation: strict positivity of the density, (to appear in Stochastic and Stochastic reports), Prépublication 576 du Laboratoire de Probabilités et Modèles Aléatoires (2000).
- [6] Dalang, R. (1999), Extending martingale measure stochastic integral with applications to spatially homogeneous S.P.D.E.'S, Electronic Journal of Probability 4, 1-29, <http://www.math.washington.edu/ejpecp/EjpVol4/paper6.abs.html>
- [7] Dalang, D. and Frangos, N. (1998), The stochastic wave equation in two-spatial dimensions, Ann. Probab. 26-1, 187-212.
- [8] Da Prato, G. and Debussche, A. (1996), Stochastic Cahn Hilliard equation, Nonlinear Anal. 26, 241-263.
- [9] Da Prato, G., Debussche, A. and Temam, R. (1994), Stochastic Burger's equation, Nonlinear Differential equations and Applications, Birkhäuser, 389-402.
- [10] Da Prato, G. and Zabczyk, J. (1992), Stochastic equations in infinite dimensions, Encyclopedia of Mathematics and its Applications, Cambridge University Press.
- [11] Debussche, A. and de Bouard, A. (1998), On the stochastic Korteweg-de Vries equation, J. Funct. Anal. 154, 215-251.

- [12] Eidelman, S.D. and Ivashin, N.V. (1970), Investigation on the Green matrix for a homogeneous parabolic boundary value problem, *Trans. Moscow Math. Soc.* 23, 179-243.
- [13] Eidelman, S.D. and Zhitarashu, N.V. (1998), *Parabolic boundary value Problems*, Birkhäuser, Basel.
- [14] Garsia, A. (1972), Continuity properties of Gaussian processes with multi-dimensional time parameter, *Proc. Sixth Berkeley Symp. Math. Statist. Probab.*, 369-374.
- [15] Gyöngy, I. (1998), Existence and uniqueness results for semi linear stochastic partial differential equations, *Stochastic Processes and their Applications* 73, 271-299.
- [16] Karkzevska, A. and Zabczyk, J. (1998), A note on stochastic wave equations, *Evolution Equations and their Applications in Physical and Life Sciences*, Proceedings of the 6th International Conference, Bad Herrenalb, Marcel Dekker.
- [17] Karkzevska, A. and Zabczyk, J. (2000), Stochastic P.D.E.s with function-valued solutions, *Infinite Dimensional Stochastic Analysis*, Proceedings of the Colloquium, Amsterdam 1999, *Verhandelingen, Afd. Natuurkunde, Vol. 52*, Royal Netherlands Academy of Arts and Sciences, Amsterdam, 197-216.
- [18] Lions, P.L. and Magenes, E., *Problèmes aux limites non linéaires et applications*, Vol. 1, Dunod 1968.
- [19] Métivier, M. (1982), *Semimartingales*, de Gruyter, Berlin.
- [20] Millet, A. and Morien, P.L., On a non linear stochastic wave equation in the plane: existence and uniqueness of the solution, *The Annals of Applied Probability* (to appear), Prépublication 00/1 de l'Équipe MODAL'X.
- [21] Millet, A. and Sanz-Solé, M. (1999), A stochastic wave equation in two space dimension: smoothness of the law, *Ann. Probab.* 27, 803-844.
- [22] Mueller, C. (1997), Long time existence for the wave equation with a noise term, *Ann. Probab.* 25, 133-151.
- [23] Nualart, D. (1995), *The Malliavin Calculus and Related Topics*, Springer Verlag, Berlin.
- [24] Peszat, S., SPDEs driven by a homogeneous Wiener process, preprint.
- [25] Peszat, S. and Zabczyk, J.(1997), Stochastic evolution equations with spatially homogeneous Wiener process, *Stoch. Proc. Appl.* 72 , 187-204.
- [26] Peszat, S. and Zabczyk, J. (2000), Nonlinear stochastic wave and heat equations, *Probab. Theory Related Fields* 116, 421-443.
- [27] Schwartz, L. (1966), *Théorie des distributions*, Hermann, Paris.
- [28] Walsh, J.B. (1986), *An introduction to stochastic partial differential equations*, Ecole d'été de Probabilités de Saint-Flour XIV, 1984, *Lecture Notes in Math* 1180, Springer Verlag.