



**HAL**  
open science

# The universal cover of a monomial triangular algebra without multiple arrows

Patrick Le Meur

► **To cite this version:**

Patrick Le Meur. The universal cover of a monomial triangular algebra without multiple arrows. 2007.  
hal-00123559v1

**HAL Id: hal-00123559**

**<https://hal.science/hal-00123559v1>**

Preprint submitted on 10 Jan 2007 (v1), last revised 6 Mar 2008 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The universal cover of a monomial triangular algebra without multiple arrows

Patrick Le Meur <sup>\*†</sup>

10th January 2007

## Abstract

Let  $A$  be a basic connected finite dimensional algebra over an algebraically closed field  $k$ . Assuming that  $A$  is monomial and that the ordinary quiver  $Q$  of  $A$  has no oriented cycle and no multiple arrows, we prove that  $A$  admits a universal cover with group the fundamental group of the underlying space of  $Q$ .

## Introduction

Let  $A$  be a finite dimensional  $k$ -algebra where  $k$  is an algebraically closed field. In order to study the category  $\text{mod}(A)$  of (left)  $A$ -modules, one may assume that  $A$  is basic and connected. When  $\mathcal{C} \rightarrow A$  is a Galois covering with  $\mathcal{C}$  locally bounded, the covering techniques introduced in [3] and [12] allow one to reduce the study of (part of)  $\text{mod}(A)$  to the study of  $\text{mod}(\mathcal{C})$  and which is easier to handle (see for example [4], [6], [7]). Hence, Galois coverings of  $A$  behave like Galois coverings of topological spaces and one can wonder if  $A$  admits a fundamental group and a universal Galois covering, i.e. a Galois covering  $\mathcal{C} \rightarrow A$  with  $\mathcal{C}$  connected and locally bounded and which is factorised by any other Galois covering. Such a universal Galois covering exists if  $A$  is of finite representation type (see [6]). Moreover, a fundamental group  $\pi_1(Q, I)$  associated with any admissible presentation  $kQ/I \simeq A$  with quiver and relations  $I$  was defined in [11]. This group is constructed using a homotopy relation  $\sim_I$  like in the topological situation and is the group of a Galois covering of  $A$  defined by the presentation  $kQ/I \simeq A$ . However, the group  $\pi_1(Q, I)$  depends on  $(Q, I)$  and there exist examples of algebras  $A$  with different presentations having non isomorphic fundamental groups. In a previous text ([9]), the author has compared the fundamental groups associated with different presentations of the same algebra. This study showed ([9, Thm. 1]) that: if  $A$  is triangular (i.e. its ordinary quiver  $Q$  has no oriented cycle) and without double bypass, and if  $k$  has characteristic zero then  $A$  admits a presentation  $kQ/I_0 \simeq A$  such that the fundamental group  $\pi_1(Q, I)$  of any other presentation  $kQ/I \simeq A$  is a quotient of  $\pi_1(Q, I_0)$ . Under the same hypotheses, this study showed ([9, Thm. 2]) that: the Galois covering with group  $\pi_1(Q, I_0)$  of  $A$  defined by the presentation  $kQ/I_0 \simeq A$  satisfies a factorisation property with respect to the Galois coverings of  $A$  similarly to the universal cover of an arcwise connected topological space. Recall ([2]) that a bypass in a quiver is a couple  $(\alpha, u)$  where  $\alpha \neq u$ ,  $\alpha$  is an arrow of  $Q$ , and  $u$  is a path parallel to  $\alpha$ . Recall also ([9]) that a double bypass is 4-tuple  $(\alpha, u, \beta, v)$  where  $(\alpha, u)$ ,  $(\beta, v)$  are bypasses such that the arrow  $\beta$  appears in the path  $u$ .

The aim of the present text is to extend [9, Thm. 2] to monomial algebras without using conditions on the characteristic of the field or on the double bypasses. Recall that  $A$  is called monomial if it admits a presentation  $kQ/I_0 \simeq A$  where  $I_0$  is an ideal of  $kQ$  generated by a set of paths. Notice that for such an algebra, [9, Thm. 1] cited above is obvious. Indeed: for any bound quiver  $(Q, I)$ , the group  $\pi_1(Q, I)$  is a quotient of the fundamental group  $\pi_1(Q)$  of the underlying graph of  $Q$ , and these two groups are isomorphic if  $I$  is generated by a set of paths. On the other hand [9, Thm. 2] is less obvious because it carries a linear setting which is not encoded in the fundamental group. With this setting, we prove the following theorem which is the main result of this text:

**Theorem 1.** *Let  $Q$  be a quiver without oriented cycle and without multiple arrows. Let  $I_0$  be a monomial ideal of  $kQ$ . Let  $\hat{\mathcal{C}} \rightarrow kQ/I_0$  be the Galois covering with group  $\pi_1(Q)$  associated to the universal Galois*

---

<sup>\*</sup> *adress:* Département de Mathématiques, Ecole normale supérieure de Cachan, 61 avenue du président Wilson, 94235 Cachan, France

<sup>†</sup> *e-mail:* plemeur@dptmaths.ens-cachan.fr

covering of  $(Q, I_0)$  (see [11]). For any Galois covering  $\mathcal{C} \rightarrow kQ/I_0$  with group  $G$  and with  $\mathcal{C}$  connected and locally bounded, there exists a commutative diagram of  $k$ -categories and  $k$ -linear functors:

$$\begin{array}{ccc}
 \widehat{\mathcal{C}} & & \mathcal{C} \\
 \downarrow & \searrow & \downarrow \\
 kQ/I_0 & \xrightarrow{\sim} & kQ/I_0
 \end{array}$$

where  $kQ/I_0 \xrightarrow{\sim} kQ/I_0$  is an isomorphism of  $k$ -algebras, restricting to the identity map on the set  $Q_0$  of vertices and where  $\widehat{\mathcal{C}} \rightarrow \mathcal{C}$  is a Galois covering with group  $N$  a normal subgroup of  $\pi_1(Q)$  such that there exists an exact sequence of groups:  $1 \rightarrow N \rightarrow \pi_1(Q) \rightarrow G \rightarrow 1$ .

As said before, the above theorem is interesting because it gives a class of algebras for which a universal Galois covering exists and which is not defined using double bypasses. Moreover, the proof of Theorem 1 allows us to prove, under the same assumptions, that the quiver  $\Gamma$  of the homotopy relations of  $A$  (see [9, Def. 2.4]) has a unique source. Recall that the unicity of the source of  $\Gamma$  was proved in [9] for algebras without double bypass over a zero characteristic field. In order to prove Theorem 1, we will compare the fundamental groups of the admissible presentations of  $kQ/I_0$  using the fact that  $I_0$  is monomial. More precisely, we will use the arguments of the proof of [9, Thm. 2]. This proof essentially uses the fact that for an admissible presentation  $kQ/I \simeq kQ/I_0$ , there exists a sequence of  $\varphi_n, \dots, \varphi_1$  of transvections of  $kQ$  such that:

- (i)  $I = \varphi_n \dots \varphi_1(I_0)$ ,
- (ii) if we set  $I_i := \varphi_i \dots \varphi_1(I_0)$ , then  $\pi_1(Q, I_i)$  is either isomorphic to or a quotient of  $\pi_1(Q, I_{i-1})$ .

Recall ([9, Sect. 1]) that a transvection is an automorphism  $\varphi_{\alpha, u, \tau}$  of  $kQ$  defined by a bypass  $(\alpha, u)$  and a scalar  $\tau \in k$ . In order to show the existence of the above sequence of transvections, we will prove the following facts:

1. There exists a suitable ordering on the set of bypasses such that if  $\psi \in \text{Aut}(kQ)$  is a product of transvections, then  $\psi$  can be written uniquely as  $\psi = \varphi_{\alpha_n, u_n, \tau_n} \dots \varphi_{\alpha_1, u_1, \tau_1}$  with  $\tau_1, \dots, \tau_n \in k^*$  and  $(\alpha_n, u_n) > \dots > (\alpha_1, u_1)$ .
2. If  $kQ/I \simeq kQ/I_0$  then there exists a unique product of transvections  $\psi_I$  verifying simple technical conditions and such that  $\psi_I(I_0) = I$  (this step will intensively use the fact that  $I_0$  is monomial).
3. If  $kQ/I \simeq kQ/I_0$ , the unique ordered sequence of transvection given by 1. and whose product equal  $\psi_I$  verify the conditions (i) and (ii) above.

The second step is particularly interesting. Indeed, in general, if  $I_0$  and  $I$  are admissible ideals (without assuming monomiality) such that  $kQ/I_0 \simeq kQ/I$ , then, there are various automorphisms  $\psi \in \text{Aut}(kQ)$  such that  $\psi(I) = I$ , whereas in the present situation, the monomiality of  $I_0$  gives us a distinguished and uniquely characterised automorphism  $\psi_I$  such that  $\psi_I(I_0) = I$ .

The text is organised as follows. In Section 1 we recall all the notions that we will need to prove Theorem 1. In Section 2, we will prove some combinatoric facts on the paths in a quiver. These will lead to the order and to the decomposition of the first step above. In Section 3 we will prove the second step above. Finally, in Section 4 we will prove the last step and Theorem 1.

## 1 Basic definitions

A  **$k$ -category** is a category  $\mathcal{C}$  whose objects class  $\mathcal{C}_0$  is a set, whose space of morphisms from  $x$  to  $y$  (denoted by  ${}_y\mathcal{C}_x$ ) is a  $k$ -vector space for any  $x, y \in \mathcal{C}_0$  and whose composition of morphisms is  $k$ -bilinear. All functors between  $k$ -categories will be assumed to be  $k$ -linear functors. In particular,  $\text{Aut}(\mathcal{C})$  will denote the group of  $k$ -linear automorphism of  $\mathcal{C}$ , and we shall denote by  $\text{Aut}_0(\mathcal{C})$  for the subgroup  $\{\psi \in \text{Aut}(\mathcal{C}) \mid \psi(x) = x \text{ for any } x \in \mathcal{C}_0\}$  of  $\text{Aut}(\mathcal{C})$ . Let  $\mathcal{C}$  be a  $k$ -category,  $\mathcal{C}$  is called **connected** if  $\mathcal{C}$  cannot be written as the disjoint union of two full subcategories. An ideal  $I$  of  $\mathcal{C}$  is the data of subspaces  ${}_yI_x \subseteq {}_y\mathcal{C}_x$  (for any  $x, y \in \mathcal{C}_0$ ) such that  $fgh \in I$  whenever  $f, g, h$  are composable morphisms in  $\mathcal{C}$  such that  $g \in I$ . We say that  $\mathcal{C}$  is **locally bounded** provided that: 1) for any  $x \in \mathcal{C}_0$ , the vector spaces  $\bigoplus_{y \in \mathcal{C}_0} {}_y\mathcal{C}_x$

and  $\bigoplus_{y \in \mathcal{C}_0} {}_x \mathcal{C}_y$  are finite dimensional, 2)  ${}_x \mathcal{C}_x$  is a local algebra for any  $x \in \mathcal{C}_0$ , 3) distinct objects are not isomorphic. Let  $A$  be a finite dimensional  $k$ -algebra and let  $\{e_1, \dots, e_n\}$  be a complete set of primitive orthogonal idempotents. Then  $A$  is also a  $k$ -category as follows:  $A_0 := \{e_1, \dots, e_n\}$ ,  $e_i A e_i := e_j A e_i$  and the composition of morphisms is induced by the product in  $A$ . Notice that different choices for the idempotents  $e_1, \dots, e_n$  give rise to isomorphic  $k$ -categories. With this setting, the  $k$ -algebra  $A$  is connected (resp. basic) if and only if the  $k$ -category  $A$  is connected (resp. locally bounded) as a  $k$ -category. In the sequel we shall make no distinction between a finite dimensional  $k$ -algebra and its associated  $k$ -category. If  $\mathcal{C}$  is a locally bounded  $k$ -category, the radical of  $\mathcal{C}$  is the ideal  $\mathcal{RC}$  of  $\mathcal{C}$  such that:  ${}_y \mathcal{RC}_x$  is the space of non-isomorphisms  $x \rightarrow y$  in  $\mathcal{C}$ , for any  $x, y \in \mathcal{C}_0$ . The ideal of  $\mathcal{C}$  generated by compositions  $gf$  where  $f$  and  $g$  lie in  $\mathcal{RC}$  will be denoted by  $\mathcal{R}^2 \mathcal{C}$ .

A **Galois covering with group**  $G$  of  $\mathcal{C}$  (by  $\mathcal{C}'$ ) is a functor  $F: \mathcal{C}' \rightarrow \mathcal{C}$  endowed with a group morphism  $G \rightarrow \text{Aut}(\mathcal{C}')$  and such that: 1) the induced action of  $G$  on  $\mathcal{C}'_0$  is free, 2)  $F \circ g = F$  for any  $g \in G$ , 3) for any  $k$ -linear functor  $F': \mathcal{C}' \rightarrow \mathcal{C}''$  such that  $F' \circ g = F'$  for any  $g \in G$ , there exists a unique  $\overline{F'}: \mathcal{C} \rightarrow \mathcal{C}''$  such that  $\overline{F'} \circ F = F'$  (in other words,  $F$  is a quotient of  $\mathcal{C}'$  by  $\mathcal{C}$  in the category of  $k$ -categories). We shall say that  $F$  is connected if  $\mathcal{C}'$  is connected and locally bounded (this implies that  $\mathcal{C}$  is connected and locally bounded). For more details on Galois coverings (in particular for the connections with representations theory), we refer the reader to [3].

**Quivers, paths, bypasses.** A quiver is a 4-tuple  $Q = (Q_1, Q_0, s, t)$  where  $Q_1$  and  $Q_0$  are sets and  $s, t: Q_1 \rightarrow Q_0$  are maps. The elements of  $Q_1$  (resp. of  $Q_0$ ) are called the arrows (resp. the vertices) of  $Q$ . If  $\alpha \in Q_1$ , the vertex  $s(\alpha)$  (resp.  $t(\alpha)$ ) is called the source (resp. the target) of  $\alpha$ . The quiver  $Q$  is called **locally finite** if and only if any vertex is the source point (resp. the target) of finitely many arrows. For example, if  $\mathcal{C}$  is a locally bounded  $k$ -category, the **ordinary quiver of  $\mathcal{C}$**  is the locally finite quiver  $Q$  such that:  $Q_0 := \mathcal{C}_0$  and for any  $x, y \in \mathcal{C}_0$ , the number of arrows starting at  $x$  and arriving at  $y$  is equal to  $\dim_k {}_y \mathcal{RC}_x / {}_y \mathcal{R}^2 \mathcal{C}_y$ . A path in  $Q$  of length  $n$  ( $n \geq 0$ ) with source  $x \in Q_0$  (or starting at  $x$ ) and target  $y \in Q_0$  (or arriving at  $y$ ) is a sequence of arrows  $\alpha_1, \dots, \alpha_n$  such that:  $x = y$  if  $n = 0$ ,  $s(\alpha_1) = x$ ,  $s(\alpha_{i+1}) = t(\alpha_i)$  for any  $i \in \{1, \dots, n-1\}$  and  $t(\alpha_n) = y$ . If  $n \geq 1$  this path will be written  $\alpha_n \dots \alpha_1$  and called non trivial. If  $n = 0$  this path will be written  $e_x$  and called stationary at  $x$ . The length of this path is  $|u| := n$ . The mappings  $s, t$  are naturally extended to paths in  $Q$ . If  $u$  and  $v$  are paths, the concatenation  $vu$  is defined if and only if  $t(u) = s(v)$  by the following rule: 1)  $vu = v$  is  $u$  is stationary, 2)  $vu = u$  is  $v$  is stationary, 3)  $vu = \beta_m \dots \beta_1 \alpha_n \dots \alpha_1$  if  $v = \beta_m \dots \beta_1$  and  $u = \alpha_n \dots \alpha_1$  (with  $\alpha_i, \beta_j \in Q_1$ ). Two paths in  $Q$  are called **parallel** whenever they have the same source and the same target. An **oriented cycle** in  $Q$  is a non trivial path whose source and target are equal. We shall say that  $Q$  has **multiple arrows** if and only if there exist in  $Q$  distinct parallel arrows. A **bypass** in  $Q$  is a couple  $(\alpha, u)$  where  $\alpha \in Q_1$  and where  $u$  is a path parallel to and different from  $\alpha$ . A **double bypass** in  $Q$  is 4-tuple  $(\alpha, u, \beta, v)$  where  $(\alpha, u)$  and  $(\beta, v)$  are bypasses and such that the arrow  $\beta$  appears in the path  $u$ . In other words, there exists paths  $u_1, u_2$  such that  $u = u_2 \beta u_1$ . When  $Q$  has no oriented cycle, the paths  $u_1, u_2$  are unique for this property, and the path  $u_2 v u_1$  will be called obtained from  $u = u_2 \beta u_1$  after replacing  $\beta$  by  $v$ .

**Admissible presentations (see [3, 2.1]).** A quiver  $Q$  defines the **path category**  $kQ$  such that  $(kQ)_0 = Q_0$ , such that  ${}_y kQ_x$  is the  $k$ -vector space with basis the family of paths starting at  $x$  and arriving at  $y$ , and the composition in  $kQ$  is induced by the concatenation of paths. If  $r \in {}_y kQ_x$ , a **normal form** for  $r$  is an equality  $r = \sum_{i=1}^n t_i u_i$  where  $t_1, \dots, t_n \in k^*$  and  $u_1, \dots, u_n$  are pairwise distinct paths in  $Q$ . With this notation, the **support** of  $r$  is the set  $\text{supp}(r) := \{u_1, \dots, u_n\}$  (with the convention  $\text{supp}(0) = \emptyset$ ). A **subexpression** of  $r$  is a linear combination  $\sum_{i \in E} t_i u_i$  with  $E \subseteq \{1, \dots, n\}$ . Later, we will need the following fact: if  $r = r_1 + \dots + r_n \in {}_y kQ_x$  is such that  $\text{supp}(r_1), \dots, \text{supp}(r_n)$  are pairwise disjoint, then  $r_{i_1} + \dots + r_{i_t}$  is a subexpression of  $r$ , for any indices  $1 \leq i_1 < \dots < i_t \leq n$ . An ideal  $I$  of  $kQ$  is called **admissible** provided that: 1) any morphism in  $I$  is a linear combination of paths of length at least 2, 2) for any vertex  $x$ , there exists an integer  $n$  such that any path starting or arriving at  $x$  and of length greater than  $n$  lies in  $I$ . A morphism in  $I$  is called a **relation** (of  $I$ ). In particular, a **minimal relation of  $I$**  (see [11]) is a non zero relation  $r$  of  $I$  such that 0 and  $r$  are the only subexpressions of  $r$  which are relations. With this definition, any relation of  $I$  is the sum of minimal relations with pairwise disjoint supports. If  $u \in I$  is a path, then  $u$  called a **monomial relation**. In particular,  $I$  is called monomial if it is generated by a set of monomial relations. A pair  $(Q, I)$  where  $Q$  is a locally finite and  $I$  is an admissible ideal of  $kQ$  is called a **bound quiver**. In such a case,  $kQ/I$  is a locally bounded  $k$ -category and is connected if and only if  $Q$  is connected (i.e. the underlying graph of  $Q$  is connected). Conversely,

if  $\mathcal{C}$  is a locally bounded  $k$ -category, then there exists an isomorphism  $kQ/I \xrightarrow{\sim} \mathcal{C}$  where  $(Q, I)$  is a bound quiver such that  $Q$  is the ordinary quiver of  $\mathcal{C}$ . Such an isomorphism is called **admissible presentation of  $\mathcal{C}$** . If the ideal  $I$  is monomial, the admissible presentation and  $\mathcal{C}$  are called monomial. Notice that  $\mathcal{C}$  may have different admissible presentations.

**Fundamental group of a presentation (see [11]).** Let  $(Q, I)$  be a bound quiver and let  $x_0 \in Q_0$ . For every arrow  $x \xrightarrow{a} y \in Q_1$  we define its formal inverse  $a^{-1}$  with source  $s(a^{-1}) = y$  and target  $t(a^{-1}) = x$ . With these notations, a **walk in  $Q$  with source  $x$  and target  $y$**  is a sequence  $a_1, \dots, a_n$  (with  $n \geq 0$ ) of arrows and formal inverses of arrows such that  $s(a_i) = t(a_{i-1})$  for any  $i$ . If  $n > 0$ , this walk is denoted by  $a_n \dots a_1$ . Hence paths are particular cases of walks and the concatenation of paths extends naturally to walks. The **homotopy relation** of  $(Q, I)$  is the equivalence relation on the set of walks in  $Q$ , denoted by  $\sim_I$  and generated by the following properties:

1.  $\alpha\alpha^{-1} \sim_I e_y$  and  $\alpha^{-1}\alpha \sim_I e_x$  for any arrow  $x \xrightarrow{\alpha} y$  in  $Q$ ,
2.  $u \sim_I v$  for any  $u, v \in \text{supp}(r)$  where  $r$  is a minimal relation of  $I$ ,
3.  $wvu \sim_I wv'u$  for any walks  $w, v, v', u$  such that  $v \sim_I v'$  and such that the concatenations  $wvu$  and  $wv'u$  are well-defined (i.e.  $\sim_I$  is compatible with the concatenation).

The  $\sim_I$ -equivalence class of a walk  $\gamma$  will be denoted by  $[\gamma]_I$ . Let  $\pi_1(Q, I, x_0)$  be the set of equivalence classes of walks in  $Q$  with source and target equal to  $x_0$ . The concatenation of walks endows this set with a group structure (with unit  $e_{x_0}$ ) and this group is called the **fundamental group of  $(Q, I)$** . If  $Q$  is connected, the isomorphism class of this group does not depend on  $x_0 \in Q_0$  and  $\pi_1(Q, I, x_0)$  is denoted by  $\pi_1(Q, I)$ . If  $\mathcal{C}$  is a connected locally bounded  $k$ -category and if  $kQ/I \simeq \mathcal{C}$  is an admissible presentation, the fundamental group  $\pi_1(Q, I)$  is called the fundamental group of this presentation.

**Dilatations, transvections (see [9]).** Let  $Q$  be a quiver. A **dilatation** of  $kQ$  is an automorphism  $D \in \text{Aut}_0(kQ)$  such that  $D(\alpha) \in k^*\alpha$  for any  $\alpha \in Q_1$ . The dilatations of  $kQ$  form a subgroup  $\mathcal{D}$  of  $\text{Aut}_0(kQ)$ . Let  $(\alpha, u)$  be a bypass in  $Q$  and let  $\tau \in k$ . This defines  $\varphi_{\alpha, u, \tau} \in \text{Aut}_0(kQ)$  as follows:  $\varphi_{\alpha, u, \tau}(\alpha) = \alpha + \tau u$  and  $\varphi_{\alpha, u, \tau}(\beta) = \beta$  for any arrow  $\beta \neq \alpha$ . The automorphism  $\varphi_{\alpha, u, \tau}$  is called a **transvection**. The composition of transvections is ruled as follows. Let  $\varphi_{\alpha, u, \tau}$  and  $\varphi_{\alpha, u, \tau'}$ , then  $\varphi_{\alpha, u, \tau}\varphi_{\alpha, u, \tau'} = \varphi_{\alpha, u, \tau+\tau'}$  and  $\varphi_{\alpha, u, \tau}^{-1} = \varphi_{\alpha, u, -\tau}$ . If  $(\alpha, u, \beta, v)$  and  $(\beta, v, \alpha, u)$  are not a double bypasses, then  $\varphi_{\alpha, u, \tau}\varphi_{\beta, v, \nu} = \varphi_{\beta, v, \nu}\varphi_{\alpha, u, \tau}$ . If  $(\alpha, u, \beta, v)$  is a double bypass and if  $Q$  has no oriented cycle, then  $\varphi_{\beta, v, \nu}\varphi_{\alpha, u, \tau} = \varphi_{\alpha, u, \tau}\varphi_{\alpha, w, \tau\nu}\varphi_{\beta, v, \nu}$ , where  $w$  is the path obtained from  $u$  after replacing  $\beta$  by  $v$ . The subgroup of  $\text{Aut}_0(kQ)$  generated by all the transvections is denoted by  $\mathcal{T}$ . Dilatations and transvections are useful to compare different admissible presentations of the same locally bounded  $k$ -category because of the following proposition:

**Proposition 1.1.** (see [9, Prop. 2.1, Prop. 2.2]) *Let  $kQ/I \simeq A$  and  $kQ/J \simeq A$  be admissible presentations of the basic finite dimensional algebra  $A$ . If  $Q$  has no oriented cycle, then there exists  $\psi \in \text{Aut}_0(kQ)$  such that  $\psi(I) = J$ . Moreover,  $\mathcal{T}$  is a normal subgroup of  $\text{Aut}_0(kQ)$  and  $\text{Aut}_0(kQ) = \mathcal{D}\mathcal{T} = \mathcal{D}\mathcal{T}$ .*

The dilatations and the transvections were introduced because they allow comparisons between the fundamental groups of presentations of the same locally bounded  $k$ -category:

**Proposition 1.2.** (see [9, Prop. 2.5]) *Let  $I$  be an admissible ideal of  $kQ$ , let  $\varphi \in \text{Aut}_0(kQ)$  and set  $J = \varphi(I)$ . If  $\varphi$  is a dilatation, then  $\sim_I$  and  $\sim_J$  coincide. If  $\varphi = \varphi_{\alpha, u, \tau}$  is a transvection, then:*

1. if  $\alpha \sim_I u$  and  $\alpha \sim_J u$  then  $\sim_I$  and  $\sim_J$  coincide.
2. if  $\alpha \not\sim_I u$  and  $\alpha \sim_J u$  then  $\sim_J$  is generated by  $\sim_I$  and  $\alpha \sim_J u$ .
3. if  $\alpha \not\sim_I u$  and  $\alpha \not\sim_J u$  then  $I = J$  and  $\sim_I$  and  $\sim_J$  coincide.

*If there exists a transvection  $\varphi$  such that  $\varphi(I) = J$  and such the second point above occurs, then we shall say that  $\sim_J$  is a direct successor of  $\sim_I$ .*

Here the expression “ $\sim_I$  is generated by  $\sim_J$  and  $\alpha \sim_I u$ ” means that  $\sim_I$  is the equivalence relation on the set of walks in  $Q$ , compatible with the concatenation and generated by the two following properties: 1)  $\gamma \sim_J \gamma' \Rightarrow \gamma \sim_I \gamma'$ , 2)  $\alpha \sim_I u$ . Following [9, Def. 2.7], if  $A$  is a basic connected finite dimensional algebra with ordinary quiver  $Q$  without oriented cycle, we define the **quiver  $\Gamma$  of the homotopy relations of  $A$**  to be the quiver such that  $\Gamma_0 = \{\sim_I \mid kQ/I \simeq A\}$  and such that there exists arrow  $\sim_I \rightarrow \sim_J$  if and only if  $\sim_J$  is a direct successor of  $\sim_I$ . Recall ([9, Rem. 5, Prop. 2.8]) that  $\Gamma$  is finite, connected, without oriented cycle and such that for any oriented path with source  $\sim_I$  and target  $\sim_J$ , the identity map on the walks in  $Q$  induces a surjective group morphism  $\pi_1(Q, I) \twoheadrightarrow \pi_1(Q, J)$ .

**Gröbner bases** Let  $E$  be a  $k$ -vector space with an ordered basis  $(e_1, \dots, e_n)$ , let  $(e_1^*, \dots, e_n^*)$  be the associated dual basis of  $E^*$ , and let  $F$  be a subspace of  $E$ . A Gröbner basis (see [1] for the usual definition) of  $F$  is a basis  $(r_1, \dots, r_d)$  such that:

1.  $r_j \in e_{i_j} + \text{Span}(e_l ; l < i_j)$  for some  $i_j$ , for any  $j \in \{1, \dots, r\}$ ,
2.  $i_1 < i_2 < \dots < i_r$ ,
3.  $e_{i_j}^*(r_{j'}) = 0$  for any  $j \neq j'$ .

It is well known that  $F$  admits a unique Gröbner basis. Also,  $r \in F$  if and only if:  $r = \sum_{j=1}^d e_{i_j}^*(r)r_j$ . In the sequel, we will use this notion in the following setting:  $E$  is the vector space with basis (for some order to be defined) the family of non trivial paths in a finite quiver  $Q$  without oriented cycles and  $F$  is the underlying subspace of  $E$  associated to an admissible ideal  $I$  of  $kQ$ . Notice that in this setting, if  $r \in E$  and if  $u$  is a non trivial path, then:  $u \in \text{supp}(r) \Leftrightarrow u^*(r) \neq 0$ .

## 2 Combinatorics on the paths in a quiver

### 2.1 Derivation of paths

**Definition 2.1.** Let  $u = \alpha_n \dots \alpha_1$  and  $v$  be paths in  $Q$ . Then  $v$  is called derived of  $u$  (of order  $t$ ) if there exist indices  $1 \leq i_1 < \dots < i_t \leq n$  and with bypasses  $(\alpha_{i_1}, v_1), \dots, (\alpha_{i_t}, v_t)$  such that  $v$  is obtained from  $u$  after replacing  $\alpha_{i_l}$  by  $v_l$  for each  $l$ :

$$v = \alpha_n \dots \alpha_{i_t+1} v_t \alpha_{i_t-1} \dots \alpha_{i_l+1} v_l \alpha_{i_l-1} \dots \alpha_{i_1+1} v_1 \alpha_{i_1-1} \dots \alpha_1$$

**Remark 2.2.** If  $\alpha \in Q_1$ , then  $u$  is derived of  $\alpha$  if and only if  $(\alpha, u)$  is a bypass.

With the above definition, the following lemma is easily verified using the fact that  $Q$  has no multiple arrows.

**Lemma 2.3.** 1. If  $v$  is derived of  $u$  with both orders  $t$  and  $t'$ , then  $t=t'$ .

2. If  $v$  is derived of  $u$  of order  $t$  then there exists a sequence of paths  $u_0 = u, u_1, \dots, u_t = v$  such that  $u_i$  is derived of  $u_{i-1}$  of order 1 for any  $i$ .
3. If  $v$  is derived of  $u$  of order  $t$ , then  $|v| \geq |u| + t$ .
4. If  $v$  is derived of  $u$  of order  $t$  and if  $w$  is derived of  $v$  of order  $t'$ , then  $w$  is derived of  $u$  of order at least  $t$ .
5. Let  $u, v, w$  be paths verifying:
  - $v$  is derived of  $u$ ,
  - $w$  is derived of  $v$ ,
  - $w$  is derived of  $u$  of order 1,

then we have:

$$u = u_2 \alpha u_1, \quad v = u_2 \theta u_1, \quad w = u_2 \theta' u_1$$

where  $u_1, u_2$  are paths,  $(\alpha, \theta)$  is a bypass and  $\theta'$  is derived of  $\theta$ .

6. If  $v$  (resp.  $v'$ ) is derived of  $u$  (resp. of  $u'$ ) of order  $t$  (resp.  $t'$ ), then  $v'v$  is derived of  $u'u$  of order  $t' + t$ , whenever these compositions of paths are well defined.

The following example shows that the inequality in the 4-th point of the preceding lemma may be an equality.

**Example 2.4.** Let  $(\alpha, u, \beta, v)$  be a double bypass. Let  $u_1, u_2$  be the paths such that  $u = u_2 \beta u_1$ . Then  $u$  is derived of  $\alpha$  of order 1,  $w := u_2 v u_1$  is derived of  $u$  of order 1 and  $w$  is derived of  $u$  of order 1.

### 2.2 Order between paths, order between bypasses

Now, we construct a total order on the set of non trivial paths in  $Q$ . This construction is a particular case of the one introduced in [5], also it depends on an arbitrary order  $\triangleleft$  on  $Q_1$ . We assume that this order  $\triangleleft$  is fixed for this subsection. We shall write  $\triangleleft$  for the lexicographical order induced by  $\triangleleft$  on the set of nontrivial paths in  $Q$ . For details on the correctness of the following definition we refer the reader to [5].

**Definition 2.5.** For  $\alpha \in Q_1$ , set:

$$W(\alpha) = \text{Card}(B(\alpha)) \quad \text{where} \quad B(\alpha) = \{(\alpha, u) \mid (\alpha, u) \text{ is a bypass in } Q\}$$

For  $u = \alpha_n \dots \alpha_1$  a path in  $Q$  (with  $\alpha_i \in Q_1$ ), let us set:

$$W(u) = W(\alpha_n) + \dots + W(\alpha_1)$$

These data define a total order  $<$  on the set of non trivial paths in  $Q$  as follows:

$$u < v \Leftrightarrow \begin{cases} W(u) < W(v) \\ \text{or} \\ W(u) = W(v) \text{ and } u \triangleleft v \end{cases}$$

We shall also write  $<$  for the lexicographical order induced by  $<$  on the set of couples of paths.

**Remark 2.6.** If  $u$  and  $v$  are (non trivial) paths such that  $vu$  is well defined, then  $W(vu) = W(u) + W(v)$ .

**Lemma 2.7.** 1. If  $u, v, u', v'$  are paths such that  $v < u$  and  $v' < u'$  then  $v'v < u'u$  whenever these compositions are well defined.

2. If  $(\alpha, u)$  is a bypass, then  $W(u) < W(\alpha)$ . So  $u < \alpha$ .

3. If  $v$  is derived of  $u$ , then  $v < u$ .

4. If  $(\alpha, u, \beta, v)$  is a double bypass and if  $w$  is the path obtained from  $u$  after replacing  $\beta$  by  $v$ , then:

$$(\beta, v) < (\alpha, w) < (\alpha, u)$$

**Proof:** 1) is a direct consequence of Definition 2.5 and Remark 2.6.

2) Let us write  $u = a_n \dots a_1$  with  $a_i \in Q_1$  for each  $i$  (hence  $a_i \neq a_j$  if  $i \neq j$  because  $Q$  has no oriented cycle). Therefore:

- .  $B(a_1), \dots, B(a_n)$  are pairwise disjoint,
- .  $W(u) = W(a_1) + \dots + W(a_n)$

Notice that if  $(a_i, v) \in B(a_i)$ , then  $(\alpha, a_n \dots a_{i+1}va_{i-1} \dots a_1) \in B(\alpha)$ . Thus, we have a well defined mapping:

$$\begin{aligned} \theta: \quad B(a_1) \sqcup \dots \sqcup B(a_n) &\longrightarrow B(\alpha) \\ (a_i, v) &\longmapsto (\alpha, a_n \dots a_{i+1}va_{i-1} \dots a_1) \end{aligned}$$

This mapping is one-to-one, indeed:

- . if  $\theta(a_i, v) = \theta(a_i, v')$  with  $(a_i, v), (a_i, v') \in B(a_i)$  then:

$$a_n \dots a_{i+1}va_{i-1} \dots a_1 = a_n \dots a_{i+1}v'a_{i-1} \dots a_1$$

and therefore  $(a_i, v) = (a_i, v')$ ,

- . if  $\theta(a_i, v) = \theta(a_j, v')$  with  $(a_i, v) \in B(a_i), (a_j, v') \in B(a_j)$  and  $j < i$ , then:

$$a_n \dots a_{i+1}va_{i-1} \dots a_1 = a_n \dots a_{j+1}v'a_{j-1} \dots a_1$$

So:

$$va_{i-1} \dots a_1 = a_i \dots a_{j+1}v'a_{j-1} \dots a_1$$

Since  $v$  and  $a_i$  are parallel and since  $Q$  has no oriented cycle, we infer that  $v = a_i$  which is impossible because  $(a_i, v) \in B(a_i)$ .

On the other hand,  $\theta$  is not onto. Indeed, if there exists  $(a_i, v) \in B(a_i)$  verifying  $\theta(a_i, v) = (\alpha, u)$ , then:

$$a_n \dots a_1 = u = a_n \dots a_{i+1}va_{i-1} \dots a_1$$

which implies  $a_i = v$ , a contradiction. Since  $\theta$  is one-to-one and not onto, we deduce that:

$$W(u) = \text{Card}(B(\alpha)) > \text{Card}(B(a_1) \sqcup \dots \sqcup B(a_n)) = W(u)$$

This proves that  $W(u) < W(\alpha)$  and that  $u < \alpha$ .

3) is a direct consequence of 1) and of 2).

4) Let us write  $u = u_2\beta u_1$  (with  $u_1, u_2$  paths) so that  $w = u_2vu_1$ . From 2), we have:

$$W(\alpha) > W(u) = W(u_1) + W(\beta) + W(u_2) \geq W(\beta)$$

So  $\beta < \alpha$  and therefore  $(\beta, v) < (\alpha, w)$ . Using 2) again, we also have:

$$W(w) = W(u_2) + W(v) + W(u_1) < W(u_2) + W(\beta) + W(u_1) = W(u)$$

So  $w < u$  and therefore  $(\alpha, w) < (\alpha, u)$  ■

## 2.3 Image of a path by a product of transvections

In this paragraph, we apply the previous constructions to find an easy way to compute  $\psi(u)$  when  $\psi \in \mathcal{T}$  and  $u$  is a path in  $Q$ . We begin with the following lemma on the description of  $\psi(\alpha)$  when  $\psi \in \mathcal{T}$  and  $\alpha \in Q_1$ . Recall that  $Q$  has no multiple arrows and no oriented cycles.

**Lemma 2.8.** *Let  $\psi \in \mathcal{T}$  and let  $\alpha \in Q_1$ . Then  $\psi(\alpha) - \alpha$  is a linear combination of paths parallel to  $\alpha$  and length greater than or equal to 2. In particular,  $\alpha \in \text{supp}(\psi(\alpha))$  and  $\alpha^*(\psi(\alpha)) = 1$ .*

**Proof:** The conclusion is immediate if  $\psi$  is a transvection because  $Q$  has no multiple arrows. The conclusion in the general case is obtained using an easy induction on the number of transvections whose product equal  $\psi$ . ■

The preceding lemma gives the following description of  $\psi(u)$  when  $\psi \in \mathcal{T}$  and  $u$  is a path. We omit the proof which is immediate thanks to Lemma 2.8 and to point 6) of Lemma 2.3.

**Proposition 2.9.** *Let  $\psi \in \mathcal{T}$  and let  $u = \alpha_n \dots \alpha_1$  be a path in  $Q$  (with  $a_i \in Q_1$  for any  $i$ ). For each  $i$ , let:*

$$\psi(\alpha_i) = \alpha_i + \sum_{j=1}^{m_i} \lambda_{i,j} u_{i_j}$$

*be a normal form for  $\psi(\alpha_i)$ . Then  $\text{supp}(\psi(u))$  is the set of the paths in  $Q$  described as follows. Let  $r \in \{0, \dots, n\}$ , let  $1 \leq i_1 < \dots < i_r \leq n$  be indices, for each  $l \in \{1, \dots, r\}$  let  $j_l \in \{1, \dots, m_{i_l}\}$ , then the following path obtained from  $u$  after replacing  $\alpha_{i_l}$  by  $u_{j_l}$  for each  $l$  belongs to  $\text{supp}(\psi(u))$ :*

$$\alpha_n \dots \alpha_{i_r+1} u_{j_r} \alpha_{i_r-1} \dots \alpha_{i_l+1} u_{j_l} \alpha_{i_l-1} \dots \alpha_{i_1+1} u_{j_1} \alpha_{i_1-1} \dots \alpha_1$$

*Moreover, this path appears in  $\psi(u)$  with coefficient:*

$$\lambda_{i_1, j_1} \dots \lambda_{i_r, j_r}$$

*As a consequence,  $\psi(u) - u$  is a linear combination of paths derived of  $u$ .*

**Remark 2.10.** *If  $(\alpha, u)$  is a bypass and if  $v \in \text{supp}(\psi(u) - u)$ , then  $(\alpha, v)$  is also a bypass and  $(\alpha, v) < (\alpha, u)$ .*

Now we are able to state the main result of this paragraph. It describes  $\psi(\alpha)$  ( $\alpha \in Q_1$ ) using a particular writing of  $\psi$  as a product of transvections.

**Proposition 2.11.** *Let  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n)$  be an increasing sequence of bypasses, let  $\tau_1, \dots, \tau_n \in k^*$  and set  $\psi = \varphi_{\alpha_n, u_n, \tau_n} \dots \varphi_{\alpha_1, u_1, \tau_1}$ . For any  $\alpha \in Q_1$ , there is a normal form for  $\psi(\alpha)$ :*

$$\psi(\alpha) = \alpha + \sum_{i \text{ such that } \alpha = \alpha_i} \tau_i u_i$$

**Proof:** Let us prove that the conclusion of the proposition is true using an induction on  $n \geq 1$ . By definition of a transvection, the proposition holds of  $n = 1$ . Assume that  $n \geq 2$  and that the conclusion of the proposition holds if we replace  $\psi = \varphi_{\alpha_n, u_n, \tau_n} \dots \varphi_{\alpha_1, u_1, \tau_1}$  by  $\varphi_{\alpha_{n-1}, u_{n-1}, \tau_{n-1}} \dots \varphi_{\alpha_1, u_1, \tau_1}$ . Therefore, for  $\alpha \in Q_1$ , we have a normal form:

$$\varphi_{\alpha_{n-1}, u_{n-1}, \tau_{n-1}} \dots \varphi_{\alpha_1, u_1, \tau_1}(\alpha) = \alpha + \sum_{i \leq n-1, \alpha = \alpha_i} \tau_i u_i$$

So:

$$\psi(\alpha) = \varphi_{\alpha_n, u_n, \tau_n}(\alpha) + \sum_{i \leq n-1, \alpha = \alpha_i} \tau_i \varphi_{\alpha_n, u_n, \tau_n}(u_i) \quad (i)$$

Let  $i \in \{1, \dots, n-1\}$ . Thanks to Lemma 2.7, the inequality  $(\alpha_i, u_i) < (\alpha_n, u_n)$  implies that  $(\alpha_i, u_i, \alpha_n, u_n)$  is not a double bypass. Thus,  $\alpha_n$  does not appear in the path  $u_i$ . This proves that:

$$(\forall i \in \{1, \dots, n-1\}) \varphi_{\alpha_n, u_n, \tau_n}(u_i) = u_i \quad (ii)$$

The definition of  $\varphi_{\alpha_n, u_n, \tau_n}$ , together with (i) and (ii), imply the equality:

$$\psi(\alpha) = \alpha + \sum_{\alpha = \alpha_i} \tau_i u_i \quad (iii)$$

It only remains to prove that the equality (iii) is a normal form. Remark that all the scalars which appear in the right-hand side of (iii) are non zero. Moreover, if  $i \in \{1, \dots, n\}$  verifies  $\alpha = \alpha_i$ , then  $\alpha \neq u_i$ , because



$(\alpha, u_i)$  is a bypass. Finally, if  $1 \leq i < j \leq n$  verify  $\alpha = \alpha_i = \alpha_j$ , then  $(\alpha, u_i) = (\alpha_i, u_i) < (\alpha_j, u_j) = (\alpha, u_j)$  so  $u_i \neq u_j$ . Therefore, (iii) is a normal form for  $\psi(\alpha)$ . ■

When  $\psi \in \mathcal{T}$  is like in Proposition 2.11, we shall say that  $\psi$  is written as a decreasing product of transvections. Later we will prove that any  $\psi \in \mathcal{T}$  can be written uniquely as a decreasing product of transvections. The description in Proposition 2.11 will be particularly useful in the sequel. We end this paragraph with two propositions concerning the description of  $\psi(r)$  when  $\psi \in \mathcal{T}$  and  $r$  is a linear combination of paths. The following proposition gives conditions for  $\psi^{-1}(r')$  to be a subexpression of  $r$  when  $r'$  is a subexpression of  $\psi(r)$ .

**Proposition 2.12.** *Let  $\psi \in \mathcal{T}$ , let  $r \in {}_y kQ_x$  and let  $r'$  be a subexpression of  $\psi(r)$ . Let  $\simeq$  be the equivalence relation on the set of paths in  $Q$  generated by:*

$$v \in \text{supp}(\psi(u)) \Rightarrow u \simeq v$$

*Assume that for any  $u, v \in \text{supp}(\psi(r))$  verifying  $u \simeq v$  we have:*

$$u \in \text{supp}(r') \Leftrightarrow v \in \text{supp}(r')$$

*Then  $\psi^{-1}(r')$  is a subexpression of  $r$ .*

**Proof:** Let  $\simeq'$  be the trace of  $\simeq$  on  $\text{supp}(r)$  and let us write  $\text{supp}(r) = c_1 \sqcup \dots \sqcup c_n$  as a disjoint union of its  $\simeq'$ -classes. This partition of  $\text{supp}(r)$  defines a decomposition of  $r = r_1 + \dots + r_n$  where  $r_i$  is the subexpression of  $r$  verifying  $\text{supp}(r_i) = c_i$ . For each  $i$ , let us fix a normal form:

$$r_i = \sum_{j=1}^{n_i} t_{i,j} u_{i,j}$$

so that we have the following normal form for  $r$ :

$$r = \sum_{i=1}^n \sum_{j=1}^{n_i} t_{i,j} u_{i,j}$$

Let us set  $r'_i := \psi(r_i)$ . In order to prove that  $\psi^{-1}(r')$  is a subexpression of  $r$ , we will prove that there exist indices  $1 \leq i_1 < \dots < i_t \leq n$  verifying  $r' = r'_{i_1} + \dots + r'_{i_t}$  (so that  $\psi^{-1}(r') = r'_{i_1} + \dots + r'_{i_t}$ ). In this purpose, we will successively prove the following facts:

- 1)  $u, v \in \text{supp}(r'_i) \Rightarrow u \simeq v$ , for any  $i$ ,
- 2)  $\text{supp}(r'_1), \dots, \text{supp}(r'_n)$  are pairwise disjoint,
- 3) for each  $i$ ,  $r'_i$  is a subexpression of  $\psi(r)$ ,
- 4) if  $i \in \{1, \dots, n\}$  verifies  $\text{supp}(r') \cap \text{supp}(r'_i) \neq \emptyset$ , then  $\text{supp}(r'_i) \subseteq \text{supp}(r')$ ,

1) Let  $i \in \{1, \dots, n\}$  and let  $u, v \in \text{supp}(r'_i)$ . So there exist  $u', v' \in \text{supp}(r_i)$  such that  $u \in \text{supp}(\psi(u'))$  and  $v \in \text{supp}(\psi(v'))$ . By definition of  $\simeq$  and of  $r_i$ , we deduce that:

$$u, v \in \text{supp}(r'_i) \Rightarrow u \simeq v \tag{i}$$

2) Let  $i, j \in \{1, \dots, n\}$  be such that there exists  $v \in \text{supp}(r'_i) \cap \text{supp}(r'_j)$ . So there exist  $u \in \text{supp}(r_i)$  and  $u' \in \text{supp}(r_j)$  such that  $v \in \text{supp}(\psi(u))$  and  $v \in \text{supp}(\psi(u'))$ . This implies that  $u \simeq v \simeq u'$ . Since  $u \in c_i = \text{supp}(r_i)$  and  $u' \in c_j = \text{supp}(r_j)$ , we deduce that  $c_i = c_j$  and therefore  $i = j$ . So:

$$i \neq j \Rightarrow \text{supp}(r'_i) \cap \text{supp}(r'_j) = \emptyset \tag{ii}$$

3) We have  $\psi(r) = r'_1 + \dots + r'_n$  so (ii) implies that:

$$r'_i \text{ is a subexpression of } \psi(r) \text{ for any } i \tag{iii}$$

4) Let  $i \in \{1, \dots, n\}$  and assume that there exists  $u \in \text{supp}(r'_i) \cap \text{supp}(r')$ . If  $v \in \text{supp}(r'_i)$  then  $u \simeq v$  thanks to (i). So, by assumption on  $r'$ , we have  $v \in \text{supp}(r')$ . This proves that:

$$\text{supp}(r'_i) \cap \text{supp}(r') \neq \emptyset \Rightarrow \text{supp}(r'_i) \subseteq \text{supp}(r') \tag{iv}$$

Now, we can prove that  $\psi^{-1}(r')$  is a subexpression of  $r$ . Thanks to (iii), the elements  $r', r'_1, \dots, r'_n$  are subexpressions of  $\psi(r)$ . So (iv) and the equality  $\psi(r) = r'_1 + \dots + r'_n$  imply that there exist indices  $1 \leq i_1 < \dots < i_t \leq n$  such that  $r' = r'_{i_1} + \dots + r'_{i_t}$ . So  $\psi^{-1}(r') = r_{i_1} + \dots + r_{i_t}$ . This proves that  $\psi^{-1}(r')$  is a subexpression of  $r$ . ■

The last proposition of this subsection gives a sufficient condition on  $u \in \text{supp}(r)$  to verify  $u \in \text{supp}(\psi(r))$ .

**Proposition 2.13.** *Let  $\psi \in \mathcal{T}$ , let  $r \in {}_y k Q_x$  and let  $u \in \text{supp}(r)$ . Then, at least one of the two following facts is verified:*

- .  $u \in \text{supp}(\psi(r))$ ,
- . there exists  $v \in \text{supp}(r)$  such that  $u \neq v$  and such that  $v \in \text{supp}(\psi(u))$ .

As a consequence, if  $u$  is not derived of  $v$  for any  $v \in \text{supp}(r)$ , then:

$$u \in \text{supp}(\psi(r)) \quad \text{and} \quad u^*(\psi(r)) = u^*(r)$$

**Proof:** Let us fix a normal form  $r = \sum_{i=1}^n t_i u_i$  where we may assume that  $u = u_1$ . Let us assume that  $u \notin \text{supp}(\psi(r))$ , i.e.  $u^*(\psi(r)) = 0$ . Recall from Proposition 2.9 that  $u^*(\psi(u)) = 1$ , so:

$$0 = u^*(\psi(r)) = t_1 + \sum_{i=2}^n t_i u^*(\psi(u_i)) \quad (i)$$

Therefore, there exists  $i_0 \in \{2, \dots, n\}$  such that  $u^*(\psi(u_{i_0})) \neq 0$ . So:

$$u_{i_0} \in \text{supp}(r), \quad u_{i_0} \neq u_1 = u \quad \text{and} \quad u_1^*(\psi(u_{i_0})) \neq 0$$

This proves the first assertion of the proposition. Now let us assume that  $u$  is not derived of  $v$  for any  $v \in \text{supp}(r)$ . Let  $i \in \{2, \dots, n\}$ . Since  $u = u_1 \neq u_i$ , Proposition 2.9 gives the following implications:

$$u \in \text{supp}(\psi(u_i)) \Rightarrow u \in \text{supp}(\psi(u_i) - u_i) \Rightarrow u \text{ is derived of } u_i$$

By assumption on  $u$ , this implies that  $u^*(\psi(u_i)) = 0$  for any  $i \geq 2$ . Using (i), we deduce the announced conclusion:  $u^*(\psi(r)) = t_1 = u^*(r) \neq 0$  ■

## 2.4 Ordering products of transvections

In Proposition 2.11 we have seen that  $\psi(\alpha)$  may be easily computed when  $\psi \in \mathcal{T}$  and  $\alpha \in Q_1$  provided that  $\psi$  is written as a decreasing product of transvections. In this subsection, we will prove that any  $\psi \in \mathcal{T}$  can be uniquely written that way. Recall that  $<$  is an order on the set of non trivial paths in  $Q$  defined in Definition 2.5. We introduce first some notations.

**Definition 2.14.** *Let  $(\alpha, u)$  be a bypass. We set  $\mathcal{T}_{<(\alpha, u)}$  and  $\mathcal{T}_{\leq(\alpha, u)}$  to be the subgroups of  $\mathcal{T}$  generated by the following sets of transvections:*

$$\begin{aligned} & \{\varphi_{\beta, v, \tau} \mid (\beta, v) < (\alpha, u) \text{ and } \tau \in k\} \quad \text{for } \mathcal{T}_{<(\alpha, u)} \\ & \{\varphi_{\beta, v, \tau} \mid (\beta, v) \leq (\alpha, u) \text{ and } \tau \in k\} \quad \text{for } \mathcal{T}_{\leq(\alpha, u)} \end{aligned}$$

Also, we define  $\mathcal{T}_{(\alpha, u)}$  to be the following subgroup of  $\mathcal{T}$ :

$$\mathcal{T}_{(\alpha, u)} = \{\varphi_{\alpha, u, \tau} \mid \tau \in k\}$$

**Remark 2.15.** .  $\mathcal{T}_{(\alpha, u)}$  is indeed a subgroup of  $\mathcal{T}$  because  $\varphi_{\alpha, u, \tau} \varphi_{\alpha, u, \tau'} = \varphi_{\alpha, u, \tau + \tau'}$  for any  $\tau, \tau' \in k$ . Actually, the following mapping is an isomorphism of abelian groups:

$$\begin{aligned} k & \longrightarrow \mathcal{T}_{(\alpha, u)} \\ \tau & \longmapsto \varphi_{\alpha, u, \tau} \end{aligned}$$

- .  $\mathcal{T}_{\leq(\alpha, u)}$  is generated by  $\mathcal{T}_{<(\alpha, u)} \cup \mathcal{T}_{(\alpha, u)}$ .
- . If  $(\alpha, u) < (\beta, v)$ , then  $\mathcal{T}_{\leq(\alpha, u)} \subseteq \mathcal{T}_{\leq(\beta, v)}$  and  $\mathcal{T}_{<(\alpha, u)} \subseteq \mathcal{T}_{<(\beta, v)}$ .
- .  $\mathcal{T} = \bigcup_{(\alpha, u)} \mathcal{T}_{\leq(\alpha, u)}$  and if  $(\alpha_m, u_m)$  is the greatest bypass in  $Q$ , then  $\mathcal{T} = \mathcal{T}_{\leq(\alpha_m, u_m)}$  (recall that  $Q$  has finitely many bypasses because it has no oriented cycle).

The following lemma proves that any  $\psi \in \mathcal{T}$  is a decreasing product of transvections.

**Lemma 2.16.** .  $\mathcal{T}_{<(\alpha, u)}$  is a normal subgroup of  $\mathcal{T}_{\leq(\alpha, u)}$ , for any bypass  $(\alpha, u)$ .

- . Let  $(a_1, v_1) < \dots < (a_N, v_N)$  be the (finite) increasing sequence of all the bypasses in  $Q$ . Then:
  - $\mathcal{T}_{\leq(a_i, v_i)} = \mathcal{T}_{<(a_{i-1}, v_{i-1})}$  if  $i \geq 1$ ,
  - $\mathcal{T}_{<(a_1, v_1)} = 1$ ,

$$- \mathcal{T}_{\leq(\alpha_i, v_i)} = \mathcal{T}_{(\alpha_i, v_i)} \mathcal{T}_{(\alpha_{i-1}, v_{i-1})} \cdots \mathcal{T}_{(\alpha_1, v_1)}.$$

**Proof:** Thanks to Remark 2.15, we only need to prove that if  $\tau, \nu \in k$  and if  $(\beta, v), (\alpha, u)$  are bypasses such that  $(\beta, v) < (\alpha, u)$ , then:

$$\varphi_{\beta, v, \nu} \varphi_{\alpha, u, \tau} \in \varphi_{\alpha, u, \tau} \mathcal{T}_{<(\alpha, u)} \quad (\star)$$

Let us distinguish two cases whether  $(\alpha, u, \beta, v)$  is a double bypass or not. Notice that thanks to Lemma 2.7 and to the inequality  $(\beta, v) < (\alpha, u)$  we know that  $(\beta, v, \alpha, u)$  is not a double bypass. If  $(\alpha, u, \beta, v)$  is a double bypass, then we have from Section 1 (recall that  $Q$  as no multiple arrows):

$$\varphi_{\beta, v, \nu} \varphi_{\alpha, u, \tau} = \varphi_{\alpha, u, \tau} \varphi_{\alpha, w, \tau} \nu \varphi_{\beta, v, \nu}$$

where  $w$  is the path obtained from  $u$  after replacing  $\beta$  by  $v$ . Moreover, Lemma 2.7 implies that  $(\beta, v) < (\alpha, w) < (\alpha, u)$ . Therefore,  $(\star)$  is satisfied when  $(\alpha, u, \beta, v)$  is a double bypass. If  $(\alpha, u, \beta, v)$  is not a double bypass, then we know from Section 1 that:

$$\varphi_{\alpha, u, \tau} \varphi_{\beta, v, \nu} = \varphi_{\beta, v, \nu} \varphi_{\alpha, u, \tau}$$

So  $(\star)$  is also satisfied when  $(\alpha, u, \beta, v)$  is not a double bypass. ■

Using the preceding lemma we are able to prove that any  $\psi \in \mathcal{T}$  can be uniquely written as a decreasing product of transvections.

**Proposition 2.17.** *Let  $(\alpha, u)$  be a bypass and let  $\psi \in \mathcal{T}_{\leq(\alpha, u)}$ . Then, there exist a non negative integer  $n$ , a sequence of bypasses  $(\alpha_1, u_1), \dots, (\alpha_n, u_n)$  and non zero scalars  $\tau_1, \dots, \tau_n \in k^*$  verifying:*

- (i)  $\psi = \varphi_{\alpha_n, u_n, \tau_n} \cdots \varphi_{\alpha_1, u_1, \tau_1}$ ,
- (ii)  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n) \leq (\alpha, u)$ .

Moreover, the integer  $n$  and the sequence  $(\alpha_1, u_1, \tau_1), \dots, (\alpha_n, u_n, \tau_n)$  are unique for these properties.

**Proof:** Thanks to Lemma 2.16 we know that there exist bypasses  $(\alpha_1, u_1), \dots, (\alpha_n, u_n)$  and non zero scalars  $\tau_1, \dots, \tau_n \in k^*$  such that (i) and (ii) are satisfied. So we only need to prove the uniqueness property. Let  $A, B$  and  $T$  be the following sets:

$$\begin{aligned} A &:= \{\alpha \in Q_1 \mid \psi(\alpha) \neq \alpha\} \\ B &:= \{(\alpha, u) \mid (\alpha, u) \text{ is a bypass, } \alpha \in A \text{ and } u \in \text{supp}(\psi(\alpha))\} \\ T &:= \{(\alpha, u, \tau) \mid (\alpha, u) \in B \text{ and } \tau = u^*(\psi(\alpha))\} \end{aligned}$$

Notice that the definition of  $A, B, T$  depend on  $\psi$  only (and not on the triples  $(\alpha_i, u_i, \tau_i)$ ). Let  $\beta \in Q_1$ . Then Proposition 2.11 gives a normal form:

$$\psi(\beta) = \beta + \sum_{i \text{ such that } \beta = \alpha_i} \tau_i u_i$$

By definition of a normal form, we deduce the following equalities:

$$\begin{aligned} A &= \{\alpha_1, \dots, \alpha_n\} \\ B &= \{(\alpha_1, u_1), \dots, (\alpha_n, u_n)\} \\ T &= \{(\alpha_1, u_1, \tau_1), \dots, (\alpha_n, u_n, \tau_n)\} \end{aligned}$$

This proves that  $n$  and  $(\alpha_1, u_1, \tau_1), \dots, (\alpha_n, u_n, \tau_n)$  are uniquely determined by the sets  $A, B, T$  (which depend on  $\psi$  only) and by the total order  $<$ . ■

### 3 Comparison of the presentations of a monomial algebra

Let  $kQ/I \simeq A$  be an admissible presentation of  $A$ . Thanks to Proposition 1.1, we know that there exists  $\psi$  equal to a product of transvections and of a dilatation and such that  $\psi(I_0) = I$ . The aim of this section is to exhibit  $\psi_I$  the “simplest” possible among all the  $\psi$ ’s verifying  $\psi(I_0) = I$ . We shall see that  $\psi_I$  verifies a property which makes it unique. In order to find  $\psi_I$  we will use specific properties of the Groebner basis of  $I$  (due to the fact that  $I_0$  is monomial). So, throughout the section,  $<$  will denote a total order on the set of non trivial paths in  $Q$ , as in Definition 2.5. We begin by giving some useful properties on the automorphisms  $\psi$  verifying  $\psi(I_0) = I_0$ .

**Lemma 3.1.** *Let  $D \in \mathcal{D}$  be a dilatation. Then  $D(I_0) = I_0$ . As a consequence, if  $kQ/I \simeq A$  is an admissible presentation, then there exists  $\psi \in \mathcal{T}$  such that  $\psi(I_0) = I$ .*

**Proof:** The first assertion is due to the fact that  $D(u) \in k^*u$  for any path  $u$  and to the fact that  $I_0$  is monomial. The second one is a consequence of the first one and of Proposition 1.1.  $\blacksquare$

**Lemma 3.2.** *Let  $(\alpha, u)$  be a bypass in  $Q$ . Then exactly one of the two following assertions is satisfied:*

- $\cdot \varphi_{\alpha, u, \tau}(I_0) = I_0$  for any  $\tau \in k$ .
- $\cdot \varphi_{\alpha, u, \tau}(I_0) \neq I_0$  for any  $\tau \in k^*$ .

**Proof:** Assume that  $\tau \in k^*$  verifies  $\varphi_{\alpha, u, \tau}(I_0) = I_0$  and let  $\mu \in k$ . Let  $v \in I_0$  be a path. If  $\alpha$  does not appear in  $v$ , then  $\varphi_{\alpha, u, \nu}(v) = v \in I_0$ . Assume that  $\alpha$  appears in  $v$ , i.e.  $v = v_2\alpha v_1$  with  $v_1, v_2$  paths in which  $\alpha$  does not appear (because  $Q$  has no oriented cycle). Therefore,  $\varphi_{\alpha, u, \tau}(v) = v + \tau v_2 u v_1$ . Since  $\varphi_{\alpha, u, \tau}(I_0) = I_0$ , we deduce that  $v_2 u v_1 \in I_0$ . This implies that  $\varphi_{\alpha, u, \nu}(v) = v + \nu v_2 u v_1 \in I_0$ . Since  $I_0$  is monomial, we infer that  $\varphi_{\alpha, u, \nu}(I_0) = I_0$ .  $\blacksquare$

**Lemma 3.3.** *Let  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n)$  be an increasing sequence of bypasses, let  $\tau_1, \dots, \tau_n \in k^*$  and set  $\psi = \varphi_{\alpha_n, u_n, \tau_n} \dots \varphi_{\alpha_1, u_1, \tau_1}$ . Then:*

$$\psi(I_0) = I_0 \Leftrightarrow \varphi_{\alpha_i, u_i, \tau_i}(I_0) = I_0 \text{ for any } i$$

**Proof:** Let us assume that  $\psi(I_0) = I_0$ . Let  $i \in \{1, \dots, n\}$ , let  $u = a_r \dots a_1 \in I_0$  be a path (with  $a_i \in Q_1$ ) and fix  $i \in \{1, \dots, n\}$ . We aim at proving that  $\varphi_{\alpha_i, u_i, \tau_i}(u) \in I_0$ . If  $a_j \neq \alpha_i$  for any  $j \in \{1, \dots, r\}$  then  $\varphi_{\alpha_i, u_i, \tau_i}(u) = u \in I_0$ . Now assume that there exists  $j \in \{1, \dots, r\}$  such that  $a_j = \alpha_i$  ( $j$  is necessarily unique because  $Q$  has no oriented cycle). Therefore:

$$\varphi_{\alpha_i, u_i, \tau_i}(u) = u + \tau_i a_r \dots a_{j+1} u_i a_{j-1} \dots a_1 \tag{i}$$

On the other hand, Proposition 2.9 and Proposition 2.11 imply that  $a_r \dots a_{j+1} u_i a_{j-1} \dots a_1 \in \text{supp}(\psi(u))$ . Thus, we have (recall that  $\psi(u) \in I_0$  and that  $I_0$  is monomial):

$$a_r \dots a_{j+1} u_i a_{j-1} \dots a_1 \in I_0 \tag{ii}$$

From (i) and (ii) we deduce that  $\varphi_{\alpha_i, u_i, \tau_i}(u) \in I_0$  for any path  $u \in I_0$ . So  $\varphi_{\alpha_i, u_i, \tau_i}(I_0) = I_0$  for any  $i$ . The remaining implication is immediate.  $\blacksquare$

**Remark 3.4.** *The three preceding lemmas imply that the group  $\text{Aut}_0(kQ, I_0)$  defined as follows:*

$$\text{Aut}_0(kQ, I_0) := \{\psi \in \text{Aut}(kQ) \mid \psi(x) = x \text{ for any } x \in Q_0, \text{ and } \psi(I_0) = I_0\}$$

*is generated by the dilatations and by all the transvections preserving  $I_0$ :*

$$\text{Aut}_0(kQ, I_0) = \langle \mathcal{D} \cup \{\varphi \mid \varphi \text{ is a transvection such that } \varphi(I_0) = I_0\} \rangle$$

Now we give some properties on the Groebner basis of the admissible ideals  $I$  of  $kQ$  such that  $kQ/I \simeq A$ . Recall that for such an  $I$ , there exists  $\psi \in \mathcal{T}$  such that  $\psi(I_0) = I$  (see Lemma 3.1).

**Proposition 3.5.** *Let  $\psi \in \mathcal{T}$  and let  $I = \psi(I_0)$ . Let  $B_0$  (resp.  $B$ ) be the Groebner basis of  $I_0$  (resp. of  $I$ ). Then  $B_0$  is made of all the paths in  $Q$  which belong to  $I_0$ . Moreover, the mapping:*

$$\begin{aligned} B &\longrightarrow B_0 \\ r &\longmapsto \max(\text{supp}(r)) \end{aligned} \tag{\star}$$

*is well defined and bijective. For  $u \in B_0$ , let us write  $r_u \in B$  for the inverse image of  $u$  under  $(\star)$ . Then  $\text{supp}(r_u - u)$  is a set of paths derived of  $u$ .*

**Proof:** Let  $u_1 < \dots < u_n$  be the increasing sequence of all the non trivial paths in  $Q$ . Let  $(r_1, \dots, r_d)$  be the Groebner basis of  $I$  and for each  $j \in \{1, \dots, d\}$ , let  $i_j \in \{1, \dots, n\}$  be such that:

$$r_j \in u_{i_j} + \text{Span}(u_l \ ; \ l < i_j)$$

Since  $I_0$  is monomial,  $B_0$  is made of all the paths in  $Q$  belonging to  $I_0$ .

Let  $j \in \{1, \dots, d\}$ . Since  $u_{i_j} = \max(\text{supp}(r_j))$ , the path  $u_{i_j}$  is not derived of  $u$  for any  $u \in \text{supp}(r_j)$  (thanks to Lemma 2.7). So Proposition 2.13 implies that  $u_{i_j} \in \text{supp}(\psi^{-1}(r_j)) \in I_0$ . Because  $I_0$  is

monomial, this proves that  $u_{i_j} \in I_0$ . Therefore, the mapping  $(\star)$  is well defined. It is also one-to-one because of the definition of the Groebner basis of  $I$ . Now let us prove that  $(\star)$  is onto. Let  $u \in B_0$ . Proposition 2.9 implies that  $u = \max(\text{supp}(\psi(u)))$ . Since  $\psi(u) \in I$ , we deduce that there exists  $j \in \{1, \dots, d\}$  such that  $u = u_{i_j} = \max(\text{supp}(r_j))$ . This proves that  $(\star)$  is onto and therefore bijective.

Let us prove the last assertion of the proposition. To do this, we will prove by induction on  $j \in \{1, \dots, d\}$  that the following assertion is true:

$$H_j : \text{"supp}(r_j - u_{i_j}) \text{ is a set of paths derived of } u_{i_j}\text{"}$$

Proposition 2.9 implies that for any  $j$ :

$$u_{i_j} = \max(\text{supp}(\psi(u_{i_j}))) \text{ and } u_{i_j}^*(\psi(u_{i_j})) = 1 \quad (i)$$

Moreover,  $\psi(u_{i_j}) \in I$  because  $(\star)$  is well defined and because  $\psi(I_0) = I$ . Let us prove that  $H_1$  is true. Both  $r_1$  and  $\psi(u_{i_1})$  lie in  $I$ . Moreover,  $u_{i_1} = \max(\text{supp}(r_1))$  by definition of  $u_{i_1}$  and  $u_{i_1} = \max(\text{supp}(\psi(u_{i_1})))$  because of Proposition 2.9. So  $H_1$  is true. Assume that  $j \geq 2$  and that  $H_1, \dots, H_{j-1}$  are true. Since  $\psi(u_{i_j}) \in I$  and because of (i), we have:

$$\psi(u_{i_j}) = r_j + \sum_{\substack{j' < j, \\ u_{i_{j'}} \in \text{supp}(\psi(u_{i_j}))}} u_{i_{j'}}^*(\psi(u_{i_j})) r_{j'}$$

So:

$$r_j - u_{i_j} = \psi(u_{i_j}) - u_{i_j} - \sum_{\substack{j' < j, \\ u_{i_{j'}} \in \text{supp}(\psi(u_{i_j}))}} u_{i_{j'}}^*(\psi(u_{i_j})) \left[ (r_{j'} - u_{i_{j'}}) + u_{i_{j'}} \right] \quad (ii)$$

Notice that in the above equality:

(iii)  $\text{supp}(\psi(u_{i_j}) - u_{i_j})$  is a set of paths derived of  $u_{i_j}$  (thanks to Proposition 2.9),

(iv) if  $j' < j$  verifies  $u_{i_{j'}} \in \text{supp}(\psi(u_{i_j}))$ , then:

(v)  $u_{i_{j'}}$  is derived of  $u_{i_j}$  (see (iii) above),

(vi)  $\text{supp}(r_{j'} - u_{i_{j'}})$  is a set of paths derived of  $u_{i_{j'}}$  (because  $H_{j'}$  is true) and therefore derived of  $u_{i_j}$  (thanks to (v) and to Lemma 2.3).

The points (ii) – (vi) prove that  $H_j$  is true. This proves that  $H_j$  is true for any  $j \in \{1, \dots, d\}$  and finishes the proof of the proposition.  $\blacksquare$

Now we can state and prove the proposition giving the existence and the uniqueness of the automorphism  $\psi_I$  mentioned at the beginning of the section.

**Proposition 3.6.** *Let  $kQ/I \simeq A$  be an admissible presentation. Then there exists a unique  $\psi_I \in \mathcal{T}$  verifying the following conditions:*

1)  $\psi_I(I_0) = I$ ,

2) if  $(\alpha, u)$  is a bypass such that  $u \in \text{supp}(\psi_I(\alpha))$  then  $\varphi_{\alpha, u, \tau}(I_0) \neq I_0$  for any  $\tau \in k^*$  (see Lemma 3.2).

**Proof:** • Let us prove the existence of  $\psi_I$ . Thanks to Lemma 3.1 we know that there exists  $\psi \in \mathcal{T}$  verifying 1). Let us set:

$$\mathcal{A} := \{\psi \in \mathcal{T} \mid \psi(I_0) = I\}$$

and let us assume that for any  $\psi \in \mathcal{A}$ , the condition 2) is not verified. So, for any  $\psi \in \mathcal{A}$ , we have a finite (recall that  $Q$  has no oriented cycle) and non empty set of bypasses (see Lemma 3.2):

$$B_\psi = \left\{ (\alpha, u) \left| \begin{array}{l} (\alpha, u) \text{ is a bypass} \\ u \in \text{supp}(\psi(\alpha)) \\ \varphi_{\alpha, u, \tau}(I_0) = I_0 \text{ for any } \tau \in k \end{array} \right. \right\}$$

For each  $\psi \in \mathcal{A}$ , let  $(\alpha_\psi, u_\psi) = \max B_\psi$  and let us fix  $\psi \in \mathcal{A}$  such that:

$$(\alpha_\psi, u_\psi) = \min \{(\alpha_{\psi'}, u_{\psi'}) \mid \psi' \in \mathcal{A}\}$$

For simplicity we shall use the following notations:

$$(\alpha, u) := (\alpha_\psi, u_\psi), \quad \tau := u^*(\psi(\alpha)) \text{ and } \psi' := \psi \varphi_{\alpha, u, -\tau}$$

Notice that  $\psi' \in \mathcal{A}$  because  $(\alpha, u) \in B_\psi$ . In order to get a contradiction, we are going to prove that  $(\alpha_{\psi'}, u_{\psi'}) < (\alpha, u)$ . To do this, let us prove first that  $(\alpha, u) \notin B_{\psi'}$ . Thanks to Proposition 2.17, we have the equality:

$$\psi = \varphi_{\alpha_n, u_n, \tau_n} \cdots \varphi_{\alpha_1, u_1, \tau_1}$$

where  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n)$  and where  $\tau_1, \dots, \tau_n \in k^*$ . On the other hand, since  $u^*(\psi(\alpha)) = \tau \neq 0$ , we know from Proposition 2.11 that:

$$(\exists i \in \{1, \dots, n\}) \quad (\alpha_i, u_i, \tau_i) = (\alpha, u, \tau)$$

Let us set:

$$\psi_1 := \varphi_{\alpha_{i-1}, u_{i-1}, \tau_{i-1}} \cdots \varphi_{\alpha_1, u_1, \tau_1} \in \mathcal{T}_{<(\alpha, u)}$$

Hence, the following equality holds:

$$\psi' = \varphi_{\alpha_n, u_n, \tau_n} \cdots \varphi_{\alpha_{i+1}, u_{i+1}, \tau_{i+1}} \varphi_{\alpha, u, \tau} \psi_1 \varphi_{\alpha, u, \tau}^{-1}$$

Since  $\psi_1 \in \mathcal{T}_{<(\alpha, u)}$ , Lemma 2.16 implies that  $\varphi_{\alpha, u, \tau} \psi_1 \varphi_{\alpha, u, \tau}^{-1} \in \mathcal{T}_{<(\alpha, u)}$ . Therefore, Proposition 2.17 gives the equality:

$$\varphi_{\alpha, u, \tau} \psi_1 \varphi_{\alpha, u, \tau}^{-1} = \varphi_{\beta_m, v_m, \nu_m} \cdots \varphi_{\beta_1, v_1, \nu_1}$$

where  $(\beta_1, v_1) < \dots < (\beta_m, v_m) < (\alpha, u)$  and  $\nu_1, \dots, \nu_m \in k^*$ . As a consequence, we have:

$$\psi' = \varphi_{\alpha_n, u_n, \tau_n} \cdots \varphi_{\alpha_{i+1}, u_{i+1}, \tau_{i+1}} \varphi_{\beta_m, v_m, \nu_m} \cdots \varphi_{\beta_1, v_1, \nu_1}$$

where  $(\beta_1, v_1) < \dots < (\beta_m, v_m) < (\alpha, u) < (\alpha_{i+1}, u_{i+1}) < \dots < (\alpha_n, u_n)$  and where  $\tau_{i+1}, \dots, \tau_n, \nu_1, \dots, \nu_m \in k^*$ . In particular, Proposition 2.11 implies that  $u \notin \text{supp}(\psi'(\alpha))$ . Therefore:

$$(\alpha, u) \notin B_{\psi'} \tag{i}$$

Hence:

$$(\alpha, u) \neq (\alpha_{\psi'}, u_{\psi'}) = \max B_{\psi'} \tag{ii}$$

Now let us prove that  $(\alpha_{\psi'}, u_{\psi'}) \leq (\alpha, u)$ . To do this, it suffices to prove that the following implication holds for any bypass  $(\beta, v)$ :

$$v \in \text{supp}(\psi'(\beta)) \text{ and } (\alpha, u) < (\beta, v) \Rightarrow \varphi_{\beta, v, t}(I_0) \neq I_0 \text{ for any } \tau \in k^* \tag{iii}$$

Let  $(\beta, v)$  be a bypass such that  $v \in \text{supp}(\psi'(\beta))$  and such that  $(\alpha, u) < (\beta, v)$ . Since  $\psi' = \psi \varphi_{\alpha, u, -\tau}$ , we have:

$$\psi'(\beta) = \begin{cases} \psi(\beta) & \text{if } \beta \neq \alpha \\ \psi(\beta) - \tau \psi(u) & \text{if } \beta = \alpha \end{cases}$$

Therefore,  $v \in \text{supp}(\psi'(\beta)) \subseteq \text{supp}(\psi(\beta)) \cup \text{supp}(\psi(u))$ . Remark that if  $v \in \text{supp}(\psi(u))$ , then Proposition 2.9 implies that  $v$  is derived of  $u$  (we have  $u \neq v$  because  $\beta = \alpha$  and  $(\alpha, u) < (\beta, v)$ ) and therefore  $(\alpha, u) > (\alpha, v) = (\beta, v)$  whereas we assumed that  $(\alpha, u) < (\beta, v)$ . This proves that  $v \in \text{supp}(\psi(\beta))$ . Since  $(\beta, v) > (\alpha, u) = (\alpha_\psi, u_\psi) = \max B_\psi$  we deduce that  $\varphi_{\beta, v, \tau}(I_0) = I_0$  for any  $\tau \in k$ . This proves that the implication (iii) is satisfied, thus:

$$(\alpha_{\psi'}, u_{\psi'}) \leq (\alpha, u) \tag{iv}$$

From (ii) and (iv) we deduce that:

$$(\alpha_{\psi'}, u_{\psi'}) < (\alpha_\psi, u_\psi) = (\alpha, u)$$

This contradicts the minimality of  $(\alpha_\psi, u_\psi)$  and proves the existence of  $\psi$ .

• Now let us prove the uniqueness of  $\psi_I$ . Assume that  $\psi, \psi' \in \mathcal{T}$  verify the conditions 1) and 2). In order to prove that  $\psi = \psi'$  we only need to prove that  $\theta^*(\psi(\alpha)) = \theta^*(\psi'(\alpha))$  for any bypass  $(\alpha, \theta)$ . Let  $\alpha \in Q_1$  and assume that there exists a minimal path  $\theta$  such that  $(\alpha, \theta)$  is bypass and such that  $\theta^*(\psi(\alpha)) \neq \theta^*(\psi'(\alpha))$ . We may assume that  $\theta^*(\psi(\alpha)) \neq 0$ , i.e.  $\theta \in \text{supp}(\psi(\alpha))$ . Since  $\psi$  verifies 2), we deduce that there exist paths  $u$  and  $v$  such that:

$$u \in I_0, v \notin I_0 \text{ and } \varphi_{\alpha, \theta, 1}(u) = u + v \notin I_0$$

Notice that Proposition 2.9 gives:

$$\begin{cases} v^*(\psi(u)) = \theta^*(\psi(\alpha)) \text{ and } u^*(\psi(u)) = 1 \\ v^*(\psi'(u)) = \theta^*(\psi'(\alpha)) \text{ and } u^*(\psi'(u)) = 1 \end{cases} \tag{i}$$

Since  $u \in I$ , we have  $\psi(u), \psi'(u) \in I_0$ . Therefore, using Proposition 3.5 and with the same notations concerning the Groebner bases, we have:

$$\begin{cases} \psi(u) = r_u + \sum_{w \in \mathcal{A}_\psi} w^*(\psi(u))r_w \\ \psi'(u) = r_u + \sum_{w \in \mathcal{A}_{\psi'}} w^*(\psi'(u))r_w \end{cases}$$

where  $\mathcal{A}_\psi$  is equal to:

$$\mathcal{A}_\psi := \{w \in \text{supp}(\psi(u)) \mid w \neq u \text{ and } w \in I_0\}$$

So:

$$\begin{cases} v^*(\psi(u)) = v^*(r_u) + \sum_{w \in \mathcal{A}_\psi} w^*(\psi(u))v^*(r_w) \\ v^*(\psi'(u)) = v^*(r_u) + \sum_{w \in \mathcal{A}_{\psi'}} w^*(\psi'(u))v^*(r_w) \end{cases} \quad (ii)$$

Let  $w \in \mathcal{A}_\psi$  be such that  $v^*(r_w) \neq 0$ , i.e.  $v \in \text{supp}(r_w)$ . Since  $v \notin I_0$  and since  $w \in I_0$  we deduce that  $v \in \text{supp}(r_w - w)$ . So:

- .  $v$  is derived of  $w$  (thanks to Proposition 3.5 and because  $v \in \text{supp}(r_w - w)$ ).
- .  $v$  is derived of  $u$  of order 1 (because  $\varphi_{\alpha, \theta, 1}(u) = u + v$ ).
- .  $w$  is derived of  $u$  (because  $w \in \mathcal{A}_\psi$  and thanks to Proposition 2.9).

Using Lemma 2.3, these three facts imply that:

$$u = u_2\alpha u_1, \quad v = u_2\theta u_1 \text{ and } w = u_2\theta' u_1 \quad (iii)$$

where  $u_1, u_2$  are paths and where  $\theta'$  is a path derived of  $\theta$ . In particular,  $(\alpha, \theta')$  is a bypass such that  $\theta' < \theta$  (see Lemma 2.7). Therefore, the minimality of  $\theta$  forces  $\theta'^*(\psi(\alpha)) = \theta'^*(\psi'(\alpha))$ . Moreover, (iii) and Proposition 2.9 imply that

$$w^*(\psi(u)) = \theta'^*(\psi(\alpha)) = \theta'^*(\psi'(\alpha)) = w^*(\psi'(u))$$

Therefore we have proved the following implication:

$$w \in \mathcal{A}_\psi \text{ and } v^*(r_w) \neq 0 \Rightarrow w^*(\psi(u))v^*(r_w) = w^*(\psi'(u))v^*(r_w) \quad (iv)$$

After exchanging the roles of  $\psi$  and  $\psi'$ , the arguments used to prove (iv) also give the following implication:

$$w \in \mathcal{A}_{\psi'} \text{ and } v^*(r_w) \neq 0 \Rightarrow w^*(\psi(u))v^*(r_w) = w^*(\psi'(u))v^*(r_w) \quad (v)$$

From (ii), (iv) and (v) we deduce that  $v^*(\psi(u)) = v^*(\psi'(u))$ , and from (i) we infer that  $\theta^*(\psi(\alpha)) = \theta^*(\psi'(\alpha))$  whereas we assumed the contrary. This proves that  $\psi = \psi'$ .  $\blacksquare$

## 4 Proof of the main theorem

The aim of this section is to prove that the quiver  $\Gamma$  of the homotopy relations of the admissible presentations of  $A$  has  $\sim_{I_0}$  as unique source. This fact will be used in order to exhibit the universal cover of  $A$ . Notice that  $\sim_{I_0}$  is a source of  $\Gamma$ . Indeed, all minimal relations in  $I_0$  are monomial relations so, for any  $\sim_I \in \Gamma_0$  we have  $\gamma \sim_{I_0} \gamma' \Rightarrow \gamma \sim_I \gamma'$ . In order to prove that  $\sim_{I_0}$  is the unique source in  $\Gamma$  we will prove that for any admissible presentation  $kQ/I \simeq A$ , the decomposition of  $\psi_I$  (given by Proposition 3.6) into a decreasing product of transvections (see Proposition 2.17) defines a path in  $\Gamma$  starting at  $\sim_{I_0}$  and ending at  $\sim_I$ . In this purpose, we begin with the following proposition.

**Proposition 4.1.** *Let  $kQ/I \simeq A$  be an admissible presentation. Then, for any bypass  $(\alpha, u)$  we have:*

$$u \in \text{supp}(\psi_I(\alpha)) \Rightarrow u \sim_I \alpha$$

**Proof:** For simplicity we shall write  $\psi$  for  $\psi_I$ . Thanks to Proposition 2.17 we have:

$$\psi = \varphi_{\alpha_n, u_n, \tau_n} \cdots \varphi_{\alpha_1, u_1, \tau_1}$$

with  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n)$  and  $\tau_1, \dots, \tau_n \in k^*$ . Notice that Proposition 2.11 implies that we only need to prove that  $\alpha_i \sim_I u_i$  for any  $i$ . We will prove this fact by a decreasing induction on  $m \in \{1, \dots, n\}$ . Let us set:

$$H_m : \text{''}\alpha_i \sim_I u_i \text{ for any } i \in \{m, m+1, \dots, n\}\text{''}$$

$H_{n+1}$  is true because  $\{i \mid n+1 \leq i \leq n\}$  is empty, so let us assume that  $m \in \{1, \dots, n\}$  is such that  $H_{m+1}$  is true and let us prove that so is  $H_m$ . From Proposition 2.11 we have  $u_m \in \text{supp}(\psi(\alpha_m))$ . So Proposition 3.6 provides a path  $u \in I_0$  such that  $\varphi_{\alpha_m, u_m, 1}(u) \notin I_0$ . Therefore, there exist paths  $v_1, v_2$  such that:

$$u = v_2 \alpha_m v_1, \quad v := v_2 u_m v_1 \notin I_0 \text{ and } \varphi_{\alpha_m, u_m, 1}(u) = u + v \quad (i)$$

Since  $\psi(u) \in I$  we have:

$$\psi(u) = r_1 + \dots + r_N$$

where  $r_1, \dots, r_N$  are minimal relations in  $I$  with pairwise disjoint supports. Remark that  $u, v \in \text{supp}(\psi(u))$  thanks to Proposition 2.9 and to Proposition 2.11. Without loss of generality, we may assume that  $v \in \text{supp}(r_1)$ . Let  $i \in \{1, \dots, N\}$  be such that  $u \in \text{supp}(r_i)$ . If  $i = 1$  then  $u \sim_I v$  hence (i) gives  $\alpha_m \sim_I u_m$ . So we can assume that  $i \neq 1$ . Since  $r_1 \in I$  we have  $\psi^{-1}(r_1) \in I_0$ . Because  $I_0$  is monomial we deduce that  $v \notin \text{supp}(\psi^{-1}(r_1))$ . Thanks to Proposition 2.13, this proves that:

$$\text{there exists } w \in \text{supp}(r_1) \text{ such that } v \text{ is derived from } w \quad (ii)$$

Therefore:

- .  $w$  is derived of  $u$  since  $w \in \text{supp}(r_1) \subseteq \text{supp}(\psi(u))$  (see Proposition 2.9,  $u \neq w$  because  $u \notin \text{supp}(r_1)$ ),
- .  $v$  is derived of  $w$  (see (ii)),
- .  $v$  is derived of  $u$  of order 1 (because of (i)).

Thanks to Lemma 2.3, these three points imply that:

$$w = v_2 \theta v_1 \text{ and } u_m \text{ is derived of } \theta \quad (iii)$$

Since  $w \in \text{supp}(\psi(u))$ , the equalities  $w = v_2 \theta v_1$ ,  $u = v_2 \alpha_m v_1$  and Proposition 2.9 imply that  $\theta \in \text{supp}(\psi(\alpha_m))$ . Hence, there exists  $j \in \{1, \dots, n\}$  such that:

$$(\alpha_m, \theta) = (\alpha_j, u_j)$$

Since  $u_m$  is derived of  $\theta$  (see (iii)), we get  $u_j = \theta > u_m$  (see Lemma 2.7) and therefore  $j > m$ . Since  $H_{m+1}$  is true, we also have:

$$\alpha_m = \alpha_j \sim_I u_j = \theta \quad (iv)$$

Finally, since  $r_1$  is a minimal relation in  $I$  such that  $v, w \in \text{supp}(r_1)$ , we have  $v \sim_I w$ . Using (i), (iii) and (iv) we deduce that:

$$\alpha_m \sim_I u_m$$

This proves that  $H_m$  is true. So  $H_1$  is true, i.e.  $\alpha_i \sim_I u_i$  for any  $i$ . ■

**Remark 4.2.** *In the preceding proposition we have proved that  $\alpha \sim_I u$  for any  $u \in \text{supp}(\psi(\alpha))$ . On the other hand,  $\sim_{I_0}$  is weaker than  $\sim_I$  (i.e.  $\gamma \sim_{I_0} \gamma' \Rightarrow \gamma \sim_I \gamma'$ ). These two properties are linked in general. Indeed, in [10, Prop. 4.2.35, Prop. 4.2.36] the author has proved that if  $I$  is an admissible ideal (non necessarily monomial) of  $kQ$  and if  $\psi \in \mathcal{T}$  is such that  $\alpha \sim_{\psi(I)} u$  for any bypass  $(\alpha, u)$  such that  $u \in \text{supp}(\psi(\alpha))$ , then  $\sim_I$  is weaker than  $\sim_{\psi(I)}$ .*

Now we can exhibit a path in  $\Gamma$  starting at  $\sim_{I_0}$  and ending at  $\sim_I$ , whenever  $kQ/I \simeq A$ .

**Proposition 4.3.** *Let  $kQ/I \simeq A$  be an admissible presentation. Let  $(\alpha_1, u_1) < \dots < (\alpha_n, u_n)$  be the bypasses and let  $\tau_1, \dots, \tau_n \in k^*$  be the scalars such that  $\psi_I = \varphi_{\alpha_n, u_n, \tau_n} \dots \varphi_{\alpha_1, u_1, \tau_1}$  (see Proposition 2.17). For each  $i \in \{1, \dots, n\}$ , set:*

$$I_i := \varphi_{\alpha_i, u_i, \tau_i} \dots \varphi_{\alpha_1, u_1, \tau_1}(I_0)$$

then, for each  $i$ , one of the two following situations occurs:

- .  $\sim_{I_{i-1}}$  and  $\sim_{I_i}$  coincide,
- .  $\varphi_{\alpha_i, u_i, \tau_i}$  induces an arrow  $\sim_{I_{i-1}} \rightarrow \sim_{I_i}$  in  $\Gamma$ .

In particular, there exists a path in  $\Gamma$  starting at  $\sim_{I_0}$  and ending at  $\sim_{I_n} = \sim_I$ .

**Proof:** Let  $i \in \{1, \dots, n\}$  and set  $\psi_i := \varphi_{\alpha_i, u_i, \tau_i} \dots \varphi_{\alpha_1, u_1, \tau_1}$ . Thus  $I_i = \psi_i(I_0)$ . Using Proposition 2.11 and Proposition 3.6 it is easily verified that:

$$\psi_i = \psi_{I_i}$$



Therefore, Proposition 4.1 applied to  $I_i$  gives:

$$\alpha_i \sim_{I_i} u_i$$

Since  $I_i = \varphi_{\alpha_i, u_i, \tau_i}(I_{i-1})$ , this proves that (see 1.2) either  $\sim_{I_{i-1}}$  and  $\sim_{I_i}$  coincide or  $\varphi_{\alpha_i, u_i, \tau_i}$  induces an arrow  $\sim_{I_{i-1}} \rightarrow \sim_{I_i}$  in  $\Gamma$ . Thus, the following vertices of  $\Gamma$ :

$$\sim_{I_0}, \sim_{I_1}, \dots, \sim_{I_n} = \sim_I$$

are the vertices of a path in  $\Gamma$  (maybe with repetitions) starting at  $\sim_{I_0}$  and ending at  $\sim_I$ . ■

The preceding proposition and the fact that  $\Gamma$  has no oriented cycle gives immediately the following corollary which was proved by the author in [9] in the case of triangular and without double bypass algebras over an algebraically closed field of characteristic zero.

**Corollary 4.4.** *Let  $Q$  be a quiver without oriented cycle and without multiple arrows, let  $I_0$  be an admissible and monomial ideal of  $kQ$  and let  $A = kQ/I_0$ . Then the quiver  $\Gamma$  of the homotopy relations of the admissible presentations of  $A$  admits  $\sim_{I_0}$  as unique source.*

Proposition 4.3 also allows us to prove the main theorem of this text. It extends [9, Thm. 2] to monomial triangular algebras without multiple arrows. Notice that in the following theorem we make no assumption on the characteristic of  $k$ . Also recall that  $\pi_1(Q, I_0) = \pi_1(Q)$ .

**Theorem 2.** *Let  $A$  be a monomial, triangular, basic and connected  $k$ -algebra. Assume that the quiver  $Q$  of  $A$  has no multiple arrows. Fix  $\varphi: kQ/I_0 \xrightarrow{\sim} A$  a monomial presentation of  $A$ . Let  $p: (\widehat{Q}, \widehat{I}_0) \rightarrow (Q, I_0)$  be the universal cover of quiver with relations (see [11]) and let  $\overline{p}: k\widehat{Q}/\widehat{I}_0 \rightarrow kQ/I_0$  be induced by  $p$ . Then, for any Galois covering  $F: C \rightarrow A$  with group  $G$  and with  $C$  connected and locally bounded, there exists a commutative diagram:*

$$\begin{array}{ccc} k\widehat{Q}/\widehat{I}_0 & \xrightarrow{F'} & C \\ \varphi \circ \overline{p} \downarrow & & \downarrow F \\ A & \xrightarrow{\sim} & A \end{array}$$

where the bottom horizontal arrow is an isomorphism which restricts to the identity map on the set of objects of  $A$ . Moreover,  $F': k\widehat{Q}/\widehat{I}_0 \rightarrow C$  is a Galois covering with group  $K$  a normal subgroup of  $\pi_1(Q)$  such that there exists an exact sequence of groups:

$$1 \rightarrow K \rightarrow \pi_1(Q) \rightarrow G \rightarrow 1$$

**Proof:** The proof of the theorem is almost identical to the proof of [8, Thm. 2]: one uses Proposition 4.3 instead of [8, Lem. 4.3]. ■

## References

- [1] W. W. Adams and P. Loustau. *An introduction to Gröbner bases*, volume 3 of *Graduate Studies in Mathematics*. American Mathematical Society, 1994.
- [2] I. Assem, D. Castonguay, E. N. Marcos, and S. Trepode. Strongly simply connected shurian algebras and multiplicative bases. *Journal of Algebra*, 283(1):161–189, 2005.
- [3] K. Bongartz and P. Gabriel. Covering spaces in representation theory. *Inventiones Mathematicae*, 65:331–378, 1982.
- [4] P. Dowbor and A. Skowroński. Galois coverings of representation infinite algebras. *Commentarii Mathematici Helvetici*, 62:311–337, 1987.
- [5] D. R. Farkas, C. D. Feustel, and E. L. Green. Synergy in the theories of Gröebner bases and path algebras. *Canadian Journal of Mathematics*, 45(4):727–739, 1993.
- [6] P. Gabriel. The universal cover of a representation finite algebra. *Lecture Notes in Mathematics.*, 903:65–105, 1981. in: Representation of algebras.
- [7] E. L. Green. Graphs with relations, coverings and group-graded algebras. *Transactions of the American Mathematical Society*, 279(1):297–310, sep 1983.

- [8] P. Le Meur. The fundamental group of a triangular algebra without double bypasses. *C. R. Acad. Sci. Paris*, Ser. I 341, 2005.
- [9] P. Le Meur. The universal cover of an algebra without double bypass. math.RT/0507513, 2005.
- [10] P. Le Meur. Revêtements galoisiens et groupe fondamental des algèbres de dimension finie. Thèse de doctorat de l'Université Montpellier 2, <http://tel.ccsd.cnrs.fr/tel-00011753>, 2006.
- [11] R. Martínez-Villa and J. A. de la Peña. The universal cover of a quiver with relations. *Journal of Pure and Applied Algebra*, 30:277–292, 1983.
- [12] Ch. Riedtmann. Algebren, darstellungsköcher ueberlagerungen und zurück. *Commentarii Mathematici Helvetici*, 55:199–224, 1980.