



HAL
open science

Integral representation of the linear Boltzmann operator for granular gas dynamics with applications

Luisa Arlotti, Bertrand Lods

► **To cite this version:**

Luisa Arlotti, Bertrand Lods. Integral representation of the linear Boltzmann operator for granular gas dynamics with applications. 2007. hal-00128370v1

HAL Id: hal-00128370

<https://hal.science/hal-00128370v1>

Preprint submitted on 31 Jan 2007 (v1), last revised 17 Aug 2007 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INTEGRAL REPRESENTATION OF THE LINEAR BOLTZMANN OPERATOR FOR GRANULAR GAS DYNAMICS WITH APPLICATIONS

L. ARLOTTI & B. LODS

ABSTRACT. We investigate the properties of the collision operator Q associated to the linear Boltzmann equation for *dissipative* hard-spheres arising in granular gas dynamics. We establish that, as in the case of non-dissipative interactions, the gain collision operator is an integral operator whose kernel is made explicit. One deduces from this result a complete picture of the spectrum of Q in an Hilbert space setting, generalizing results from T. Carleman [5] to granular gases. In the same way, we obtain from this integral representation of Q^+ that the semigroup in $L^1(\mathbb{R}^3 \times \mathbb{R}^3, dx \otimes dv)$ associated to the linear Boltzmann equation for dissipative hard spheres is *honest* generalizing known results from [1].

KEYWORDS. Granular gas dynamics, linear Boltzmann equation, detailed balance law, spectral theory, C_0 -semigroup.

1. INTRODUCTION

We deal in this paper with the linear Boltzmann equation for *dissipative interactions* modeling the evolution of a granular gas, undergoing *inelastic collisions* with its underlying medium. Actually, we shall see in the sequel that there is no contrast, *in the linear description*, between the kinetic theory of granular gases and that of classical (*elastic*) gases. This may seem quite surprising if one has in mind the fundamental differences that may be emphasized between the *nonlinear kinetic theory* of granular gases and that of classical gases, as briefly recalled in the next lines.

1.1. Granular gas dynamics: linear and nonlinear descriptions. Let us begin by recalling the general features of the kinetic description of granular gas dynamics that can be recovered from the monograph [4] or the more mathematically oriented survey [17]. If $f(x, v, t)$ denotes the distribution function of granular particles with position $x \in \mathbb{R}^3$ and velocity $v \in \mathbb{R}^3$ at time $t \geq 0$, then the evolution of $f(x, v, t)$ is governed by the following generalization of Boltzmann equation

$$\partial_t f(x, v, t) + v \cdot \nabla_x f(x, v, t) = C(f)(x, v, t), \quad (1.1)$$

with initial condition

$$f(x, v, 0) = f_0(x, v) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3, dx \otimes dv)$$

where the right-hand side $C(f)$ models the collision phenomena and depends on the description we adopt.

In the **nonlinear description**, the collision operator $C(f) =: \mathcal{B}[f, f]$ is a quadratic operator modeling the collision phenomena between particles. For *hard-spheres* interactions, it reads

$$\mathcal{B}[f, f](v) = \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| \left[\frac{1}{\epsilon^2} f(x, v_\star, t) f(x, w_\star, t) - f(x, v, t) f(x, w, t) \right] dw dn,$$

where q is the relative velocity, $q = v - w$. The microscopic velocities (v_\star, w_\star) are the pre-collisional velocities of the so-called inverse collision, which results in (v, w) as post-collisional velocities. The main peculiarity of the kinetic description of granular gas is the inelastic character of the microscopic collision mechanism which induces that *part of the total kinetic energy is dissipated*. This energy dissipation might be due to the roughness of the surface or just to a non-perfect restitution and is measured through a restitution coefficient $0 < \epsilon < 1$ (which we assume here to be constant for simplicity). As a consequence, the collision phenomenon is a *non microreversible process*. Generally, we assume that the energy dissipation does not affect the conservation of momentum. Therefore, in the homogeneous setting, i.e. when $f_0(x, v) = f_0(v)$ is independent of the position, the number density of the gas is constant while the bulk velocity is conserved if initially zero. However, the temperature of the gas

$$\vartheta(t) = \frac{1}{3} \int_{\mathbb{R}^3} |v|^2 f(t, v) dv$$

continuously decreases (*cooling of granular gas*). As a consequence, the stationary state of the inelastic collision operator \mathcal{B} is a given Dirac mass. However, the homogeneous Boltzmann equation for granular gases exhibits self-similar solution (*homogeneous cooling state*) [7, 12]. Note the important contrast with the classical kinetic theory, i.e. when $\epsilon = 1$, for which it is well-known that the steady state of the collision operator is a Maxwellian distribution.

The **linear description** of granular gases concerns dilute particles (test particles with negligible mutual interactions) immersed in a fluid at thermal equilibrium [10, 11, 16]. The dilute suspension dissipates its kinetic energy colliding with particles of the host fluid. In such a case, the collision operator $\mathcal{C}(f) = \mathcal{Q}(f)$ is a linear operator given by

$$\mathcal{Q}(f) = \mathcal{B}[f, \mathcal{M}_1] = \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| \left[\frac{1}{\epsilon^2} f(x, v_\star, t) \mathcal{M}_1(w_\star) - f(x, v, t) \mathcal{M}_1(w) \right] dw dn \quad (1.2)$$

where \mathcal{M}_1 stands for the distribution function of the host fluid. Note that in such a description, the microscopic masses of the dilute particles m and that of the host particles m_1 can be different. We will assume throughout this paper that the distribution function of the host fluid is a given normalized Maxwellian function:

$$\mathcal{M}_1(v) = \left(\frac{m_1}{2\pi\vartheta_1} \right)^{3/2} \exp \left\{ -\frac{m_1(v - u_1)^2}{2\vartheta_1} \right\}, \quad v \in \mathbb{R}^3,$$

where $u_1 \in \mathbb{R}^3$ is the given bulk velocity and $\vartheta_1 > 0$ is the given effective temperature of the host fluid. It can be shown in this case that the number density and bulk velocity of the dilute gas are still conserved. The temperature is still not conserved but it does not decrease anymore. This strongly contrasts to the nonlinear description and suggests that, in the linear setting, granular gases behave like classical ones. The first mathematical result in this direction is the following derived in [10] (see also [11]) according to which, as in the classical case, the steady state of \mathcal{Q} remains Gaussian:

Theorem 1.1. *The Maxwellian velocity distribution:*

$$\mathcal{M}(v) = \left(\frac{m}{2\pi\vartheta^\#} \right)^{3/2} \exp \left\{ -\frac{m(v - u_1)^2}{2\vartheta^\#} \right\} \quad v \in \mathbb{R}^3,$$

with $\vartheta^\# = \frac{(1 + \epsilon)m}{2m + (3 + \epsilon)m_1} \vartheta_1$ is the unique equilibrium state of Q with unit mass.

The existence and uniqueness of such an equilibrium state allows to establish a linear version of the famous H -Theorem. Precisely, for any convex C^1 -function $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}$, one can define the associated entropy functional as

$$H_\Phi(f|\mathcal{M}) = \int_{\mathbb{R}^3} \mathcal{M}(v) \Phi\left(\frac{f(v)}{\mathcal{M}(v)}\right) dv, \quad (1.3)$$

Theorem 1.2 (H -Theorem [10, 14]). *Let $f_0(v)$ be a space homogeneous distribution function with unit mass and finite entropy, i.e. $H_\Phi(f_0|\mathcal{M}) < \infty$. Then,*

$$\frac{d}{dt} H_\Phi(f(t)|\mathcal{M}) \leq 0 \quad (t \geq 0), \quad (1.4)$$

where $f(t)$ stands for the (unique) solution to (1.1) in $L^1(\mathbb{R}^3, dv)$.

Here again, this linear version of H -Theorem contrasts very much with the situation in the nonlinear setting where such a Lyapunov functional does not exist. As an important consequence, it can be shown by suitable compactness arguments that any solution to the Boltzmann equation (1.1) (with unit mass) converges towards the Maxwellian equilibrium \mathcal{M} .

To summarize, the steady state of the linear collision operator for dissipative interactions has the same nature (a Maxwellian distribution) as the one corresponding to non-dissipative interactions. Moreover, as in the classical case, by virtue of the H -Theorem, such a steady state attracts any solution to the space homogeneous Boltzmann equation (1.1). This seems to indicate that *the linear description of granular gases does not contrast too much with that of classical gases*. It is the main subject of this paper to make precise and confirm such an indication and the key ingredient will be the derivation of an integral representation of the gain part of the collision operator.

1.2. Main results. The main concern of our paper is the derivation of a suitable representation of the gain part of the collision operator Q as an integral operator with *explicit kernel*. Precisely, the linear collision operator Q can be split into $Q(f) = Q^+(f) - Q^-(f)$, where the gain part is

$$Q^+(f)(v) = \epsilon^{-2} \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| f(v_\star) \mathcal{M}_1(w_\star) dw dn$$

while

$$Q^-(f)(v) = \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| f(v) \mathcal{M}_1(w) dw dn = \sigma(v) f(v)$$

where the collision frequency $\sigma(v)$ is given by $\sigma(v) = \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| \mathcal{M}_1(w) dw dn$. It is well-known that, for non-dissipative interactions, i.e. when $\epsilon = 1$, the gain part Q^+ can be written as an integral operator with explicit kernel [5, 9] (see also [8, 6] for similar results for the *linearized* Boltzmann equation). We prove that such a representation is still valid in the dissipative case:

Theorem 1.3. *If $f \geq 0$ is such that $\sigma(v)f(v) \in L^1(\mathbb{R}^3, dv)$, then*

$$Q^+(f)(v) = \int_{\mathbb{R}^3} k(v, v') f(v') dv'$$

where the integral kernel $k(v, v')$ can be made explicit (see (2.2)).

Actually, most important is the fact that the integral kernel $k(v, v')$ turns out to be very similar to that obtained in the classical case (see for instance [9, 5]), the only changes standing in some explicit numerical constants. Moreover, as we shall see, the kernel $k(v, v')$ and the Maxwellian distribution \mathcal{M} satisfy the following **detailed balance law**:

$$k(v, v')\mathcal{M}(v') = k(v', v)\mathcal{M}(v), \quad v, v' \in \mathbb{R}^3,$$

that allows us to recover Theorem 1.1 in a direct way. Recall that, in [10], the Gaussian nature of the steady state of Q was obtained by replacing Q by its grazing collision limit.

We derive from these two results some important consequences on the linear Boltzmann equation (1.1) with $C = Q$. The applications are dealing with space dependent version of (1.1) as well as with space homogeneous version of it. The first one concerns the spectral properties of the Boltzmann collision operator in its natural Hilbert space setting.

1.3. Spectral properties of the Boltzmann operator in $L^2(\mathcal{M}^{-1})$. Applying the above H -Theorem 1.2 with the quadratic convex function $\Phi(x) = (x - 1)^2$, one sees that a natural function space for the study of the *homogeneous linear Boltzmann equation* is the weighted space $L^2(\mathcal{M}^{-1})$. Now, from Theorem 1.3, it is possible to prove that the gain collision operator Q^+ is *compact* in $L^2(\mathcal{M}^{-1})$. This compactness result has important consequences on the structure of the spectrum of Q as an operator in $L^2(\mathcal{M}^{-1})$. Precisely, from Weyl's Theorem, the spectrum of Q in this space is given by the (essential) range of the collision frequency $\sigma(\cdot)$ and of isolated eigenvalues with finite algebraic multiplicities. Since $\lambda = 0$ is a simple eigenvalue of Q (its associated null space is spanned by \mathcal{M}), this leads to the existence of a positive spectral gap. In turns, one proves that any solution to the space-homogeneous linear Boltzmann equation (1.1) converges at an exponential rate towards the equilibrium. These spectral results are technical generalizations of some of the fundamental results of T. Carleman [5], but are new in the context of granular gas dynamics.

1.4. Honest solutions for hard-spheres model. It is easily seen that, for any nonnegative f ,

$$\int_{\mathbb{R}^3} Q^+(f)(v)dv = \int_{\mathbb{R}^3} \sigma(v)f(v)dv, \quad (1.5)$$

i.e. the collision operator Q is *conservative*. Then, formally, any nonnegative solution $f(x, v, t)$ to (1.1) (with $C = Q$) should satisfy the following *mass conservation equation*:

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} f(x, v, t)dx dv = \int_{\mathbb{R}^3 \times \mathbb{R}^3} f(x, v, 0)dx dv, \quad \forall t > 0. \quad (1.6)$$

It is the main concern of Section 4 to prove that such a formal mass conservation property holds true for any nonnegative initial datum $f(x, v, 0) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3)$. As well documented in the monograph [3], this is strongly related to the honesty of the C_0 -semigroup governing Eq. (1.1). More precisely, if we denote by T_0 the streaming operator:

$$\mathcal{D}(T_0) = \{f \in X, v \cdot \nabla_x f \in X\}, \quad T_0 f = -v \cdot \nabla_x f,$$

it is not difficult to see that there exists some extension G of $T_0 + Q$ that generates a C_0 -semigroup of contractions $(V(t))_{t \geq 0}$ in $X = L^1(\mathbb{R}^3 \times \mathbb{R}^3)$. According to the so-called "sub-stochastic perturbation" theory, developed in [1, 3, 19], it can be proved that

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} V(t)f(x, v)dx dv = \int_{\mathbb{R}^3 \times \mathbb{R}^3} f(x, v)dx dv, \quad \forall f \in X, f \geq 0$$

if and only if $\mathcal{D}(G) = \overline{\mathcal{D}(T_0 + Q)}$. We show in Section 4 that the latter holds. To do so, we shall use the integral representation (Theorem 1.3) in order to apply some of the results of [1] (see also [3, Chapter 10]) dealing with the classical linear Boltzmann equation.

1.5. Organization of the paper. We derive in Section 2 the integral representation of Q^+ (Theorem 2.1) as well as some of its immediate consequences concerning the explicit expression of the collision frequency. We also recover Theorem 1.1 through a detailed balance law. Section 3 is devoted to the study of the collision operator Q in the narrow space $L^2(\mathcal{M}^{-1}(v)dv)$ and its spectral consequences. In Section 4 we applied the results of Section 2 as well as some known facts about the classical linear Boltzmann equation [1, 3] to the honesty of the solutions to the Boltzmann equation for dissipative hard-spheres.

2. INTEGRAL REPRESENTATION OF THE GAIN OPERATOR

Let us consider the gain operator for dissipative hard-spheres:

$$Q^+(f)(v) = \epsilon^{-2} \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| f(v_\star) \mathcal{M}_1(w_\star) dw dn$$

and let $\sigma(v)$ be the corresponding collision frequency:

$$\sigma(v) = \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| \mathcal{M}_1(w) dw dn, \quad v \in \mathbb{R}^3.$$

Recall that \mathcal{M}_1 is a Maxwellian distribution function with bulk velocity u_1 and effective temperature ϑ_1 . We recall here the general microscopic description of the pre-collisional velocities (v_\star, w_\star) which result in (v, w) after collision. For a constant restitution coefficient $0 < \epsilon < 1$, one has [4, 17]

$$\begin{cases} v_\star = v - 2\alpha \frac{1-\beta}{1-2\beta} [q \cdot n] n, \\ w_\star = w + 2(1-\alpha) \frac{1-\beta}{1-2\beta} [q \cdot n] n; \end{cases}$$

where $q = v - w$, α is the mass ratio and β denotes the inelasticity parameter

$$\alpha = \frac{m_1}{m + m_1}, \quad \beta = \frac{1-\epsilon}{2}.$$

We show in this section that, as it occurs for the classical Boltzmann equation, Q^+ turns out to be an integral operator with explicit kernel. Precisely, we prove the following whose proof is inspired by similar calculations in [9].

Theorem 2.1 (Integral representation of Q^+). For any $f \in L^1(\mathbb{R}^3 \times \mathbb{R}^3, dx \otimes \sigma(v)dv)$,

$$Q^+ f(x, v) = \frac{1}{2\epsilon^2 \gamma^2} \int_{\mathbb{R}^3} f(x, v') k(v, v') dv', \quad (2.1)$$

where

$$k(v, v') = \left(\frac{m_1}{2\pi \vartheta_1} \right)^{1/2} |v - v'|^{-1} \exp \left\{ -\frac{m_1}{8\vartheta_1} \left((1 + \mu)|v - v'| + \frac{|v - u_1|^2 - |v' - u_1|^2}{|v - v'|} \right)^2 \right\} \quad (2.2)$$

with $\mu = \frac{2\alpha(1 - \beta) - 1}{\alpha(1 - \beta)} > 0$ and $\gamma = \alpha \frac{1 - \beta}{1 - 2\beta}$.

Proof. The local (in x) nature of Q^+ is obvious and we can restrict ourselves to prove the result for a function $f \in L^1(\mathbb{R}^3, \sigma(v)dv)$ that does not depend on x . Set $\gamma = \alpha \frac{1 - \beta}{1 - 2\beta}$ and $\bar{\gamma} = (1 - \alpha) \frac{1 - \beta}{1 - 2\beta}$ so that

$$v_\star = v - 2\gamma[q \cdot n]n \quad \text{and} \quad w_\star = w + 2\bar{\gamma}[q \cdot n]n.$$

The following formula, for smooth φ :

$$\int_{\mathbb{S}_+} (q \cdot n) \varphi((q \cdot n)n) dn = \frac{|q|}{4} \int_{\mathbb{S}^2} \varphi\left(\frac{q - |q|\sigma}{2}\right) d\sigma,$$

applied to

$$\varphi(x) = f(v - 2\gamma x) \mathcal{M}_1(w + 2\bar{\gamma}x)$$

yields

$$Q^+ f(v) = \frac{1}{2\epsilon^2} \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q| f(v') \mathcal{M}_1(w') dw d\sigma,$$

where

$$\begin{cases} v' &= v + \gamma(|q|\sigma - q), \\ w' &= w - \bar{\gamma}(|q|\sigma - q). \end{cases} \quad (2.3)$$

Now, let us recall the following formula valid for sufficiently smooth Φ :

$$\frac{1}{2} \int_{\mathbb{S}^2} \Phi(|q|\sigma - q) d\sigma = \frac{1}{|q|} \int_{\mathbb{R}^3} \delta(2x \cdot q + x^2) \Phi(x) dx. \quad (2.4)$$

We have, according to (2.3)–(2.4),

$$Q^+ f(v) = \epsilon^{-2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} \delta(2x \cdot q + x^2) f(v + \gamma x) \mathcal{M}_1(w - \bar{\gamma}x) dw dx.$$

The change of variables $x \mapsto v' = v + \gamma x$ leads to

$$Q^+ f(v) = \epsilon^{-2} \gamma^{-3} \int_{\mathbb{R}^3 \times \mathbb{R}^3} \delta(2\gamma^{-1}(v' - v) \cdot q + \gamma^{-2}|v' - v|^2) f(v') \mathcal{M}_1(w - \frac{\bar{\gamma}}{\gamma}(v' - v)) dw dv'.$$

Now, keeping v and v' fixed, we perform the change of variables $w \mapsto w' = w - \frac{\bar{\gamma}}{\gamma}(v' - v)$, which leads to

$$Q^+ f(v) = \epsilon^{-2} \gamma^{-3} \int_{\mathbb{R}^3 \times \mathbb{R}^3} \delta \left(2\gamma^{-1}(v' - v) \cdot [v - w' - \frac{\bar{\gamma}}{\gamma}(v' - v)] + \gamma^{-2}|v' - v|^2 \right) \times \\ \times f(v') \mathcal{M}_1(w') dw' dv'.$$

Writing $w' = v + \lambda_1 n + V_2$ with $\lambda_1 = (w' - v) \cdot n \in \mathbb{R}$, $n = (v' - v)/|v' - v|$ and $V_2 \cdot n = 0$, we get, noting that $dw' = dV_2 d\lambda_1$,

$$Q^+ f(v) = \epsilon^{-2} \gamma^{-3} \int_{\mathbb{R}^3} f(v') dv' \int_{\mathbb{R}} d\lambda_1 \int_{V_2 \cdot n=0} \mathcal{M}_1(v + V_2 + \lambda_1 n) dV_2 \times \\ \times \delta \left(\gamma^{-2}|v' - v|^2 - 2\bar{\gamma}\gamma^{-2}|v' - v|^2 - 2\gamma^{-1}\lambda_1|v' - v| \right).$$

Thanks to the change of variables $\lambda_1 \mapsto 2\gamma^{-1}|v' - v|\lambda_1$, one can evaluate the Dirac mass as

$$\int_{\mathbb{R}} \delta \left(\gamma^{-2}|v' - v|^2 - 2\bar{\gamma}\gamma^{-2}|v' - v|^2 - 2\gamma^{-1}\lambda_1|v' - v| \right) \mathcal{M}_1(v + V_2 + \lambda_1 n) d\lambda_1 \\ = \frac{\gamma}{2|v' - v|} \mathcal{M}_1 \left(v + V_2 + \frac{1 - 2\bar{\gamma}}{2\gamma}(v' - v) \right)$$

where we used that $n = (v' - v)/|v' - v|$. Consequently,

$$Q^+ f(v) = \frac{1}{2\epsilon^2 \gamma^2} \int_{\mathbb{R}^3} k(v, v') f(v') dv'$$

where

$$k(v, v') = \frac{1}{|v - v'|} \int_{V_2 \cdot (v' - v) = 0} \mathcal{M}_1 \left(v + V_2 + \frac{1 - 2\bar{\gamma}}{2\gamma}(v' - v) \right) dV_2.$$

It remains now to explicit $k(v, v')$. We will use the approach of [9]. Let us assume v, v' to be fixed. Let \mathbf{P} be the hyperplan orthogonal to $(v' - v)$. For any $V_2 \in \mathbf{P}$, set

$$z = v + \frac{1 - 2\bar{\gamma}}{2\gamma}(v' - v) + V_2 - u_1$$

so that

$$k(v, v') = \left(\frac{\varrho_1}{\pi} \right)^{3/2} |v - v'|^{-1} \int_{V_2 \in \mathbf{P}} \exp\{-\varrho_1 z^2\} dV_2.$$

where $\varrho_1 = \frac{m_1}{2\vartheta_1}$. Denoting for simplicity $u = \frac{v + v'}{2} - u_1$ and $\mu = -\frac{1 - 2\bar{\gamma}}{\gamma}$, one has

$$z^2 = \left(u + \frac{v - v'}{2} + \frac{\mu}{2}(v - v') + V_2 \right)^2 \\ = |u + V_2|^2 + \frac{(1 + \mu)^2}{4} |v - v'|^2 + \frac{1 + \mu}{2} (|v - u_1|^2 - |v' - u_1|^2)$$

where we used the fact that V_2 is orthogonal to $(v' - v)$. Splitting u as

$$u = u_0 + u_\perp$$

where u_0 is parallel to $v - v'$ while u_\perp is orthogonal to $v - v'$ (i.e. $u_\perp \in \mathbf{P}$), we see that

$$|u + V_2|^2 = |u_0|^2 + |u_\perp + V_2|^2 \quad \text{and} \quad |u_0|^2 = \frac{[|v - u_1|^2 - |v' - u_1|^2]^2}{4|v - v'|^2},$$

so that

$$k(v, v') = |v - v'|^{-1} \left(\frac{\varrho_1}{\pi} \right)^{3/2} \int_{\mathbf{P}} \exp(-\varrho_1 |u_\perp + V_2|^2) dV_2 \\ \exp \left\{ -\frac{\varrho_1}{4} \left((1 + \mu)^2 |v - v'|^2 + 2(1 + \mu)(|v - u_1|^2 - |v' - u_1|^2) + \frac{[|v - u_1|^2 - |v' - u_1|^2]^2}{|v - v'|^2} \right) \right\}.$$

Finally, since $u_\perp \in \mathbf{P}$,

$$\int_{\mathbf{P}} \exp(-\varrho_1 |u_\perp + V_2|^2) dV_2 = \int_{\mathbb{R}^2} \exp(-\varrho_1 x^2) dx = \frac{\pi}{\varrho_1},$$

one obtains the desired expression for $k(v, v')$. \square

The very important fact to be noticed out is that the expression of $k(v, v')$ is very similar to that one obtains in the elastic case [9], the only change being the expression of the constant μ . In particular, in the elastic case $\epsilon = 1$, we recover the expression of the kernel obtained in [5] for particles of same mass (i.e. $m = m_1$) and in [9] for particles with different masses.

Another fundamental property of the kernel $k(v, v')$ is that it allows us to recover the steady state of Q through some *microscopic detailed balance law*. Precisely,

Theorem 2.2. *With the notations of the Theorem 2.1, the following detailed balance law:*

$$k(v, v') \exp \left\{ -\frac{m_1}{2\vartheta_1} (1 + \mu)(v' - u_1)^2 \right\} = k(v', v) \exp \left\{ -\frac{m_1}{2\vartheta_1} (1 + \mu)(v - u_1)^2 \right\}, \quad (2.5)$$

holds for any $v, v' \in \mathbb{R}^3$. As a consequence, the Maxwellian velocity distribution:

$$\mathcal{M}(v) = \left(\frac{m}{2\pi\vartheta^\#} \right)^{3/2} \exp \left\{ -\frac{m(v - u_1)^2}{2\vartheta^\#} \right\} \quad v \in \mathbb{R}^3,$$

with $\vartheta^\# = \frac{(1 - \alpha)(1 - \beta)}{1 - \alpha(1 - \beta)} \vartheta_1$ is the unique equilibrium state of Q with unit mass.

Proof. According to Eq. (2.2), it is easily seen that

$$k(v', v) = k(v, v') \exp \left\{ \frac{m_1}{2\vartheta_1} (1 + \mu) (|v - u_1|^2 - |v' - u_1|^2) \right\}, \quad v, v' \in \mathbb{R}^3$$

which is nothing but (2.5). Now, writing $\frac{m_1}{2\vartheta_1}(1 + \mu) = \frac{m}{2\vartheta^\sharp}$, straightforward calculations lead to the desired expression for the equilibrium temperature ϑ^\sharp . The fact that \mathcal{M} is an equilibrium solution with unit mass follows then from the fact that

$$Q(\mathcal{M}) = \int_{\mathbb{R}^3} k(v, v') \mathcal{M}(v') dv' - \sigma(v) \mathcal{M}(v) = \int_{\mathbb{R}^3} [k(v, v') \mathcal{M}(v') - k(v', v) \mathcal{M}(v)] dv'$$

and from the detailed balance law (2.5). To prove that the steady state is unique, we adopt the strategy of [15, Theorem 1]. Precisely, consider the equation

$$\sigma(v) f(v) = Q^+ f(v), \quad \forall v \in \mathbb{R}^3 \quad (2.6)$$

which admits at least the solution $f = \mathcal{M}_1$. Since $\sigma(v)$ does not vanish, any solution f to (2.6) is such that

$$f(v) = \frac{1}{\sigma(v)} Q^+(f)(v), \quad \forall v \in \mathbb{R}^3.$$

Since Q^+ is an integral operator with *nonnegative* kernel, it is clear that $|f(v)| \leq \frac{1}{\sigma(v)} Q^+(|f|)(v)$ for any $v \in \mathbb{R}^3$, i.e.

$$\sigma(v) |f(v)| \leq Q^+(|f|)(v), \quad \forall v \in \mathbb{R}^3.$$

Now, from the positivity of both σ and Q^+ , one sees that the conservation of mass (1.5) reads:

$$\|\sigma f\|_X = \|\sigma |f|\|_X = \|Q^+(|f|)\|_X.$$

This shows that, actually,

$$|Q^+(f)(v)| = \sigma(v) |f(v)| = Q^+(|f|)(v) \quad \forall v \in \mathbb{R}^3.$$

Again, since Q^+ is a positive operator, one obtains that

$$f = \pm |f|.$$

Now, assume that (2.6) admits two solutions f_1, f_2 with $\int_{\mathbb{R}^3} f_1(v) dv = \int_{\mathbb{R}^3} f_2(v) dv = 1$. Then, $f_1 - f_2$ is again a solution to (2.6) so that, $f_1 - f_2 = \pm |f_1 - f_2|$. Thus,

$$\pm \int_{\mathbb{R}^3} |f_1(v) - f_2(v)| dv = \int_{\mathbb{R}^3} f_1(v) dv - \int_{\mathbb{R}^3} f_2(v) dv = 0$$

and the uniqueness follows. \square

The above result allows to derive the explicit expression of the collision frequency $\sigma(v)$:

Corollary 2.3. *The collision frequency $\sigma(v)$ for dissipative hard-spheres interactions is given by*

$$\sigma(v) = \frac{2\pi}{(2 + \mu)^2} \sqrt{\frac{m_1}{2\pi\vartheta_1}} \left\{ \frac{4\vartheta_1}{m_1} \exp\left(-\frac{m_1}{2\vartheta_1}|v - u_1|^2\right) + \left(2|v - u_1| + \frac{2\vartheta_1}{m_1|v - u_1|}\right) \int_0^{2|v - u_1|} \exp\left(-\frac{m_1}{8\vartheta_1}t^2\right) dt \right\}. \quad (2.7)$$

Consequently, there exist positive constants ν_0, ν_1 such that

$$\nu_0(1 + |v - u_1|) \leq \sigma(v) \leq \nu_1(1 + |v - u_1|), \quad \forall v \in \mathbb{R}^3.$$

Proof. Set $C = \sqrt{\frac{m_1}{2\pi\vartheta_1}}$. Noting that $\sigma(v) = \int_{\mathbb{R}^3} k(v', v) dv'$ for any $v \in \mathbb{R}^3$, one has, with the change of variable $z = v' - v$,

$$\begin{aligned}\sigma(v) &= C \int_{\mathbb{R}^3} \exp \left\{ -\frac{m_1}{8\vartheta_1} \left((1 + \mu)|z| - \frac{|v - u_1|^2 - |z + v - u_1|^2}{|z|} \right)^2 \right\} |z|^{-1} dz \\ &= C \int_{\mathbb{R}^3} \exp \left\{ -\frac{m_1}{8\vartheta_1} ((2 + \mu)|z| + 2|v - u_1| \cos \varphi)^2 \right\} |z|^{-1} dz \\ &= 2\pi C \int_0^\infty d\rho \int_0^\pi \rho \exp \left\{ -\frac{m_1}{8\vartheta_1} ((2 + \mu)\rho + 2|v - u_1| \cos \varphi)^2 \right\} \sin \varphi d\varphi.\end{aligned}$$

The computation of this last integral leads to the desired expression for $\sigma(v)$. The estimates are then straightforward [9]. \square

3. APPLICATION TO THE BOLTZMANN OPERATOR IN $L^2(\mathcal{M}^{-1})$.

We investigate in this section the properties of the Boltzmann operator Q in the weighted space

$$\mathcal{H} = L^2(\mathbb{R}^3; \mathcal{M}^{-1}(v)dv)$$

We shall denote by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ the inner product in \mathcal{H} . The introduction of such an Hilbert space setting is motivated by the application of the H -Theorem 1.2 with the convex function

$$\Phi(x) = (x - 1)^2.$$

In this case, one sees that, if $f_0 \geq 0$ is a space homogeneous initial distribution such that

$$\int_{\mathbb{R}^3} f_0(v) dv = 1, \quad \int_{\mathbb{R}^3} |f_0(v)|^2 \mathcal{M}^{-1}(v) dv < \infty,$$

then any solution $f(t, v)$ to the space homogeneous equation

$$\partial_t f(t, v) = Q(f)(t, v), \quad f(0, v) = f_0(v) \in \mathcal{H}, \quad (3.1)$$

satisfies the following estimate:

$$\frac{d}{dt} \int_{\mathbb{R}^3} |f(t, v) - \mathcal{M}(v)|^2 \mathcal{M}(v)^{-1} dv \leq 0, \quad t \geq 0.$$

In other words, the mapping $t \mapsto \|f(t, \cdot) - \mathcal{M}\|_{\mathcal{H}}$ is *nonincreasing*. For these reasons, the study of the properties of the collision operator Q in \mathcal{H} is of particular relevance for the asymptotic behavior of the solution

$$\partial_t f(t, v) = Q(f)(t, v), \quad f(0, v) \in \mathcal{H}. \quad (3.2)$$

The material of this section borrows some techniques already employed by T. Carleman [5] in the study of non-dissipative gas dynamics (see also, e.g. [6] or [8] for similar results in the context of the *linearized* Boltzmann equation). Let \mathcal{L} be the realization of the operator Q in \mathcal{H} , i.e.

$$\mathcal{D}(\mathcal{L}) = \left\{ f \in \mathcal{H}; \int_{\mathbb{R}^3} |f(v)|^2 \sigma(v) \mathcal{M}^{-1}(v) dv < \infty \right\}.$$

and, for any $f \in \mathcal{D}(\mathcal{L})$, $\mathcal{L}f(v) = Q(f)(v)$ is given by (1.2). As previously, one can use the following splitting of \mathcal{L} as a gain operator and a loss (multiplication) operator, $\mathcal{L} = \mathcal{L}^+ - \mathcal{L}^-$ with

$$\mathcal{L}^+(f)(v) = \int_{\mathbb{R}^3} k(v, v') f(v') dv' \quad \text{and} \quad \mathcal{L}^-(f) = \sigma(v) f(v), \quad f \in \mathcal{D}(\mathcal{L}).$$

We shall show, as in the classical case, that \mathcal{L}^+ is actually a bounded operator in \mathcal{H} . Precisely, let \mathcal{J} define the natural bijection operator from $L^2(\mathbb{R}^3, dv)$ to \mathcal{H} :

$$\left\{ \begin{array}{l} \mathcal{J} : L^2(\mathbb{R}^3, dv) \longrightarrow \mathcal{H} \\ f \longmapsto \mathcal{J} f(v) = \mathcal{M}^{1/2}(v) f(v) \end{array} \right.$$

It is clear that \mathcal{J} is a bounded bijective operator whose inverse is given by

$$\mathcal{J}^{-1} g(v) = \mathcal{M}^{-1/2}(v) g(v) \in L^2(\mathbb{R}^3, dv), \quad \forall g \in \mathcal{H}.$$

Now, let us define

$$G(v, v') = \mathcal{M}^{-1/2}(v) k(v, v') \mathcal{M}^{1/2}(v'), \quad v, v' \in \mathbb{R}^3,$$

i.e.

$$G(v, v') = \left(\frac{m_1}{2\pi\vartheta_1} \right)^{1/2} |v - v'|^{-1} \exp \left\{ -\frac{m_1}{8\vartheta_1} \left((1 + \mu)^2 |v - v'|^2 + \frac{(|v - u_1|^2 - |v' - u_1|^2)^2}{|v - v'|^2} \right) \right\}. \quad (3.3)$$

From the detailed balance law (2.5), one easily checks that $G(v, v') = G(v', v)$ for any $v, v' \in \mathbb{R}^3$. Therefore, defining \mathcal{G} as the integral operator in $L^2(\mathbb{R}^3, dv)$ with kernel $G(v, v')$, i.e.

$$\mathcal{G}f(v) = \int_{\mathbb{R}^3} G(v, v') f(v') dv',$$

one can prove the following:

Proposition 3.1. *\mathcal{G} is a bounded symmetric operator in $L^2(\mathbb{R}^3, dv)$ and $\mathcal{L}^+ = \mathcal{J} \mathcal{G} \mathcal{J}^{-1}$. Consequently, \mathcal{L}^+ is a bounded symmetric operator in \mathcal{H} .*

Proof. It is clear that \mathcal{G} is symmetric since $G(v, v') = G(v', v)$. Now, to prove the boundedness of \mathcal{G} , one adopts a strategy already used in the non-dissipative case by T. Carleman [5, p. 75]. Precisely, from (3.3), the use of spherical coordinates leads to

$$\begin{aligned} \int_{\mathbb{R}^3} G(v, v') dv' &= \left(\frac{m_1}{2\pi\vartheta_1} \right)^{1/2} \int_0^{2\pi} d\xi \int_0^\pi \sin \varphi d\varphi \times \\ &\quad \times \int_0^\infty \varrho \exp \left\{ -\frac{m_1}{8\vartheta_1} \left((1 + \mu)^2 \varrho^2 + (\varrho + 2|v - u_1| \cos \varphi)^2 \right) \right\} d\varrho \\ &\leq 4\pi \left(\frac{m_1}{2\pi\vartheta_1} \right)^{1/2} \int_0^\infty \varrho \exp \left(-\frac{m_1}{8\vartheta_1} (1 + \mu)^2 \varrho^2 \right) d\varrho < \infty. \end{aligned}$$

Therefore,

$$C := \sup_{v \in \mathbb{R}^3} \int_{\mathbb{R}^3} G(v, v') dv' < \infty.$$

Since $G(\cdot, \cdot)$ is symmetric, one also has $\sup_{v' \in \mathbb{R}^3} \int_{\mathbb{R}^3} G(v, v') dv = C < \infty$. Denoting by $\langle \cdot, \cdot \rangle$ the usual inner product of $L^2(\mathbb{R}^3, dv)$, one deduces from Cauchy-Schwarz identity,

$$\langle \mathcal{G}f, g \rangle \leq \frac{C}{2} \left(\int_{\mathbb{R}^3} |f(v)|^2 dv + \int_{\mathbb{R}^3} |g(v')|^2 dv' \right), \quad \forall f, g \in L^2(\mathbb{R}^3, dv),$$

which leads to the boundedness of \mathcal{G} . Now, since $G(v, v') = \mathcal{M}^{-1/2}(v)k(v, v')\mathcal{M}^{1/2}(v')$ for any $v, v' \in \mathbb{R}^3$, one gets easily that $\mathcal{L}^+ = \mathcal{J}\mathcal{G}\mathcal{J}^{-1}$ and the conclusion follows. \square

In Proposition 3.1, we proved that the gain operator \mathcal{L}^+ is bounded in \mathcal{H} , i.e. $\mathcal{L}^+ \in \mathcal{B}(\mathcal{H})$. Actually, we have much better and it is possible, as in the non-dissipative case, to prove that \mathcal{L}^+ is a compact operator in \mathcal{H} . Indeed, arguing as in [5, p. 70–75], one can prove the following:

Lemma 3.2. *For any $0 < p < 3$ and any $q \geq 0$, there exists $C(p, q) > 0$ such that*

$$\int_{\mathbb{R}^3} |G(v, v')|^p \frac{dv'}{(1 + |v' - u_1|)^q} \leq \frac{C(p, q)}{(1 + |v - u_1|)^{q+1}}, \quad \forall v \in \mathbb{R}^3.$$

Proof. The proof is a technical generalization of a similar result due to T. Carleman [5] in the classical case (i.e. when $m = m_1$ and $\varepsilon = 1$). Let us fix $0 < p < 3$ and $q \geq 0$ and set

$$I(v) = \int_{\mathbb{R}^3} |G(v, v')|^p \frac{dv'}{(1 + |v' - u_1|)^q}.$$

Then, one sees easily that

$$\begin{aligned} I(v) &= \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \int_0^{2\pi} d\xi \int_0^\pi \sin \varphi d\varphi \int_0^\infty \varrho^{-p} \exp \left\{ -\frac{m_1 p}{8\vartheta_1} \left((1 + \mu)^2 \varrho^2 + (\varrho + 2|v - u_1| \cos \varphi)^2 \right) \right\} \times \\ &\quad \times \frac{\varrho^2 d\varrho}{\left(1 + \sqrt{\varrho^2 + |v - u_1|^2 + 2\varrho|v - u_1| \cos \varphi} \right)^q} \\ &= 2\pi \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \int_0^\pi \sin \varphi d\varphi \int_0^\infty \varrho^{2-p} \frac{\exp \left\{ -\frac{m_1 p}{8\vartheta_1} \left((1 + \mu)^2 \varrho^2 + (\varrho + 2|v - u_1| \cos \varphi)^2 \right) \right\}}{\left(1 + \sqrt{\varrho^2 + |v - u_1|^2 + 2\varrho|v - u_1| \cos \varphi} \right)^q} d\varrho. \end{aligned}$$

Note that, since $0 < p < 3$,

$$\sup_{v \in \mathbb{R}^3} I(v) \leq 4\pi \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \int_0^\infty \varrho^{2-p} \exp \left\{ -\frac{m_1 p}{8\vartheta_1} (1 + \mu)^2 \varrho^2 \right\} d\varrho < \infty. \quad (3.4)$$

Performing the change of variable $x = \varrho/|v - u_1| + 2 \cos \varphi$, $y = \varrho/|v - u_1|$, one has $(x, y) \in \Omega$ where

$$\Omega = \{(x, y) \in \mathbb{R}^2; y > 0, |x - y| \leq 2\}$$

and

$$I(v) = \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \pi |v - u_1|^{3-p} \int_\Omega \exp \left\{ -\frac{m_1 p |v - u_1|^2}{8\vartheta_1} \left((1 + \mu)^2 y^2 + x^2 \right) \right\} \frac{dx dy}{y^{p-2} \left(1 + |v - u_1| \sqrt{1 + xy} \right)^q}.$$

We split Ω into $\Omega = \Omega_1 \cup \Omega_2$ where Ω_1 is the half-ellipse

$$\Omega_1 = \{(x, y) \in \mathbb{R}^2 : y > 0, (1 + \mu)^2 y^2 + x^2 < 1/4\} \quad \text{while} \quad \Omega_2 = \Omega \setminus \Omega_1.$$

Note that, since $1 + \mu \geq 1$, one has $\Omega_1 \subset \Omega$. One define correspondingly $I_1(v)$ and $I_2(v)$ as the above integral over Ω_1 and Ω_2 respectively. One notes first that, if $(x, y) \in \Omega_1$ then $xy > -\frac{1}{8(1+\mu)}$ so that

$$I_1(v) \leq \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \frac{\pi|v - u_1|^{3-p}}{(1 + a|v - u_1|)^q} \int_{\Omega_1} \exp \left\{ -\frac{m_1 p |v - u_1|^2}{8\vartheta_1} \left((1 + \mu)^2 y^2 + x^2 \right) \right\} \frac{dx dy}{y^{p-2}}$$

where $a = \sqrt{1 - \frac{1}{8(1+\mu)}}$, $0 < a < 1$. Letting $R = \left(\frac{m_1 p}{8\vartheta_1} \right)^{1/2}$ and setting $t = R|v - u_1|x$, $u = R(1 + \mu)|v - u_1|y$, it is easy to check that $-R|v - u_1|/2 \leq t \leq R|v - u_1|/2$, while $0 \leq u \leq R|v - u_1|/2$, so that

$$I_1(v) \leq \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \frac{\pi R^{p-4} (1 + \mu)^{p-3}}{|v - u_1| (1 + a|v - u_1|)^q} \int_{\mathbb{R}} dt \int_0^\infty \frac{\exp\{-(t^2 + u^2)\}}{u^{p-2}} du.$$

Thus, there exists a constant $C_1(p, q) > 0$ such that

$$I_1(v) \leq \frac{C_1(p, q)}{|v - u_1| (1 + a|v - u_1|)^q}, \quad \forall v \in \mathbb{R}^3. \quad (3.5)$$

Let us now deal with $I_2(v)$. Arguing as above,

$$\begin{aligned} I_2(v) &= \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \pi |v - u_1|^{3-p} \int_{\Omega_2} \exp \left\{ -\frac{m_1 p |v - u_1|^2}{8\vartheta_1} \left((1 + \mu)^2 y^2 + x^2 \right) \right\} \\ &\quad \times \frac{dx dy}{y^{p-2} \left(1 + |v - u_1| \sqrt{1 + xy} \right)^q} \\ &= \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \pi |v - u_1|^{3-p} \int_{\Omega_2} \exp \left\{ -\frac{R^2 |v - u_1|^2}{2} \left((1 + \mu)^2 y^2 + x^2 \right) \right\} \times \\ &\quad \times \frac{\exp \left\{ -\frac{R^2 |v - u_1|^2}{2} \left((1 + \mu)^2 y^2 + x^2 \right) \right\}}{y^{p-2} \left(1 + |v - u_1| \sqrt{1 + xy} \right)^q} dx dy. \end{aligned}$$

Clearly, since $(1 + \mu)^2 y^2 + x^2 > 1/4$ for any $(x, y) \in \Omega_2$, then

$$\begin{aligned} I_2(v) &\leq \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \pi |v - u_1|^{3-p} \int_{\Omega_2} \exp \left(-\frac{R^2 |v - u_1|^2}{8} \right) \exp \left\{ -\frac{R^2 |v - u_1|^2}{2} \left((1 + \mu)^2 y^2 + x^2 \right) \right\} \frac{dx dy}{y^{p-2}} \\ &\leq \left(\frac{m_1}{2\pi\vartheta_1} \right)^{p/2} \pi |v - u_1|^{3-p} \exp \left(-\frac{R^2 |v - u_1|^2}{8} \right) \int_0^\infty \exp \left(-\frac{R^2 |v - u_1|^2}{2} (1 + \mu)^2 y^2 \right) \frac{dy}{y^{p-2}} \int_{y-2}^{y+2} dx. \end{aligned}$$

Hence, there is some constant $C_2(p, q)$ such that

$$I_2(v) \leq C_2(p, q) \exp \left(-\frac{R^2 |v - u_1|^2}{8} \right), \quad v \in \mathbb{R}^3. \quad (3.6)$$

Combining (3.5) and (3.6), one sees that

$$I(v) \leq \frac{C_1(p, q)}{|v - u_1|(1 + a|v - u_1|)^q} + C_2(p, q) \exp\left(-\frac{R^2|v - u_1|^2}{8}\right), \quad v \in \mathbb{R}^3.$$

According to (3.4), $\limsup_{|v - u_1| \rightarrow 0} I(v) < \infty$, from which one can conclude that there exists some constant $C(p, q) > 0$ with

$$I(v) \leq \frac{C(p, q)}{(1 + |v - u_1|)^q}, \quad v \in \mathbb{R}^3$$

which ends the proof. \square

From the above Lemma, one has the following compactness result:

Proposition 3.3. *\mathcal{G} is compact in $L^2(\mathbb{R}^3, dv)$. Consequently, \mathcal{L} is a compact operator in \mathcal{H} .*

Proof. Applying arguments already used in [5], the above Lemma implies that the third iterate of \mathcal{G} is an Hilbert–Schmidt operator in $L^2(\mathbb{R}^3, dv)$, i.e. the kernel of \mathcal{G}^3 is square summable over $\mathbb{R}^3 \times \mathbb{R}^3$. The compactness of \mathcal{G} follows then from standard arguments and that of \mathcal{L} is deduced from the identity $\mathcal{L}^+ = \mathcal{J} \mathcal{G} \mathcal{J}^{-1}$ (see Proposition 3.1). \square

The following, which generalizes a known result from classical kinetic theory, proves that \mathcal{L} is a negative symmetric operator in \mathcal{H} :

Proposition 3.4. *The operator $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ is a negative self-adjoint operator of \mathcal{H} . Precisely,*

$$\langle \mathcal{L}f, f \rangle_{\mathcal{H}} = -\frac{1}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} k(v, v') \mathcal{M}(v') \left[\mathcal{M}^{-1}(v) f(v) - f(v') \mathcal{M}^{-1}(v') \right]^2 dv dv' \leq 0$$

for any $f \in \mathcal{D}(\mathcal{L})$.

Proof. The fact that $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ is self-adjoint is a direct consequence of Proposition 3.1 since \mathcal{L}^- is clearly symmetric. Now, it is a classical feature, from the detailed balance law (2.5), that

$$\begin{aligned} \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &= \int_{\mathbb{R}^3 \times \mathbb{R}^3} f(v) \mathcal{M}^{-1}(v) \left[k(v, v') f(v') - k(v', v) f(v) \right] dv dv' \\ &= \int_{\mathbb{R}^3 \times \mathbb{R}^3} k(v, v') \mathcal{M}(v') \left[\mathcal{M}^{-1}(v') f(v') - f(v) \mathcal{M}^{-1}(v) \right] f(v) \mathcal{M}^{-1}(v) dv dv'. \end{aligned}$$

Exchanging v and v' and using again the detailed balance law (2.5), one sees that

$$\begin{aligned} \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &= \int_{\mathbb{R}^3 \times \mathbb{R}^3} k(v', v) \mathcal{M}(v) \left[\mathcal{M}^{-1}(v) f(v) - f(v') \mathcal{M}^{-1}(v') \right] f(v') \mathcal{M}^{-1}(v') dv dv' \\ &= \int_{\mathbb{R}^3 \times \mathbb{R}^3} k(v, v') \mathcal{M}(v') \left[\mathcal{M}^{-1}(v) f(v) - f(v') \mathcal{M}^{-1}(v') \right] f(v') \mathcal{M}^{-1}(v') dv dv' \end{aligned}$$

so that, taking the mean of the two quantities,

$$\langle \mathcal{L}f, f \rangle_{\mathcal{H}} = -\frac{1}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3} k(v, v') \mathcal{M}(v') \left[\mathcal{M}^{-1}(v) f(v) - f(v') \mathcal{M}^{-1}(v') \right]^2 dv dv' \leq 0$$

which ends the proof. \square

Remark 3.5. From the above result, the spectrum $\mathfrak{S}(\mathcal{L})$ of \mathcal{L} lies in \mathbb{R}_- , i.e. $\mathfrak{S}(\mathcal{L}) \subset (-\infty, 0]$. It is clear that $\lambda = 0$ lies in $\mathfrak{S}(\mathcal{L})$. Precisely 0 is a simple eigenvalue of \mathcal{L} since \mathcal{M} is the unique (up to a multiplication factor) steady state of \mathcal{L} .

Combining the above results with Proposition 3.1 leads to a precise description of the spectrum of \mathcal{L} :

Theorem 3.6. The spectrum of \mathcal{L} (as an operator in \mathcal{H}) consists of the spectrum of \mathcal{L}^- and of, at most, eigenvalues of finite multiplicities. Precisely, setting $\nu_0 = \inf_{v \in \mathbb{R}^3} \sigma(v) > 0$,

$$\mathfrak{S}(\mathcal{L}) = \{\lambda \in \mathbb{R}; \lambda \leq -\nu_0\} \cup \{\lambda_n; n \in I\}$$

where $I \subset \mathbb{N}$ and $(\lambda_n)_n$ is a decreasing sequence of real eigenvalues of \mathcal{L} with finite algebraic multiplicities: $\lambda_0 = 0 > \lambda_1 > \lambda_2 \dots > \lambda_n > \dots$, whose unique possible cluster point is $-\nu_0$.

Proof. From Proposition 3.3, \mathcal{L} is nothing but a compact perturbation of the loss operator $-\mathcal{L}^-$. Henceforth, Weyl's Theorem asserts that $\mathfrak{S}(\mathcal{L}) \setminus \mathfrak{S}(-\mathcal{L}^-)$ consists of, at most, eigenvalues of finite algebraic multiplicities whose unique cluster point is $\sup\{\lambda, \lambda \in \mathfrak{S}(-\mathcal{L}^+)\}$. In particular, up to a rearrangement, one can write $\mathfrak{S}(\mathcal{L}) \setminus \mathfrak{S}(-\mathcal{L}^-) = \{\lambda_n, n \in I\}$ with $\lambda_0 > \lambda_1 > \lambda_2 \dots > \lambda_n \geq \dots$. We already saw that $\lambda_0 = 0$ since \mathcal{M} is a steady state of Q and $\mathcal{M} \in \mathcal{H}$. Now, since $-\mathcal{L}^-$ is a multiplication operator by the collision frequency $-\sigma(\cdot)$, its spectrum $\mathfrak{S}(-\mathcal{L}^-)$ is given by the essential range $R_{\text{ess}}(-\sigma(\cdot))$ of the collision frequency. From Corollary 2.7, one sees without difficulty that

$$R_{\text{ess}}(-\sigma(\cdot)) = (-\infty, -\nu_0]$$

where $\nu_0 = \inf_{v \in \mathbb{R}^3} \sigma(v) = \lim_{|v-u_1| \rightarrow 0} \sigma(v) = \frac{8}{(2+\mu)^2} \sqrt{\frac{2\pi g_1}{m_1}}$ is positive. \square

The above result provides a complete picture of the spectrum of Q as an operator in \mathcal{H} and shows, in particular, the existence of a positive spectral gap $|\lambda_1|$ of \mathcal{L} . Note that such a result, combined with Proposition 3.4, has important consequence on the entropy production, since it can be shown in an easy way that the H -Theorem reads as

$$\frac{d}{dt} \|f(t) - \mathcal{M}\|_{\mathcal{H}}^2 = \langle \mathcal{L}f(t), f(t) \rangle_{\mathcal{H}}.$$

Consequently, the Dirichlet form $B(f) = -\langle \mathcal{L}f, f \rangle_{\mathcal{H}}$ plays the role of entropy-dissipation functional and the existence of a spectral gap $|\lambda_1|$ is equivalent to the following coercivity estimate for $B(f)$:

$$B(f) \geq -|\lambda_1| \|f\|_{\mathcal{H}}^2 \quad \forall f \perp \text{span}(\mathcal{M}).$$

One deduces easily the following corollary on the exponential trend towards equilibrium:

Corollary 3.7. Let $f_0(v) \in \mathcal{H}$ and let $f(t, v)$ be the unique solution to the linear homogeneous Boltzmann equation (3.2). Then, there is some constant $C \geq 0$ such that

$$\|f(t, \cdot) - \mathcal{M}\|_{\mathcal{H}} \leq C \exp(-|\lambda_1|t) \|f_0 - \mathcal{M}\|_{\mathcal{H}}, \quad \text{for any } t \geq 0,$$

where $0 < |\lambda_1| \leq \nu_0$ is provided by Theorem 3.6.

We refer the reader to [13] for details on the matter, and in particular, for an explicit estimate of the spectral gap $|\lambda_1|$.

4. APPLICATION TO THE HONEST SOLUTIONS OF THE BOLTZMANN EQUATION

4.1. Conservative solutions. We are interested in this section in applying the result of the previous section to prove the existence of honest solutions to the linear Boltzmann equation for dissipative hard-spheres

$$\partial_t f(x, v, t) + v \cdot \nabla_x f(x, v, t) = Q(f)(x, v, t), \quad (4.1)$$

with initial condition

$$f(x, v, 0) = f_0(x, v) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3, dx \otimes dv),$$

where the collision operator Q is given by Eq. (1.2). Recall that the streaming operator T_0 is defined by

$$\mathcal{D}(T_0) = \{f \in \mathcal{X}, v \cdot \nabla_x f \in \mathcal{X}\}, \quad T_0 f = -v \cdot \nabla_x f$$

where $\mathcal{X} = L^1(\mathbb{R}^3 \times \mathbb{R}^3, dx \otimes dv)$. One can define then the multiplication operator Σ by

$$\mathcal{D}(\Sigma) = \{f \in \mathcal{X}, \sigma f \in \mathcal{X}\}, \quad \Sigma f(x, v) = -\sigma(v) f(x, v)$$

where, as in the previous Section, $\sigma(v)$ is the collision frequency corresponding to *dissipative hard spheres* interactions and given by Eq. (2.7). The following generation result is well-known [3]

Theorem 4.1. *The operator T_0 generates a C_0 -semigroup of isometries $(U(t))_{t \geq 0}$ of \mathcal{X} given by:*

$$U(t)f(x, v) = f(x - tv, v), \quad t \geq 0.$$

The operator $A = T_0 + \Sigma$ with domain $\mathcal{D}(A) = \mathcal{D}(T_0) \cap \mathcal{D}(\Sigma)$ is the generator of a contractions C_0 -semigroup $(V(t))_{t \geq 0}$ given by

$$V(t)f(x, v) = \exp(-\sigma(v)t)f(x - tv, v), \quad t \geq 0.$$

Let us define now K as the gain operator Q^+ endowed with the domain of A :

$$\mathcal{D}(K) = \mathcal{D}(A), \quad Kf(x, v) = Q^+(f)(x, v) = \epsilon^{-2} \int_{\mathbb{R}^3 \times \mathbb{S}^2} |q \cdot n| f(x, v_\star) \mathcal{M}_1(w_\star) dw dn.$$

It is clear from (1.5) that, for any $f \in \mathcal{D}(K)$,

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} (Af + Kf) dx dv = 0, \quad (4.2)$$

while $Kf \geq 0$ for any $f \in \mathcal{D}(K)$, $f \geq 0$. Then, the following generation result is a direct consequence of [1, 19]:

Theorem 4.2. *There exists a positive contractions semigroup $(Z(t))_{t \geq 0}$ in \mathcal{X} whose generator G is an extension of $A + K$. Moreover, $(Z(t))_{t \geq 0}$ is minimal, i.e. if $(T(t))_{t \geq 0}$ is a positive C_0 -semigroup generated by an extension of $A + K$, then $T(t) \geq Z(t)$ for any $t \geq 0$.*

The natural question is now to determine whether the "formal" mass conservation identity (1.6) can be made rigorous. Namely, one aims to prove that, for any nonnegative $f \in \mathcal{X}$, the following holds:

$$\|Z(t)f\| = \|f\|, \quad \forall t > 0.$$

The important point to be noticed is the following. If $G = \overline{A + K}$, then any function $\varphi \in \mathcal{D}(G)$ can be approximated by a sequence $(\varphi_n)_n \subset \mathcal{D}(A + K) = \mathcal{D}(A)$ such that $\varphi_n \rightarrow \varphi$ and $(A + K)\varphi_n \rightarrow G\varphi$ as $n \rightarrow \infty$. In particular, (4.2) implies that

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} G\varphi \, dx dv = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^3 \times \mathbb{R}^3} (A + K)\varphi_n \, dx dv = 0, \quad \forall \varphi \in \mathcal{D}(G).$$

Now, for any given initial datum $f_0 \in \mathcal{D}(G)$, $f_0 \geq 0$, the solution $f(t) = Z(t)f_0$ of (4.1) is such that

$$\frac{d}{dt} \|f(t)\| = \int_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{d}{dt} f(t) \, dx dv = \int_{\mathbb{R}^3 \times \mathbb{R}^3} Gf(t) \, dx dv = 0,$$

i.e.

$$\|f(t)\| = \|f_0\|, \quad \forall t \geq 0.$$

This means that, if $G = \overline{A + K}$, then the solutions to the linear Boltzmann equation (4.1) are conservative. On the other hand, if G is a larger extension of $A + K$ than $\overline{A + K}$, then there may be a loss of particles in the evolution (see [3] for the matter as well as [2] for examples of transport equation for which such a loss of particles occurs because of boundary conditions). Precisely, if $G \neq \overline{A + K}$ then there exists $f_0 \in \mathcal{X}$, $f_0 \geq 0$ such that

$$\|Z(t)f_0\| < \|f_0\| \quad \text{for some } t > 0.$$

This shows that the determination of the domain $\mathcal{D}(G)$ of G is of primary importance in the study of the Boltzmann equation. This is the main concern of the so-called *substochastic perturbation theory of C_0 -semigroups* [3].

To prove the honesty of the C_0 -semigroup $(Z(t))_{t \geq 0}$, we will adopt the strategy developed first in [1] and systematized in [3]. More precisely, we will show that the gain operator K fullfils the assumption of [1]:

Proposition 4.3. *There exists $C > 0$ such that, for any fixed $\varrho > 0$,*

$$\operatorname{ess\,sup}_{|v-u_1| \leq \varrho} \int_{|v'-u_1| \geq \varrho} k(v', v) \, dv' \leq C.$$

Proof. Since the kernel $k(v, v')$ differs from the corresponding one for classical gas, except from numerical constants, one can apply *mutatis mutandis* the technical calculations of [1, Section 4.1] (see also [3, p. 329-330]) to get the desired estimate. \square

As a consequence, one deduces immediately from [1], the main result of this section:

Theorem 4.4. *The domain $\mathcal{D}(G)$ of G is given by*

$$\mathcal{D}(G) = \overline{\mathcal{D}(A + K)}.$$

In particular, the C_0 -semigroup $(Z(t))_{t \geq 0}$ is honest and

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} Z(t)f(x, v) \, dx dv = \int_{\mathbb{R}^3 \times \mathbb{R}^3} f(x, v) \, dx dv \quad \text{for any } f \in \mathcal{X} \text{ and any } t > 0.$$

4.2. Consequence on the entropy production. The above result (Theorem 4.4) allows to give a rigorous proof of the linear H -Theorem of [10]. In order to stay in the formalism of [10], we shall restrict ourselves to the space-homogenous case. Precisely, let Y denote the set of functions depending only on the velocity and integrable with respect to velocities:

$$Y = L^1(\mathbb{R}^3, dv),$$

equipped with its natural norm $\|\cdot\|_Y$. For any nonnegative f and g in Y , we define the *information* of f with respect to g by

$$\mathcal{I}(f|g) = \int_{\mathbb{R}^3} \left(f(v) \ln f(v) - f(v) \ln g(v) \right) dv$$

with the conventions $0 \ln 0 = 0$ and $x \ln 0 = -\infty$ for any $x > 0$. This means that the information is nothing but the entropy functional H_Φ for the particular choice of $\Phi(s) = s \ln s$. One recalls the main result of [18]:

Theorem 4.5. *Let U be a stochastic operator of Y , i.e. U is a positive operator such that $\|Uf\|_Y = \|f\|_Y$ for any $f \in Y, f \geq 0$. Then,*

$$\mathcal{I}(Uf|Ug) \leq \mathcal{I}(f|g)$$

for any nonnegative f, g in Y . In particular, if $g \in Y$ is a nonnegative fixed point of U then,

$$\mathcal{I}(Uf|g) \leq \mathcal{I}(f|g), \quad \forall f \in Y, f \geq 0.$$

According to the results of the previous section, it is not difficult to see that the restriction of $(Z(t))_{t \geq 0}$ to Y is a C_0 -semigroup of stochastic operators of Y . Since the unique equilibrium state $\mathcal{M} \in Y$ is space independent, one sees that $(T_1 + K)\mathcal{M} = 0$ and, in particular,

$$Z_Y(t)\mathcal{M} = \mathcal{M}, \quad \forall t \geq 0.$$

Combining this with Theorem 4.5, one obtains a rigorous proof of the H -Theorem [10, Theorem 5.1]:

Theorem 4.6. *Let $f_0 \in Y$ be a given nonnegative (space homogeneous) distribution function with unit mass, i.e. $\|f_0\|_Y = 1$. Assume that $\mathcal{I}(f_0|\mathcal{M}) < \infty$, then*

$$\frac{d}{dt} \mathcal{I}(f(t)|\mathcal{M}) \leq 0, \quad (t \geq 0),$$

where $f(t) = Z_Y(t)f_0 = Z(t)f_0$ is the unique solution to (4.1) with $f(0) = f_0$.

REFERENCES

- [1] L. Arlotti, A perturbation theorem for positive contraction semigroups on L^1 -spaces with applications to transport equations and Kolmogorov's differential equations. *Acta Appl. Math.* **23**, 129–144 (1991).
- [2] L. Arlotti, B. Lods, Substochastic semigroups for transport equations with conservative boundary conditions, *J. Evol. Equations* **5** 485–508, 2005.
- [3] J. Banasiak, L. Arlotti, **Perturbations of positive semigroups with applications**, Springer-Verlag, 2005.
- [4] N. V. Brilliantov, T. Pöschel, **Kinetic theory of granular gases**, Oxford University Press, 2004.
- [5] T. Carleman, Problèmes mathématiques dans la théorie cinétique des gaz. *Publications Scientifiques de l'Institut Mittag-Leffler*, **2** (1957).

- [6] C. Cercignani, R. Illner and M. Pulvirenti, **The mathematical theory of dilute gases**, Springer–Verlag, New York (1994).
- [7] M. H. Ernst, R. Brito. Scaling solutions of inelastic Boltzmann equation with over–populated high energy tails. *J. Statist. Phys.* **109**, 407–432 (2002).
- [8] H. Grad, Asymptotic theory of the Boltzmann equation. II. *Rarefied Gas Dynamics (Proc. 3rd Internat. Sympos., Palais de l’UNESCO, Paris, 1962)*, Vol. I, 26–59 (1963).
- [9] F. A. Molinet, Existence, uniqueness and properties of the solutions of the Boltzmann kinetic equation for a weakly ionized gas, *J. Math. Phys.* **18**, 984–996 (1977).
- [10] B. Lods, G. Toscani, The dissipative linear Boltzmann equation for hard spheres, *J. Statist. Phys.* **117**, 635–664 (2004).
- [11] P. A. Martin, J. Piacieski, Thermalization of a particle by dissipative collisions, *Europhys. Lett.* **46**, 613–616 (1999).
- [12] S. Mischler, C. Mouhot, Cooling process for inelastic Boltzmann equations for hard spheres, Part II: Self-similar solutions and tail behavior, *J. Statist. Phys.* **124**, 703–746 (2006).
- [13] C. Mouhot, G. Toscani, Relaxation rate and diffusive limit for inelastic scattering Boltzmann models, Work in progress.
- [14] R. Pettersson, On solutions to the linear Boltzmann equation for granular gases, *Transp. Theory Statist. Phys.* **33**, 527–543 (2004).
- [15] F. Poupaud, Runaway phenomena and fluid approximation under high fields in semiconductor kinetic theory, *Z. Angew. Math. Mech.* **72**, 359–372 (1992).
- [16] G. Spiga, G. Toscani, The dissipative linear Boltzmann equation, *Appl. Math. Lett.* **17**, 255–301 (2004).
- [17] C. Villani, Mathematics of granular materials, *J. Statist. Phys.* **124**, 781–822 (2006).
- [18] J. Voigt, Stochastic operators, information, and entropy, *Comm. Math. Phys.* **81**, 31–38 (1981).
- [19] J. Voigt, On substochastic C_0 -semigroups and their generators, *Transp. Theory Stat. Phys.* **16**, 453–466, (1987).

LUISA ARLOTTI, DIPARTIMENTO DI INGEGNERIA CIVILE, UNIVERSITÀ DI UDINE, VIA DELLE SCIENZE 208, 33100 UDINE, ITALY.

E-mail address: luisa.arlotti@uniud.it

BERTRAND LODS, LABORATOIRE DE MATHÉMATIQUES, CNRS UMR 6620, UNIVERSITÉ BLAISE PASCAL (CLERMONT–FERRAND 2), 63177 AUBIÈRE CEDEX, FRANCE.

E-mail address: lods@math.univ-bpclermont.fr