

## First archaeomagnetic secular variation curve for the Iberian peninsula: Comparison with other data from Western Europe and with global geomagnetic field models

Miriam Gomez-Paccard, Annick Chauvin, Philippe Lanos, G. Mcintosh, Maria-Luisa. Osete, Gianluca Catanzariti, V.C. Ruiz-Martinez, J.I. Nuñez

### ▶ To cite this version:

Miriam Gomez-Paccard, Annick Chauvin, Philippe Lanos, G. Mcintosh, Maria-Luisa. Osete, et al.. First archaeomagnetic secular variation curve for the Iberian peninsula: Comparison with other data from Western Europe and with global geomagnetic field models. Geochemistry, Geophysics, Geosystems, 2006, 7 (12), pp.Q12001. 10.1029/2006GC001476. hal-00128411

## HAL Id: hal-00128411 https://hal.science/hal-00128411

Submitted on 9 Jun2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## First archaeomagnetic secular variation curve for the Iberian Peninsula: Comparison with other data from western Europe and with global geomagnetic field models

#### M. Gómez-Paccard

Géosciences-Rennes, CNRS, UMR 6118, Université de Rennes 1, Rennes, France

Facultad de Ciencias Físicas, Universidad Complutense de Madrid, Madrid, Spain

Civilisations Atlantiques et Archéosciences, CNRS, UMR 6566, Université de Rennes 1, Rennes, France

Now at Facultad de Geología, Universidad de Barcelona, Zona Universitaria de Pedralbes, E-08028 Barcelona, Spain. (miriam.gomez@univ-rennes1.fr)

#### A. Chauvin

Géosciences-Rennes, CNRS, UMR 6118, Université de Rennes 1, Rennes, France (annick.chauvin@univ-rennes1.fr)

#### P. Lanos

Civilisations Atlantiques et Archéosciences, CNRS, UMR 6566, Université de Rennes 1, Rennes, France (philippe.lanos@univ-rennes1.fr)

#### G. McIntosh, M. L. Osete, G. Catanzariti, V. C. Ruiz-Martínez, and J. I. Núñez

Facultad de Ciencias Físicas, Universidad Complutense de Madrid, Ciudad Universitaria, E-28040 Madrid, Spain (gregc@fis.ucm.es; mlosete@fis.ucm.es; gcatanza@fis.ucm.es; vcarlos@fis.ucm.es; jnuneza.hgugm@salud.madrid.org)

[1] A first secular variation (SV) curve for the Iberian Peninsula was computed by hierarchical Bayesian method using a total of 134 archaeomagnetic directions with ages ranging from -775 to 1959 A.D. A general agreement is observed between the Iberian curve and the French and German SV curves, although some interesting differences were found, such as the occurrence of lower inclinations between the 11th and 14th centuries in the Iberian curve. The analysis of these three reference curves indicates that SV in western Europe is characterized by three major directional changes at -125, 200, and 1350 A.D. It is suggested that these cusps are regional features of the geomagnetic field. The Iberian curve has been compared with the predictions of the Jackson, CALSK7K.2, and Hongre global models. Despite large differences recognized between these models, even for the dipolar terms, they predict reasonably well the Iberian archaeomagnetic SV.

Components: 9126 words, 11 figures, 2 tables.

Keywords: archaeomagnetism; geomagnetic secular variation; Bayesian statistics; Iberian Peninsula.

**Index Terms:** 1503 Geomagnetism and Paleomagnetism: Archeomagnetism; 1522 Geomagnetism and Paleomagnetism: Paleomagnetic secular variation; 1532 Geomagnetism and Paleomagnetism: Reference fields: regional, global.

Received 11 September 2006; Accepted 25 September 2006; Published 5 December 2006.



Gómez-Paccard, M., A. Chauvin, P. Lanos, G. McIntosh, M. L. Osete, G. Catanzariti, V. C. Ruiz-Martínez, and J. I. Núñez (2006), First archaeomagnetic secular variation curve for the Iberian Peninsula: Comparison with other data from western Europe and with global geomagnetic field models, *Geochem. Geophys. Geosyst.*, 7, Q12001, doi:10.1029/2006GC001476.

#### 1. Introduction

[2] The variation of the geomagnetic field in the archaeological past can be obtained from paleomagnetic directions determined from heated and well dated archaeological structures. Archaeomagnetic data are presently the most precise highresolution paleomagnetic data for the last millennia (data from lakes and sediments may be smoothed and/or offset due to the magnetization acquisition process and to sedimentation gaps). These data are used to build secular variation (SV) reference curves that provide knowledge of geomagnetic field variations at regional and global scales for periods covering the last few millennia. Along with historical data, detailed SV curves can also be used for dating purposes. Several SV curves are now available for Europe, including Great Britain [Batt, 1997], Bulgaria [Kovacheva et al., 1998], France [Gallet et al., 2002], Hungary [Márton, 2003], Germany [Schnepp and Lanos, 2005] and Austria [Schnepp and Lanos, 2006]. Despite the high number of archaeological excavations carried out in Spain only limited data are available for this region. In a recent work, Gómez-Paccard et al. [2006] provide a compilation of 5 previous results and present 58 new archaeomagnetic directions. In the present study this data set is used, along with data from southern France and northern Morocco, to propose a first SV curve for the Iberian Peninsula. The French and Moroccan data can be used in constructing the SV curve due to their geographical proximity to the Iberian Peninsula. The hierarchical bivariate method by moving average technique [Le Goff, 1990; Le Goff et al., 1992; Daly and Le Goff, 1996; Lanos et al., 2005] and the hierarchical Bayesian modeling [Lanos, 2004] are implemented in order to determine the best way to calculate the Iberian curve. To determine the characteristics of the geomagnetic SV in western Europe over the last two millennia, the Iberian curve has been compared with the French and German archaeomagnetic reference curves. Finally, a comparison between available geomagnetic field global models [Hongre et al., 1998; Jackson et al., 2000; Korte and Constable, 2005] has been conducted in order to test the reliability of their predictions.

# 2. Available Data From Spain, France, and Morocco

#### 2.1. Spanish Data

[3] Gómez-Paccard et al. [2006] compiled the available data for Spain and presented 58 new archaeomagnetic directions, carefully describing the sampling, experimental procedures and dating constraints of the new directions. Samples were taken as oriented blocks or drilled directly from the archaeological heated structures. Classical thermal, alternating field demagnetization or Thellier method were applied in order to determine the characteristic directions for each structure. In most cases, high-quality, single-component remanence vectors were obtained. For some structures, which were heated to relatively low temperatures, more than 1 component could be recognized and great circle analysis was required. The characteristic directions and their statistical parameters were calculated using principal component [Kirschvink, 1980] or great circle [McFadden and McElhinny, 1988] analysis and Fisher [1953] statistics. As proposed by Lanos et al. [2005], a hierarchical structure was followed in the calculation of the 58 new directions. Together with 5 previous data from Spain, a total of 63 archaeomagnetic directions are available from Spain (see Table 1 and Figure 1).

#### 2.2. French and Moroccan Data

[4] Archaeomagnetic research started in France with the work of E. Thellier [*Thellier*, 1938], who later published the first SV curve for France for the last two millennia [*Thellier*, 1981]. This curve has been revised, completed and extended by *Bucur* [1994]. Since then, new archaeomagnetic directional results have been collected by *Moutmir* [1995], *Chauvin et al.* [2000], and *Gallet et al.* [2002], who presented a new SV curve for France. Together they provide a high-quality collection of archaeomagnetic data. In this study, only the data from sites with a geographical proximity to the Iberian Peninsula (within a 900 km radius of

Table 1. Archaeomagnetic Direction	ons Froi	n Spa	nish, Fre	nch, and	More	occan	Sites <sup>a</sup>					
Name	tmin	t <sub>max</sub>	tmean	t <sub>bayesian</sub>	I <sub>s</sub> , deg	D <sub>s</sub> , deg	N	35 k	Lat. °N	, Long., °E	Stratigraphic Constraints	Source
Dlaza da Morros (DI M)	150	50	100	08 7	6 85	Sit A O	es (Spo	1 197	305	0 7 00		
Ampurias (AMP)	-200	100	-50	-98.7	63.5	-8.5	0 4 ⊦ –	5 259	2 42.1	2 3.13		Thellier [1981]
El Monastil (MON)	-50	35	-7.5	-5.6	57.3	0.3	9 2.	2 54(	5 38.4	7 -0.79		, L
El Gallinero (GA)	40	50	45	45.0	53.4	3.8	7.4.	9 15(	36.5	3 -6.19		
Villa del Pañuelo I (VILI)	0	100	50	51.4	57.1	-5.5	25 2.	6 11(	5 40.3	0 - 3.40		Oyamburu et al. [1996]
Calahorra, La Maja (LMA)	0	100	50	54.9	58.4	-0.8	29 2.	1 128	8 42.2	7 -2.02		Parés et al. [1992]
Costalita (COS)	0	100	50	46.8	54.3	3.7	14 6.	0 46	36.4	2 -5.15		1
Venta del Carmen (VC)	80	90	85	85.3	46.7	2.1	27 2.	3 14(	5 36.1	8 -5.49		
Villares Andujar (VIA)	50	150	100	109.2	50.5	1.4	9.3.	2 25	9 38.0	6 -4.04		
Cartuja I (CAR-HI)	50	150	100	86.3	55.3	-1.7	7 3.	5 293	3 37.1	8 -3.10		
Cartuja II (CAR-HII)	50	150	100	100.1	52.2	-3.1	7 5.	4 12	5 37.1	8 -3.10		
Cartuja III (CAR-HIII)	50	150	100	96.5	52.3	2.9	5 3	9 392	2 37.1	8 -3.10		
Patio Cardenal I (PARI)	90	130	110	109.0	53.0	-2.4	10 2.	2 482	4 37.3	8 -5.98		
Patio Cardenal III (PAR3)	90	130	110	108.6	54.1	-4.1	63.	7 324	4 37.3	8 -5.98		
Patio Cardenal IV (PAR4)	90	130	110	109.1	53.3	-2.7	7 1.	9 102	6 37.3	8 -5.98		
Patio Cardenal V (PAR5)	90	130	110	109.0	53.5	-4.3	17 2.	6 193	3 37.3	8 -5.98		
Baelo Claudia (BC)	100	150	125	128.2	46.7	-2.2	14 3.	6 14	5 36.0	3 -5.62		
Gallineras (GAL)	100	220	160	140.5	55.8	-0.1	7 4	6 17(	5 36.4	7 -6.18		
Arva (ARV)	150	250	200	199.1	51.8	-3.7	17 1.	3 68.	2 37.6	0 -5.50		Evans, private communication
Setla Mirarosa Miraflor (DENA)	220	250	235	234.9	52.6	-4.5	10 1.	99 66	9 38.8	6 0.02		
Hyppolytus (HIP)	225	325	275	274.4	52.6	-4.0	18 3.	4 102	4 40.4	8 -3.32		
Villa del Pañuelo II (VIL2)	250	350	300	296.6	51.7	-2.7	31 2.	2 13	1 40.3	0 -3.40		Oyamburu et al. [1996]
Valeria (VAL)	270	330	300	299.7	48.0	-5.4	7 9.	7 58	39.7	7 -2.13		
Puente Grande I (PG1)	400	410	405	404.9	46.4	-14.1	13 7.	333	36.1	8 -5.49		
Puente Grande II (PG2)	400	410	405	405.0	52.5	0.3	15 3.	4 12	36.1	8 -5.49		
Ramon Ortega II (RO2)	1000	1050	1025	1027.7	50.9	17.3	14 4	0 100	38.8	6 0.12		
Murcia c/Sagasti (MURG)	1000	1100	1050	1033.2	51.3	21.7	8.	5 143	7 37.9	8 -1.12		
Ramon Ortega I (RO1)	1050	1100	1075	1079.0	47.9	15.8	16 1.	9 38(	38.8	6 0.12		
Cabrera d'Anoia (CDAP)	1010	1220	1115	1174.4	50.6	3.6	ж	6 23(	5 41.5	0 1.50		
Cabrera d'Anoia (CDAU)	1029	1203	1116	1180.0	45.7	9.6	6 1.	3 276	5 41.5	0 1.50		
Murcia c/Puxmarina (MURO)	1100	1200	1150	1119.6	44.7	13.6	63.	8 308	3 37.9	8 -1.12 b	efore MURN	
Murcia c/Puxmarina (MURN)	1100	1200	1150	1135.8	45.1	14.0	6 5	1 103	8 37.9	8 -1.12 a	fter MURO and before MURM	
Murcia c/Puxmarina (MURM)	1100	1200	1150	1151.7	45.5	14.9	7 2.	1 81	1 37.9	8 -1.12 8	fter MURN and before MURL	
Murcia c/Puxmarina (MURL)	1100	1200	1150	1170.8	44.2	16.2	9	2 184	2 37.9	8 -1.12 8	fter MURM	
Murcia c/Puxmarina (MURI)	1100	1200	1150	1157.4	41.7	15.2	7 4	1 21:	5 37.9	8 -1.12		
Murcia c/Puxmarina (MURK)	1100	1200	1150	1146.4	45.2	14.1	9	2 54'	7 37.9	8 -1.12		
Murcia c/Puxmarina (MURH)	1100	1200	1150	1134.3	46.4	17.3	10 2.	7 33(	37.9	8 -1.12		
Cabrera d'Anoia (CDAJ)	1056	1262	1159	1217.0	45.2	9.0	8 13	7 41(	5 41.5	0 1.50		
Cabrera d'Anoia (CDAH)	1043	1281	1162	1215.1	46.4	10.7	5 1.	3 346	1 41.5	0 1.50		
Magisterio I (MAGI)	1250	1300	1275	1270.6	42.6	11.5	5. 4.	8 50(	9.04	3 -3.16		
Magisterio II (MAGII)	1250	1300	1275	1275.9	47.3	4.1	6 6	1 179	9 40.6	3 -3.16		
Calatrava la Vieja (CALA)	1275	1300	1287.5	1286.6	44.9	11.4	8 8	9 35(	5 39.0	2 -3.82		

3

]]

Num         L         D <thd< th="">         D         D         D</thd<>	Table 1. (continued)															
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Name	tmin	t <sub>max</sub>	t <sub>mean</sub>	t <sub>bayesian</sub>	I <sub>s</sub> , deg	Ds, deg	Z	<sup>65</sup> 1	J C	tt., Lor N	ıg., E Stratig	straphic Constraints	Source		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Calatrava la Vieia (CALB)	1275	1300	1287.5	1287.4	44.4	7.1	10 2	9 28	39	02 - 3	.82				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Valancia Vallutare (VALN)	1738	1350	1204	1310.6	46.6	с -	0	с 18	50 30	47 _0	27				
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Guadalaise (GIIA1)	1275	1275	1200	hot need	8 2 3	- 7 - 7	- c \ _	i & 5 E		60 3	00				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Castillo de San Romialdo (CSR)	1100	1500	1300	1334 0	46 0		1 (~	2 C	26.20	30 -6	10				
Weils (MUL)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (56)         (57)         (56)         (57)         (56)         (57)	Valencia Velluters (VALD	1238	1400	1319	1329.6	46.4	4	,		5 39	47 -0	37				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Av. Blas de Infante (BI)	1369	1369	1369	1369.0	41.9	0.4	10	. 7	)6 36 36	$\frac{13}{-5}$	45				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Valencia Velluters (VALM)	1300	1450	1375	1342.8	47.0	7.2	6	0 27	33 39	47 -0	.37 before VA	LK			
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Valencia Velluters (VALK)	1300	1450	1375	1383.5	44.2	3.0	9	.3 16	06 39	47 -0	.37 after VAL	Μ			
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Llano las Damas (LLD)	1400	1415	1407.5	1407.2	37.0	9.6	14	0.0	8 35	89 -5	.30				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Huerta Rufino (HR)	1400	1415	1407.5	1407.0	35.9	7.8	5	.1 4	17 35	89 -5	.30				
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Calatrava la Vieja (CALC)	1400	1420	1410	1410.3	47.0	3.0	8	.0	90 39	02 -3	.82				
$ \begin{array}{c} \mbox{Termer} \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$	Paterna c/Huertos (PATA)	1450	1500	1475	1487.6	56.4	3.6	10	52	39.	50 -0	.43 before PA	ΓB			
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Paterna Testar del Moli (TMO)	1490	1540	1515	1520.1	55.7	7.5	16	.0 32	t3 39	50 -0	.43				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Paterna Testar del Moli (PATJ)	1429	1611	1520	1593.6	62.2	6.6	16	.1 12	01 39	50 -0	.43				
	Paterna Testar del Moli (PATH)	1450	1600	1525	1515.7	53.5	7.3	10	.7 8	31 39	50 -0	.43				
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Paterna c/Huertos (PATB)	1525	1650	1587.5	1625.6	64.1	5.8	=	.6 8	27 39	50 -0	.43 after PAT/	~			
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Valencia Velluters (VALL)	1575	1625	1600	1591.5	56.6	9.1	1	.9 5.	57 39	47 -0	.37				
Huerts del Carmen (GUA2)         1825         1845         53.3         60.0 $= 3.20$ $= 2.93$ $= 3.00$	Monasterv at Yuste (YUS1)	1784	1814	1799	1835.2	55.5	-14.3	7	.6	73 40	10 -5	.70				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Huertas del Carmen (GUA2)	1825	1845	1835	1800.4	615	- 1 1 1	. 4	с с	88 40	60 -3	20				
$ \begin{array}{c cccc} \mbox{worres} \mb$	Palacio de Perales (AI)	1830	1910	1870	1865 5	63.0	-14.7		- i -	0 70	10 - 3	10				
	Monasterv at Vilste (VUS2)	1959	1959	1959	1959.0	58.2	-11.1	s v s v	5 K	40	10 - 5	01				
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	(2001) mont in finantions					1		2	;	-						
					Site	S T < 0	600 km	(Fra	nce ai	to Mor	occo)					
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Saint Florence	-325	-275	-300.0	-299.4	66.4	-2.8	1	.2	36 44	.8 .0	0		Bucur [1994]		
Aguillon F1+3+3 $-120$ $-80$ $-1000$ $-995$ $624$ $-24$ $45$ $11$ $1825$ $443$ $0.3$ $Bucw$ $1994$ Aguillon F4 $-100$ $-900$ $633$ $-26$ $19$ $08$ $1607$ $443$ $0.3$ $Bucw$ $1994$ Montans 4 $-100$ $-900$ $633$ $-26$ $10$ $1096$ $433$ $0.3$ $Bucw$ $1994$ Montans 6 $-80$ $-900$ $-900$ $633$ $-212$ $610$ $-900$ $633$ $216$ $634$ $319$ $Bucw$ Montans 6 $-900$ $-900$ $530$ $-903$ $633$ $-12$ $7$ $12$ $634$ $394$ Montans 6 $-900$ $-900$ $530$ $233$ $433$ $2.3$ $360$ $355$ $600$ $Bucw$ $1994$ Montans 6 $-100$ $-500$ $-900$ $530$ $243$ $613$ $133$ $2.3$ $Bucw$ $1994$ Montans 16 $-1100$ $-500$ $230$ $543$ $613$ $333$ $52$ $600$ $Bucw$ $1994$ Montans 6 $-100$ $-900$ $580$ $-933$ $543$ $-12$ $712$ $209$ $343$ $2.3$ $209$ Montans 6 $-100$ $900$ $580$ $230$ $100$ $333$ $610$ $333$ $610$ $1994$ Montans 7 $-100$ $175$ $220$ $293$ $513$ $133$ $225$ $100$ Montans 7 $175$ $225$ $2000$ $2$	Dchar Jdid. the citadel	-300	-100	-200.0	-186.9	53.0	1.8	10 5	.1	5 35	.5 –	5.0		Kovacheva [1984]		
Ağuillon F4 $-120 - 80 -1000 -997$ $64.1 - 2.6 19$ $0.8 1607$ $44.3 0.3$ $Bucw$ [1994]Maguillon F4 $-100 - 80 -900$ $-900 -997$ $64.1 - 2.6 11$ $100 166 4.33$ $1.8$ $Bucw$ [1994]MagreeLimoux II (bur II) $-20 - 3$ $-37 + 21.6 652$ $-3.7 + 4.0 + 0.2$ $Bucw$ [1994]Limoux II (bur II) $-20 - 5 - 12.5 - 12.6 620$ $-0.3 7 + 21.6 652$ $43.9 1.8$ $Bucw$ [1994]Montans 6 $-200 - 5 - 12.5 - 12.6 620$ $-0.3 7 + 21.6 653$ $43.9 1.8$ $Bucw$ [1994]Montans 6 $-100 0 - 500 - 550 - 24.9 62.3 -11.2 7 1.2 2054 43.9 1.8$ $Bucw$ [1994]Montans 1 $-20 - 30 - 25.0 - 24.9 62.3 -11.2 7 1.2 2054 43.9 1.8$ $Bucw$ [1994]Montans 1 $-100 0 - 500 - 900 56.8 -0.1 12 1.6 673 43.9 1.8$ $Bucw$ [1994]Montans 1 $-100 0 -500 0 -500 0 56.8 -0.1 12 1.1 031 43.3 2.8$ $Bucw$ [1994]Montans 1 $-100 0 -500 0 -500 0 56.8 -0.1 12 1.1 031 43.3 2.8$ $Bucw$ [1994]Montans 1 $-100 0 -500 0 -900 56.8 -0.1 13 1.1 031 43.3 0.16$ $Bucw$ [1994]Montans 1 $-100 0 0 -500 0 56.8 -0.1 13 1.1 031 43.5 -1.0$ $Bucw$ [1994]Barat de Vin $375 425 400 0 3977 53.4 1.0 9 0.6 575 44.8 -0.5$ $Bucw$ [1994]Sadirac SB4 $1350 1400 1375.6 11.0 9 0.8 2865 44.8 -0.5$ $Bucw$ [1994]Sadirac SB3 $1500 0 3977 53.4 1.0 9 0.6 575 64.3 -0.5$ $Bucw$ [1994]Sadirac SB4 $1350 1375.6 11.0 9 0.8 2865 44.8 -0.5$ $Bucw$ [1994]Sadirac SB4 $1350 1375.6 11.9 9 0.8 2865 44.8 -0.5$ $Bucw$ [1994]Sadirac SB3 $150$	Aiguillon F1+2+3	-120	-80	-100.0	-99.5	62.4	-2.4	45	.1 18	25 44	.3	3		Bucur $[1994]$		
	Aiguillon F4	-120	-80	-100.0	-99.7	64.1	2.6	19 (	8 16	07 44	.3	3		Bucur $[1994]$		
Lagrere $-80 - 40 - 60.0 - 599 6.29 - 3.7 14 0.9 1797 4.4 0.2$ Bucur [1994]Linoux II (four II) $-20 - 5 - 12.5 - 12.6 6.20 - 0.3 7 2.1 625 4.31 2.3$ $Bucur [1994]$ Linoux II (four II) $-20 - 5 - 12.5 - 12.6 6.20 - 0.3 7 2.1 6.52 4.31 2.3$ $Bucur [1994]$ Montans I $-100 0 - 50.0 - 49.3 5.43 - 6.10 7 1.2 205 4.3.9 1.8$ $Bucur [1994]$ Al Kouss (Chkakra) $-100 0 - 50.0 - 49.3 5.43 - 6.1 2 7 1.2 205 4.3.9 1.8$ $Bucur [1994]$ Montans I $-100 0 - 50.0 - 49.3 5.43 - 6.1 3 1.3 205 4.3.9 1.8$ $Bucur [1994]$ Abrens (Laure Minervois) $175 2.25 20.00 2000 5.68 - 0.3 16 1.1 1031 4.33 2.5$ $Bucur [1994]$ Abrens (Laure Minervois) $375 4.5 - 4.0 0 2000 5.68 - 0.3 16 1.1 1031 4.33 2.5$ $Bucur [1994]$ Barat de Vin $375 4.25 200.0 2000 5.68 - 1.0 8 1.5 1079 4.3.5 - 1.0$ $Bucur [1994]$ Barat de Vin $375 4.5 - 4.0 0 0 8.7 5.3.4 1.0 9 0.6 5769 4.3.6 1.4$ $Bucur [1994]$ Sadirac SB4 $1350 1400 13750 13775 6.51.0 1.9 10 0.8 2865 4.48 - 0.5$ $Bucur [1994]$ Barat de Vin $Bucur [1994]$ $Bucur [1994]$ Sadirac SB3 $1500 1525 1512.5 57.9 10.0 8 1.0 2.311 4.48 - 0.5$ $Bucur [1994]$ Bucur [1994] $Bucur [1994]$ $Bucur [1994]$ Sadirac SB4 $1500 1525 1512.5 57.9 10.0 8 1.0 2.317 4.48 - 0.5$ $Bucur [1994]$ Bucur [1994] $Bucur [1994]$ $Bucur [1994]$ Sadirac SB4 $1500 0 5525 1512.5 57.9 10.0 8 1.0 2.9 237 4.62 3.3$ $Aumir [1995]$ Bucur [1994] $Bucur [1994]$ $Bucur [1994]$ Sadirac SB4 $-7750 - 7750 - 7752 0 571 2.44 2.4 14.8 5.8 3.3$	Montans 4	-100	-80	-90.0	-90.0	63.5	-2.6	1	.0 16	96 43	.9	8		Bucur [1994]		
$ \begin{array}{llllllllllllllllllllllllllllllllllll$	Lagrere	-80	-40	-60.0	-59.9	62.9	-3.7	14 0	.9 17	97 44	.4	2		Bucur [1994]		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Limoux II (four II)	-20	-5	-12.5	-12.6	62.0	-0.3	5	.1	25 43	.1	3		Bucur [1994]		
Al Kouass (Chkakra) $-100 \ 0 \ -50.0 \ -49.3 \ 54.3 \ -6.1 \ 5 \ 3.3 \ 56 \ 0 \ 57.5 \ -6.0 \ 673 \ 43.9 \ 1.8 \ Bucur [1994]$ Montans I 140 160 150.0 150.1 56.0 $-0.1$ 12 1.6 673 43.9 1.8 Bucur [1994] Abrens (Laure Minervois) 175 225 200.0 56.8 $-0.3$ 16 1.1 1031 43.3 2.5 $Bucur [1994]$ Barat de Vin 375 425 400.0 397.9 57.5 $-4.3 \ 20 \ 1.7 \ 311 \ 43.9 \ 0.6 \ Bucur [1994]$ Lectoure 450 500.0 508.7 $6_{3.4} \ 1.0 \ 9 \ 0.6 \ 5769 \ 4.6 \ 1.4 \ Bucur [1994]$ Sadiras SB4 1.0 9 0.6 5769 4.4.8 $-0.5 \ Bucur [1994]$ Sadiras SB3 150 1375.0 1375.0 1373.6 51.0 1.9 10 0.8 2865 44.8 $-0.5 \ Bucur [1994]$ Bucur [1994] Sadiras CB4 1.0 9 0.6 5769 4.4.8 $-0.5 \ Bucur [1994]$ Sadiras CB4 1.0 0 2.2 231 44.8 $-0.5 \ Bucur [1994]$ Sadiras CB4 1.0 0 2.2 231 44.8 $-0.5 \ Bucur [1994]$ Sadiras CB4 1.0 0 2.9 235 44.8 $-0.5 \ Bucur [1994]$ Sadiras CB4 1.0 0 2.9 237 46.2 3.3 Bucur [1994] Sadiras CB4 1.0 0 2.9 237 46.2 3.3 Bucur [1994] Sucur [1995] Moutmir [1955] Steps r > 900 km (France and Morocco) Moutmir [1995] Moutmir [1995] Steps r > 00.0 -662.7 691 2.1.6 2.1 1.4 961 45.5 3.3 Moutmir [1955] Steps r > 00.0 -662.7 691 2.1.6 2.1 1.4 961 45.5 3.3 Moutmir [1955] Moutmir [1995] Steps r > 00.0 -662.7 691 2.1.6 2.1 1.4 961 45.5 3.3 Moutmir [1955] Moutmir [1955] Moutmir [1955] Steps r > 00.0 -662.7 691 2.1.6 2.1 1.4 961 45.5 3.3 Moutmir [1955] Moutmir [1955] Moutmir [1955] Moutmir [1955] Moutmir [1955] Moutmir [1955] Steps r > 00.0 -662.7 691 2.1.6 2.1 1.4 961 45.5 3.3 Moutmir [1955] Moutmir [1955] Mo	Montans 6	20	30	25.0	24.9	62.3	-1.2	7	.2 20	54 43	.9	8		Bucur [1994]		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Al Kouass (Chkakra)	-100	0	-50.0	-49.3	54.3	-6.1	<b>5</b> 60	.3	50 35	5 –(	5.0		Kovacheva [1984]		
Abrens (Laure Minervois)175225200.0260.056.8 $-0.3$ 161.1103143.32.5BucurBucur[1994]Barat de Vin375425400.0397.957.5 $-4.3$ 201.731143.90.6Bucur[1994]Lectoure375425400.0397.957.5 $-4.3$ 201.731143.90.6Bucur[1994]Lectoure375425400.0397.957.5 $-4.3$ 201.731143.90.6Sadirac SIA135014001375.051.01.9100.8236544.8 $-0.5$ Bucur[1994]Sadirac SI3150015251512.51512.257.910.081.0233144.8 $-0.5$ Bucur[1994]Sadirac SI3150015251512.51512.257.910.081.0233144.8 $-0.5$ Bucur150015251512.51512.257.910.081.0233746.23.3Sadirac SI3 $-850$ $-700$ 775.0 $-744.2$ $65.1$ 14.0102.923746.23.3Gannat $-850$ $-700$ 775.0 $-744.2$ $65.1$ 14.010.22355 $43.5$ 5.0Gannat $-850$ $-700$ $-775.0$ $-744.2$ $65.1$ $10.0$ $2.92$ $2.4$ $2.4$ $2.4$ $2.$	Montans I	140	160	150.0	150.1	56.0	-0.1	12	.6	73 43	.9	8		Bucur $[1994]$		
Barat de Vin400400400.0400.559.6-1.081.51079 $4.5.$ -1.0Bucur [1994]Lectoure375425400.0397.957.5-4.3201.731143.90.6Bucur [1994]Lectoure375550500.0508.763.41.090.6576943.61.4Bucur [1994]Sadirac SB413501375.01373.651.01.9100.8286544.8-0.5Bucur [1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8-0.5Bucur [1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8-0.5Bucur [1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8-0.5Sadirac SB3150015251512.51512.257.910.081.0233144.8-0.5Sadirac SB3150015251512.61512.257.910.081.0233144.8-0.5Satirac SB3-860-700-775.0-744.265.114.0102.923746.23.3Gannat-850-700-775.0-762.967.729.42.414.645.83.3Subat-555-550-562.5-561	Abrens (Laure Minervois)	175	225	200.0	200.0	56.8	-0.3	16	.1 10	31 43	.3 .2	5		Bucur [1994]		
Lectoure $375$ $425$ $400.0$ $397.9$ $57.5$ $-4.3$ $20$ $1.7$ $311$ $43.9$ $0.6$ $Bucur$ $[1994]$ Tolouse Place St. Etienne $450$ $550$ $500.0$ $508.7$ $63.4$ $1.0$ $9$ $0.6$ $5769$ $43.6$ $1.4$ $Bucur$ $[1994]$ Sadirac SB4 $1350$ $1375.0$ $1373.6$ $51.0$ $1.9$ $10$ $0.8$ $2865$ $44.8$ $-0.5$ $Bucur$ $[1994]$ Sadirac SB3 $1500$ $1525$ $1512.5$ $1512.2$ $57.9$ $10.0$ $8$ $1.0$ $2331$ $44.8$ $-0.5$ $Bucur$ $[1994]$ Sadirac SB3 $1500$ $1525$ $1512.5$ $1512.2$ $57.9$ $10.0$ $8$ $1.0$ $2331$ $44.8$ $-0.5$ $Bucur$ $[1994]$ Sadirac SB3 $1500$ $1525$ $1512.5$ $1512.2$ $57.9$ $10.0$ $8$ $1.0$ $2331$ $44.8$ $-0.5$ Sadirac SB3 $1500$ $1525$ $1512.5$ $1512.5$ $1512.6$ $124.0$ $0.2$ $237$ $46.2$ $3.3$ Gannat $-850$ $-700$ $-775.0$ $-744.2$ $65.1$ $14.0$ $10$ $29.2$ $45.2$ $3.3$ Moutmir $1995$ $170$ $-775.0$ $-762.9$ $67.7$ $29.4$ $24$ $24$ $24.5$ $3.5$ Size $-550$ $-560.6$ $-560.0$ $-662.7$ $69.1$ $21.6$ $21.6$ $45.5$ $3.3$ $Moutmir<[1995]$ Light	Barat de Vin	400	400	400.0	400.5	59.6	-1.0	8	.5 10	79 43	.5	0.1		Bucur [1994]		
Tolouse Place St. Etienne450550500.0508.763.41.090.6576943.61.4BucurBucurBucurBucurBucurBucurBucurBucurBucur[1994]Sadirac SB413501375.01375.01373.651.01.9100.8286544.8 $-0.5$ BucurBucur[1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8 $-0.5$ Bucur[1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8 $-0.5$ Bucur[1994]Sadirac SB3150015251512.51512.257.910.081.0233144.8 $-0.5$ Bucur[1994]Gannat $-850$ $-700$ $-775.0$ $-744.2$ 65.114.0102.923746.23.3Moutmir<[1995]	Lectoure	375	425	400.0	397.9	57.5	-4.3	20	.7	11 43	.0	9		Bucur [1994]		
Sadirac SB4135014001375.01373.651.01.9100.82865 $44.8$ $-0.5$ $Bucur$ $Bucur$ $[1994]$ Sadirac SB3150015251512.51512.257.910.081.02331 $44.8$ $-0.5$ $Bucur$ $[1994]$ Sadirac SB3510015251512.51512.257.910.081.02331 $44.8$ $-0.5$ $Bucur$ $[1994]$ Satirac SB351015251512.51512.257.910.081.02331 $44.8$ $-0.5$ $Bucur$ $[1994]$ Gannat $-850$ $-700$ $-775.0$ $-744.2$ $65.1$ 14.0102.9237 $46.2$ $3.3$ $Moutmir$ $[1995]$ Lignat $-850$ $-700$ $-775.0$ $-762.9$ $67.7$ $29.4$ $24$ $24.4$ $46.2$ $3.3$ $Moutmir$ $[1995]$ St. Blaise $-750$ $-550$ $-560.0$ $-662.7$ $69.1$ $21.6$ $21.4$ $45.5$ $3.3$ $Moutmir$ $[1995]$ Isoire 1 $-750$ $-250$ $-500.0$ $-662.7$ $69.1$ $21.6$ $21.4$ $45.5$ $3.3$ $Moutmir$ $[1995]$	Tolouse Place St. Etienne	450	550	500.0	508.7	63.4	1.0	9	.6 57	69 43	.6 1.	4		Bucur [1994]		
Sadirac SB3150015251512.257.910.081.0233144.8 $-0.5$ BucurBucur[1994]Gannat1995]Gannat1995]GannatGannatSt. BlaiseIsonice 1Isonice 1St. BlaiseIsonice 1 <td solution<="" t<="" td=""><td>Sadirac SB4</td><td>1350</td><td>1400</td><td>1375.0</td><td>1373.6</td><td>51.0</td><td>1.9</td><td>10</td><td>8 28</td><td>65 44</td><td>)  </td><td>.5</td><td></td><td>Bucur [1994]</td><td></td></td>	<td>Sadirac SB4</td> <td>1350</td> <td>1400</td> <td>1375.0</td> <td>1373.6</td> <td>51.0</td> <td>1.9</td> <td>10</td> <td>8 28</td> <td>65 44</td> <td>)  </td> <td>.5</td> <td></td> <td>Bucur [1994]</td> <td></td>	Sadirac SB4	1350	1400	1375.0	1373.6	51.0	1.9	10	8 28	65 44	)	.5		Bucur [1994]	
Sites $r < 900 \ km$ (France and Morocco)Sites $r < 900 \ km$ (France and Morocco)Moutmir [1995]Lignat $-850 -700 -775.0 -744.2 \ 65.1 \ 14.0 \ 10 \ 2.9 \ 237 \ 46.2 \ 3.3 \ Moutmir [1995]St. Blaise-575 -550 -562.5 -561.6 \ 65.6 \ 0.5 \ 11 \ 0.9 \ 2355 \ 43.5 \ 5.0 \ Moutmir [1994]Issoire 1-750 -250 -500.0 -662.7 \ 69.1 \ 21.6 \ 21 \ 1.4 \ 961 \ 45.5 \ 3.3 \ Moutmir [1995]$	Sadirac SB3	1500	1525	1512.5	1512.2	57.9	10.0	~~~	0 23	31 44		).5		Bucur [1994]		
GannatSites $r < 900 \ km$ (France and Morocco)Moutmir [1995]Gannat $-850 -700 -775.0 -775.0 -744.2 \ 65.1 \ 14.0 \ 10 \ 2.9 \ 237 \ 46.2 \ 3.3 \ Moutmir [1995]Moutmir [1995]Lignat-850 -700 -775.0 -775.0 -762.9 \ 67.7 \ 29.4 \ 24 \ 24 \ 24 \ 24 \ 24 \ 3.5 \ 3.3 \ Moutmir [1995]Moutmir [1995]St. Blaise-575 -550 -562.5 -561.6 \ 65.6 \ 0.5 \ 11 \ 0.9 \ 2355 \ 43.5 \ 5.0 \ Moutmir [1994]Bucur [1994]Issoire 1-750 -250 -500.0 -662.7 \ 69.1 \ 21.6 \ 21 \ 1.4 \ 961 \ 45.5 \ 3.3 \ Moutmir [1995]$									2	-						
Lignat $-850 -700 -775.0 -762.9 67.7 29.4 24 2.4 146 45.8 3.3$ $Moutmir [1995]$ Moutmir [1995]       St. Blaise $-575 -550 -562.5 -561.6 65.6 0.5 11 0.9 2355 43.5 5.0$ $Bucur [1994]$ Issoire I $-750 -250 -500.0 -662.7 69.1 21.6 21 1.4 961 45.5 3.3$ $Aoutmir [1995]$	Gannat	-850	-700	-775.0	Site -744.2	s r < 9	00 kn 14 0	(Fra 10 2	nce an	1d Mor 37 46	occo)	"		Moutmir [1995]		
St. Blaise $-755 - 550 - 562.5 - 561.6 \ 65.6 \ 0.5 \ 11 \ 0.9 \ 2355 \ 43.5 \ 5.0 \ Bucur [1994]         Issoire I       -750 - 250 - 500.0 - 662.7 \ 69.1 \ 21.6 \ 21 \ 1.4 \ 961 \ 45.5 \ 3.3 \ Moutmir [1995]   $	Lionat	-850	-700	-775.0	-762.9	67.7	20.4	27	1 <del>-</del>	16 45	i «			Moutmir [1995]		
Issoire 1 -750 -250 -500.0 -662.7 69.1 21.6 21 1.4 961 45.5 3.3 Moutmir [1995]	St Blaise	-575	-550	-562 5	-561.6	65.6	- Y O	; =	9 23	55 43	o v	0				
	Issoire 1	-750	-250	-500.0	-662.7	69.1	21.6	21	; 4: ; 9	54 15	5 . 2	s m		Moutmir [1995]		

Geochemistry Geophysics Geosystems

3

]]

Table 1. (continued)												
Name	t <sub>min</sub>	t <sub>max</sub>	tmean	t <sub>bayesian</sub>	I <sub>s</sub> , deg	$\mathrm{D}_{\mathrm{s}}^{\mathrm{s}},$ deg	α N	95 k	°N. °N	Long., °E	Stratigraphic Constraints	Source
Aspiran. AFR 6186	-525	-475	-500.0	-501.1	69.5	9.9	17 1	8 398	43.6	3.6		Gallet et al. [2002]
Aspiran. AFR 6416	-525	-475	-500.0	-501.4	69.4	9.6	13 1	5 728	43.6	3.6		Gallet et al. [2002]
Issoire 2	-700	-600	-650.0	-648.0	73.0	22.0	8	.6 483	45.5	3.3		Moutmir [1995]
Loupiac. Loup-01	-850	-700	-775.0	-777.6	64.5	27.0	14	3 889	44.9	1.5		Gallet et al. [2002]
Loupiac. Loup-03	-850	-700	-775.0	-814.6	62.4	36.2	14 2	6 242	44.9	1.5		Gallet et al. [2002]
Loupiac. Loup-12	-850	-700	-775.0	-742.0	66.7	19.7	93	4 236	44.9	1.5		Gallet et al. [2002]
Loupiac. Loup-16	-850	-700	-775.0	-794.2	62.4	27.4	17 3	2 123	44.9	1.5		Gallet et al. [2002]
Petit et grand Lezat	-100	-50	-75.0	-75.9	65.7	6.7-	10	5 931	46.0	3.3		Bucur [1994]
Sallèles d'Aude F10	-30	30	0.0	-1.3	62.5	0.2	8	8 756	43.2	3.0		Chauvin et al. [2000]
Marseille	-30	40	5.0	-2.3	63.7	-2.4	46 0	6 121	5 43.3	5.4		Bucur [1994]
Lezoux F3	-25	50	12.5	6.8	64.5	-2.4	14 1	.1 124	3 45.8	3.4		Bucur [1994]
Lezoux LAS65	15	20	17.5	17.5	63.5	-1.3	22 1	.3 684	. 45.8	3.4		Bucur [1994]
Gievres II sol brûlé	40	40	40.0	40.5	64.1	-0.5	6 0	8 504	7 47.3	1.7		Bucur [1994]
Mougon II	40	60	50.0	50.9	61.4	-3.3	20 1	.1 169	8 47.1	0.5		Bucur [1994]
Sallèles d'Aude F12	50	90	70.0	71.6	57.8	-1.6	18 1	.2 756	43.2	3.0		Chauvin et al. [2000]
Yseure (Le Pavillon)	79	80	80.0	80.0	61.0	-1.5	17 0	.7 226	5 46.6	3.4		Bucur [1994]
Amboise	75	125	100.0	103.0	60.2	-5.5	14 1	.7 474	47.4	1.0		Bucur [1994]
Pouillé les Bordes (four 5)	75	150	112.5	101.4	64.6	-6.3	8	.8 771	47.3	1.3		Bucur [1994]
Pouillé les Bordes (four 6)	75	150	112.5	116.2	60.4	-1.1	7 1	5 124	8 47.3	1.3		$Bucur \ [1994]$
Sallèles d'Aude F13	100	160	130.0	130.4	56.7	-5.1	8	.0 245	0 43.2	3.0		Chauvin et al. [2000]
Sallèles d'Aude F15	120	180	150.0	150.9	55.9	-5.6	10 1	.7 678	43.2	3.0		Chauvin et al. [2000]
La Croiselle/Briance	150	200	175.0	176.1	56.9	1.1	10 2	.3 372	45.6	1.6		Bucur [1994]
Varennes/Allier	150	210	180.0	180.6	58.8	-1.5	8	.8 745	46.3	3.4		$Bucur \ [1994]$
Lezoux (Hôpital)	160	200	180.0	181.3	53.7	-2.5	11 1	.7 548	45.8	3.4		$Bucur \ [1994]$
Toulon/Allier	180	180	180.0	180.5	57.9	-2.5	17 1	0 401	2 46.6	3.3		$Bucur \ [1994]$
Lezoux 85	175	190	182.5	182.5	58.2	-5.3	15 0	.6 320	6 45.8	3.4		$Bucur \ [1994]$
Sallèles d'Aude F3	190	230	210.0	210.2	56.1	0.2	20 1	.7 335	43.2	3.0		Chauvin et al. [2000]
Sallèles d'Aude F9	195	235	215.0	215.2	56.9	0.3	16 2	.0 306	43.2	3.0		Chauvin et al. [2000]
Volubilis. the big furnace	200	300	250.0	252.6	50.3	-7.8	10 3	.6 151	34.0	-5.5		Kovacheva [1984]
Sallèles d'Aude F7	235	295	265.0	264.5	58.8	1.2	15 2	.3 907	43.2	3.0		Chauvin et al. [2000]
Volubilis. NE	200	390	295.0	283.2	44.7	0.1	ω 1	8 201	6 34.1	-5.5		Najid [1986]
Volubilis. palais Justice	200	390	295.0	295.7	47.2	-8.4	4 8	.5 68	34.1	-5.5		Najid [1986]
Sallèles d'Aude F8	280	340	310.0	309.9	56.5	-2.8	10 2	.8 250	43.2	3.0		Chauvin et al. [2000]
Sallèles d'Aude F14	280	340	310.0	310.0	56.5	-3.3	9	.1 494	43.2	3.0		Chauvin et al. [2000]
Volubilis. S thermal bath	250	390	320.0	320.0	47.5	-1.8	9	.1 227	34.1	-5.5		Najid [1986]
Lezoux III Jardin de l'Hospice	385	450	417.5	418.8	61.1	-2.0	7 1	.3 162	2 45.8	3.4		Bucur [1994]
Le Thoronet (Abbaye)	400	500	450.0	443.9	58.5	2.0	7 1	5 133	3 43.5	6.3		Bucur $[1994]$
Volubilis. the furnace in the S of the site	400	500	450.0	444.7	51.7	-6.7	5 2	5 627	34.0	-5.5		Kovacheva [1984]
Masmolene I	500	600	550.0	547.1	62.9	4. 4.	11 0	.6 436	0 44.0	4.5		$Bucur \ [1994]$
El Basra	009	1000	800.0	864.8	51.9	22.5	28 2	.3 132	34.8	-5.9		Najid [1986]
El Basra	600	1000	800.0	858.2	51.5	18.1	8	4 212	34.8	-5.9		Najid [1986]
Peran en Pledran	910	930	920.0	919.8	65.6	23.3	12 1	0 177	6 48.5	-2.8		Bucur [1994]



$\sim$
ed
inu
ont
ğ
:
le
ab
Ë

Nama	+	+	÷	+	וs, לפת	Чs, Дел	Ž	<u>~</u>	°N.	, LUIIB., °F	Strationaphic Constraints	Source
INALLIC	hmin	hnax	umean	ubayesian	aca	ncg	ที่กับ	95 N	5	1	Suaugraphic Consuants	201100
Douè la Fontaine (cellule II)	006	1000	950.0	949.7	63.1	20.3	10 1.	8 607	47.2	0.3		Bucur [1994]
Moulins/Cephons	1032	1068	1050.0	1050.5	59.3	15.7	15 1.	1 1133	8 47.0	1.6		Bucur [1994]
Cabasse	1100	1150	1125.0	1115.4	58.5	15.8	10 1.	1 1603	8 43.4	6.2		Bucur [1994]
Planier	1250	1300	1275.0	1277.2	53.7	6.9	14 1.	0 137.	2 43.2	: 5.9		Bucur [1994]
Sainte Barbe (Marseille)	1280	1320	1300.0	1301.7	49.2	4.6	15 1.	7 460	43.3	5.4		Bucur [1994]
Cadrix	1365	1385	1375.0	1374.7	48.9	7.2	14 1.	3 841	43.5	5.9		Bucur [1994]
Tintaine (four à chaux)	1550	1590	1570.0	1568.9	60.5	9.0	7 1.	8 985	43.1	3.1		Bucur [1994]
Blois	1578	1578	1578.0	1578.5	60.4	6.7	15 0.	7 256.	2 47.6	5 1.3		Bucur [1994]
La Madrague (St. Cyr./Mer)	1680	1720	1700.0	1702.8	67.1	-11.5	7 1.	3 179	6 43.2	: 5.6		Bucur [1994]
Thoronet Four de l'âne	1829	1831	1830.0	1830.0	62.9	-18.3	6 1.	4 162	6 43.5	5 6.3		Bucur $[1994]$

Geochemistry

Geophysics Geosystems

> t<sub>mean</sub>, mean age of the interval proposed for the abandon of the structures in years A.D.; t<sub>hayesian</sub>, time at which the maximum of probability is obtained (using Bayesian modeling); I<sub>s</sub> and D<sub>s</sub>, inclination and declination of the mean site direction in situ, N, number of samples used for the calculation of the archaeomagnetic mean direction;  $\alpha_{95}$  and k, 95% confidence limit and precision parameter from Fisher statistics; Lat. and Long., site latitude and longitude; Stratigraphic Constraints between the structures; Source for each archaeomagnetic direction (for the Spanish [2006]) al. entry indicates Gómez-Paccard et abandon of the structure; blank data.

Madrid) have been considered (see Figure 1). This includes data from *Gallet et al.* [2002] (which itself includes a selection of well-dated results (called PC) from *Bucur* [1994], data from *Moutmir* [1995], and some new archaeomagnetic directions), together with the directional results of *Chauvin et al.* [2000]. A total of 63 archaeomagnetic directions from France have been compiled (see Table 1).

[5] In contrast, very few archaeomagnetic studies have been performed in Morocco. The first concerned the study of six structures coming from three localities [*Kovacheva*, 1984], in which four reliable archaeomagnetic directions were given. Five directions from *Najid* [1986] are also available in the archaeomagnetic database of *Korte et al.* [2005], giving a total of 9 archaeomagnetic directions from Morocco (Figure 1). They correspond to sites within a 900 km radius of Madrid (see Table 1).

### 3. Reference Curve Data Set

[6] The reference curves have been established assuming that within a restricted region the SV curves can be represented by a single curve assigned to a fixed reference site. Madrid is close to the geographic centre of the Iberian Peninsula  $(40.4^{\circ}N, 3.7^{\circ}W)$  and has been chosen as the reference point. A total of 63 Spanish archaeomagnetic directions are available from within the 600 km radius, with a further 16 from Morocco and southern France. One of the Spanish directions (GUA1 in Table 1) is offset from the general trend displayed by the other data. Recent results of radiocarbon dating cast doubt on the age assigned to the structure and so the direction has not been used in calculating SV curves.

[7] The data set (called the 600 km data set) contains 78 directions, which are shown in Figure 2, after relocation to Madrid via the virtual geomagnetic pole (VGP) method. They are generally very well defined. Around 60% have  $\alpha_{95}$  values less than 3°, and only 5 have values bigger than 5°. Uncertainties related to the ages of the structures can differ depending on the archaeological information available, but for the Spanish data set it is generally around 50 years. The compiled data span approximately 2000 years, from -100 to 1959 A.D. Throughout most of the record several directions per century are available, although there is a gap in the data between the 6th and 10th centuries.



**Figure 1.** Map showing locations of archaeomagnetic sites (plotted with different symbols for the different age ranges) compiled for the construction of the secular variation curve for the Iberian Peninsula. Madrid (40.4°N, 3.7°W) has been chosen as the reference site. Two circles with radii of 600 km and 900 km are shown.

[8] In southern France or northern Morocco several data can help to cover this gap, but they are from localities more than 600 km from Madrid (see Figure 1). Using data from such distal sources may introduce systematic errors due to the process of relocation to the central reference point. To investigate this question the angular errors introduced during the relocation process via the VGP method have been estimated in the same way as by Noel and Batt [1990]. The IGRF values have been calculated for an area described by 30°N, 50°N and 15°W, 10°E, relocated to Madrid, and the results are compared with the Madrid IGRF values. For an area of radius 900 km, inclination errors are less than  $2^{\circ}$  and declination errors are less than  $3^{\circ}$  (see Figure 3). Assuming that the harmonic content of the field has been similar throughout archaeological time, then similar error values can be expected in the past. Angular relocation errors of  $2^{\circ}-3^{\circ}$  are of the same order as the typical  $\alpha_{95}$  values of the Spanish archaeomagnetic directions, and if the error margins of the estimated SV curve are bigger, then these errors would not be significant. The assumption of almost constant harmonic content of the field during archaeological times has been tested

Geochemistry

Geophysics Geosystems

using the global model CALSK3.2 of *Korte and Constable* [2005]. The results indicate that the ratio of the magnetic energy of the nondipole (from degree 2 up to degree 10) to the energy of the dipole, at the Earth surface, has remained almost constant during the last 3 millennia with a mean value of  $0.02 \pm 0.005$ .

[9] Two archaeomagnetic directional data sets have been compiled. The first one includes archaeomagnetic directions corresponding to sites which fall within a 600 km radius circle centered on Madrid, and the second includes an additional 56 ("foreign") archaeomagnetic directions corresponding to sites situated between 600 and 900 km from Madrid, giving a total of 134 data (see Table 1). It can be seen (Figure 2) that the relocated directions from the two data sets are similar over the same time periods, and that there is no observable systematic error associated with the data between 600 and 900 km from Madrid.

[10] As the accuracy of any calculated SV curve depends principally on the number of reference points and on their associated dating errors, the use of the 900 km data set should increase the reliabil-



Geochemistry

Geophysics Geosystems

**Figure 2.** Declination and inclination of the archaeomagnetic directions available for the two reference areas, relocated to Madrid. Data coming from localities less than 600 km from Madrid are plotted in red, and data coming from localities between 600 and 900 km are plotted in blue.

ity of a curve obtained for the Iberian Peninsula. Furthermore, the 900 km data set includes data that partially fill the gap between the 6th and 10th centuries A.D. in the 600 km data set. It also includes data from the -6th to -8th centuries, which are not represented in the 600 km data set, thus extending coverage over a longer time period.

#### 4. First SV Curve for Iberia

[11] In order to determine the best SV curve for Iberia the two compiled data sets (600 and 900 km) were treated using the hierarchical bivariate [*Lanos et al.*, 2005] and the hierarchical Bayesian modeling based on roughness penalty [*Lanos*, 2004]. A comparison between the classical stratified bivariate [*Le Goff*, 1990; *Le Goff et al.*, 1992; *Daly and Le Goff*, 1996] and the hierarchical bivariate approaches has been presented by *Lanos et al.*  [2005]. The principal difference found was in the estimation of the error margins. In the stratified approach these errors are divided by the total number of samples contributing to the window, while in the hierarchical approach these errors are divided by the number of sites contributing to the window. As demonstrated by *Lanos et al.* [2005], a hierarchical approach should be used in archaeomagnetic curve estimation. Therefore this approach has been retained here.

[12] In order to determine the best window widths to use with hierarchical bivariate statistics several values were tested (50, 80, 100 and 120 years). No significant differences were found, therefore it was decided to use a window width of 80 years, plotted in 25 year steps, with at least four reference points per window.

[13] Bayesian modeling [Lanos, 2004] puts some prior knowledge on the global nature of the curve to be estimated, i.e., it is assumed that the studied physical phenomena vary in a smooth way, and it allows the window width to be automatically adapted to the density of points along the time axis, making the points movable within their respective dating error ranges. This approach allows the fitting of a spherical spline function based on roughness penalty to the data in three dimensions (declination, inclination and time), calculating the weight on the construction of the SV curve of all of the possible three-dimensional data. The results are expressed as a mean curve and an envelope (error) at a 95% confidence level. The "real" curve will lie somewhere inside the error band. In addition, this method takes into account stratigraphic constraints provided by archaeological investigations, some of which are known for the Spanish data (Table 1) [Gómez-Paccard et al., 2006].

[14] Figures 4a and 4b show the curves obtained for the 600 and 900 km data sets calculated using Bayesian methods and the comparison between the hierarchical bivariate and Bayesian modeling for the 900 km data set. The hierarchical bivariate curves are not continuous because some time windows contain less than four reference points (Figure 4b). It is noticeable that the mean values obtained with the two methods are very similar, although the hierarchical bivariate mean curve and corresponding errors seem a little choppy.

[15] The archaeomagnetic data sets are unevenly distributed in time. For this reason, Bayesian modeling has been retained in order to calculate





**Figure 3.** Isolines of the errors in declination and inclination introduced during the VGP relocation between any point on the plot and Madrid. The errors have been calculated using the IGRF 1900 and IGRF 2000 models. Two circles centered at Madrid with 600 and 900 km radii are shown.

the best SV curve for Iberia. The Bayesian curves, and their error envelopes, calculated with the 600 and 900 km data sets are very similar between -350 and 2000 A.D. (Figure 4a). As previously discussed, the maximum error due to the VGP relocation process from any point inside a reference area of 900 km (Figure 3) to Madrid is approximately 3°. This error is lower than the errors of declination and inclination at the 95% confidence level obtained for the Bayesian curve (of about 5°, Figure 4a). Therefore it is considered that the error introduced during relocation is included in the error inherent to the archaeomagnetic data.

[16] Taking into account all of the previous discussions, Bayesian modeling of the 900 km data set has been used to calculate the curve most representative of the SV for the Iberian Peninsula. It is shown in Figure 5. Table 2 gives the results obtained for inclination and declination (in steps of approximately 25 years), together with their 95%



**Figure 4.** Comparison between the hierarchical bivariate and the Bayesian modeling: (a) the results from Bayesian modeling for the 600 and 900 km data set and (b) a comparison between the results obtained by hierarchical bivariate (calculated using 25 year steps with a least four reference points per window) and Bayesian modeling for the 900 km data set. All curves have been calculated at Madrid and are plotted with their 95% confidence level error bands.

error envelopes. In Table 1, t<sub>bayesian</sub> indicates the most probable ages (at the 95% confidence level) obtained by Bayesian modeling for each structure. In order to check the influence of data with larger errors on the construction of the SV curve, the Iberian curve has also been calculated rejecting data with dating errors  $\geq$  75 years or  $\alpha_{95} \geq$  3.5. A total of 106 points has been considered. The results obtained are very similar to those obtained using the complete 900 km data set (134 points).

[17] The age of the reference archaeomagnetic data used (134 points) ranges from -775 to 1959 A.D. Throughout most of the record several directions per century are available, although there is a need to extend the data, especially between the 6th and 10th centuries A.D. and prior to 0 A.D., where the

majority of the data come from neighboring countries.

# 5. Comparison With French and German Secular Variation Curves

[18] In order to compare the Iberian SV curve with the French and German curves, the Bayesian curves for these two countries have been computed. The German archaeomagnetic curve was published by *Schnepp and Lanos* [2005], relocated to Göttingen. It includes a total of 166 archaeomagnetic directions. The French SV curve was calculated at Paris, by Bayesian modeling, the reference data set used is the same as that of *Gallet et al.* [2002], together with the 12 archaeomagnetic directions determined by *Chauvin et al.*  GÓMEZ-PACCARD ET AL.: FIRST ARCHAEOMAGNETIC SV CURVE

10.1029/2006GC001476



**Figure 5.** First SV curve for Iberia obtained by Bayesian modeling of the 900 km data set. (a) Stereoplot of declination and inclination, where orange and red indicate century timescales (thin line indicates where the curve is based on a small number of data), (b) declination versus time, and (c) inclination versus time. In Figures 5b and 5c, data coming from localities less than 600 km from Madrid are plotted in red, and data located between 600 and 900 km from Madrid are plotted in blue.

[2000]. The three curves were recalculated at Lyon (45.46°N, 4.50°E, Figure 1), chosen because of its central location between Madrid and Göttingen (Figure 6 and 7). No systematic relocation effect was observed related to this process. However, and in order to avoid relocation errors, the velocities and changes in curvatures of the curves have been calculated directly from the spherical splines obtained for Madrid (Iberian curve), Paris (French curve) and Göttingen (German curve). It is worth pointing out that the Iberian and French curves have 61 archaeomagnetic directions in common, while the French and German curves

Geochemistry

Geophysics Geosystems

have 25 common data. In contrast, the Iberian and the German curves have no points in common.

[19] The three SV curves look similar during most of the past three millennia, especially the French and Iberian curves. However, closer inspection reveals some significant differences in the details of the curves. The observed differences cannot be explained by the effect of relocation processes alone. Lower inclinations are seen in the Iberian curve between the 11th and 14th century and between the 16th and 19th century (Figures 6 and 7). These two differences are constrained by



Table	2.	Secular	Variation	Curve	for	the	Iberian
Penins	ula :	at Madrid	a				

Table 2. (continued)

I CIIIIIS	ula at 1	viauriu					t,	I,	I <sub>min</sub> ,	I <sub>max</sub> ,	D,	D <sub>min</sub> ,	D <sub>max</sub> ,
t.	I.	Imin.	Imax.	D.	D <sub>min</sub> ,	D <sub>max</sub> .	years	deg	deg	deg	deg	deg	deg
years	deg	deg	deg	deg	deg	deg	5(2	50.0	57.0	(2.(	4.4	0.0	0.0
	0	U	0	0	U		505	59.9	57.2	62.6	4.4	-0.9	9.9
-815	58.5	55.4	61.6	27.2	21.3	33.1	288 612	60.5	57.5	62.6	0.1	0.4	12.0
-790	59.6	56.8	62.5	25.6	19.8	31.3	615	60.0	57.0	62.0	7.9	1.9	14.1
-767	60.6	57.9	63.4	24.1	18.4	29.6	038	60.8	57.0	03.9	9.7	5.4	10.2
-745	61.5	58.9	64.2	22.5	16.9	28.1	003	60.8	57.0	04.0	11.5	5.0	18.1
-723	62.4	59.8	65.0	21.0	15.3	26.6	088	60.8	57.5	04.1	13.2	0.5	20.0
-698	63.2	60.7	65.9	19.3	13.5	25.0	/13	60.7	57.5	04.0	14.8	8.1 0.5	21.0
-673	64.0	61.5	66.6	17.6	11.6	23.4	/38	60.5	5/.1	03.8	10.2	9.5	23.0
-650	64.6	62.1	67.1	16.0	10.0	21.8	/03	60.2	50.9	03.5	1/.5	10.9	24.2
-626	65.1	62.6	67.6	14.4	8.5	20.1	/88	59.8	50.0	03.1	18.0	12.2	25.1
-601	65.4	63.1	67.8	12.8	6.9	18.4	813	59.4	55.0	02.0	19.4	13.3	25.7
-576	65.7	63.4	68.0	11.2	5.5	16.7	838	59.0	55.9	62.0	20.1	14.2	26.0
-556	65.8	63.5	68.1	10.0	4.5	15.5	861	58.6	55.0	61.4	20.4	14.9	26.0
-531	65.9	63.6	68.1	8.7	3.2	14.2	883	58.1	55.4	60.8	20.6	15.5	25.7
-506	65.8	63.5	68.1	7.5	1.9	13.0	908	57.5	55.0	60.0 50.2	20.5	15.8	25.2
-490	65.7	63.4	68.0	6.8	1.1	12.4	931	56.9	52.0	59.2	20.2	15.9	24.5
-465	65.5	63.1	67.9	5.6	-0.2	11.4	952	56.2	53.9	58.4	19.8	15.8	23.8
-440	65.3	62.7	67.8	4.5	-1.6	10.5	9//	55.Z	53.0	5/.5	19.2	15.5	22.9
-415	64.9	62.3	67.6	3.5	-2.9	9.6	1002	54.1	52.0	50.1	18.5	15.0	21.9
-390	64.6	61.8	67.3	2.5	-4.0	8.9	1027	53.0	51.0	54.8	17.0	14.4	20.8
-365	64.2	61.2	67.0	1.5	-5.1	8.1	1044	52.1	50.2 40.2	54.0	1/.0	13.9	20.0
-340	63.7	60.7	66.7	0.7	-6.0	7.4	1007	51.0	49.2	52.8	10.1	13.2	18.9
-315	63.3	60.2	66.3	0.0	-6.7	6.7	1091	49.9	48.1	50.5	15.2	12.4	1/.8
-293	62.9	59.9	65.9	-0.6	-7.2	6.0	1110	40.0	47.0	30.3 40.7	14.1	11.5	10./
-268	62.5	59.5	65.4	-1.1	-7.5	5.3	1157	48.0	40.2	49.7	13.2	10.5	15.0
-243	62.1	59.2	64.9	-1.5	-7.6	4.5	1133	4/.4	45.0	49.2	12.4	9.7	13.0
-218	61.7	59.1	64.4	-1.9	-7.5	3.7	11/1	40.9	43.0	40.7	10.5	0.0 7 7	14.1
-193	61.4	59.0	63.9	-2.2	-7.3	2.9	1216	40.4	44.0	40.2	0.5	6.6	11.7
-171	61.2	58.9	63.4	-2.4	-7.1	2.2	1210	45.0	14.2	47.6	9.2	5.6	10.4
-146	61.0	59.0	63.0	-2.6	-6.7	1.5	1259	45.9	44.3	47.0	6.0	2.0 4.6	9.2
-121	60.8	59.1	62.5	-2.7	-6.3	0.8	1204	45.9	1/1 3	47. <del>4</del> 17.1	6.4	4.0	9.2 8.6
-100	60.6	59.1	62.2	-2.7	-5.9	0.5	1298	45.9	44.3	47.5	57	33	79
-65	60.4	59.0	61.7	-2.5	-5.2	0.2	1321	46.0	44.3	47.5	5.0	2.6	7.9
-49	60.2	58.9	61.5	-2.3	-4.9	0.3	1338	46.1	44.4	47.8	4 7	2.0	7.1
-24	59.8	58.5	61.0	-2.0	-4.5	0.5	1362	46.3	44 7	48.0	4 5	2.2	6.8
-5	59.3	58.1	60.5	-1./	-4.2	0.7	1381	46.7	45.1	48.3	4.5	2.2	6.8
13	58.8	57.5	60.0	-1.6	-4.0	0.8	1402	47.3	45.8	48.9	4.7	2.4	7.0
31	58.2	56.9	59.4	-1.5	-3.9	0.8	1416	47.9	46.4	49.5	4.9	2.6	7.2
48	57.5	50.2	38./ 59.0	-1.0	-3.9	0.7	1441	49.2	47.7	50.9	5.4	3.0	7.9
65	56.8	53.5 54.7	58.0	-1./	-4.0	0.5	1466	50.8	49.2	52.5	6.0	3.4	8.7
80 110	55.1	52.0	56.2	-2.0	-4.2	0.2	1488	52.3	50.6	54.1	6.5	3.7	9.4
121	54.5	52.9	55.6	-2.5	-4.4	-0.2	1513	54.0	52.3	55.9	7.0	3.9	10.1
151	53.0	52.5	55.0	-2.3	-4.5	-0.0	1531	55.3	53.5	57.2	7.1	3.9	10.4
191	53.9	52.0	54.5	-2.7	-4.0	-0.8	1556	57.0	55.2	58.8	7.1	3.6	10.4
200	53.4	52.4	54.5	-2.8 -2.9	-4.0 -4.7	-1.0	1580	58.4	56.7	60.4	6.5	2.9	10.0
200	53.3	52.2	54.4	-2.9	7	-1.1	1600	59.7	57.8	61.7	5.6	1.6	9.3
236	53.3	52.2	54.5	-2.9	_4 9	-1.0	1625	61.0	59.0	63.1	3.8	-0.5	7.9
258	53.4	52.2	54 7	-3.0	-5.1	-0.9	1647	62.0	59.9	64.1	1.7	-2.9	6.1
280	53.6	52.2	54.9	-3.1	-53	-0.9	1672	62.8	60.6	65.0	-1.1	-6.0	3.6
311	53.9	52.5	55.3	-3.2	-5.6	-0.9	1697	63.3	61.0	65.5	-4.1	-9.1	0.7
336	54.2	52.8	55.6	-3.3	-57	-0.9	1719	63.4	61.2	65.6	-6.8	-11.7	-1.9
361	54 7	53.2	56.1	-33	-5.8	-0.8	1744	63.3	61.2	65.4	-9.6	-14.4	-4.9
386	55.2	53.8	56.7	-3.2	-5.8	-0.5	1769	63.0	60.9	65.1	-12.0	-16.7	-7.5
406	55.2	54.2	573	-2.9	-57	-0.1	1794	62.5	60.4	64.6	-14.0	-18.6	-9.6
424	563	54.6	57.9	-2.5	-55	0.1	1817	62.0	59.9	64.1	-15.4	-20.0	-10.9
446	56.9	55.1	58.8	-1.9	-5.2	1.5	1836	61.6	59.4	63.8	-16.2	-20.9	-11.6
466	57.5	55.6	59.5	-1.1	-4.8	2.6	1861	61.1	58.5	63.5	-16.9	-22.0	-11.8
491	58.2	56.1	60.4	0.0	-4.0	4.2	1882	60.6	57.8	63.4	-17.2	-22.9	-11.5
515	58.9	56.6	61.3	1.4	-3.2	6.0	1907	60.1	56.7	63.3	-17.4	-24.0	-10.7
540	59.5	56.9	62.0	2.9	-2.1	8.0	1932	59.5	55.6	63.4	-17.4	-25.0	-9.7



 Table 2. (continued)

t,	I,	I <sub>min</sub> ,	I <sub>max</sub> ,	D,	D <sub>min</sub> ,	D <sub>max</sub> ,
years	deg	deg	deg	deg	deg	deg
1959	59.0	54.5	63.5	-17.3	-26.0	-8.5

<sup>a</sup>Here t, age in years A.D.; I (D), inclination (declination);  $I_{min} - I_{max}$  (D<sub>min</sub> - D<sub>max</sub>), error margins for inclination (declination).

a relatively large number of Spanish data and may be related to features of the local geomagnetic field. If the Iberian and German SV curves are characterized by a period of two centuries (from 1200 to 1400 A.D.) with low inclinations, this period of low inclinations seems shorter in the French SV curve. The trend of the Iberian SV curve between the 10th and 15th centuries is very similar to that obtained for Italy recorded in historical lava flows [*Tanguy et al.*, 1999; *Arrighi et al.*, 2004]. It is noticeable that the lowest inclinations observed for France and Iberia do not have exactly the same values as for Germany. The differences between the three curves observed



**Figure 6.** SV curves for (a) Iberia and France and (b) Iberia and Germany at Lyon (45.46°N, 4.50°E). All have been calculated by Bayesian modeling. Ages are plotted in black for the French and Germany SV curves and in gray for the Iberian SV curve. Thin lines indicates where the Iberian curve is based on a small number of data.



**Figure 7.** Declination and inclination reference curves together with their 95% error envelope versus time for the Iberian, French, and German SV curves at Lyon. All results are obtained by Bayesian modeling.

during the 6th–9th centuries can be explained by the small number of data available for this period in the Iberian curve. It is important to notice that the French and German curves are also less well constrained during this period (6th–9th) because of a sparse number of data.

[20] The same analysis can be done taking into account the error estimates of the curves (Figure 7). They are generally lower for the French SV curve than for the Iberian or German curves, due to the larger number of archaeomagnetic data available for France. However, the error margins of the Iberian curve for the period between the 10th and 17th centuries are small and the differences discussed before are well constrained for this region.

[21] Figure 8 shows the rate of the secular variation (velocity) and curvature values computed using Iberian, French and German Bayesian curves. The general trend of the obtained values of velocity and curvature is very similar. Three major directional cusps, related with velocity minima, are

observed in each of the curves, the first around -125 A.D., the second around 200 A.D. and the third around 1350 A.D. Two of these major directional changes (at 200 and 1350 A.D.) are in agreement with those found by Gallet et al. [2003] (200 and 1400 A.D.) and seem to occur a little later in the German curve. A further change in curvature is seen at around 600-700 A.D. for the German curve, which is suggested in the French curve but poorly expressed in the Iberian curve (Figure 8). This last directional cusp could be related to the one derived by Gallet et al. [2003] and identified at around 800 A.D. The weak temporal resolution of this last directional change could be explained by the lack of data in the European records between the 5th and the 10th centuries (the so-called Dark Ages).

[22] It is concluded that the SV for western Europe over the last three millennia is characterized by 3 major directional changes associated with low secular variation rates. In order to see if these characteristics have a regional or a global signifi-



**Iberian Peninsula** 



Figure 8. Comparison of the velocity and changes in curvature of the Iberian, French, and German SV curves, calculated by Bayesian modeling.

cance, archaeomagnetic data from other regions have been analyzed. In the global archaeomagnetic data set built by *Korte et al.* [2005] the records from Japan and southwest United States are the most continuous (outside Europe), both in declination and inclination, and are also the less scattered over the last 1500 years. After relocation of the data to a common point (35.3°N, 136.8°E for Japan and 39.5°N, 250.7°E for southwest United States), SV curves for these two regions have been calculated using Bayesian modeling (Figure 9a). The only difference with the previous analysis is that the number of samples per site used to calculate the mean directional data is not taken into account, because this information is not reported in the data set published by *Korte et al.* [2005].

[23] The secular variation rates and changes in curvature obtained are shown in Figure 9b. Three (for southwest United States) and four (for Japan) major directional changes can be observed, occurring at different time periods. Moreover, the ages of

#### a) Bayesian curves

Geochemistry Geophysics Geosystems



**Figure 9.** (a) Declination and inclination values together with the error margins versus time for Japan and southwest United States and (b) corresponding velocities and changes in curvature. All results are obtained by Bayesian modeling. N indicates the number of archaeomagnetic directions used to obtain the Bayesian curves.



**Figure 10.** Location of the north geomagnetic pole as computed from CALS3K.2 [*Korte and Constable*, 2005], from *Hongre et al.* [1998] global models, and from the model of *Jackson et al.* [2000]. The results obtained from DRGF-1950, DRGF-1970, and DRGF-1990 are also given.

these cusps are not in agreement with those observed in Europe, except perhaps the one present for the period 650–800 A.D. This seems to indicate that major directional changes are a regional feature of the geomagnetic field, even though more detailed and longer records are clearly needed.

#### 6. Comparison With Global Models

[24] The Iberian SV curve presented here has been compared with the CALS3K.2 and CALS7K.2 global models of Korte and Constable [2005], the geomagnetic model of Jackson et al. [2000] and the model of Hongre et al. [1998]. The Hongre et al. [1998] model is derived with Gauss coefficients up to degree 2 plus degree 3 order 3. The models of Korte and Constable [2005] are derived with Gauss coefficients up to order 10. This is a significant difference between these models. As a smoothing constraint is used by Korte and Constable [2005], their resolution is probably in the order of degree 4 or 5. For the last two millennia there is little difference between the CALS3K.2 and CALS7K.2 models, so the CALS3K.2 model has been used here. The Jackson et al.'s model, based on massive 400 years historical records, is derived with Gauss coefficients up to degree 14.

[25] Differences between the available global models can be seen in Figure 10. This gives the corresponding positions of the north geomagnetic pole. It was expected that the dipole terms would be the most robust and should be similar for the three models, but this is clearly not the case. It can be noticed that DRGF and Jackson models are in close agreement over the past five decades.

[26] Model predictions of declination and inclination at Madrid have been calculated. The results are display in Figure 11, together with the Iberian SV curve. The directional changes expected at Madrid when only dipole terms are included (Figure 11a), when terms up to order 2 (Figure 11b) and 3 (Figure 11c) are used and when all Gauss coefficients are used (Figure 11d) are shown. It can be seen how the results for the dipole directional changes are quite different for both models, the curves obtained with *Hongre et al.*'s [1998] model being smoother. However, the dipole part predictions of declination and inclination show the same overall shape and features as the archaeomagnetic



Geochemistry

Geophysics Geosystems



**Figure 11.** Comparison between the declination and inclination of the SV reference curve obtained for Iberia (in gray) and declination and inclination values predicted by *Korte and Constable* [2005] and *Hongre et al.* [1998] and from *Jackson et al.*'s [2000] models using the Gauss coefficients up to different degrees, calculated at Madrid: (a) degree 1, (b) degree 2, (c) degree 3, and (d) all Gauss coefficients provided by the models. Results from CALS7.K2 are very similar to those from CALSK3.2 and are not plotted.

SV curve for Iberia, although the amplitudes are very different. When non dipole terms are included, the amplitudes of the fluctuations of the predicted SV curves increase considerably and the models fit reasonably well the archaeomagnetic curve, showing the same fluctuations with time. However, there is a discrepancy between some of the amplitudes of these fluctuations (in both declination and inclination). The declination results obtained using the models from Korte and Constable [2005] agree very well with the results from Hongre et al. [1998], except before 600 A.D. where the declination values are generally more easterly for the CALS3K.2 model. In contrast, for inclination, the amplitudes are clearly more important for the Hongre et al. [1998] model when all the coefficients are used. The best fit is obtained with the Hongre et al. [1998] model, up to degree 2 (Figure 11b). As can be seen in Figure 11d, Jackson et al.'s [2000] results provide higher inclinations between 1590 and 1800 A.D. than Korte and Constable's [2005] and Hongre et al.'s [1998] models and the Bayesian curve.

Geochemistry

Geophysics Geosystems

[27] In spite of their differences the Korte and Constable [2005] and Hongre et al. [1998] models predict the Iberian archaeomagnetic SV reasonably well for the last two millennia. This is not very surprising since the majority of the available archaeomagnetic data used to build these models (especially CALS3K.2 and Hongre models) comes from Europe. The good fit between the SV curve and the global model predictions is interpreted as an indicator of the good agreement between the new data and the previous European archaeomagnetic data. To improve knowledge of the evolution of the dipole, more data are clearly needed in order to increase the accuracy of global models. The Iberian data used in this article could be used for future global modeling and will increase their resolution.

#### 7. Conclusions

[28] This study presents the first archaeomagnetic SV reference curve for Iberia. It has been calculated using Bayesian modeling. The age of the reference archaeomagnetic data used ranges from -775 to 1959 A.D. Throughout most of the record several directions per century are available, although there is a need to extend the data, especially before 0 A.D. and between the 6th and 10th centuries (where the majority of the data comes from neighboring countries). However, the first SV for Iberia is now available.

[29] Compared to French and German SV curves the same general patterns are found but significant differences can be observed between the 11th and 14th centuries A.D. and the last few centuries. where lower inclinations are found for the Iberian curve. Analysis of the Iberian, French and German reference curves indicates that SV in western Europe is characterized by three cusps or major directional changes associated with low SV rates (at -125, 200, and 1350 A.D.). The SV curves and corresponding rates and changes in curvature for Japan and southwest United States have also been calculated using Bayesian modeling. The results suggest that the observed cusps are representing a regional feature of the geomagnetic field. Longer and more detailed records are clearly needed in order to test this idea.

[30] Differences between the available global models [*Korte and Constable*, 2005; *Hongre et al.*, 1998; *Jackson et al.*, 2000] are clearly observed. Even the lowest terms (i.e., the dipole) are quite different for these models. Despite these differences, all models predict reasonably well the Iberian archaeomagnetic SV curve.

[31] The SV curve obtained extends the archaeomagnetic dating technique to Spain and Portugal. This provides a valuable tool for archaeologists working in the region, which has a rich and increasingly studied archaeological past. Finally, the Iberian curve contributes to the increase in the knowledge of the Earth's magnetic field, and the data can be used for future modeling of geomagnetic field behavior.

#### Acknowledgments

[32] M. Gómez-Paccard and G. Catanzariti acknowledge their fellowships from the AARCH Network (Archaeomagnetic Applications for the Rescue of Cultural Heritage), contract EU:HPRN-CT-2002-00219. This research was also funded by the CGL2005-00211 project from the Spanish ministry of Education and Science. Two anonymous referees provided careful reviews which helped to greatly improved the paper.

#### References

- Arrighi, S., M. Rosi, J. C. Tanguy, and V. Courtillot (2004), Recent eruptive history of Stromboli (Aeolian Islands, Italy) determined from high-accuracy archeomagnetic dating, *Geophys. Res. Lett.*, 31, L19603, doi:10.1029/2004GL020627.
- Batt, C. M. (1997), The British archaeomagnetic calibration curve: An objective treatment, *Archaeometry*, *39*, 153–168.
- Bucur, I. (1994), The direction of the terrestrial magnetic field in France during the last 21 centuries, *Phys. Earth Planet. Inter.*, 87, 95–109.

- Geochemistry Geophysics Geosystems
- Chauvin, A., Y. Garcia, P. Lanos, and F. Laubenheimer (2000), Paleointensity of the geomagnetic field recovered on archaeomagnetic sites from France, *Phys. Earth Planet. Inter.*, *120*, 111–136.
- Daly, L., and M. Le Goff (1996), An updated and homogeneous world secular variation data base. 1. Smoothing of the archaeomagnetic results, *Phys. Earth Planet. Inter.*, *93*, 159– 190.
- Fisher, R. A. (1953), Dispersion on a sphere, Proc. R. Soc. London, Ser. A, 217, 295–305.
- Gallet, Y., A. Genevey, and M. Le Goff (2002), Three millennia of directional variations of the Earth's magnetic field in western Europe as revealed by archaeological artefacts, *Phys. Earth Planet. Inter.*, 131, 81–89.
- Gallet, Y., A. Genevey, and V. Courtillot (2003), On the possible occurrence of "archaeomagnetic jerks" in the geomagnetic field over the past three millennia, *Earth Planet. Sci. Lett.*, 214, 237–242.
- Gómez-Paccard, M., et al. (2006), A catalogue of Spanish archaeomagnetic data, *Geophys. J. Int.*, *166*, 1125–1143.
- Hongre, L., G. Hulot, and A. Khokhlov (1998), An analysis of the geomagnetic field over the past 2000 years, *Phys. Earth Planet. Inter.*, 106, 311–335.
- Jackson, A., A. Jonkers, and M. Walker (2000), Four centuries of geomagnetic secular variation from historical records, *Philos. Trans. R. Soc. London, Ser. A*, 358, 957–990.
- Kirschvink, J. L. (1980), The least-squares line and plane and the analysis of palaeomagnetic data, *Geophys. J. R. Astron. Soc.*, 62, 699–718.
- Korte, M., and C. G. Constable (2005), Continuous geomagnetic field models for the past 7 millenia: 2. CALS7K, *Geochem. Geophys. Geosyst.*, 6, Q02H16, doi:10.1029/ 2004GC000801.
- Korte, M., A. Genevey, C. G. Constable, U. Frank, and E. Schnepp (2005), Continuous geomagnetic field models for the past 7 millennia: 1. A new global data compilation, *Geochem. Geophys. Geosyst.*, 6, Q02H15, doi:10.1029/ 2004GC000800.
- Kovacheva, M. (1984), Some archaeomagnetic conclusions from three archaeological localities in north-west Africa, *C. R. Acad. Sci. Bulgaria*, *37*, 171–174.
- Kovacheva, M., N. Jordanova, and V. Karloukovski (1998), Geomagnetic field variations as determined from Bulgaria archaeomagnetic data. Part II: The last 8000 years, *Surv. Geophys.*, 19, 431–460.
- Lanos, P. (2004), Bayesian inference of calibration curves: Application to archaeomagnetism, in *Tools for Constructing Chronologies: Crossing Disciplinary Boundaries, Lect. Notes Stat.*, vol. 177, edited by C. Buck and A. Millard, pp. 43–82, Springer, New York.
- Lanos, P., M. Le Goff, M. Kovacheva, and E. Schnepp (2005), Hierarchical modelling of archaeomagnetic data and curve

estimation by moving average technique, Geophys. J. Int., 160, 440-476.

- Le Goff, M. (1990), Lissage et limites d'incertitude des courbes de migration polaire: Pondération des données et extension bivariate de la statistique de Fisher, *C. R. Acad. Sci. Paris, Ser. II, 311*, 1191–1198.
- Le Goff, M., B. Henry, and L. Daly (1992), Practical method for drawing VGP path, *Phys. Earth Planet. Inter.*, 70, 201– 204.
- Márton, P. (2003), Recent achievements in archaeomagnetism in Hungary, *Geophys. J. Int.*, 153, 675–690.
- McFadden, P. L., and M. W. McElhinny (1988), The combined analysis of remagnetization circles and direct observations in paleomagnetism, *Earth Planet. Sci. Lett.*, 87, 161–172.
- Moutmir, M. (1995), Analyses magnétiques de terres cuites protohistoriques en France. Apports en archéomagnétisme (Premier millénaire avant J.C.) et en archéologie, Ph.D. thesis, Mus. Natl. d'hist. Nat., Paris.
- Najid, D. (1986), Palaeomagnetic studies in Morocco, Ph.D. thesis, Univ. of Newcastle upon Tyne, Newcastle upon Tyne, U. K.
- Noel, M., and C. M. Batt (1990), A method for correcting geographically separated remanence directions for the purpose of archeomagnetic dating, *Geophys. J. Int.*, 102, 753–756.
- Oyamburu, I., J. J. Villalain, M. L. Osete, M. Zarzalejos, and C. Blasco (1996), Estudio paleomagnético del yacimiento de Villa del Pañuelo (Villamanta, Madrid), *Geogaceta*, 20, 1044–1046.
- Parés, J. M., R. De Jonge, J. O. Pascual, A. Bermúdez, C. J. Tovar, R. A. Luezas, and N. Maestro (1992), Archaeomagnetic evidence for the age of a Roman pottery kiln from Calahorra (Spain), *Geophys. J. Int.*, 112, 533–537.
- Schnepp, E., and P. Lanos (2005), Archaeomagnetic secular variation in Germany during the past 2500 years, *Geophys. J. Int.*, 163, 479–490.
- Schnepp, E., and P. Lanos (2006), A preliminary secular variation reference curve for archaeomagnetic dating in Austria, *Geophys. J. Int.*, 166, 91–96.
- Tanguy, J. C., M. Le Goff, V. Chillemi, A. Paiotti, C. Principe, S. La Delfa, and G. Patanè (1999), Variation séculaire de la direction du champ géomagnétique enregistrée par les laves de l'Etna et du Vésuve pendant les deux derniers millénaires, *C. R. Acad. Sci Paris., Ser. IIa, Terre Planetes*, 329, 557– 564.
- Thellier, E. (1938), Sur l'aimantation des terres cuites et ses applications géophysiques, *Ann. Inst. Phys. Globe Univ. Paris, 16,* 157–302.
- Thellier, E. (1981), Sur la direction du champ magnétique terrestre en France durant les deux derniers millénaires, *Phys. Earth Planet. Inter.*, *24*, 89–132.