



**HAL**  
open science

## On Magid's approach to the inverse problem in differential Galois theory.

Jerry Kovacic, Claude Mitschi, Michael F. Singer

► **To cite this version:**

Jerry Kovacic, Claude Mitschi, Michael F. Singer. On Magid's approach to the inverse problem in differential Galois theory.. 1996. hal-00129562

**HAL Id: hal-00129562**

**<https://hal.science/hal-00129562>**

Preprint submitted on 8 Feb 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On Magid's Approach to the Inverse Problem in Differential Galois Theory

J. Kovacic\* C. Mitschi† M.F. Singer‡

March 25, 1996

The aim of this note is to expand on a footnote contained in our paper [4]. This footnote alluded to a counterexample (due to Kovacic) to the following Theorem (Theorem 7.13 of [3]):

*Let  $D_1, \dots, D_n$  be a  $C$ -basis of  $\text{Lie}(G)$  and let  $f_1, \dots, f_n \in F = C(x)$  be linearly independent over  $C$ . With  $D_F = \frac{d}{dx}$  let  $D = D_F \otimes 1 + \sum f_i \otimes D_i$  be the corresponding derivation of  $F[G_F]$ . Let  $P$  be a maximal  $D$ -stable ideal of  $F[G_F]$  and let  $E$  be the fraction field of  $F[G_F]/P$ . Then  $E \supset F$  is a Picard-Vessiot extension with  $G(E/F) \leq G$ , where  $\text{codim}_G(G(E/F)) \leq 1$ .*

*In particular, if  $G$  has no algebraic subgroups of codimension 1, then  $G(E/F) = G$  and  $F(G_F) \supset F$  is a Picard-Vessiot extension with  $G(F(G_F)/F) = G$ .*

This statement is incorrect as the following examples show [2]. Note that Theorem 7.3 of [3] is incorrect as well since the ideals presented in these examples have height larger than 1.

**Example 1:** Let  $G$  be  $C \times C$ , under addition. If we wish, we may consider  $G$  to be a subgroup of  $\text{GL}(3)$  by:

$$(a, b) \mapsto \begin{pmatrix} 1 & a & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The ring of regular functions on  $G$  is  $C[G] = C[X, Y]$  where  $X$  and  $Y$  are the coordinate functions:  $X(a, b) = a$  and  $Y(a, b) = b$ . It is nothing more than the ring of polynomials in the indeterminates  $X$  and  $Y$ . The Lie algebra of right-invariant derivations (over  $C$ ) has

---

\*JYACC, 116 John Street, New York, N.Y. 10038

†Institut de Recherche Mathématique Avancée, Université Louis Pasteur et C.N.R.S., 7 rue René Descartes, 67084 Strasbourg Cedex, France.

‡North Carolina State University, Department of Mathematics, Box 8205, Raleigh, NC 27695-8205.

basis  $\frac{\partial}{\partial X}$  and  $\frac{\partial}{\partial Y}$ , the usual partial derivatives with respect to the variables.

Define  $f_1 = 1$  and  $f_2 = 2x$ . These are in  $F$  and are linearly independent over  $C$ . Let  $D$  be a derivation on  $F[G_F] = F \otimes_C C[G]$  defined as follows:

$$D = \frac{d}{dx} \otimes 1 + 1 \otimes \frac{\partial}{\partial X} + 2x \otimes \frac{\partial}{\partial Y}$$

Consider the ideal  $I = (x \otimes 1 - 1 \otimes X, x^2 \otimes 1 - 1 \otimes Y)$  of  $F[G]$ . Then

$$D(x \otimes 1 - 1 \otimes X) = 1 \otimes 1 - 1 \otimes 1 = 0$$

and

$$D(x^2 \otimes 1 - 1 \otimes Y) = 2x \otimes 1 - 2x \otimes 1 = 0$$

Thus  $I$  is a  $D$ -differential ideal of  $F[G]$ . Note that if we identify  $F[G]$  with  $F[X, Y]$  then  $I = (X - x, Y - x^2)$ .

Now consider the substitution homomorphism  $\pi : X \mapsto x, Y \mapsto x^2$  over  $F$ . In other words,  $\pi : F[G] = F \otimes C[G] \rightarrow F$  is given by  $\sum(k_i \otimes h_i) \mapsto \sum(k_i h_i(x, x^2))$ . One can show that this is a  $D$  to  $d/dx$  differential homomorphism whose kernel is  $I$ . Since the image of  $\pi$  is a field (*i.e.*,  $F$ ),  $I$  is a maximal ideal, and therefore a maximal differential ideal. But  $E = F[G]/I$  is isomorphic to  $F$ . The Galois group  $G(E/F)$  is the identity, which is of codimension 2 in  $G$ . We note that arbitrary codimension can be achieved by using  $G = C^n$  and elements  $f_i = ix^{i-1}, i = 1, \dots, n$ . ■

Analyzing this example further shows where the argument in the proof of Theorem 7.3 is not valid. With notations as before, let  $V = \text{Spec } C[x] = \mathbb{A}_C^1$  as in the proof of Theorem 7.3 in [3] (the  $f_i$  here have no denominators). Then  $V \times G = \text{Spec } C[x, X, Y]$  and the irreducible variety defined by the ideal  $I$  is the rational curve  $W = \text{Spec } C[x, X, Y]/(X - x, Y - x^2)$ . The generic point  $\text{Spec } F$  of  $V$  (as before,  $F = C(x)$ ) becomes an  $F$ -point of  $V \times G$  via the natural homomorphism  $C[x, X, Y] \rightarrow C[x, X, Y]/(X, Y) \rightarrow F$ . It extends to an  $\overline{F}$ -point of  $V \times G$  which is *not* an  $\overline{F}$ -point of  $W$ , since the ideal  $I$  is not contained in the kernel  $(X, Y)$  of  $C[x, X, Y] \rightarrow \overline{F}$ . On line 24 of [3], an implicit assumption is made that (using the notation of [3])  $(\overline{\eta}, e)$  is a point of  $W_{\overline{\eta}}$ , that is,  $(t, e)$  is a point of  $W(\overline{F})$  for all  $t \in \overline{F}$ . In the above example  $(t, e) = (t, 0, 0)$  and  $W(\overline{F}) = \{(t, t, t^2), t \in \overline{F}\}$  and so we see that this is not the case.

One can also construct a counterexample for a reductive group.

**Example 2:** Let  $G = \text{GL}(n)$ . Let  $\xi_{ij}$  be the coordinate functions on  $G$ . Thus  $\xi_{ij}(g) = g_{ij}$  whenever  $g \in \text{GL}(n)$ . The ring of regular functions  $C[G]$  is  $C[\xi_{ij}, \frac{1}{\det \xi}]$ . The derivations  $D_{ij} = \sum_k \xi_{jk} \frac{\partial}{\partial \xi_{ik}}$  form a basis of  $\text{Lie}(G)$  (*cf.* [1], p. 329). As before, let  $F = C(x)$ .

One wishes to find  $f_{11}, \dots, f_{nn} \in F$  and  $p_{11}, \dots, p_{nn} \in F$  such that:

- 1) the  $f_{ij}$  are linearly independent over  $C$ ,

- 2) the ideal  $P$  generated by  $p_{ij} \otimes 1 - 1 \otimes \xi_{ij}$  is a maximal ideal of  $F[G_F]$ ,  
3) if  $D = \frac{d}{dx} \otimes 1 + \sum_{ij} f_{ij} \otimes D_{ij}$  then  $P$  is a  $D$ -stable ideal.

We will then have a counterexample to the theorem.

Leave condition 1 aside for the moment. Condition 2 is satisfied so long as  $\det(p)$  is not 0. This is equivalent to saying that  $p \in G_F$ . Condition 3 is more interesting.

$$\begin{aligned}
D(p_{\mu\nu} \otimes 1 - 1 \otimes \xi_{\mu\nu}) &= p'_{\mu\nu} \otimes 1 - \sum_{ij} f_{ij} \otimes D_{ij} \xi_{\mu\nu} \\
&= p'_{\mu\nu} \otimes 1 - \sum_{ijk} f_{ij} \otimes \xi_{jk} \frac{\partial \xi_{\mu\nu}}{\partial \xi_{ik}} \\
&= p'_{\mu\nu} \otimes 1 - \sum_j f_{\mu j} \otimes \xi_{j\nu} \\
&\equiv p'_{\mu\nu} \otimes 1 - \sum_j f_{\mu j} p_{j\nu} \otimes 1 \quad (\text{mod } P).
\end{aligned}$$

So it suffices to find a matrix  $p = (p_{\mu\nu})$  such that  $p'_{\mu\nu} = \sum_j f_{\mu j} p_{j\nu}$ , or, in matrix form,  $f = p'p^{-1}$ .

In order that all the conditions be satisfied (including condition 1), we need to find examples of  $p \in G_F$  such that the coordinates of its logarithmic derivative are linearly independent over  $C$ . We now do this for  $\text{GL}(2)$ . Let

$$\begin{aligned}
a &= 2^u \\
b &= 2^v \\
c &= 2^w \\
d &= 2^x
\end{aligned}$$

where the  $u, v, w, x$  are positive integers chosen so that with  $u < v < w < x$  (in fact  $a = 2, b = 2^2, c = 2^3, d = 2^4$  will work). Let  $p = (p_{\mu\nu}) =$

$$\begin{pmatrix} \frac{1}{x^a} & \frac{1}{x^b} \\ \frac{1}{x^c} & \frac{1}{x^d} \end{pmatrix}$$

We then have that  $p'p^{-1} =$

$$\begin{pmatrix} \frac{1}{x^a} & \frac{1}{x^b} \\ \frac{1}{x^c} & \frac{1}{x^d} \end{pmatrix}' \begin{pmatrix} \frac{1}{x^a} & -\frac{1}{x^b} \\ -\frac{1}{x^c} & \frac{1}{x^d} \end{pmatrix} \frac{x^{a+b+c+d}}{x^{b+c} - x^{a+d}}$$

To show that the entries of this matrix are linearly independent over the constants, it is enough to show that the entries of the following matrix are linearly independent

$$\begin{aligned}
\begin{pmatrix} \frac{1}{x^a} & \frac{1}{x^b} \\ \frac{1}{x^c} & \frac{1}{x^d} \end{pmatrix}' \begin{pmatrix} \frac{1}{x^d} & -\frac{1}{x^b} \\ -\frac{1}{x^c} & \frac{1}{x^a} \end{pmatrix} &= \begin{pmatrix} \frac{-a}{x^{a+1}} & \frac{-b}{x^{b+1}} \\ \frac{-c}{x^{c+1}} & \frac{-d}{x^{d+1}} \end{pmatrix} \begin{pmatrix} \frac{1}{x^d} & -\frac{1}{x^b} \\ -\frac{1}{x^c} & \frac{1}{x^a} \end{pmatrix} \\
&= \begin{pmatrix} \frac{-a}{x^{a+d+1}} + \frac{b}{x^{c+b+1}} & \frac{a}{x^{a+b+1}} - \frac{b}{x^{a+b+1}} \\ \frac{-c}{x^{c+d+1}} + \frac{d}{x^{c+d+1}} & \frac{c}{x^{b+c+1}} - \frac{d}{x^{a+d+1}} \end{pmatrix}
\end{aligned}$$

Assume we have a relation  $0 =$

$$A\left(\frac{-a}{x^{a+d+1}} + \frac{b}{x^{c+b+1}}\right) + B\left(\frac{a}{x^{a+b+1}} - \frac{b}{x^{a+b+1}}\right) + C\left(\frac{-c}{x^{c+d+1}} + \frac{d}{x^{c+d+1}}\right) + D\left(\frac{c}{x^{b+c+1}} - \frac{d}{x^{a+d+1}}\right)$$

for some constants  $A, B, C, D$ . Since  $d + c + 1 > d + a + 1 > c + b + 1 > a + b + 1$  (these are 2-adic expansions) we must have that

$$\begin{aligned} C(d - c) &= 0 \\ A(-a) + D(-d) &= 0 \\ A(b) + D(c) &= 0 \\ B(a - b) &= 0 \end{aligned}$$

Since  $d > c$  and  $b > a$ , we have that  $C = B = 0$ . Since  $ac > bd$  we have that  $-ac + bd \neq 0$  so  $A = D = 0$ . Therefore the entries of  $p'p^{-1}$  are linearly independent. A similar idea should work for  $\text{GL}(n)$ ,  $n > 2$  as well.

## References

- [1] E. R. Kolchin *Differential Algebra and Algebraic Groups*, Academic Press, 1973.
- [2] J. Kovacic, *E-mail communication to A. Magid, C. Mitschi, M.F. Singer, et. al.* March 2, 1995.
- [3] A. Magid, *Lectures on Differential Galois Theory*, 7, *University Lecture Series*, Vol. 7, American Mathematical Society, 1994.
- [4] C. Mitschi, M.F. Singer, *Connected linear groups as differential Galois groups over  $C(x)$* , to appear in *Journal of Algebra*.