



HAL
open science

Sequential estimation of the parameters in unstable AR(2)

Leonid Galtchouk, Victor Konev

► **To cite this version:**

Leonid Galtchouk, Victor Konev. Sequential estimation of the parameters in unstable AR(2). 2004.
hal-00129705

HAL Id: hal-00129705

<https://hal.science/hal-00129705v1>

Preprint submitted on 8 Feb 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sequential estimation of the parameters in unstable AR(2). *

L. Galtchouk

IRMA, Department of Mathematics,
Strasbourg University
7, st. René Descartes
Strasbourg Cedex France
e-mail: galtchou@math.u-strasbg.fr

and

V.Konev

Department of Applied Mathematics and Cybernetics,
Tomsk University
Lenin str. 36, 634050 Tomsk, Russia
e-mail: vvkonev@vmm.tsu.ru

Abstract

For a second order non-explosive autoregressive process with unknown vector parameter $\theta = (\theta_1, \theta_2)'$, it is shown that the sequential least squares estimate with a particular stopping time is asymptotically normally distributed uniformly in θ belonging to any compact set in the stability region of the process supplemented with the part of its boundary corresponding to complex roots of the characteristic polynomial.

AMS 1991 Subject Classification : 62L10, 62L12.

Key words and phrases: Autoregressive process, least squares estimator, sequential estimation, uniform asymptotic normality.

*This research was carried out at the Department of Mathematics, the Strasbourg University, France. The second author is partially supported by the RFFI-DFG-Grant 02-01-04001.

1 Introduction

Consider the autoregressive $AR(2)$ model

$$x_n = \theta_1 x_{n-1} + \theta_2 x_{n-2} + \varepsilon_n, \quad n = 1, 2, \dots, \quad (1.1)$$

where (x_n) is the observation, (ε_n) is a sequence of independent identically distributed (i.i.d.) random variables with $\mathbf{E}\varepsilon_1 = 0$ and $0 < \mathbf{E}\varepsilon_1^2 = \sigma^2 < \infty$, σ^2 is known, $x_0 = x_{-1} = 0$. The process (1.1) is assumed to be unstable, that is both roots of the characteristic polynomial

$$\mathcal{P}(z) = z^2 - \theta_1 z - \theta_2 \quad (1.2)$$

lie on or inside the unit circle. The model (1.1) is a particular case of unstable autoregressive process $AR(p)$ which have been studied by many authors due to their applications in automatic control, identification and in modeling economic and financial time series (we refer the reader to Anderson (1971), Ahtola and Tiao (1987), Dickey and Fuller (1979), Chan and Wei (1988), Rao (1978) for details and further references).

A commonly used estimate of parameter vector $\theta = (\theta_1, \theta_2)'$ is the least squares estimate (LSE)

$$\theta(n) = (\theta_1(n), \theta_2(n))' = M_n^{-1} \sum_{k=1}^n X_{k-1} x_k, \quad M_n = \sum_{k=1}^n X_{k-1} X_{k-1}', \quad (1.3)$$

where $X_k = (x_k, x_{k-1})'$; the prime denotes the transposition; M_n^{-1} denotes the inverse of matrix M_n if $\det M_n > 0$ and $M_n^{-1} = 0$ otherwise.

It is well known that

$$\sqrt{n}(\theta(n) - \theta) \xrightarrow{\mathcal{L}} \mathcal{N}(0, F), \quad \text{as } n \rightarrow \infty,$$

for all $\theta \in \Lambda$, where Λ is the stability region of process (1.1), that is

$$\Lambda = \{\theta = (\theta_1, \theta_2)' : -1 + \theta_2 < \theta_1 < 1 - \theta_2, |\theta_2| < 1\}, \quad (1.4)$$

$F = F(\theta)$ is a positive definite matrix (see, e.g. Anderson (1971), Th. 5.5.7), $\xrightarrow{\mathcal{L}}$ indicates convergence in law. If θ belongs to the boundary $\partial\Lambda$ of the stability region Λ , the limiting distribution is no longer normal. Moreover, there is no one universal limiting distribution for all $\theta \in \partial\Lambda$ and the corresponding set of limiting distributions counts 6 different types depending on the values of roots z_1 and z_2 of polynomial (1.2). Each of the limiting distribution coincides with that of the ratio of certain Brownian functionals

(see Chan and Wei (1988) for general results and details). For example, for a pair of conjugate complex roots $z_1 = e^{i\varphi}$, $z_2 = e^{-i\varphi}$ one has

$$n \cdot (\theta_1(n) - 2 \cos \varphi) \xrightarrow{\mathcal{L}} \frac{(W_1^2(1) - W_2^2(1)) \sin \varphi + (W_1^2(1) + W_2^2(1) - 2) \cos \varphi}{\int_0^1 [W_1^2(s) + W_2^2(s)] ds},$$

$$n \cdot (\theta_2(n) + 1) \xrightarrow{\mathcal{L}} (2 - W_1^2(1) - W_2^2(1)) / \int_0^1 [W_1^2(s) + W_2^2(s)] ds,$$

where $(W_1(t), 0 \leq t \leq 1)$ and $(W_2(t), 0 \leq t \leq 1)$ are independent standard Brownian motion processes. It is well-known that a similar situation takes place in case of $AR(1)$ process

$$x_n = \theta x_{n-1} + \varepsilon_n,$$

for which the limiting distributions of the least squares estimate are not normal at the end-points $\theta = \pm 1$ of stability interval $(-1, 1)$ (see, e.g. White (1958)).

Lai and Siegmund (1983) for a first order non-explosive autoregressive process proposed to use a sequential sampling scheme and proved that the sequential least squares estimate for θ with the stopping time based on the observed Fisher information is asymptotically normal uniformly in $\theta \in [-1, 1]$ in contrast with the usual LSE.

In this paper we apply a sequential sampling scheme for estimating parameter vector $\theta = (\theta_1, \theta_2)'$ in (1.1). The sequential least squares estimate is defined by replacing sample size n in (1.3) with the stopping time

$$\tau(h) = \inf \left\{ n \geq 1 : \sum_{k=1}^n (x_{k-1}^2 + x_{k-2}^2) \geq h \right\}, \quad \inf \{\emptyset\} = +\infty, \quad (1.5)$$

where h is a positive number (threshold). Our main result (Theorems 2.1) claims that as $h \rightarrow \infty$

$$M_{\tau(h)}^{1/2}(\theta(\tau(h)) - \theta) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma^2 I), \quad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (1.6)$$

uniformly in $\theta \in K$, K is any compact set in the stability region (1.4) supplemented with the part of its boundary

$$\{\theta = (\theta_1, \theta_2)' : -1 + \theta_2 < \theta_1 < 1 - \theta_2, \theta_2 = -1\}, \quad (1.7)$$

It will be observed that this part of the boundary includes the values of θ corresponding to the case when roots of polynomial (1.2) are $e^{i\phi}$ and $e^{-i\phi}$.

Note that for stable autoregressive processes of order 1 with two unknown parameters and of order $p, p \geq 1$, the uniform asymptotic normality have been proved in author's papers (2003_a, 2003_b).

The remainder of this paper is arranged as follows. Section 2 gives proofs of (1.6) and some properties of stopping time (1.5). In Section 3 some properties of unstable $AR(2)$ process needed to prove the main results are established. The appendix contains some technical results.

2 Sequential least squares estimate. Uniform asymptotic normality.

In this section we consider the sequential least squares estimate

$$\theta(\tau(h)) = M_{\tau(h)}^{-1} \sum_{k=1}^{\tau(h)} X_{k-1} x_k, \quad (2.1)$$

$$\tau(h) = \inf \left\{ n \geq 1 : \sum_{k=1}^n \|X_{k-1}\|^2 \geq h \right\}, \quad \inf\{\emptyset\} = +\infty, \quad (2.2)$$

and study its properties. First we will establish the property of uniform asymptotic normality (1.6).

Theorem 2.1 *Let $(\varepsilon_n)_{n \geq 1}$ be a sequence of independent identically distributed random variables with $\mathbf{E}\varepsilon_n = 0$ and $0 < \mathbf{E}\varepsilon_n^2 = \sigma^2 < \infty$. Then for any compact set $K \subset \Lambda_1$*

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} \sup_{t \in R^2} |\mathbf{P}_\theta \left(M_{\tau(h)}^{1/2} (\theta(\tau(h)) - \theta) \leq t \right) - \Phi_2(t/\sigma)| = 0,$$

where $\Phi_2(t) = \Phi(t_1)\Phi(t_2)$, Φ is the standard normal distribution function,

$$\Lambda_1 = \{\theta = (\theta_1, \theta_2) : -1 + \theta_2 < \theta_1 < 1 - \theta_2, -1 \leq \theta_2 < 1\}.$$

Proof. Substituting (1.1) in (2.1) yields

$$M_{\tau(h)}^{1/2} (\theta(\tau(h)) - \theta) = M_{\tau(h)}^{-1/2} \sum_{k=1}^{\tau(h)} X_{k-1} \varepsilon_k = \sqrt{h} M_{\tau(h)}^{-1/2} L^{1/2}(\theta_1, \theta_2) Y_h, \quad (2.3)$$

where $L(\theta_1, \theta_2)$ is given in Lemma 3.4,

$$Y_h = \frac{1}{\sqrt{h}} \sum_{k=1}^{\tau(h)} L^{-1/2}(\theta_1, \theta_2) X_{k-1} \varepsilon_k. \quad (2.4)$$

To prove Theorem 2.1 it suffices to establish the following Lemmas.

Lemma 2.1 *Under conditions of Theorem 2.1 for any compact set $K \subset \Lambda_1$ and $\delta > 0$*

$$\limsup_{h \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta \left(\|\sqrt{h} M_{\tau(h)}^{-1/2} L^{1/2}(\theta_1, \theta_2) - I_2\| > \delta \right) = 0,$$

where I_2 is the unit matrix of order 2.

Lemma 2.2 *Under conditions of Theorem 2.1 for any compact set $K \subset \Lambda_1$ and for each constant vector $v \in R^2$ with $\|v\| = 1$*

$$\limsup_{h \rightarrow \infty} \sup_{\theta \in K} \sup_{t \in R} |\mathbf{P}_\theta(v' Y_h \leq t) - \Phi(t)| = 0.$$

The proofs of these results are given in the Appendix.

By Lemma 2.2 vector Y_h in (2.4) is asymptotically normal uniformly in $\theta \in K$. In view of Lemma 2.1 this completes the proof of Theorem 2.1.

Now we will study the properties of the stopping time $\tau(h)$ defined by (2.2). Denote

$$\Gamma_1 = \{\theta = (\theta_1, \theta_2) : -\theta_1 + \theta_2 = 1, -2 < \theta_1 < 0\},$$

$$\Gamma_2 = \{\theta = (\theta_1, \theta_2) : \theta_1 + \theta_2 = 1, 0 < \theta_1 < 2\},$$

$$\Gamma_3 = \{\theta = (\theta_1, -1) : -2 < \theta_1 < 2\},$$

$$A = \begin{pmatrix} \theta_1 & \theta_2 \\ 1 & 0 \end{pmatrix}; \quad B = \begin{pmatrix} \sigma^2 & 0 \\ 0 & 0 \end{pmatrix}.$$

Theorem 2.2 *Let $(\varepsilon_n)_{n \geq 1}$ in (1.1) be a sequence of independent identically distributed random variables with $E\varepsilon_n = 0$, $E\varepsilon_n^2 = \sigma^2$ and $\tau(h)$ be defined by (2.2). Let z_1 and z_2 be roots of polynomial (1.2). Then*

for each $\theta \in \Lambda$

$$(2.5) \quad \mathbf{P}_\theta - \lim_{h \rightarrow \infty} \frac{\tau(h)}{h} = \frac{1}{\text{tr} F}, \quad F - AFA' = B;$$

for each $\theta \in \Gamma_1$

$$(2.6) \quad \frac{\tau(h)}{(1 + z_2)\sqrt{h/2}} \xrightarrow{\mathcal{L}} \nu_1, \quad \text{as } h \rightarrow \infty, \quad \nu_1 = \inf \left\{ t \geq 0 : \int_0^t W^2(s) ds = 1 \right\},$$

for each $\theta \in \Gamma_2$

$$(2.7) \quad \frac{\tau(h)}{(1 - z_1)\sqrt{h/2}} \xrightarrow{\mathcal{L}} \nu_2, \quad \text{as } h \rightarrow \infty, \quad \nu_2 = \inf \left\{ t \geq 0 : \int_0^t W_1^2(s) ds = 1 \right\},$$

for each $\theta = (2 \cos \varphi, -1) \in \Gamma_3$

$$(2.8) \quad \frac{\tau(h)}{\sqrt{2h \sin \varphi}} \xrightarrow{\mathcal{L}} \nu_3, \text{ as } h \rightarrow \infty, \nu_3 = \inf \left\{ t \geq 0 : \int_0^t (W^2(s) + W_1^2(s)) ds \geq 1 \right\},$$

where $W(t)$ and $W_1(t)$ are independent standard Brownian motions, Λ is defined in (1.4).

Proof Assertion (2.5) easily follows from Lemma 3.12 in Galtchouk and Konev (2003_b).

Let $\theta \in \Gamma_1$. In this case polynomial (1.2) takes the form

$$\mathcal{P}(z) = (z + 1)(z - a)$$

where $-1 < a < 1$. By applying the backshift operator q^{-1} one can write down equation (1.1) as

$$(2.8) \quad q^{-2}(q + 1)(q - a)x_k = \varepsilon_k.$$

Denote

$$u_k = q^{-1}(q - a)x_k, \quad v_k = q^{-1}(q + 1)x_k,$$

that is

$$x_k - ax_{k-1} = u_k, \quad x_k + x_{k-1} = v_k,$$

or in the vector form

$$(2.9) \quad QX_k = \begin{pmatrix} u_k \\ v_k \end{pmatrix}; \quad Q = \begin{pmatrix} 1 & -a \\ 1 & 1 \end{pmatrix}.$$

The processes u_k and v_k satisfy the equations

$$u_k = -u_{k-1} + \varepsilon_k, \quad v_k = av_{k-1} + \varepsilon_k, \quad u_0 = 0, \quad v_0 = 0.$$

From here one gets

$$u_k = -\sum_{j=1}^k (-1)^j \varepsilon_j, \quad v_k = \sum_{j=1}^k a^{k-j} \varepsilon_j.$$

The identity

$$(2.10) \quad \sum_{k=1}^n x_{k-1}^2 = (1/2) \sum_{k=1}^n \|X_{k-1}\|^2 + \frac{x_{n-1}^2}{2}$$

and (2.9) imply

(2.11)

$$\sum_{k=1}^n x_{k-1}^2 = \frac{1}{2}x_{n-1}^2 + \frac{1}{(a+1)^2} \sum_{k=1}^n u_{k-1}^2 + \frac{a-1}{a+1} \sum_{k=1}^n u_{k-1}v_{k-1} + \frac{a^2+1}{2(a+1)^2} \sum_{k=1}^n v_{k-1}^2.$$

It is known that (Lai and Wei (1983), Chan and Wei (1988))

$$\liminf_{n \rightarrow \infty} \frac{\sum_{k=1}^n u_{k-1}^2}{n^2 / \log \log n} = \sigma^2 / 4 \quad \mathbf{P}_\theta - \text{a.s.},$$

$$P_\theta - \lim_{n \rightarrow \infty} n^{-3/2} \sum_{k=1}^n u_{k-1}v_{k-1} = 0.$$

This and (2.11) yield

$$(2.12) \quad P_\theta - \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n x_{k-1}^2}{\sum_{k=1}^n u_{k-1}^2} = (a+1)^{-2}.$$

Further we have

$$P_\theta \left(\frac{\tau(h)}{b\sqrt{h}} \leq t \right) = P_\theta \left(\sum_{k=1}^{\lfloor tb\sqrt{h} \rfloor} \|X_{k-1}\|^2 \geq h \right) = P_\theta \left(2 \sum_{k=1}^{\lfloor tb\sqrt{h} \rfloor} x_{k-1}^2 - x_{\lfloor tb\sqrt{h} \rfloor - 1} \geq h \right),$$

where $b = (a+1)/\sqrt{2}$. Applying herein Lemma 3.2 and (2.10) yields

$$P_\theta \left(\frac{\tau(h)}{b\sqrt{h}} \leq t \right) = P_\theta \left(\frac{1}{b^2 h} \sum_{k=1}^{\lfloor tb\sqrt{h} \rfloor} u_{k-1}^2 \mu_{\lfloor tb\sqrt{h} \rfloor} \geq 1 \right),$$

where $P_\theta - \lim_{h \rightarrow \infty} \mu_{\lfloor tb\sqrt{h} \rfloor} = 1$. From here by Donsker's Theorem we come to (2.6). By a similar argument one can show (2.7).

Assume that $\theta \in \Gamma_3$. Then in view of (2.10) one has

(2.13)

$$P_\theta \left(\frac{\tau(h)}{bh^{1/2}} \leq t \right) = P_\theta \left(\sum_{k=1}^{\lfloor tbh^{1/2} \rfloor} \|X_{k-1}\|^2 \geq h \right) = P_\theta \left(2b^2 Z_{\lfloor tbh^{1/2} \rfloor} \tilde{\mu}_{\lfloor tb\sqrt{h} \rfloor} \geq 1 \right),$$

where

$$Z_{\lfloor tb\sqrt{h} \rfloor} = \frac{1}{b^2 h} \sum_{k=1}^{\lfloor tbh^{1/2} \rfloor} x_{k-1}^2,$$

and

$$P_\theta - \lim_{h \rightarrow \infty} \tilde{\mu}_{\lfloor tb\sqrt{h} \rfloor} = 1.$$

Now we apply Theorem (3.3.4) from the paper of Chan and Wei (1988), which yields

$$Z_{[tb\sqrt{h}]} \xrightarrow{\mathcal{L}} \frac{1}{4 \sin^2 \phi} \int_0^t (W^2(s) + W_1^2(s)) ds.$$

Putting $b = \sqrt{2} \sin \phi$ in (2.13) and taking this into account we come to (2.8). This completes the proof.

3 Auxiliary propositions.

In this section we establish some properties of the process (1.1) and the observed Fisher information matrix M_n .

In the sequel we will need the following two probabilistic results for martingales from the paper of Lai and Siegmund (1983).

Proposition 3.1 *Let $x_n, \varepsilon_n, n = 0, 1, \dots$ be random variables adapted to the increasing sequence of σ -algebras $(\mathcal{F}_n)_{n \geq 0}$. Let $\{\mathbf{P}_\theta, \theta \in \Theta\}$ be a family of probability measures such that under every \mathbf{P}_θ*

- A_1 : $\varepsilon_1, \varepsilon_2, \dots$ are i.i.d. with $\mathbf{E}_\theta \varepsilon_1 = 0, \mathbf{E}_\theta \varepsilon_1^2 = 1$;
- A_2 : $\sup_\theta \mathbf{E}_\theta \{\varepsilon_1^2; |\varepsilon_1| > a\} \rightarrow 0$ as $a \rightarrow \infty$;
- A_3 : ε_n is independent of \mathcal{F}_{n-1} for each $n \geq 1$;
- A_4 : $\mathbf{P}_\theta(\sum_{i=0}^\infty x_i^2 = \infty) = 1$;
- A_5 : $\sup_\theta \mathbf{P}_\theta(x_n^2 > a) \rightarrow 0$ as $a \rightarrow \infty$ for each $n \geq 0$;
- A_6 : $\lim_{m \rightarrow \infty} [\sup_\theta \mathbf{P}_\theta(x_n^2 \geq \delta \sum_{i=0}^{n-1} x_i^2 \text{ for some } n \geq m)] = 0$ for each $\delta > 0$.

For $c > 0$ let $T_c = \inf\{n : \sum_{i=1}^n x_{i-1}^2 \geq c\}, \inf\{\emptyset\} = +\infty$. Then uniformly in $\theta \in \Theta$ and $-\infty < t < \infty$

$$\mathbf{P}_\theta \left\{ c^{-1/2} \sum_{i=1}^{T_c} x_{i-1} \varepsilon_i \leq t \right\} \rightarrow \Phi(t) \text{ as } c \rightarrow \infty,$$

where Φ is the standard normal distribution function.

Lemma 3.1 *Suppose that the measurability conditions of Proposition 3.1, A_1, A_3 are satisfied. Then for each $\gamma > 1/2, \delta > 0$, and increasing sequence of positive constants $c_n \rightarrow \infty$,*

$$\sup_{\theta \in \Theta} \mathbf{P}_\theta \left\{ \left| \sum_{k=1}^n x_{k-1} \varepsilon_k \right| \geq \delta \max(c_n, \left(\sum_{k=1}^n x_{k-1}^2 \right)^\gamma) \text{ for some } n \geq m \right\} \rightarrow 0$$

as $m \rightarrow \infty$.

Now we show the following result.

Lemma 3.2 *Let $(\varepsilon_n)_{n \geq 1}$ in (1.1) be a sequence of i.i.d. random variables with $\mathbf{E}\varepsilon_n = 0$ and $\mathbf{E}\varepsilon_n^2 = \sigma^2 < \infty$ and roots of the characteristic polynomial (1.2) lie on or inside the unit circle, i.e. $|z_1| \leq 1$, $|z_2| \leq 1$. Then, for any compact set $K \subset \Lambda_0$ and $\delta > 0$,*

$$\lim_{m \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta \left(\max(x_n^2, x_{n-1}^2) \geq \delta \sum_{k=1}^n x_{k-1}^2 \text{ for some } n \geq m \right) = 0,$$

where

$$\Lambda_0 = [\Lambda] \setminus \{(-2, -1), (2, -1)\}, \quad (3.1)$$

$[\Lambda]$ is the closure of the stability region (1.4).

Proof. In view of the equality

$$\sum_{k=1}^n x_{k-1}^2 = (1/2) \sum_{k=1}^n \|X_{k-1}\|^2 + \frac{x_{n-1}^2}{2} \quad (3.2)$$

it suffices to show that, for any compact set $K \subset \Lambda_0$ and $\delta > 0$,

$$\lim_{m \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta(B_m(\delta)) = 0, \quad (3.3)$$

where

$$B_m(\delta) = \{ \|X_n\|^2 \geq \delta \sum_{k=1}^n \|X_{k-1}\|^2 \text{ for some } n \geq m \}.$$

By making use of the vector form of equation (1.1)

$$X_n = AX_{n-1} + \xi_n, \quad (3.4)$$

where $X_n = (x_n, x_{n-1})'$, $\xi_n = (\varepsilon_n, 0)'$,

$$A = \begin{pmatrix} \theta_1 & \theta_2 \\ 1 & 0 \end{pmatrix}, \quad \xi_n = \begin{pmatrix} \varepsilon_n \\ 0 \end{pmatrix}. \quad (3.5)$$

one obtains

$$\sum_{k=1}^n X_{k-1} X_{k-1}' \geq \sum_{k=1}^n \xi_k \xi_k' - X_n X_n' + \sum_{k=1}^n \xi_k X_{k-1}' A' + A \sum_{k=1}^n X_{k-1} \xi_k.$$

This implies

$$\sum_{k=1}^n \|X_{k-1}\|^2 \geq \sum_{k=1}^n \varepsilon_k^2 - \|X_n\|^2 - 2 \left| \sum_{k=1}^n \langle AX_{k-1} \rangle_1 \varepsilon_k \right|, \quad (3.6)$$

where $\langle AX_{k-1} \rangle_1$ is the first coordinate of vector AX_{k-1} .

For each $0 < \lambda < \sigma^2/4$ and $m \geq 1$ we introduce the sets

$$\Omega_{m,\lambda} = \left\{ \left| n^{-1} \sum_{k=1}^n \varepsilon_k^2 - \sigma^2 \right| < \lambda \right\} \cap \left\{ \left| \sum_{k=1}^n \langle AX_{k-1} \rangle_1 \varepsilon_k \right| < \max \left(\lambda n, \left(\sum_{k=1}^n \langle AX_{k-1} \rangle_1^2 \right)^{2/3} \right) \text{ for all } n \geq m \right\}.$$

It is easily seen that

$$\Omega_{m,\lambda} \subset \left\{ \sum_{k=1}^m \|X_{k-1}\|^2 \leq 1 \right\} \cap \Omega'_{m,\lambda}, \quad (3.7)$$

where

$$\begin{aligned} \Omega'_{m,\lambda} &\subset \cap_{n \geq m} (\mathcal{D}_n \cup \{\|X_n\|^2 > \lambda n\}), \quad (3.8) \\ \mathcal{D}_n &= \left\{ \left| n^{-1} \sum_{k=1}^n \varepsilon_k^2 - \sigma^2 \right| < \lambda \right\} \cap \\ &\left\{ \left| \sum_{k=1}^n \langle AX_{k-1} \rangle_1 \varepsilon_k \right| < \max \left(\lambda n, \left(\sum_{k=1}^n \langle AX_{k-1} \rangle_1^2 \right)^{2/3} \right), \right. \\ &\left. \|X_n\|^2 \leq \lambda n, \sum_{k=1}^m \|X_{k-1}\|^2 > 1 \right\}. \end{aligned}$$

By (3.6), on the set \mathcal{D}_n , one has the inequality

$$\begin{aligned} \sum_{k=1}^n \|X_{k-1}\|^2 &\geq (\sigma^2 - \lambda)n - \lambda n - 2\lambda n - 2 \left(\sum_{k=1}^n \|AX_{k-1}\|^2 \right)^{2/3} \\ &\geq (\sigma^2 - 4\lambda)n - 2\|A\|^{4/3} \left(\sum_{k=1}^n \|X_{k-1}\|^2 \right)^{2/3}, \end{aligned}$$

which implies

$$(1 + 2\|A\|^{4/3}) \sum_{k=1}^n \|X_{k-1}\|^2 \geq \left(\frac{\sigma^2}{\lambda} - 4\right)\lambda n \geq \left(\frac{\sigma^2}{\lambda} - 4\right)\|X_n\|^2.$$

Given $\delta > 0$, by choosing λ so that

$$(1 + 2 \sup_{\theta \in K} \|A\|^{4/3}) \left(\frac{\sigma^2}{\lambda} - 4\right)^{-1} < \delta$$

one has, on the set \mathcal{D}_n , the inequality

$$\|X_n\|^2 < \delta \sum_{k=1}^n \|X_{k-1}\|^2, \quad n \geq m.$$

This and (3.8) lead to the inclusion

$$\Omega'_{m,\lambda} \subset \bigcap_{n \geq m} \left(\left\{ \|X_n\|^2 < \delta \sum_{k=1}^n \|X_{k-1}\|^2 \right\} \cup \{ \|X_n\|^2 > \lambda n \} \right). \quad (3.9)$$

Now assume that $\|X_n\|^2 \geq \lambda n$. By (3.4) one gets

$$X_n = AX_{n-k} + \sum_{j=0}^{k-1} A^j \xi_{n-j}. \quad (3.10)$$

Further we will show that for every compact set $K \subset \Lambda_0$ there exists a positive number κ such that

$$\sup_{\theta \in K} \|A^n\| \leq \kappa, \quad n \geq 1. \quad (3.11)$$

Without loss of generality we can consider a compact set of the form

$$K_d = [-2+d, 2-d] \times [-1, -1+d] \cup \{\theta : -1+\theta_2 \leq \theta_1 \leq 1-\theta_2, -1+d \leq \theta_2 \leq 1\},$$

where $0 < d \leq 2$. One can verify that if roots z_1 and z_2 of the characteristic polynomial (1.2) are real then

$$A^n = \frac{1}{z_1 - z_2} \begin{pmatrix} z_1^{n+1} - z_2^{n+1} & -(z_1^n - z_2^n)z_1 z_2 \\ z_1^n - z_2^n & -(z_1^{n-1} - z_2^{n-1})z_1 z_2 \end{pmatrix}, \quad -1 \leq z_1 < z_2 \leq 1,$$

and, if the roots are complex, that is $z_1 = ae^{i\phi}$ and $z_2 = ae^{-i\phi}$, then

$$A^n = \frac{a^{n-1}}{\sin \phi} \begin{pmatrix} \sin(n+1)\phi & -a^2 \sin(n\phi) \\ \sin n\phi & -\sin(n-1)\phi \end{pmatrix}.$$

By making use of these formulas respectively on the subsets

$$K'_d = [-2+d, 2-d] \times [-1, -1+d] \cup \{\theta : -2+d \leq \theta_1 \leq 2-d, -1 \leq \theta_2 \leq -\frac{1}{4}\theta_1^2\},$$

$$K''_d = \{\theta : -2+d \leq \theta_1 \leq 0, -\frac{1}{4}\theta_1^2 < \theta_2 \leq 1+\theta_1\}$$

$$\cup \{\theta : 0 \leq \theta_1 \leq 2-d, -\frac{1}{4}\theta_1^2 < \theta_2 \leq 1-\theta_1\},$$

we establish that (3.9) is satisfied with $K = K_d$. From (3.10) in view of (3.11) one obtains

$$\kappa \|X_{n-i}\| \geq \|A^i\| \cdot \|X_{n-i}\| \geq \|A^i X_{n-i}\| \geq \|X_n\| - \sum_{j=0}^{i-1} \|A^j\| \cdot |\varepsilon_{n-j}|.$$

This implies that for $k \leq n$

$$\begin{aligned} \kappa \min_{1 \leq i \leq k} \|X_{n-i}\| &\geq \|X_n\| - \sum_{j=0}^{k-1} \|A^j\| \cdot |\varepsilon_{n-j}| \\ &\geq \|X_n\| - \zeta_{n,k} \geq \|X_n\| (1 - (\lambda n)^{-1/2} \zeta_{n,k}), \end{aligned}$$

where

$$\zeta_{n,k} = \kappa \sum_{j=0}^{k-1} |\varepsilon_{n-j}|.$$

Therefore

$$\sum_{i=1}^n \|X_{i-1}\|^2 \geq \sum_{i=1}^n \|X_{n-i}\|^2 \geq \|X_n\|^2 k (1 - (\lambda n)^{-1/2} \zeta_{n,k})^2 / \kappa, \quad k \leq n.$$

This inequality yields the inclusions

$$\{\|X_n\|^2 > \lambda n\} \subset \left\{ \sum_{i=1}^n \|X_{i-1}\|^2 \geq \|X_n\|^2 \frac{k}{\kappa} (1 - \Delta)^2 \right\} \cup \left\{ \sup_{n \geq m} (\lambda n)^{-1/2} \zeta_{n,k} > \Delta \right\}$$

for $n \geq m$, where $0 < \Delta < 1$. Choosing k so that

$$k(1 - \Delta)^2 / \kappa^2 > \delta^{-1}$$

we obtain

$$\{\|X_n\|^2 > \lambda n\} \supset \left\{ \|X_n\|^2 < \delta \sum_{k=1}^n \|X_{k-1}\|^2 \right\} \subset \left\{ \sup_{n \geq m} (\lambda n)^{-1/2} \zeta_{n,k} > \Delta \right\}, \quad n \geq m.$$

From here and (3.7), (3.9) it follows that

$$\Omega_{m,\lambda} \subset \left\{ \sum_{k=1}^m \|X_{k-1}\|^2 \leq 1 \right\} \bigcup \left\{ \sup_{n \geq m} (\lambda n)^{-1/2} \zeta_{n,k} > \Delta \right\} \\ \bigcup \left\{ \|X_n\|^2 < \delta \sum_{k=1}^n \|X_{k-1}\|^2 \right\}.$$

This implies

$$\Omega_{m,\lambda}^c \supset \left\{ \sum_{k=1}^m \|X_{k-1}\|^2 > 1 \right\} \supset \left\{ \sup_{n \geq m} (\lambda n)^{-1/2} \zeta_{n,k} \leq \Delta \right\} \supset B_m(\delta).$$

Finally we get the inclusion

$$B_m(\delta) \supset \Omega_{m,\lambda}^c \bigcup \left\{ \sum_{k=1}^m \|X_{k-1}\|^2 \leq 1 \right\} \bigcup \left\{ \sup_{n \geq m} \frac{\zeta_{n,k}}{\sqrt{\lambda n}} > \Delta \right\}.$$

Hence

$$\sup_{\Theta \in K} P_{\Theta}(B_m(\delta)) \leq \sup_{\Theta \in K} P_{\Theta}(\Omega_{m,\lambda}^c) + \sup_{\Theta \in K} P_{\Theta} \left\{ \sum_{k=1}^m \|X_{k-1}\|^2 \leq 1 \right\} \\ + \sup_{\Theta \in K} P_{\Theta} \left\{ \sup_{n \geq m} (\lambda n)^{-1/2} \zeta_{n,k} > \Delta \right\}.$$

By Lemma 3.1 with $x_{i-1} = \langle AX_{i-1} \rangle_1$ and $\gamma = 2/3$ the first term in the right-hand side of this inequality vanishes as $m \rightarrow \infty$. The last two summands tend to zero by the law of large numbers.

This completes the proof of Lemma 3.2.

Lemma 3.3 *Let $(\varepsilon_n)_{n \geq 1}$ in (1.1) be a sequence of i.i.d. random variables with $\mathbf{E}\varepsilon_n = 0$ and $\mathbf{E}\varepsilon_n^2 = \sigma^2 < \infty$. Then for any compact set $K \subset \Lambda_0$*

$$\lim_{m \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_{\theta} \left(\left| \sum_{k=1}^n x_{k-1} \varepsilon_k \right| > \delta \sum_{k=1}^n x_{k-1}^2 \text{ for some } n \geq m \right) = 0, \quad (3.12)$$

where Λ_0 is given in (3.1).

Proof. Let $c_n = n^{3/4}$. For the set of interest one has the following inclusions

$$\left\{ \left| \sum_{k=1}^n x_{k-1} \varepsilon_k \right| > \delta \sum_{k=1}^n x_{k-1}^2 \text{ for some } n \geq m \right\}$$

$$\begin{aligned}
&= \left\{ \frac{|\sum_{k=1}^n x_{k-1}\varepsilon_k|}{(\sum_{k=1}^n x_{k-1}^2)^{2/3} \vee c_n} \cdot \frac{(\sum_{k=1}^n x_{k-1}^2)^{2/3} \vee c_n}{\sum_{k=1}^n x_{k-1}^2} > \delta \text{ for some } n \geq m \right\} \\
&\subset \left\{ \frac{|\sum_{k=1}^n x_{k-1}\varepsilon_k|}{(\sum_{k=1}^n x_{k-1}^2)^{2/3} \vee c_n} > \sqrt{\delta} \text{ for some } n \geq m \right\} \cup \\
&\quad \left\{ \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1/3} \vee c_n \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \sqrt{\delta} \text{ for some } n \geq m \right\} \\
&\subset \left\{ \left| \sum_{k=1}^n x_{k-1}\varepsilon_k \right| \left(\left(\sum_{k=1}^n x_{k-1}^2 \right)^{2/3} \vee c_n \right)^{-1} > \sqrt{\delta} \text{ for some } n \geq m \right\} \cup \\
&\quad \cup \left\{ \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \delta^{3/2} \text{ for some } n \geq m \right\} \cup \\
&\quad \cup \left\{ c_n \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \sqrt{\delta} \text{ for some } n \geq m \right\} \\
&\subset \left\{ \left| \sum_{k=1}^n x_{k-1}\varepsilon_k \right| \left(\left(\sum_{k=1}^n x_{k-1}^2 \right)^{2/3} \vee c_n \right)^{-1} > \sqrt{\delta} \text{ for some } n \geq m \right\} \\
&\quad \cup \left\{ c_n \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \sqrt{\delta} \wedge \delta^{3/2} \text{ for some } n \geq m \right\}.
\end{aligned}$$

From here it follows that

$$\begin{aligned}
&\mathbf{P}_\theta \left(\left| \sum_{k=1}^n x_{k-1}\varepsilon_k \right| > \delta \sum_{k=1}^n x_{k-1}^2 \text{ for some } n \geq m \right) \\
&\leq \mathbf{P}_\theta \left\{ \left| \sum_{k=1}^n x_{k-1}\varepsilon_k \right| \left(\left(\sum_{k=1}^n x_{k-1}^2 \right)^{2/3} \vee c_n \right)^{-1} > \sqrt{\delta} \text{ for some } n \geq m \right\} \\
&\quad + \mathbf{P}_\theta \left\{ n^{3/4} \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \sqrt{\delta} \wedge \delta^{3/2} \text{ for some } n \geq m \right\}.
\end{aligned} \tag{3.13}$$

By making use of (1.1) and the elementary inequalities we obtain

$$\sum_{k=1}^n \varepsilon_k^2 = \sum_{k=1}^n (x_k - \theta_1 x_{k-1} - \theta_2 x_{k-2})^2$$

$$\begin{aligned}
&\leq 3 \left(\sum_{k=1}^n x_k^2 + \theta_1^2 \sum_{k=1}^n x_{k-1}^2 + \theta_2^2 \sum_{k=1}^n x_{k-2}^2 \right) \\
&\leq 15(x_n^2 + \sum_{k=1}^n x_{k-1}^2) = 15 \sum_{k=1}^n x_{k-1}^2 \left(1 + \frac{x_n^2}{\sum_{k=1}^n x_{k-1}^2} \right).
\end{aligned}$$

Therefore the second summand in the right-hand side of (3.13) can be estimated as

$$\begin{aligned}
&\mathbf{P}_\theta \left\{ n^{3/4} \left(\sum_{k=1}^n x_{k-1}^2 \right)^{-1} > \sqrt{\delta} \wedge \delta^{3/2} \text{ for some } n \geq m \right\} \\
&\leq \mathbf{P}_\theta \left\{ 30n^{3/4} \left(\sum_{k=1}^n \varepsilon_k^2 \right)^{-1} > \sqrt{\delta} \wedge \delta^{3/2} \text{ for some } n \geq m \right\} \\
&\quad + \mathbf{P}_\theta \left\{ x_n^2 / \sum_{k=1}^n x_{k-1}^2 \geq 1 \text{ for some } n \geq m \right\}.
\end{aligned}$$

Combining this and (3.13) and applying Lemmas 3.1, 3.2 lead to (3.12).

This completes the proof of Lemma 3.3.

Lemma 3.4 *Let $(\varepsilon_n)_{n \geq 1}$ in (1.1) be a sequence of i.i.d. random variables with $\mathbf{E}\varepsilon_n = 0$ and $\mathbf{E}\varepsilon_n^2 = \sigma^2 < \infty$ and M_n be given by (1.3).*

Then, for any compact set $K \subset \overset{\circ}{\Lambda}$ and $\delta > 0$,

$$\lim_{m \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta \left(\left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| \geq \delta \text{ for some } n \geq m \right) = 0, \quad (3.14)$$

where

$$L(\theta_1, \theta_2) = \begin{pmatrix} 1 & \frac{\theta_1}{1-\theta_2} \\ \frac{\theta_1}{1-\theta_2} & 1 \end{pmatrix}, \quad \overset{\circ}{\Lambda} = [\Lambda] \setminus \{(0, 1), (-2, -1), (2, -1)\}. \quad (3.15)$$

Proof. Multiplying (1.1) by x_{k-1} and summing yield

$$\sum_{k=1}^n x_{k-1}x_k = \theta_1 \sum_{k=1}^n x_{k-1}^2 + \theta_2 \sum_{k=1}^n x_{k-1}x_{k-2} + \sum_{k=1}^n x_{k-1}\varepsilon_k.$$

From here it follows that

$$\sum_{k=1}^n x_{k-1}x_{k-2} = -\frac{1}{1-\theta_2} x_n x_{n-1} + \frac{\theta_1}{1-\theta_2} \sum_{k=1}^n x_{k-1}^2 + \frac{1}{1-\theta_2} \sum_{k=1}^n x_{k-1}\varepsilon_k.$$

Substituting this in M_n yields

$$M_n = L(\theta_1, \theta_2) \sum_{k=1}^n x_{k-1}^2 + r_n,$$

where

$$r_n = \frac{1}{1 - \theta_2} \begin{pmatrix} 0; & x_n x_{n-1} + \sum_{k=1}^n x_{k-1} \varepsilon_k \\ -x_n x_{n-1} + \sum_{k=1}^n x_{k-1} \varepsilon_k; & -x_{n-1}^2 \end{pmatrix}.$$

By applying Lemmas 3.3, 3.4 to $r_n / \sum_{k=1}^n x_{k-1}^2$ we come to (3.14). This completes the proof of Lemma 3.4.

Lemma 3.5 *Let $M_n, \tau(h)$ and $L(\theta_1, \theta_2)$ be given by (1.3), (1.5) and (3.15) respectively. Assume that $\{\varepsilon_n\}$ in (1.1) be a sequence of independent identically distributed random variables with $E\varepsilon_n = 0, E\varepsilon_n^2 = \sigma^2$. Then, for any compact set $K \subset \overset{\circ}{\Lambda}$ and $\delta > 0$,*

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta \left(\left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\| > \delta \right) = 0, \quad (3.16)$$

where $\overset{\circ}{\Lambda}$ is the same as in (3.15).

Proof. By making use of the equality

$$\frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) = \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} - L(\theta_1, \theta_2) + M_{\tau(h)} \left(\frac{1}{h} - \frac{1}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} \right)$$

one gets

$$\left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\| \leq \left\| \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} - L(\theta_1, \theta_2) \right\| + \frac{\|M_{\tau(h)}\|}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} \frac{\left(\sum_{k=1}^{\tau(h)} x_{k-1}^2 - h \right)}{h}.$$

From here it follows that

$$\begin{aligned} \left\{ \left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\| > \delta \right\} &\subset \left\{ \left\| \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \delta/2 \right\} \\ &\cup \left\{ \left\| \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} \right\| \frac{\left(\sum_{k=1}^{\tau(h)} x_{k-1}^2 - h \right)}{h} > \delta/2 \right\}. \end{aligned} \quad (3.17)$$

Further we have the inclusions

$$\begin{aligned}
& \left\{ \left\| \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \delta/2 \right\} \subseteq \{ \tau(h) \leq m \} \quad (3.18) \\
& \cup \left\{ \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \delta/2 \text{ for some } n \geq m \right\}, \\
& \left\{ \left\| \frac{M_{\tau(h)}}{\sum_{k=1}^{\tau(h)} x_{k-1}^2} \right\| \frac{\left(\sum_{k=1}^{\tau(h)} x_{k-1}^2 - h \right)}{h} > \delta/2 \right\} \subset \{ \tau(h) \leq m \} \\
& \cup \left\{ \frac{\|M_n\|}{\sum_{k=1}^n x_{k-1}^2} \cdot \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta/2 \text{ for some } n \geq m \right\}.
\end{aligned}$$

From here and (3.17) one obtains

$$\begin{aligned}
& \mathbf{P}_\theta \left\{ \left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\| > \delta \right\} \leq 2\mathbf{P}_\theta \{ \tau(h) \leq m \} \quad (3.19) \\
& + \mathbf{P}_\theta \left\{ \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \delta/2 \text{ for some } n \geq m \right\} \\
& + \mathbf{P}_\theta \left\{ \frac{\|M_n\|}{\sum_{k=1}^n x_{k-1}^2} \cdot \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta/2 \text{ for some } n \geq m \right\}.
\end{aligned}$$

By the definition of $\tau(h)$ in (1.5)

$$\begin{aligned}
& \{ \tau(h) < m \} = \left\{ \sum_{k=1}^m (x_{k-1}^2 + x_{k-2}^2) > h \right\} \\
& = \left\{ \sum_{k=1}^m (x_{k-1}^2 + x_{k-2}^2) > h, \max_{1 \leq j \leq m} (x_{j-1}^2 + x_{j-2}^2) < l \right\} \\
& + \left\{ \sum_{k=1}^m (x_{k-1}^2 + x_{k-2}^2) > h, \max_{1 \leq j \leq m} (x_{j-1}^2 + x_{j-2}^2) \geq l \right\} \\
& \subset \{ ml > h \} \cup \cup_{j=1}^m \{ (x_{j-1}^2 + x_{j-2}^2) \geq l \}.
\end{aligned}$$

Therefore

$$\mathbf{P}_\theta \{ \tau(h) < m \} \leq I_{(ml > h)} + \sum_{k=1}^m \mathbf{P}_\theta \{ x_{k-1}^2 + x_{k-2}^2 \geq l \}. \quad (3.20)$$

It remains to estimate the last term in the right-hand side of (3.17).

By the inequality

$$\frac{\|M_n\|}{\sum_{k=1}^n x_{k-1}^2} \leq \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| + \|L(\theta_1, \theta_2)\|$$

one has

$$\begin{aligned} & \mathbf{P}_\theta \left\{ \frac{\|M_n\|}{\sum_{k=1}^n x_{k-1}^2} \cdot \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta/2 \text{ for some } n \geq m \right\} \\ & \leq \mathbf{P}_\theta \left\{ \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \sqrt{\delta/4} \text{ for some } n \geq m \right\} \\ & \quad + \mathbf{P}_\theta \left\{ L_k^* \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \sqrt{\delta/4} \text{ for some } n \geq m \right\} \\ & \quad + \mathbf{P}_\theta \left\{ L_k^* \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta/4 \text{ for some } n \geq m \right\}, \end{aligned} \tag{3.21}$$

where $L_k^* = \sup_{\theta \in K} \|L(\theta_1, \theta_2)\|$.

Combining (3.19)–(3.21) yields

$$\begin{aligned} & \sup_{\theta \in K} \mathbf{P}_\theta \left(\left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\| > \delta \right) \\ & \leq 2I_{(ml>h)} + 2 \sum_{k=1}^m \sup_{\theta \in K} \mathbf{P}_\theta \{x_{k-1}^2 + x_{k-2}^2 \geq l\} \\ & + \mathbf{P}_\theta \left\{ \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L(\theta_1, \theta_2) \right\| > \frac{1}{2}(\delta \wedge \sqrt{\delta}) \text{ for some } n \geq m \right\} \\ & + \mathbf{P}_\theta \left\{ L_k^* \frac{x_{n-1}^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \frac{1}{2}(\delta \wedge \sqrt{\delta}) \text{ for some } n \geq m \right\}. \end{aligned}$$

Limiting $h \rightarrow \infty$, $l \rightarrow \infty$, $m \rightarrow \infty$ and taking into account Lemma 3.3 we come to (3.16). Hence Lemma 3.5.

Lemma 3.6 *Let x_k and $\tau(h)$ be defined by (1.1) and (1.5). Then for any compact set $K \subset \Lambda_0$ and $\delta > 0$*

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} P_\theta \left(\frac{x_{\tau-1}^2}{\sum_{k=1}^{\tau-1} x_{k-1}^2} > \delta \right) = 0. \tag{3.22}$$

Proof. One has the inclusion

$$\left\{ \frac{x_{\tau-1}^2}{\sum_{k=1}^{\tau-1} x_{k-1}^2} > \delta \right\} \subset \{\tau(h) \leq m\} \cup \left\{ \frac{x_n^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta \text{ for some } n \geq m \right\}.$$

From here and (3.20) it follows that

$$\begin{aligned} \sup_{\Theta \in K} P_{\Theta} \left\{ \frac{x_{\tau-1}^2}{\sum_{k=1}^{\tau-1} x_{k-1}^2} > \delta \right\} &\leq I_{(ml > h)} + \sum_{j=1}^m \sup_{\Theta \in K} P_{\Theta} \{x_{j-1}^2 + x_{j-2}^2 \geq l\} \\ &+ \sup_{\Theta \in K} P_{\Theta} \left\{ \frac{x_n^2}{\sum_{k=1}^{n-1} x_{k-1}^2} > \delta \text{ for some } n \geq m \right\}. \end{aligned}$$

Limiting $h \rightarrow \infty$, $l \rightarrow \infty$, $m \rightarrow \infty$ and applying Lemma 3.2 lead to (3.22). This completes the proof of Lemma 3.6.

4 Appendix.

In this Section we prove some results used in the paper.

1. *Proof of Lemma 2.1.* Denote

$$G_{\tau(h)} = L^{-1/2}(\theta_1, \theta_2) M_{\tau(h)} L^{-1/2}(\theta_1, \theta_2).$$

One can easily verify that

$$\begin{aligned} \left\| L^{-1/2}(\theta_1, \theta_2) \frac{M_{\tau(h)}^{1/2}}{\sqrt{h}} - I_2 \right\|^2 &= \left\| \frac{1}{\sqrt{h}} G_{\tau(h)}^{1/2} - I_2 \right\|^2 \\ &\leq \left\| h^{-1} G_{\tau(h)} - I_2 \right\|^2 \leq \text{tr} L^{-1}(\theta_1, \theta_2) \left\| \frac{M_{\tau(h)}}{h} - L(\theta_1, \theta_2) \right\|^2. \end{aligned}$$

From here by Lemma 3.5 we come to the desired result.

Hence Lemma 2.1.

2. *Proof of Lemma 2.2.* We have

$$v'Y_h = \frac{1}{\sqrt{h}} \sum_{k=1}^{\tau(h)} g_{k-1} \varepsilon_k, \quad g_{k-1} = v' L^{-1/2}(\theta_1, \theta_2) X_{k-1}.$$

For each $h > 0$ we define the stopping time as

$$\tau_0 = \tau_0(h) = \inf\{n \geq 1 : \sum_{k=1}^n g_{k-1}^2 \geq h\}, \quad \inf\{\emptyset\} = +\infty.$$

Further we use the representation

$$v'Y_h = \frac{1}{\sqrt{h}} \sum_{k=1}^{\tau_0(h)} g_{k-1} \varepsilon_k + \eta(h) + \Delta(h),$$

where $\Delta(h) = \Delta_1(h) + \dots + \Delta_4(h)$,

$$\Delta_1(h) = h^{-1/2} I_{(\tau(h)=1)} g_0 \varepsilon_1, \quad \Delta_2(h) = h^{-1/2} g_{\tau(h)-1} \varepsilon_{\tau(h)},$$

$$\Delta_3(h) = -h^{-1/2} I_{(\tau_0(h)=1)} g_0 \varepsilon_1, \quad \Delta_4(h) = -h^{-1/2} g_{\tau_0(h)-1} \varepsilon_{\tau_0(h)},$$

$$\eta(h) = \frac{1}{\sqrt{h}} \sum_{k=1}^{\tau(h)-1} g_{k-1} \varepsilon_k - \frac{1}{\sqrt{h}} \sum_{k=1}^{\tau_0(h)-1} I_{(\tau_0 > 1)} g_{k-1} \varepsilon_k.$$

Now we show that

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} \sup_{t \in R} |\mathbf{P}_\theta \left(\frac{1}{\sqrt{h}} \sum_{k=1}^{\tau_0(h)} g_{k-1} \varepsilon_k \leq t \right) - \Phi(t)| = 0, \quad (4.1)$$

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta (|\eta(h)| > \delta) = 0, \quad (4.2)$$

$$\lim_{h \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta (|\Delta(h)| > \delta) = 0. \quad (4.3)$$

The proof of (4.1) is based on Proposition 3.1 in Lai and Siegmund (1983). To this end we have to verify only the condition A_δ , that is for each $\delta > 0$

$$\lim_{m \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta \left(g_n^2 \geq \delta \sum_{k=1}^n g_{k-1}^2 \text{ for some } n \geq m \right) = 0. \quad (4.4)$$

Conditions $A_1 - A_5$ are evidently satisfied.

First we note that

$$\sum_{k=1}^n g_{k-1}^2 = \left(v' L^{-1/2} \left(\frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L \right) L^{-1/2} v + 1 \right) \sum_{k=1}^n x_{k-1}^2.$$

Therefore we have

$$\begin{aligned} & \left\{ g_n^2 \geq \delta \sum_{k=1}^n g_{k-1}^2 \text{ for some } n \geq m \right\} \\ & \subseteq \left\{ \|X_n\|^2 \geq \delta_1 \sum_{k=1}^n g_{k-1}^2 \text{ for some } n \geq m \right\} \end{aligned}$$

$$\begin{aligned}
&= \left\{ \|X_n\|^2 \geq \delta_1 \sum_{k=1}^n x_{k-1}^2 \left[1 + v' L^{-1/2} \left(\frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L \right) L^{-1/2} v \right] \text{ for some } n \geq m \right\} \\
&\subseteq \left\{ \|X_n\|^2 \geq \delta_1 \sum_{k=1}^n x_{k-1}^2 \left[1 - \|L^{-1/2} v\|^2 \cdot \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L \right\| \right] \text{ for some } n \geq m \right\} \\
&\subseteq \left\{ \|X_n\|^2 \geq \delta_1 \sum_{k=1}^n x_{k-1}^2 \left[1 - a^* \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L \right\| \right] \text{ for some } n \geq m \right\} \\
&\subset \left\{ \left\| \frac{M_n}{\sum_{k=1}^n x_{k-1}^2} - L \right\| \geq \frac{1}{2a^*} \text{ for some } n \geq m \right\} \\
&\quad \cup \left\{ \|X_n\|^2 > \frac{\delta_1}{2} \sum_{k=1}^n x_{k-1}^2 \text{ for some } n \geq m \right\},
\end{aligned}$$

where $\delta_1 = \delta/a^*$, $a^* = \sup_{\theta \in K} \|v' L^{-1/2}\|^2$. This in view of Lemmas 2.5, 3.1 leads to (4.1).

It will be observed that (4.4) enables one to prove (by the same argument as in Lemma 3.6) that, for any compact set $K \subset \Lambda_1$ and $\delta > 0$,

$$\limsup_{h \rightarrow \infty} \sup_{\theta \in K} P_\theta \left(g_{\tau_0-1}^2 / \sum_{k=1}^{\tau_0-1} g_{k-1}^2 > \delta \right) = 0. \quad (4.5)$$

Now we check (4.2). One can easily verify that

$$\mathbf{E}_\theta \eta^2(h) = \mathbf{E}_\theta u(h), \quad u(h) = \frac{1}{h} \left| \sum_{k=1}^{\tau(h)-1} g_{k-1}^2 - \sum_{k=1}^{\tau_0(h)-1} g_{k-1}^2 \right|.$$

The random variable $u(h)$ is uniformly bounded from above uniformly in $\theta \in K$ because

$$\begin{aligned}
u(h) &\leq \frac{1}{h} \sum_{k=1}^{\tau(h)-1} g_{k-1}^2 + 1 = \frac{1}{h} v' L^{-1/2} M_{\tau(h)-1} L^{-1/2} v + 1 \\
&\leq \frac{a^*}{h} \sum_{k=1}^{\tau(h)-1} \|X_{k-1}\|^2 + 1 \leq a^* + 1.
\end{aligned}$$

Therefore it suffices to establish that for each $\delta > 0$

$$\limsup_{h \rightarrow \infty} \sup_{\theta \in K} \mathbf{P}_\theta(u(h) > \delta) = 0. \quad (4.6)$$

To this end one can use the following estimate

$$\begin{aligned}
u(h) &= \frac{1}{h} |v' L^{-1/2} M_{\tau(h)-1} L^{-1/2} v - \sum_{k=1}^{\tau_0(h)-1} g_{k-1}^2| \\
&= \left| \frac{\sum_{k=1}^{\tau(h)-1} x_{k-1}^2}{h} v' L^{-1/2} \left(\frac{M_{\tau(h)-1}}{\sum_{k=1}^{\tau(h)-1} x_{k-1}^2} - L \right) L^{-1/2} v \right. \\
&\quad \left. + \frac{\sum_{k=1}^{\tau(h)-1} x_{k-1}^2}{h} - \frac{\sum_{k=1}^{\tau_0(h)-1} g_{k-1}^2}{h} \right| \\
&\leq a^* \left\| \frac{M_{\tau(h)-1}}{\sum_{k=1}^{\tau(h)-1} x_{k-1}^2} - L \right\| + \frac{x_{\tau(h)-1}^2}{\sum_{k=1}^{\tau(h)-1} x_{k-1}^2} + \frac{g_{\tau_0(h)-1}^2}{\sum_{k=1}^{\tau_0(h)-1} g_{k-1}^2}.
\end{aligned}$$

From here by making use of (3.16), (3.22) and (4.5) we come to (4.6) which in its turn implies (4.2).

By a similar argument one can check (4.3). This completes the proof of Lemma 2.2.

References

- [1] Ahtola, J.A. and Tiao, G.C. (1987) Distribution of least squares estimators of autoregressive parameters for a process with complex roots on the unit circle. *J. Time Ser. Anal.* **8**, p. 1-14.
- [2] Anderson, T.W. (1971) The statistical analysis of time series. *John Wiley Sons*, New York.
- [3] Anscombe, F. (1953) Sequential estimation. *J. Roy. Statist. Soc., Ser. B*, **15**, p. 1-21.
- [4] Chan, N.H. and Wei, C.Z. (1988) Limiting distribution of least squares estimates of unstable autoregressive processes. *Ann. Statist.*, **16**, p. 367-401.
- [5] Dickey, D.A. and Fuller, W.A. (1979) Distribution of the estimators for autoregressive time series with a unit root. *J. Amer. Statist. Assoc.*, **74**, p. 427-431.
- [6] Galtchouk, L. and Konev V. (2003_a) On uniform asymptotic normality of sequential least squares estimators for the parameters in a stable AR(1). *Sequential Anal.* **22**, p. 31-54.
- [7] Galtchouk, L. and Konev V. (2003_b) On uniform asymptotic normality of sequential least squares estimators for the parameters in a stable AR(p). *J. Multivariate Anal.* **?**, p. ?.
- [8] Greenwood, P.E. and Shiryaev, A.N. (1992) Asymptotic minimaxity of a sequential estimator for a first order autoregressive model. *Stochastic and Stochastics Rep.*, **38**, 49-65.
- [9] Konev, V.V. and Pergamenshchikov, S.M. (1996) Guaranteed parameter estimation on the basis of a generalized least-squares method. *Theory Probab. Appl.* **41**, p. 765-784.
- [10] Lai, T.L. and Siegmund, D. (1983) Fixed-accuracy estimation of an autoregressive parameter. *Ann.Statist.* **11**, p. 478-485.
- [11] Lai, T.L. and Wei, C.Z. (1983) Asymptotic properties of general autoregressive models and strong consistency of least squares estimates of their parameters. *J. Multivariate Anal.* **13**, p. 1-23.
- [12] Nyblon, J. (1985) On the least squares estimators for the unstable AR(2) models. *Contributed Papers, 45th Session, ISI*, p. 385-386.

- [13] Rao, M.M. (1978) Asymptotic distribution of an estimator of the boundary parameter of an unstable process. *Ann.Statist.* **6**, p. 185-190.
- [14] Shiryaev, A.N. and Spokoiny, V.G. (1997) On sequential estimation of an autoregressive parameter. *Stochastic and Stochastics Rep.*, **60**, p. 219-240.
- [15] Sriram, T.N. (1987) Sequential estimation of the mean of a first order stationary autoregressive process. *Ann.Statist.*, **15**, p. 1079-1090.
- [16] Sriram, T.N. (1988) Sequential estimation of the autoregressive parameter in a first order autoregressive process. *Sequential Anal.*, **7**, p. 53-74.
- [17] Stein, C. (1945) A two-sample test for a linear hypothesis whose power is independent of the variance. *Ann. Math. Statist.* **16**, p. 243-258.
- [18] White, J.S. (1958) The limiting distribution of the serial correlation coefficient in the explosive case. *Ann. Math. Statist.* **29**, p. 1188-1197.