



**HAL**  
open science

**A new proof of the Borel-de Siebenthal Theorem, the classification of equi-rank groups, and some compact and semi-compact dual pairs.**

Hubert Rubenthaler

► **To cite this version:**

Hubert Rubenthaler. A new proof of the Borel-de Siebenthal Theorem, the classification of equi-rank groups, and some compact and semi-compact dual pairs.. 1998. hal-00129720

**HAL Id: hal-00129720**

**<https://hal.science/hal-00129720>**

Preprint submitted on 8 Feb 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A new proof of the Borel-de Siebenthal Theorem, the classification of equi-rank groups, and some compact and semi-compact dual pairs.

HUBERT RUBENTHALER

## 1. Introduction

Let  $G$  be a connected simple real Lie group with finite center and let  $K$  be a maximal compact subgroup of  $G$ . By a well known result of Harish-Chandra ([HC]) the group  $G$  has a discrete series of representations if and only if the rank of  $K$  is equal to the rank of  $G$ .

The list of such  $G$ 's, which we will call *equi-rank groups* was known since a long time, mostly by case by case computation from the list of real simple Lie algebras (see for example [0-V], [War], [He]).

But there exists an intrinsic classification which is essentially due to Borel and de Siebenthal [B-dS] (more precisely what is now called the Borel-de Siebenthal Theorem is Remarque 1, p 218 of [B-dS]). Their theorem however was given in terms of compact groups and the transcription in terms of equi-rank groups was given by Murakami who called these equi-rank groups *groups of interior type* ([M], Théorème 1 p. 295) (see also the paper by Wallach [Wal]).

Recently A. Knapp [K1], gave a quick proof of the classification of real semi-simple Lie algebras which relies also on the Borel-de Siebenthal Theorem. In his book ([K2]), A. Knapp gave a new proof of this theorem (Thm 6.96. p.350) which uses together the Lie algebra  $\mathfrak{g}_{\mathbf{R}}$  of  $G$  and its complexification  $\mathfrak{g}$ , and the complex, imaginary, compact, non-compact roots.

We propose here a third proof of the Borel-de Siebenthal Theorem, or to be more precise, a new classification of equi-rank groups, which is totally complex (i.e. we work only at the  $\mathfrak{g}$  level) and is based on the work of Dynkin on subalgebras of semi-simple Lie algebras ([D], see [T] for a nice résumé).

Our proof underlines what one could call the *parabolic* and *Z-graded* aspects of the involved structures which were never, as far as we know, noticed before.

Moreover our results give some extra information about the different "parabolic realizations" of the complexified Lie algebra  $\mathfrak{k}$  of  $K$  (paragraph 5) and lead naturally to the description of some families of dual pairs  $(\mathfrak{k}_1, \mathfrak{k}_2)$  inside  $\mathfrak{g}_{\mathbf{R}}$ , where at least one of these subalgebras is compact (paragraph 6).

**Acknowledgements.** The author would like to thank Nicole Bopp and Robert J. Stanton for several very useful discussions.

## 2. Maximal compact subalgebras and complex involutions.

Let  $\mathfrak{g}_{\mathbf{R}} = \mathfrak{k}_{\mathbf{R}} \oplus \mathfrak{p}_{\mathbf{R}}$  be a Cartan decomposition of  $\mathfrak{g}_{\mathbf{R}}$  relatively to a Cartan involution  $\theta$ . Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  be the decomposition of  $\mathfrak{g} = \mathfrak{g}_{\mathbf{R}} \otimes \mathbf{C}$  obtained by complexification. We will denote by  $\sigma$  the complex linear extension of  $\theta$ . Conversely let us show the following ( certainly well known) result :

**Proposition 2.1.** — *If one has a decomposition  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  where  $\mathfrak{k}$  (resp.  $\mathfrak{p}$ ) is the  $+1$  (resp.  $-1$ ) eigenspace of a complex involution  $\sigma$  of  $\mathfrak{g}$ , then there exists a real form  $\mathfrak{g}_{\mathbf{R}}$  of  $\mathfrak{g}$  whose maximal compact subalgebra  $\mathfrak{k}_{\mathbf{R}}$  is a real form of  $\mathfrak{k}$ . Moreover all such real forms are conjugate under the group  $\exp \mathfrak{k}$ .*

Proof. First we need the following lemma.

**Lemma 2.2.** — *Let  $\theta_1$  and  $\theta_2$  be two Cartan involutions of the complex Lie algebra  $\mathfrak{g}$  which commute with a given complex linear involution  $\sigma$  whose fixed point set is denoted by  $\mathfrak{k}$ . Then there exists an element  $X \in \mathfrak{k}$  such that  $\theta_2 = (\exp X)\theta_1(\exp -X)$ .*

Let us prove the lemma. Put  $N = \theta_2\theta_1$ . One shows easily that if  $B_{\theta_1}$  is defined by  $B_{\theta_1}(X, Y) = -B(X, \theta_1(Y))$ , then the linear operator  $N$  is symmetric with respect to  $B_{\theta_1}$ . But  $B_{\theta_1}$  is hermitian positive definite, therefore  $N$  is diagonalizable with real eigenvalues  $\lambda_i$ . Hence, for any  $t \in \mathbf{R}$ , one can define an operator  $P^t$  whose eigenvalues are  $(\lambda_i^2)^t$ . In particular  $P = N^2$ . Of course  $P^t$  is a one parameter subgroup inside  $Aut(\mathfrak{g})$ , therefore there exists  $X \in \mathfrak{g}$  such that  $P^t = \exp(t \operatorname{ad} X)$ . Moreover this one parameter subgroup commutes with  $\sigma$  (because it is the case for  $N$ ) :  $\sigma P^t \sigma^{-1} = \exp(t \operatorname{ad} \sigma(X)) = P^t = \exp(t \operatorname{ad}(X))$  for all  $t \in \mathbf{R}$ . Hence  $\operatorname{ad} X = \operatorname{ad} \sigma(X)$  and one gets  $X = \sigma(X)$ . Therefore  $X$  is in  $\mathfrak{k}$ .

Set  $\theta' = P^{\frac{1}{4}}\theta_1 P^{-\frac{1}{4}}$ , this again is a Cartan involution of  $\mathfrak{g}$ . Let us show that  $\theta'$  and  $\theta_2$  commute. Notice first that  $\theta_1 N \theta_1^{-1} = \theta_1 \theta_2 \theta_1 \theta_1^{-1} = \theta_1 \theta_2 = N^{-1}$ , hence  $\theta_1 P^t \theta_1^{-1} = P^{-t}$ . Then we have

$$\theta' \theta_2 = P^{\frac{1}{4}} \theta_1 P^{-\frac{1}{4}} \theta_1 \theta_2 = P^{\frac{1}{4}} P^{\frac{1}{4}} \theta_1 \theta_2 = P^{\frac{1}{2}} N^{-1},$$

on the other hand we have

$$\theta_2 \theta' = \theta_2 \theta_1 \theta_1 P^{\frac{1}{4}} \theta_1 P^{-\frac{1}{4}} = N P^{-\frac{1}{2}} = P^{\frac{1}{2}} N^{-1} \quad (\text{because } P = N^2).$$

So we get  $\theta' \theta_2 = \theta_2 \theta'$ . But two Cartan involutions which commute are equal, therefore  $\theta_2 = P^{\frac{1}{4}} \theta_1 P^{-\frac{1}{4}} = \exp(\operatorname{ad} \frac{1}{4} X) \theta_1 \exp(-\operatorname{ad} \frac{1}{4} X)$ , with  $X \in \mathfrak{k}$ .  $\square$

A second lemma is needed.

**Lemma 2.3.** — *Take the same notations as in the preceding Lemma. Let  $\mathfrak{g} = \mathfrak{u}_1 \oplus i\mathfrak{u}_1 = \mathfrak{u}_2 \oplus i\mathfrak{u}_2$  be the Cartan decompositions of  $\mathfrak{g}$  relatively to the Cartan involutions  $\theta_1$  and  $\theta_2$ . Let  $\mathfrak{g}_{\mathbb{R}}^1$  (resp.  $\mathfrak{g}_{\mathbb{R}}^2$ ) be the fixed point algebra of  $\sigma\theta_1$  (resp.  $\sigma\theta_2$ ). Then  $\mathfrak{g}_{\mathbb{R}}^1$  and  $\mathfrak{g}_{\mathbb{R}}^2$  are real forms of  $\mathfrak{g}$  and there exists  $k \in \exp(\text{ad}\mathfrak{k})$  such that  $\mathfrak{g}_2 = k\mathfrak{g}_1$ .*

Proof of the Lemma. As  $\sigma$  commutes with  $\theta_1$  and  $\theta_2$  one has the decompositions  $\mathfrak{g}_{\mathbb{R}}^1 = \mathfrak{k} \cap \mathfrak{u}_1 \oplus \mathfrak{p} \cap i\mathfrak{u}_1$  and  $\mathfrak{g}_{\mathbb{R}}^2 = \mathfrak{k} \cap \mathfrak{u}_2 \oplus \mathfrak{p} \cap i\mathfrak{u}_2$ . The fact that  $\mathfrak{g}_{\mathbb{R}}^1$  and  $\mathfrak{g}_{\mathbb{R}}^2$  are real forms is now obvious. From the preceding Lemma one has  $\theta_2 = k\theta_1k^{-1}$  where  $k = \exp(\text{ad}\frac{1}{4}X)$  ( $X \in \mathfrak{k}$ ). This implies that  $ku_1 = u_2$ . As  $k\mathfrak{k} = \mathfrak{k}$  and  $k\mathfrak{p} = \mathfrak{p}$ , one gets  $k\mathfrak{g}_{\mathbb{R}}^1 = \mathfrak{g}_{\mathbb{R}}^2$ .  $\square$

Let us now prove Proposition 2.1. There always exists a Cartan involution  $\theta$  of  $\mathfrak{g}$  which commutes with  $\sigma$ . Let  $\mathfrak{g} = \mathfrak{u} \oplus i\mathfrak{u}$  be the corresponding Cartan decomposition and  $\mathfrak{g}_{\mathbb{R}}$  be the fixed point algebra of the involution  $\sigma\theta$ . One has  $\mathfrak{g}_{\mathbb{R}} = \mathfrak{k} \cap \mathfrak{u} \oplus \mathfrak{p} \cap i\mathfrak{u}$  and this is a Cartan decomposition of  $\mathfrak{g}_{\mathbb{R}}$ . The Lie algebra  $\mathfrak{g}_{\mathbb{R}}$  is a real form of  $\mathfrak{g}$  and  $\mathfrak{k}_{\mathbb{R}} = \mathfrak{k} \cap \mathfrak{u}$  is a real form of  $\mathfrak{k}$ . This proves the first assertion of the proposition.

Let now  $\mathfrak{g}'_{\mathbb{R}}$  be another real form of  $\mathfrak{g}$  with Cartan decomposition  $\mathfrak{g}'_{\mathbb{R}} = \mathfrak{k}'_{\mathbb{R}} \oplus \mathfrak{p}'_{\mathbb{R}}$  such that  $\mathfrak{k} = \mathfrak{k}'_{\mathbb{R}} \oplus i\mathfrak{k}'_{\mathbb{R}}$ . Then automatically  $\mathfrak{p} = \mathfrak{p}'_{\mathbb{R}} \oplus i\mathfrak{p}'_{\mathbb{R}}$ . Set  $\mathfrak{u}' = \mathfrak{k}'_{\mathbb{R}} \oplus i\mathfrak{p}'_{\mathbb{R}}$  then  $\mathfrak{g} = \mathfrak{u}' \oplus i\mathfrak{u}'$  is a Cartan decomposition and the corresponding Cartan involution  $\theta'$  is easily verified to commute with  $\sigma$ . By the preceding lemma there exists  $k \in \exp(\text{ad}\mathfrak{k})$  such that  $\mathfrak{g}'_{\mathbb{R}} = k\mathfrak{g}_{\mathbb{R}}$ .  $\square$

### 3. The classification of equi-rank complex simple Lie algebras.

Proposition 2.1. reduces the problem of classifying all equi-rank groups to the classification of all  $\mathbb{Z}_2$ -graduations  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  ( $[\mathfrak{k}, \mathfrak{k}] \subseteq \mathfrak{k}$ ,  $[\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{k}$ ,  $[\mathfrak{k}, \mathfrak{p}] \subseteq \mathfrak{p}$ ) of the complex simple Lie algebra  $\mathfrak{g}$  having the property  $\text{rank}(\mathfrak{g}) = \text{rank}(\mathfrak{k})$ , up to conjugacy by an element of the adjoint group of  $\mathfrak{g}$ . The required Lie algebras are simple because a simple equi-rank group is never a complex group.

From now on  $\mathfrak{g}$  will always denote a simple complex Lie algebra.

The data consisting of a simple complex Lie algebra together with a  $\mathbb{Z}_2$ -graduation having the above mentioned properties is called an *equi-rank algebra*.

Our main result is the as follows.

**Theorem 3.1.** — *Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  ( $\text{rank}(\mathfrak{g}) = \text{rank}(\mathfrak{k})$ ) be an equi-rank algebra, with  $\mathfrak{k} \neq \mathfrak{g}$ . Then either*

1)  $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$ , the parabolic subalgebra  $\mathfrak{g}_0 \oplus \mathfrak{g}_1$  is maximal, and  $\mathfrak{k} = \mathfrak{g}_0$ ,  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$  is the corresponding  $\mathbb{Z}_2$ -graduation. This corresponds exactly to the case where  $\mathfrak{k}$  is reductive not semi-simple, i.e. has a no non trivial center necessarily one dimensional. This is also exactly the case where  $G/K$  is hermitian.

or 2)  $\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$ , the parabolic subalgebra  $\mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$  is maximal, and then  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2$  and  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$  is the corresponding  $\mathbb{Z}_2$ -graduation. This corresponds exactly to the case where  $\mathfrak{k}$  is semi-simple.

Conversely if  $\mathfrak{g}$  is given together with a graduation of one of the preceding types, then the corresponding  $\mathfrak{k}$  and  $\mathfrak{p}$  (given as is 1) or 2)) determine an equi-rank algebra.

Let us now prove this theorem. First of all we need to recall the following definition due to Dynkin ([D]).

A subalgebra  $\mathfrak{r}$  of  $\mathfrak{g}$  is called a *regular* subalgebra if there exists a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$  such that  $[\mathfrak{h}, \mathfrak{r}] \subseteq \mathfrak{r}$ .

**Lemma 3.2.** — *Let  $\mathfrak{r} = \mathfrak{z} \oplus \mathfrak{r}'$  be a reductive subalgebra of  $\mathfrak{g}$  where  $\mathfrak{z}$  is the center of  $\mathfrak{r}$  and  $\mathfrak{r}' = [\mathfrak{r}, \mathfrak{r}]$ . Suppose moreover that the rank of  $\mathfrak{r}$  is equal to the rank of  $\mathfrak{g}$ . Then  $\mathfrak{z}$  is the center of a Levi subalgebra  $\mathfrak{l} = \mathfrak{z} \oplus \mathfrak{l}'$  of  $\mathfrak{g}$  and  $\mathfrak{r}'$  is a regular subalgebra of  $\mathfrak{l}'$  of maximal rank, i.e.  $\text{rank}(\mathfrak{r}') = \text{rank}(\mathfrak{l}')$ .*

Proof.

As  $\text{rank}(\mathfrak{r}) = \text{rank}(\mathfrak{g})$  one can choose a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$  of the form

$$\mathfrak{h} = \mathfrak{z} \oplus \mathfrak{h}_{\mathfrak{r}'}$$

where  $\mathfrak{h}_{\mathfrak{r}'}$  is a Cartan subalgebra of  $\mathfrak{r}'$ . Let  $\mathfrak{l} = \mathcal{Z}_{\mathfrak{g}}(\mathfrak{z})$  be the centralizer of  $\mathfrak{z}$ . The Lie subalgebra  $\mathfrak{l}$  is a Levi subalgebra of  $\mathfrak{g}$  and therefore is of the form :

$$\mathfrak{l} = \mathcal{Z}_{\mathfrak{g}}(\mathfrak{z}) = \mathfrak{c} \oplus \mathfrak{l}'$$

where  $\mathfrak{c}$  is the center of  $\mathfrak{l}$  and  $\mathfrak{l}' = [\mathfrak{l}, \mathfrak{l}]$ . Moreover  $\mathfrak{r}' \subseteq \mathfrak{r} \subseteq \mathfrak{l}$ .

From the fact that  $\mathfrak{c}$  is the centralizer of  $\mathfrak{l}$  in  $\mathfrak{g}$ , one deduces that  $\mathfrak{z} \subseteq \mathfrak{c}$ .

Let  $B$  be the Killing form of  $\mathfrak{g}$ . It is well known that  $\mathfrak{c}^{\perp} \cap \mathfrak{l} = \mathfrak{l}'$  where  $\mathfrak{c}^{\perp}$  is the orthogonal of  $\mathfrak{c}$  with respect to  $B$ . Let  $\mathfrak{h}_{\mathfrak{l}'} = \mathfrak{c}^{\perp} \cap \mathfrak{h}$ . Then

$$\mathfrak{h} = \mathfrak{c} \oplus \mathfrak{h}_{\mathfrak{l}'}$$

One has  $\mathfrak{h}_{\mathfrak{l}'} \subseteq \mathfrak{h}_{\mathfrak{r}'}$ , and therefore  $\text{rank}(\mathfrak{l}') \leq \text{rank}(\mathfrak{r}')$ .

Let  $\Sigma(\mathfrak{g}, \mathfrak{h})$  be the root system of the pair  $(\mathfrak{g}, \mathfrak{h})$ . For  $\alpha \in \Sigma(\mathfrak{g}, \mathfrak{h})$ , let  $\mathfrak{g}^{\alpha}$  be the corresponding root space. Then  $\mathfrak{g}^{\alpha} \perp \mathfrak{h}$  and hence  $\mathfrak{g}^{\alpha} \perp \mathfrak{c}$ . Therefore, as  $\mathfrak{r}' \subseteq \mathfrak{l}$ , any root space in  $\mathfrak{r}'$  is in fact in  $\mathfrak{l}'$ , this implies that  $\mathfrak{r}' \subseteq \mathfrak{l}'$ , and consequently  $\text{rank}(\mathfrak{r}') \leq \text{rank}(\mathfrak{l}')$ .

As the inverse inequality was proved before we get  $\text{rank}(\mathfrak{r}') = \text{rank}(\mathfrak{l}')$ .

Recall that  $\mathfrak{z} \subseteq \mathfrak{c}$ . From the equality

$$\dim \mathfrak{h} = \dim \mathfrak{z} + \dim \mathfrak{h}_{\mathfrak{r}'} = \dim \mathfrak{c} + \dim \mathfrak{h}_{\mathfrak{l}'}$$

one gets  $\mathfrak{c} = \mathfrak{z}$ . The lemma is proved.  $\square$

Let us return to the proof of Theorem 3.1.

**First case :**  $\mathfrak{k}$  is reductive non semi-simple.

By the preceding lemma  $\mathfrak{k} = \mathfrak{z} \oplus \mathfrak{k}'$  where  $\mathfrak{k}'$  is the derived algebra of  $\mathfrak{k}$  and where  $\mathfrak{z}$  is the center of a Levi subalgebra  $\mathfrak{l} = \mathfrak{z} \oplus \mathfrak{l}'$  and  $\mathfrak{k}' \subseteq \mathfrak{l}'$  with  $\text{rank}(\mathfrak{k}') = \text{rank}(\mathfrak{l}')$ .

Let  $\mathfrak{P}$  be a parabolic subalgebra of  $\mathfrak{g}$  with Levi factor  $\mathfrak{l}$ . One has  $\mathfrak{P} = \mathfrak{l} \oplus \mathfrak{n}$ , where  $\mathfrak{n}$  is the nilradical of  $\mathfrak{P}$ . It is well known ([Bou2]) that one can choose a system of simple roots  $\Psi \subseteq R = \sum(\mathfrak{g}, \mathfrak{h})$ , and a subsystem  $\Gamma \subseteq \Psi$ , such that the set of roots  $\langle \Gamma \rangle$  which are linear combinations of elements of  $\Gamma$  is the root system of  $(\mathfrak{l}, \mathfrak{h})$ , and such that

$$\mathfrak{n} = \sum_{\alpha \in R^+ \setminus \langle \Gamma \rangle^+} \mathfrak{g}^\alpha$$

where the sets of positive roots  $R^+$  and  $\langle \Gamma \rangle^+$  are defined by  $\Psi$  and  $\Gamma$  respectively.

Let  $h_\Gamma$  be the unique element of  $\mathfrak{z}$  defined by the equations

$$\begin{aligned} \alpha(h_\Gamma) &= 1 & \text{if } \alpha \in \Psi \setminus \Gamma \\ \alpha(h_\Gamma) &= 0 & \text{if } \alpha \in \Gamma \end{aligned} \tag{3-1}$$

and put, for  $i \in \mathbf{Z}$ ,  $\mathfrak{g}_i = \{X \in \mathfrak{g} \mid [h_\Gamma, X] = iX\}$ . Then one has the  $\mathbf{Z}$ -graduation  $\mathfrak{g} = \bigoplus_{i \in \mathbf{Z}} \mathfrak{g}_i$ , and moreover  $\mathfrak{l} = \mathfrak{g}_0$  and  $\mathfrak{n} = \bigoplus_{i > 0} \mathfrak{g}_i$ .

Let  $n$  be the greatest integer such that  $\mathfrak{g}_n \neq \{0\}$ .

**Lemma 3.3.** — *(Notations as before) Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  be a simple equi-rank algebra with  $\mathfrak{k}$  reductive non-semi-simple. Then  $n = 1$  (i.e. the nilradical  $\mathfrak{n}$  is commutative) and  $\dim \mathfrak{z} = 1$ .*

Proof of the Lemma.

The assertion  $\dim \mathfrak{z} = 1$  is an easy consequence of  $n = 1$ .

Suppose that  $n \geq 2$ . It is well known that then  $[\mathfrak{g}_1, \mathfrak{g}_1] \neq \{0\}$  (in fact  $[\mathfrak{g}_1, \mathfrak{g}_1] = \mathfrak{g}_2$ ). As  $\mathfrak{p}^\perp = \mathfrak{k}$  and  $\mathfrak{l}^\perp = \mathfrak{g}_{-n} \oplus \cdots \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_n \subseteq \mathfrak{p}$ , one should have  $[\mathfrak{g}_1, \mathfrak{g}_1] \subseteq [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{k} \subseteq \mathfrak{l} = \mathfrak{g}_0$ . This is not true if  $\mathfrak{g}_2 \neq \{0\}$ . Therefore  $n = 1$ .  $\square$

Note that the condition  $n = 1$  we just proved, implies that the parabolic subalgebra  $\mathfrak{P} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$  is maximal (because  $\mathfrak{g}$  is simple).

In order to prove Theorem 3.1. in the case where  $\mathfrak{k}$  is reductive non semi-simple, it is now sufficient to prove that  $\mathfrak{k} = \mathfrak{g}_0$ .

Let  $\mathfrak{l} = \mathfrak{k} \oplus \mathfrak{p}_1$  where  $\mathfrak{p}_1 = \mathfrak{l} \cap \mathfrak{p}$ . Recall that then  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}_1 \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$ .

Put  $\mathfrak{a} = \mathfrak{g}_{-1} \oplus [\mathfrak{g}_{-1}, \mathfrak{g}_1] \oplus \mathfrak{g}_1$ . It is easily seen that  $\mathfrak{a}$  is an ideal of  $\mathfrak{g}$ . As  $\mathfrak{g}$  is simple, one has  $\mathfrak{g} = \mathfrak{a}$ .

Consider now the bracket space  $[\mathfrak{p}_1, \mathfrak{g}_1]$ . As  $\mathfrak{p}_1 \subseteq \mathfrak{p}$  and  $\mathfrak{g}_1 \subseteq \mathfrak{p}$ , one has  $[\mathfrak{p}_1, \mathfrak{g}_1] \subseteq \mathfrak{k}$ . But  $\mathfrak{p}_1 \subseteq \mathfrak{l} = \mathfrak{g}_0$ , therefore  $[\mathfrak{p}_1, \mathfrak{g}_1] \subseteq \mathfrak{g}_1 \subseteq \mathfrak{p}$ . Hence  $[\mathfrak{p}_1, \mathfrak{g}_1] = \{0\}$ . The same arguments shows that  $[\mathfrak{p}_1, \mathfrak{g}_{-1}] = \{0\}$ , and hence  $[\mathfrak{p}_1, [\mathfrak{p}_1, \mathfrak{g}_1]] = \{0\}$ , and finally  $\mathfrak{p}_1$  is central in  $\mathfrak{g}$ , therefore  $\mathfrak{p}_1 = \{0\}$  and  $\mathfrak{k} = \mathfrak{g}_0$  and  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$ .

Conversely suppose that one has a  $\mathbf{Z}$ -graduation

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1.$$

Then the decomposition  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  where  $\mathfrak{k} = \mathfrak{g}_0$  and  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$  determines an equi-rank algebra.

Theorem 3.1. is proved in the first case.

**Second case :**  $\mathfrak{k}$  is semi-simple.

Lets us recall what an *elementary operation* in the sense of Dynkin is.

Let  $D$  be the Dynkin diagram of a simple Lie algebra  $\mathfrak{g}$ . Once a Cartan subalgebra  $\mathfrak{h}$  is chosen, as well as a set of simple roots  $\Psi$ , the vertices of  $D$  are associated to the elements of  $\Psi$ , and the different vertices are connected by edges according to well known rules. Let  $\omega$  be the highest root with respect to  $\Psi$ . Then one gets the so-called *extended* Dynkin diagram  $\tilde{D}$  by adding a vertex associated to  $-\omega$  and connecting this new vertex to the other vertices by using the same rules.

It is a matter of fact that  $\tilde{D}$  is *not* a Dynkin Diagram, but if one removes any vertex from it, as well as the edges connected to this vertex, then one gets a, possibly non connected, new Dynkin diagram  $D_1$  (with of course the same number of vertices as  $D$ ).

This new Dynkin diagram  $D_1$  is said to be obtained from  $D$  by an elementary operation.

Let now  $(X_{-\alpha}, H_\alpha, X_\alpha)_{\alpha \in D_1}$  be the collection of the usual  $\mathfrak{sl}_2$ -triples in  $\mathfrak{g}$  associated to the roots corresponding to the vertices in  $D_1$ .

This collection of triples generates a semi-simple Lie subalgebra  $\mathfrak{a}_1$  of  $\mathfrak{g}$  whose Dynkin diagram is precisely  $D_1$  ( $[D], [T]$ ). The Lie algebra  $\mathfrak{a}_1$  is said to be obtained from  $\mathfrak{g}$  by an elementary operation.

Another way to understand this algebra  $\mathfrak{a}_1$  is as follows. Let  $\alpha_0$  be the root which is removed by the elementary operation, and let  $\Gamma = \Psi \setminus \{\alpha_0\}$ . This subset  $\Gamma$  defines, as in the proof of the first case, a (maximal) parabolic subalgebra and therefore a  $\mathbf{Z}$ -graduation

$$\mathfrak{g} = \bigoplus_{i=-k}^{i=k} \mathfrak{g}_i. \quad (3-2)$$

One can prove that

$$\mathfrak{a}_1 = \mathfrak{g}_{-k} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_k$$

(see [Bou1] Exercice 4, Chap. VI, par. 4, p. 229 and [Bou2] Exercice 2, Chap VIII par. 3, p. 223).

But now, one can make a new elementary operation on any of the connected components of  $D_1$ . Of course each time that a diagram of type  $A_n$  occurs, any elementary operation does not change the algebra.

Any semi-simple Lie algebra of maximal rank of a simple Lie algebra is obtained by a finite number of elementary operations (see [D], Th. 5.3. page 145 or [T]).

Therefore  $\mathfrak{k}$  is obtained in such a way : one has a chain of subalgebras

$$\mathfrak{g} = \mathfrak{a}_0 \supseteq \mathfrak{a}_1 \supseteq \cdots \supseteq \mathfrak{a}_m = \mathfrak{k},$$

where each  $\mathfrak{a}_i$  is obtained from  $\mathfrak{a}_{i-1}$  by an elementary operation.

Let us focus on  $\mathfrak{a}_1$ . As we explained before it exists a  $\mathbf{Z}$ -graduation of  $\mathfrak{g}$  as in (3-2) coming from a maximal parabolic subalgebra such that

$$\mathfrak{a}_1 = \mathfrak{g}_{-k} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_k.$$

As  $\mathfrak{k} \subseteq \mathfrak{a}_1$  one has

$$\mathfrak{p} = \mathfrak{k}^\perp \supseteq \mathfrak{g}_{-k+1} \oplus \cdots \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_{k-1}.$$

If  $k \geq 3$ , that is if  $k-1 \geq 2$ , one should have  $[\mathfrak{g}_1, \mathfrak{g}_1] \subseteq \mathfrak{g}_2 \subseteq \mathfrak{p}$  and also  $[\mathfrak{g}_1, \mathfrak{g}_1] \subseteq \mathfrak{k} \subseteq \mathfrak{a}_1 = \mathfrak{g}_{-k} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_k$ . This is impossible since  $[\mathfrak{g}_1, \mathfrak{g}_1] \neq \{0\}$ .

Hence  $k \leq 2$ . This means that in fact  $k = 2$ , because the case  $k = 1$  would lead to  $\mathfrak{a}_1 = \mathfrak{g}$  which is excluded since  $\mathfrak{k} \neq \mathfrak{g}$ .

Now we are in the following situation :



$$\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2,$$

$$\mathfrak{k} \subseteq \mathfrak{a}_1 = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2, \quad \mathfrak{g}_{-1} \oplus \mathfrak{g}_1 \subseteq \mathfrak{p}.$$

As  $\mathfrak{g}$  is simple one has  $[\mathfrak{g}_{-1}, \mathfrak{g}_1] = \mathfrak{g}_0$  and hence  $\mathfrak{g}_0 \subseteq \mathfrak{k}$ .

Moreover the graduation is defined by the eigenvalues of an element  $h_\Gamma$  as in (3-1) where  $\Gamma = \Psi \setminus \{\alpha_0\}$  for some simple root  $\alpha_0$  (the parabolic subalgebra  $\mathfrak{P} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$  is maximal).

This implies that

$$\mathfrak{k} = \mathfrak{g}_2 \cap \mathfrak{k} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2 \cap \mathfrak{k} \tag{3-3}$$

The following lemma is needed.

**Lemma 3.4.** — *Let  $\mathfrak{P}$  be a maximal parabolic subalgebra of  $\mathfrak{g}$ . Let  $\mathfrak{g} = \mathfrak{g}_{-n} \oplus \cdots \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_n$  be the associated  $\mathbf{Z}$ -graduation (as in (3-2)). Then for any  $i = 1, \dots, n$  the space  $\mathfrak{g}_i$  is an irreducible  $\mathfrak{g}_0$ -module.*

Proof of the Lemma.

It is well known that the result is true for  $i = 1$  (see for example [Ru1]). As  $\mathfrak{P}$  is maximal, the center of  $\mathfrak{g}_0$  is one dimensional. Let us consider  $\mathfrak{g}' = \bigoplus_{p \in \mathbf{Z}} \mathfrak{g}_{pi}$ . This algebra is certainly reductive. If it has a non trivial center, then this center is the one dimensional center of  $\mathfrak{g}_0$ , which does not act trivially on the  $\mathfrak{g}_i$ 's. Therefore  $\mathfrak{g}'$  is semi-simple.

On the other hand  $\mathfrak{P}' = \bigoplus_{p > 0} \mathfrak{g}_{pi}$  is a parabolic subalgebra of  $\mathfrak{g}'$  which is maximal again because the center of the Levi factor  $\mathfrak{g}_0$  is one dimensional.

Then by the result cited at the beginning of the proof, the ad-action of  $\mathfrak{g}_0$  on the first step (here  $\mathfrak{g}_i$ ) is irreducible.  $\square$

The Lemma implies that the ad-representation of  $\mathfrak{g}_0$  on  $\mathfrak{g}_{-2}$  and  $\mathfrak{g}_2$  is irreducible. Therefore one has

$$\begin{aligned} \mathfrak{g}_2 \cap \mathfrak{k} &= \{0\} \quad \text{or} \quad \mathfrak{g}_2 \\ \mathfrak{g}_{-2} \cap \mathfrak{k} &= \{0\} \quad \text{or} \quad \mathfrak{g}_{-2} \end{aligned}$$

The case  $\mathfrak{g}_{-2} \cap \mathfrak{k} = \mathfrak{g}_2 \cap \mathfrak{k} = \{0\}$  is impossible because (3-3) would imply that  $\mathfrak{k} = \mathfrak{g}_0$  and here  $\mathfrak{k}$  is semi-simple.

The case  $\mathfrak{g}_2 \cap \mathfrak{k} = \{0\}$  and  $\mathfrak{g}_{-2} \cap \mathfrak{k} = \mathfrak{g}_{-2}$  (as well as the symmetric case) is also impossible because it would imply that  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0$  and hence non semi-simple.

Finally  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2$  and hence  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$ .

Conversely suppose that one has a  $\mathbb{Z}$ -graduation of the following type :

$$\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2,$$


then it is easy to see that the decomposition  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  where  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2$  and  $\mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1$  corresponds to an equi-rank subalgebra. The proof of Theorem 3.1. is completed.  $\square$

**Definition 3.5.** — *Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  be an equi-rank algebra. Then any  $\mathbb{Z}$ -graduation  $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$  (if  $\mathfrak{k} = \mathfrak{g}_0$  is reductive non-semi-simple) or  $\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$  (if  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2$  is semi-simple) as described in Theorem 3.1. is called a parabolic realization of the equi-rank algebra. (This implies in particular that the parabolic subalgebras  $\mathfrak{g}_0 \oplus \mathfrak{g}_1$  or  $\mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$  are maximal).*

#### 4. The list of equi-rank complex simple Lie algebras.

Let us first recall how parabolic subalgebras of a simple  $\mathfrak{g}$  may be described by weighted Dynkin diagrams. Each parabolic subalgebra is uniquely defined (up to conjugacy) by a subset  $\Gamma$  of the set  $\Psi$  of simple roots. We make the convention that such a data is described by the Dynkin diagram of  $\mathfrak{g}$  where the roots in  $\Psi \setminus \Gamma$  are circled.


For example consider the following diagram :




This diagram describes a parabolic subalgebra of  $\mathfrak{g} \simeq D_9$  where the Levi factor is of type  $A_1 \times A_1 \times D_4 \times \mathbb{C}^3$ .

We will now give a diagrammatical description of an elementary operation. The vertex corresponding to  $-\omega$  where  $\omega$  is the highest root will be denoted by  $\nabla$ , and the vertex which corresponds to the removed root will be marked by a cross.

For example the diagram



describes an elementary operation in  $E_7$  where the resulting algebra is of type  $A_7$ . Remember from the previous paragraph that this  $A_7$ -algebra is also obtained by considering the parabolic subalgebra associated to the diagram



More precisely as the circled root has coefficient 2 in the highest root the corresponding parabolic realization is of type

$$E_7 = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$$

and

$$A_7 = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2.$$

As we explained in the previous paragraph this gives  $E_7$  the structure of an equi-rank algebra where  $\mathfrak{k} = A_7$ .

Theorem 3.1. now gives a very easy way to classify all equi-rank complex Lie algebras. They split into two distinct families.

The first family is obtained by taking for  $\mathfrak{k}$  the Levi factor of a maximal parabolic subalgebra with an abelian nilradical. This family corresponds to weighted Dynkin diagrams where the unique circled root has coefficient 1 in the highest root. This is the hermitian symmetric case.

The second family corresponds to elementary operations where the removed root (marked by a cross on the extended Dynkin diagram) has coefficient 2 in the highest root.

These remarks lead very easily to the following table. In this table we have already taken into account the isomorphisms of the (extended or not) Dynkin diagram.

**TABLE 1**

**First Type (Hermitian Case)**

$$(1-1) \quad \bullet \text{---} \bullet \text{---} \dots \text{---} \bullet \text{---} \circ \text{---} \bullet \text{---} \dots \text{---} \bullet \text{---} \bullet \quad A_n \quad \mathfrak{k} = A_{p-1} \times A_{n-p} \times \mathbf{C}$$

$\alpha_1 \qquad \qquad \qquad \alpha_p \qquad \qquad \qquad \alpha_n$

(1-2)		$B_n$	$\mathfrak{k} = B_{n-1} \times \mathbb{C}$
(1-3)		$C_n$	$\mathfrak{k} = A_{n-1} \times \mathbb{C}$
(1-4)		$D_n$	$\mathfrak{k} = D_{n-1} \times \mathbb{C}$
(1-5)		$D_n$	$\mathfrak{k} = A_{n-1} \times \mathbb{C}$
(1-6)		$E_6$	$\mathfrak{k} = D_5 \times \mathbb{C}$
(1-7)		$E_7$	$\mathfrak{k} = E_6 \times \mathbb{C}$

### Second Type (Non Hermitian Case)

(2-1)		$B_n$	$p \geq 2$	$\mathfrak{k} = D_p \times B_{n-p}$
(2-2)		$C_n$	$p \leq n-1$	$\mathfrak{k} = C_p \times C_{n-p}$
(2-3)		$D_n$	$2 \leq p \leq n-2$	$\mathfrak{k} = D_p \times D_{n-p}$
(2-4)		$E_6$		$\mathfrak{k} = A_1 \times A_5$
(2-5)		$E_7$		$\mathfrak{k} = A_1 \times D_6$
(2-6)		$E_7$		$\mathfrak{k} = A_7$
(2-7)		$E_8$		$\mathfrak{k} = A_1 \times E_7$

$$(2-8) \quad \begin{array}{c} \bullet \\ \bullet \\ \bullet \\ \bullet \\ \bullet \\ \bullet \\ \bullet \\ \bullet \\ \nabla \end{array} \quad E_8 \quad \mathfrak{k} = D_8$$

$$(2-9) \quad \nabla \leftarrow \bullet \rightleftarrows \bullet \bullet \bullet \quad F_4 \quad \mathfrak{k} = A_1 \times C_3$$

$$(2-10) \quad \nabla \leftarrow \bullet \rightleftarrows \bullet \bullet \bullet \rightarrow \bullet \quad F_4 \quad \mathfrak{k} = B_4$$


$$(2-11) \quad \nabla \leftarrow \bullet \rightleftarrows \bullet \bullet \quad G_2 \quad \mathfrak{k} = A_1 \times A_1$$

### 5. Distinct parabolic realizations of the same equi-rank algebra.

In this paragraph we will give some examples of distinct parabolic realizations of the same equi-rank algebra.

**5.1.** This first example will show that the description by means of a maximal parabolic subalgebra and the associate  $\mathbf{Z}$ -graduation given in theorem 3.1. may be rather different for the same equi-rank algebra.

Let us consider the following three elementary operations in  $E_6$



These three diagrams are obviously conjugate under an isomorphism of the extended Dynkin diagram, which induces an isomorphism of  $E_6$ . Therefore the corresponding equi-rank algebras are conjugate. In each case  $\mathfrak{k} = A_1 \times A_5$  and the corresponding equi-rank Lie algebra is the case (2-4) of the table in the previous paragraph.

The first two diagrams are in fact conjugate under an automorphism of the Dynkin diagram itself and therefore the parabolic realizations basically only differ by a distinct numbering of the simple roots.

Let us consider the parabolic subalgebras corresponding to the first and third case :



Of course in each case the circled root has coefficient 2 in the highest root.  
 Consider now the associate parabolic realizations :

$$(a) \quad E_6 = \mathfrak{g}_{-2}^1 \oplus \mathfrak{g}_{-1}^1 \oplus \mathfrak{g}_0^1 \oplus \mathfrak{g}_1^1 \oplus \mathfrak{g}_2^1 \quad (b) \quad E_6 = \mathfrak{g}_{-2}^2 \oplus \mathfrak{g}_{-1}^2 \oplus \mathfrak{g}_0^2 \oplus \mathfrak{g}_1^2 \oplus \mathfrak{g}_2^2$$

From Theorem 3.1. one knows that

$$\mathfrak{k} \simeq A_1 \times A_5 \simeq \mathfrak{g}_{-2}^1 \oplus \mathfrak{g}_0^1 \oplus \mathfrak{g}_2^1 \simeq \mathfrak{g}_{-2}^2 \oplus \mathfrak{g}_0^2 \oplus \mathfrak{g}_2^2$$

and the corresponding  $\mathbf{Z}$ -graduations of  $\mathfrak{k}$  are described by the following diagrams :



In the (a) case, one has  $\dim \mathfrak{g}_2^1 = 5$  ("natural" representation of  $A_4 = \mathfrak{sl}_5$ ) and  $\mathfrak{g}_0^1 \simeq \mathbb{C} \times A_1 \times A_4$  whereas in the (b) case  $\dim \mathfrak{g}_2^2 = 1$  (this is a general result if the removed root in the elementary operation is the one connected to  $-\omega$ , in fact the subalgebra  $\mathfrak{g}_1^2 \oplus \mathfrak{g}_2^2$  is known to be a Heisenberg algebra (see for example Corollary 2.3. in [J])) and  $\mathfrak{g}_0^2 \simeq \mathbb{C} \times A_5$ .

Of course the spaces  $\mathfrak{g}_1^1$  and  $\mathfrak{g}_1^2$  have the same dimension ( $= \frac{1}{2} \dim \mathfrak{p} = 20$ ) but they are distinct in  $E_6$ .

**5.2.** Consider the following two elementary operations in  $E_7$  :



As the removed roots have coefficient 2 in the highest root, these diagrams correspond to parabolic realizations of two equi-rank algebra structure on  $E_7$  which are obviously conjugate under an extended diagram automorphism.

As in 5.1. the associate parabolic realizations

$$(a) \quad E_7 = \mathfrak{g}_{-2}^1 \oplus \mathfrak{g}_{-1}^1 \oplus \mathfrak{g}_0^1 \oplus \mathfrak{g}_1^1 \oplus \mathfrak{g}_2^1 \quad (b) \quad E_7 = \mathfrak{g}_{-2}^2 \oplus \mathfrak{g}_{-1}^2 \oplus \mathfrak{g}_0^2 \oplus \mathfrak{g}_1^2 \oplus \mathfrak{g}_2^2$$

are different.

For example the corresponding graduation of  $\mathfrak{k} \simeq A_1 \times D_6$  are given by the following diagrams



This shows that in the (a) case  $\dim \mathfrak{g}_2^1 = 1$  whereas in the (b) case  $\dim \mathfrak{g}_2^2 = 10$ .

**5.3.** The proof of the converse part of Theorem 3.1. 2) shows that if one has *any*  $\mathbb{Z}$ -graduation of the form

$$\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2 \quad (5-3-1)$$

then, even if the parabolic subalgebra  $\mathfrak{P} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2$  is *not* maximal, that is even if (5-3-1) is *not* a parabolic realization, one can define an equi-rank algebra structure on  $\mathfrak{g}$  by taking

$$\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2 \quad \mathfrak{p} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_1.$$

This situation occurs if  $\Psi \setminus \Gamma = \{\alpha_1, \alpha_2\}$  (notations as in paragraph 3) where  $\alpha_1$  and  $\alpha_2$  have coefficient 1 in the highest root.

This is for example the case in the following diagram :



In this case it is easy to see that  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2 \simeq \mathbb{C} \times D_5$  (think in terms of the extended diagram). Hence the Lie algebra  $\mathfrak{k}$  is reductive non semi-simple. Therefore by Theorem 3.1. there exists a parabolic realization of the type

$$E_6 = \mathfrak{g}'_{-1} \oplus \mathfrak{g}'_0 \oplus \mathfrak{g}'_1$$

where  $\mathfrak{k} = \mathfrak{g}'_0$ .

The only possibilities to get a maximal Levi factor of type  $\mathbb{C} \times D_5$  correspond to the following diagrams



which are conjugate.

Of course the vector spaces  $\mathfrak{g}_1$  and  $\mathfrak{g}'_1$  are isomorphic (of dimension  $2^4 = 16$ ), but  $\mathfrak{g}'_1$  is  $\mathfrak{k}$ -stable (as a representation of  $D_5$  it is the Spin representation) whereas  $\mathfrak{g}_1$  is not  $\mathfrak{k}$ -stable.

Let  $\Omega_1$  (resp.  $\Omega_2$ ) be the set of positive roots  $\alpha$  which can be expressed in the form  $\alpha = \alpha_1 \bmod(\Gamma)$  (resp.  $\alpha = \alpha_2 \bmod(\Gamma)$ ) where  $\alpha_1$  (resp.  $\alpha_2$ ) is the first (resp. second root) circled in (5-3-2).

Let

$$\mathfrak{g}_{\pm 1}^1 = \sum_{\alpha \in \pm \Omega_1} \mathfrak{g}^\alpha \quad \text{and} \quad \mathfrak{g}_{\pm 1}^2 = \sum_{\alpha \in \pm \Omega_2} \mathfrak{g}^\alpha.$$

Then

$$\mathfrak{g}_1 = \mathfrak{g}_1^1 \oplus \mathfrak{g}_1^2 \quad \text{and} \quad \mathfrak{g}_{-1} = \mathfrak{g}_{-1}^1 \oplus \mathfrak{g}_{-1}^2$$

and in this picture there are the spaces  $\mathfrak{g}_1^1 \oplus \mathfrak{g}_{-1}^1$  and  $\mathfrak{g}_1^2 \oplus \mathfrak{g}_{-1}^2$  which are  $\mathfrak{k}$ -invariant (here  $\mathfrak{k} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_2$ ).

## 6. Associated compact and semi-compact dual pairs

Let  $\mathfrak{g}_{\mathbf{R}}$  be a real form of  $\mathfrak{g}$  whose maximal compact subalgebra  $\mathfrak{k}_{\mathbf{R}}$  is a real form of  $\mathfrak{k}$ . Recall from Proposition 2.1. that such a real form exists and is unique up to conjugacy by  $\exp \mathfrak{k}$ .

Let us also recall that a *dual pair* in  $\mathfrak{g}_{\mathbf{R}}$  is a pair  $(\mathfrak{a}_{\mathbf{R}}, \mathfrak{b}_{\mathbf{R}})$  of reductive subalgebras of  $\mathfrak{g}_{\mathbf{R}}$ , such that  $\mathfrak{b}_{\mathbf{R}} = \mathcal{Z}_{\mathfrak{g}_{\mathbf{R}}}(\mathfrak{a}_{\mathbf{R}})$  and  $\mathfrak{a}_{\mathbf{R}} = \mathcal{Z}_{\mathfrak{g}_{\mathbf{R}}}(\mathfrak{b}_{\mathbf{R}})$ , where  $\mathcal{Z}_{\mathfrak{g}_{\mathbf{R}}}(\mathfrak{a})$  (resp.  $\mathcal{Z}_{\mathfrak{g}_{\mathbf{R}}}(\mathfrak{b})$ ) denotes the centralizer of  $\mathfrak{a}$  (resp.  $\mathfrak{b}$ ) in  $\mathfrak{g}_{\mathbf{R}}$ .

In this paragraph, we will describe three families of compact or semi-compact dual pairs (this means that the two algebras  $\mathfrak{a}_{\mathbf{R}}$  and  $\mathfrak{b}_{\mathbf{R}}$  are compact or one of them) inside  $\mathfrak{g}_{\mathbf{R}}$  which are naturally associated to the corresponding equi-rank algebra.

**6.1.** This first family is associated to the first type of equi-rank algebras (described in the first part of Theorem 3.1. and listed as the first type in Table 1). The obvious remark to do is that as  $\mathfrak{k}$  is a Levi subalgebra of  $\mathfrak{g}$  then  $(\mathfrak{k}, \mathcal{Z}(\mathfrak{k}))$  (where  $\mathcal{Z}(\mathfrak{k})$  is the center of  $\mathfrak{k}$ ) is a dual pair in  $\mathfrak{g}$ . Therefore  $(\mathfrak{k}_{\mathbf{R}}, \mathcal{Z}(\mathfrak{k}_{\mathbf{R}}))$  is a dual pair in  $\mathfrak{g}_{\mathbf{R}}$ . Here  $\mathcal{Z}(\mathfrak{k}_{\mathbf{R}})$  is one dimensional because  $\mathfrak{k}$  is the Levi factor of a maximal parabolic subalgebra. Hence  $\mathcal{Z}(\mathfrak{k}_{\mathbf{R}})$  is isomorphic to  $\mathfrak{so}(2)$ . This leads easily to the following table.

**Table 2**

(The numbering of the different cases is as in Table 1, the notations for the real simple Lie algebras are the same as in Helgason's book, Table V, page 518 [He])



$$(1 - 1) \quad (\mathfrak{so}(2) \times \mathfrak{su}(p) \times \mathfrak{su}(n+1-p), \mathfrak{so}(2)) \subset \mathfrak{su}(p, n+1-p)$$

$$(1 - 2) \quad (\mathfrak{so}(2) \times \mathfrak{so}(2n-1), \mathfrak{so}(2)) \subset \mathfrak{so}(2, 2n-1)$$

$$(1 - 3) \quad (\mathfrak{so}(2) \times \mathfrak{su}(n), \mathfrak{so}(2)) \subset \mathfrak{sp}(n, \mathbf{R})$$

$$(1 - 4) \quad (\mathfrak{so}(2) \times \mathfrak{so}(2n-2), \mathfrak{so}(2)) \subset \mathfrak{so}(2, 2n-2)$$

$$(1 - 5) \quad (\mathfrak{so}(2) \times \mathfrak{su}(n), \mathfrak{so}(2)) \subset \mathfrak{so}^*(2n)$$

$$(1 - 6) \quad (\mathfrak{so}(2) \times \mathfrak{so}(10), \mathfrak{so}(2)) \subset \mathfrak{e}_{6(-14)}$$

$$(1 - 7) \quad (\mathfrak{so}(2) \times \mathfrak{e}_{6(-78)}, \mathfrak{so}(2)) \subset \mathfrak{e}_{7(-25)}$$

**6.2.** The second family is a family of semi-compact dual pairs, which is also associated to the first type of equi-rank algebras, or to be more precise to those equi-rank algebras which correspond to hermitian symmetric spaces of *tube type*.

But the definition of these pairs is more involved as the preceding one. In fact these pairs are real forms of very basic complex dual pairs which have been used by the author as "primitive objects" from which all complex dual pairs in  $\mathfrak{g}$  can be built ([Ru 2]).

Here we shall only consider equi-rank algebras of the first type such that the underlying hermitian pair  $(\mathfrak{g}_{\mathbf{R}}, \mathfrak{k}_{\mathbf{R}})$  is of *tube type*.

Let  $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$  be a parabolic realization of such an algebra. Let  $H$  be the element of  $\mathfrak{g}_0$  which defines the graduation (i.e. which has eigenvalue  $i$  on  $\mathfrak{g}_i$ , see (3-1)). It is known that the tube type condition is equivalent to the fact that the element  $2H$  can be put into an  $\mathfrak{sl}_2$ -triple. More precisely there exists

$X \in \mathfrak{g}_1$  and  $Y \in \mathfrak{g}_{-1}$  such that  $(Y, H, X)$  is an  $\mathfrak{sl}_2$ -triple ( $[Y, X] = H$ ,  $[H, X] = 2X$ ,  $[H, Y] = -2Y$ ). For a proof see for example [K-W].

Moreover the elements  $H, X, Y$  can be described in the following way. Let  $\alpha_1, \alpha_2, \dots, \alpha_k$  be a maximal set of strongly orthogonal roots such that the corresponding root spaces  $\mathfrak{g}^{\alpha_i}$  are in  $\mathfrak{g}_1$ . Let  $(X_{-\alpha_i}, H_{\alpha_i}, X_{\alpha_i})$  be a classical  $\mathfrak{sl}_2$ -triple associated to each root. Here  $H_{\alpha_i}$  is the co-root of  $\alpha_i$ ,  $X_{\alpha_i} \in \mathfrak{g}^{\alpha_i}$  and  $X_{-\alpha_i} \in \mathfrak{g}^{-\alpha_i}$ . Then  $2H = \sum_{i=1}^k H_{\alpha_i}$  and one can take  $X = \sum_{i=1}^k X_{\alpha_i}$  and  $Y = \sum_{i=1}^k X_{-\alpha_i}$  (see for example [M-R-S]).

The algebra  $\mathfrak{k}_{\mathbf{R}}$  has a non trivial center, which is necessarily  $i\mathbf{R}H$  because the restriction of the Killing form must be negative definite. As there exists a Cartan subalgebra included in  $\mathfrak{k}_{\mathbf{R}}$  on which all the roots take purely imaginary values, one has for any root  $\overline{\mathfrak{g}^{\alpha}} = \mathfrak{g}^{-\alpha}$ , where bar denotes the conjugation with respect to  $\mathfrak{g}_{\mathbf{R}}$ . Therefore there exists a non zero constant  $c_i$  such that  $\overline{X_{\alpha_i}} = c_i X_{-\alpha_i}$ . As  $X_{\alpha_i} \in \mathfrak{p}$ , one has  $B(X_{-\alpha_i}, X_{\alpha_i}) > 0$ , where  $B$  is the Killing form. On the other hand, from the invariance of the Killing form, one gets  $B(X_{-\alpha_i}, X_{\alpha_i}) = -\frac{1}{2}B(H_{\alpha_i}, H_{\alpha_i})$ . Hence  $B(\overline{X_{\alpha_i}}, X_{\alpha_i}) = c_i B(X_{-\alpha_i}, X_{\alpha_i}) = -\frac{1}{2}B(H_{\alpha_i}, H_{\alpha_i})c_i > 0$ . This implies  $c_i < 0$ .

Let  $E_{\alpha_i} = |c_i|^{-\frac{1}{2}} X_{\alpha_i}$ . Then  $\overline{E_{\alpha_i}} = |c_i|^{-\frac{1}{2}} \overline{X_{\alpha_i}}$  and  $(-\overline{E_{\alpha_i}}, H_{\alpha_i}, E_{\alpha_i})$  is an  $\mathfrak{sl}_2$ -triple. Define  $\tilde{X} = \sum_{i=1}^k E_{\alpha_i}$  and  $\tilde{Y} = \sum_{i=1}^k -\overline{E_{\alpha_i}}$ . Then  $(\tilde{Y}, 2H, \tilde{X})$  is again an  $\mathfrak{sl}_2$ -triple. This triple generates an  $\mathfrak{sl}_2$  subalgebra  $\mathfrak{a}$  of  $\mathfrak{g}$  which is stable under conjugation with respect to  $\mathfrak{g}_{\mathbf{R}}$ , therefore  $\mathfrak{a}_{\mathbf{R}} = \mathfrak{g}_{\mathbf{R}} \cap \mathfrak{a}$  is a real form of  $\mathfrak{a}$ . One can take as a basis of  $\mathfrak{a}_{\mathbf{R}}$  the elements

$$U = \sum_{i=1}^k (E_{\alpha_i} + \overline{E_{\alpha_i}}), \quad V = -i \sum_{i=1}^k (E_{\alpha_i} - \overline{E_{\alpha_i}}), \quad T = i \sum_{i=1}^k H_{\alpha_i}.$$

The relations are then  $[T, U] = -2V$ ,  $[T, V] = 2U$  and  $[V, U] = -2T$  and this proves that  $\mathfrak{a}_{\mathbf{R}}$  is isomorphic to  $\mathfrak{sl}_2(\mathbf{R})$ . The existence of this  $\mathfrak{sl}_2(\mathbf{R})$  subalgebra in the tube type case is due to Koranyi and Wolf ([K-W], Prop. 3.12.)

Recall now the following result.

**Theorem 6.2.1..** — ([Ru 2], Th. 4.3.)

Let  $\mathfrak{b} = \mathcal{Z}_{\mathfrak{g}}(\mathfrak{a})$  be the centralizer of  $\mathfrak{a}$  in  $\mathfrak{g}$ . Then  $(\mathfrak{a}, \mathfrak{b})$  is a dual pair in  $\mathfrak{g}$ .

As  $\mathfrak{a}$  is split relatively to  $\mathfrak{g}_{\mathbf{R}}$ , the Lie algebra  $\mathfrak{b}$  is also split relatively to  $\mathfrak{g}_{\mathbf{R}}$ , i.e.  $\mathfrak{b}_{\mathbf{R}} = \mathfrak{b} \cap \mathfrak{g}$  is a real form of  $\mathfrak{b}$ .

The following corollary is then straightforward.

**Corollary 6.2.2..** —

The pair  $(\mathfrak{a}_{\mathbf{R}}, \mathfrak{b}_{\mathbf{R}})$  is a dual pair in  $\mathfrak{g}_{\mathbf{R}}$ . The Lie algebra  $\mathfrak{a}_{\mathbf{R}}$  is isomorphic to  $\mathfrak{sl}_2(\mathbf{R})$  and  $\mathfrak{b}_{\mathbf{R}}$  is a subalgebra of  $\mathfrak{k}_{\mathbf{R}}$ , hence is compact.

This leads to the following table of dual pairs.

**Table 3**

(The numbering of the different cases is as in table 1, the notations for the real simple Lie algebras are the same as in Helgason's book, Table V, page 518 [He], the type of the compact algebra  $\mathfrak{b}_{\mathbf{R}}$  is deduced from [Ru2], 6.13.Table 5.)

$$(1 - 1) \quad (n = 2p - 1) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{su}(p)) \subset \mathfrak{su}(p, p)$$

$$(1 - 2) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{so}(2n - 2)) \subset \mathfrak{so}(2, 2n - 1)$$

$$(1 - 3) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{so}(n)) \subset \mathfrak{sp}(n, \mathbf{R})$$

$$(1 - 4) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{so}(2n - 3)) \subset \mathfrak{so}(2, 2n - 2)$$

$$(1 - 5) \quad (n \text{ even}) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{sp}(n)) \subset \mathfrak{so}^*(2n)$$

$$(1 - 7) \quad (\mathfrak{sl}_2(\mathbf{R}), \mathfrak{f}_{4(-52)}) \subset \mathfrak{e}_{7(-25)}$$

**6.3.** The dual pairs in the third family are compact dual pairs associated the second type of equi-rank algebras.

Recall the following result.

**Proposition 6.3.1.** — ([Ru 2], Prop. 5.15.)

*Let  $\mathfrak{g}$  be a simple Lie algebra over  $\mathbf{C}$ . Let  $\mathfrak{a}$  and  $\mathfrak{b}$  be two semi-simple subalgebras of  $\mathfrak{g}$  such that  $\mathfrak{a} \cap \mathfrak{b} = \{0\}$  and such that  $\mathfrak{a} \times \mathfrak{b}$  is a regular maximal subalgebra of  $\mathfrak{g}$  (this implies that  $\mathfrak{a} \times \mathfrak{b}$  is obtained by a single elementary operation). Then  $(\mathfrak{a}, \mathfrak{b})$  is a dual pair in  $\mathfrak{g}$ .*

**Remark 6.3.2.** —

As noticed in the preceding proposition, all maximal semi-simple regular subalgebras are obtained by a single elementary operation, the converse is true only if the coefficient of the root which is removed has a prime coefficient in the highest root (see the remark following Theorem 3.5. in [T], exercise 4, par. 4, p. 229 in [Bou 1] and exercise 2, par. 3, p. 222 in [Bou 2]).

**Corollary 6.3.3.** —

*In all equi-rank algebras of the second type where the subalgebra  $\mathfrak{k}$  can be written as a product  $\mathfrak{g} = \mathfrak{a} \times \mathfrak{b}$  with  $\mathfrak{a}$  and  $\mathfrak{b}$  simple algebras, the pair  $(\mathfrak{a}, \mathfrak{b})$  is a dual pair in  $\mathfrak{g}$ .*

Proof. As explained in paragraph 3 and 4, the equi-rank algebras of the second type are obtained by a single elementary operation where the removed root has coefficient 2 in the highest root.  $\square$

**Corollary 6.3.4.** —

*Let  $\mathfrak{g}$  be an equi-rank algebra of the second type where  $\mathfrak{k}$  can be written as a product  $\mathfrak{k} = \mathfrak{a} \times \mathfrak{b}$  with  $\mathfrak{a}$  and  $\mathfrak{b}$  simple algebras. Let  $\mathfrak{a}_R = \mathfrak{g}_R \cap \mathfrak{a}$  and  $\mathfrak{b}_R = \mathfrak{g}_R \cap \mathfrak{b}$ , then  $(\mathfrak{a}_R, \mathfrak{b}_R)$  is a dual pair in  $\mathfrak{g}_R$ .*

This leads easily to the following table.

**Table 4**

(The numbering of the different cases is as in table 1, the notations for the real simple Lie algebras are the same as in Helgason's book, Table V, page 518 [He])

$$(2-1) \quad p \geq 2 \quad (\mathfrak{so}(2p), \mathfrak{so}(2n-2p+1)) \subset \mathfrak{so}(2p, 2n-2p+1)$$

$$(2-2) \quad p \leq n-1 \quad (\mathfrak{sp}(p), \mathfrak{sp}(n-p)) \subset \mathfrak{sp}(p, n-p)$$

$$(2-3) \quad 2 \leq p \leq n-2 \quad (\mathfrak{so}(2p), \mathfrak{so}(2n-2p)) \subset \mathfrak{so}(2p, 2n-2p)$$

$$(2-4) \quad (\mathfrak{su}(2), \mathfrak{su}(6)) \subset \mathfrak{e}_{6(2)}$$

$$(2 - 5) \quad (\mathfrak{su}(2), \mathfrak{so}(12)) \subset \mathfrak{e}_{7(-5)}$$

$$(2 - 7) \quad (\mathfrak{su}(2), \mathfrak{e}_{7(-133)}) \subset \mathfrak{e}_{8(-24)}$$

$$(2 - 9) \quad (\mathfrak{su}(2), \mathfrak{sp}(3)) \subset \mathfrak{f}_{4(4)}$$

$$(2 - 11) \quad (\mathfrak{su}(2), \mathfrak{su}(2)) \subset \mathfrak{g}_{2(2)}$$

**Remark 6.3.5.** —

In the cases from the preceding table where  $\mathfrak{k}_{\mathbf{R}}$  has an  $\mathfrak{su}(2)$  factor, Gross and Wallach have studied the discrete series of some groups corresponding to  $\mathfrak{g}_{\mathbf{R}}$  ([G-W]).

### Bibliography

- [B-dS] A. BOREL and J. de SIEBENTHAL. — *Les sous-groupes fermés de rang maximum des groupes de Lie clos*, Comment. Math. Helv. 23 (1949), 200-221.
- [Bou1] N. BOURBAKI. — *Groupes et Algèbres de Lie Ch. 4,5 et 6*, Masson, Paris (1981).
- [Bou2] N. BOURBAKI. — *Groupes et Algèbres de Lie Ch. 7 et 8*, Hermann, Paris (1975).
- [D] E.B. DYNKIN. — *Semi-simple subalgebras of semi-simple Lie algebras*, A.M.S. Tanslations 6 (1957), 111-244.
- [G-W] B. H. GROSS and N. R. WALLACH. — *On quaternionic discrete series representations, and their continuations.*, J. reine angew. Math. 481 (1996), 73-123.
- [HC] HARISH-CHANDRA . — *Discrete series for semisimple Lie groups I,II* , Acta Math. 113 (1965), 241-318, Acta Math. 116 (1966), 1-111.
- [He] S. HELGASON. — *Differential Geometry, Lie Groups, and Symmetric spaces*, Academic Press, New york (1978).

- [J] A. JOSEPH. — *The minimal orbit in a simple Lie algebra and its associated maximal ideal*, Ann. Scient. Ec. Norm. Sup. t.9. (1976), 1-30.
- [K-W] A. KORANYI and J. A. WOLF. — *Realization of Hermitian Symmetric Spaces as Generalized Half-planes*, Annals of Math. 81 (1965), 265-288.
- [K1] A.W. KNAPP. — *A quick proof of the classification of simple real Lie algebras*, Proc. Amer. Math. Soc. 124 (1996) 3257-3259.
- [K2] A.W. KNAPP. — *Lie Groups Beyond an Introduction*, Birkhäuser, Basel (1996).
- [M-R-S] I. MULLER, H. RUBENTHALER and G. SCHIFFMANN. — *Structure des espaces préhomogènes associés à certaines algèbres de Lie graduées*, Math. Ann. 274 (1986), 95-123.
- [M] S. MURAKAMI. — *Sur la classification des algèbres de Lie réelles et simples*, Osaka J. Math. 2 (1965), 291-307.
- [O-V] A.L. ONISHCHIK and E.B. VINBERG. — *Lie Groups and Algebraic Groups* Springer-Verlag, Berlin (1990).
- [Ru1] H. RUBENTHALER. — *Espaces préhomogènes de type parabolique*, Lect. Math. Kyoto Univ. 14 (1982), 189-221.
- [Ru 2] H. RUBENTHALER. — *Les paires duales dans les algèbres de Lie réductives*, Astérisque 219 (1994).
- [T] J. TITS. — *Sous-algèbres des algèbres de Lie semi-simples*, Séminaire Bourbaki, exposé 119 (1955).
- [Wal] N. R. WALLACH. — *On maximal subsystems of roots systems*, Canad. J. Math. 20 (1968), 555-574.
- [War] G. WARNER. — *Harmonic Analysis on Semi-Simple Lie Groups, Vol. 1* Springer-Verlag, New-York (1972).

HUBERT RUBENTHALER  
 IRMA et Département de Mathématiques  
 Université Louis Pasteur  
 7, rue René Descartes  
 67084 Strasbourg Cedex  
 France  
 e-mail : rubenth@math.u-strasbg.fr

December, 1998