



**HAL**  
open science

## **-trees and laminations for free groups II: The dual lamination of an -tree**

Thierry Coulbois, Arnaud Hilion, Martin Lustig

► **To cite this version:**

Thierry Coulbois, Arnaud Hilion, Martin Lustig. -trees and laminations for free groups II: The dual lamination of an -tree. 2007. hal-00130239v1

**HAL Id: hal-00130239**

**<https://hal.science/hal-00130239v1>**

Preprint submitted on 9 Feb 2007 (v1), last revised 9 Jun 2007 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# $\mathbb{R}$ -trees and laminations for free groups II: The dual lamination of an $\mathbb{R}$ -tree

Thierry Coulbois, Arnaud Hilion, Martin Lustig

February 10, 2007

## 1 Introduction

A geodesic lamination  $\mathfrak{L}$  on a hyperbolic surface  $S$ , provided with a transverse measure, defines (via the lift  $\tilde{\mathfrak{L}}$  of  $\mathfrak{L}$  to the universal covering of  $S$ ) an action of  $\pi_1 S$  on an  $\mathbb{R}$ -tree  $T$  which is often called *dual* to the lamination  $\tilde{\mathfrak{L}}$ . Conversely, every small action of  $\pi_1 S$  on an  $\mathbb{R}$ -tree  $T$  comes from this construction, provided the surface is closed and the action on  $T$  is *small* [Sko96].

A generalization of this concept occurs first in the work of E. Rips, and is since then widely used. A particular kind of  $\mathbb{R}$ -tree actions can be defined as spaces dual to measured foliations on finite 2-complexes. If the latter is not a surface (and can not be made into a surface by certain elementary moves), then the resulting  $\mathbb{R}$ -tree is qualitatively different from the ones dual to surface laminations, see [GLP94, BF95, LP97].

A third occurrence of laminations in direct relation to  $\mathbb{R}$ -trees takes place in the context of free group automorphisms, specifically for *irreducible automorphisms with irreducible powers* (= *iwip automorphisms*). For such an  $\alpha \in \text{Aut}(F_N)$  every (non-periodic) conjugacy class of elements in  $F_N$  converges to a collection of biinfinite *legal* paths on a train track representing  $\alpha$ , see [BFH97]. On the other hand, the train track itself converges towards an  $\mathbb{R}$ -tree  $T_\alpha$  which is projectively fixed under the induced action of  $\alpha$  on the set of  $\mathbb{R}$ -tree actions of  $F_N$ . In the particular case where  $\alpha$  is induced by a pseudo-Anosov homeomorphism  $\varphi$  of some surface  $S$  (with one boundary component), then the above collection of biinfinite legal paths is in precise 1 - 1 correspondence with the leaves of the stable lamination of  $\varphi$ .

Finally, in [LL03] a collection of one-sided infinite words reminiscent to half-leaves of a lamination was associated to an arbitrary very small  $F_N$ -action on an  $\mathbb{R}$ -tree  $T$ , as a tool to prove that iwip automorphisms of  $F_N$  have a North-South dynamics on the space  $\overline{CV}_N$  of very small  $\mathbb{R}$ -tree actions of  $F_N$ .

This puzzle of coinciding and consistent observations induced the authors to set out for a general theory, in the realm of free (and perhaps later word-hyperbolic) groups. As a first step, in [CHL-I] algebraic laminations were defined, generalizing at the same time geodesic laminations on surfaces, as well as symbolic flows as known from discrete dynamics. They come in three equivalent languages, group theoretic, dynamic and combinatorial, and passing from one to the other turns out to be rather helpful. In [CHL-I] these “translations” were established with some care, and the topology, the partial order, as well as the natural  $\text{Out}(F_N)$ -action were studied.

In the present paper we use these tools to define, for any isometric  $F_N$ -action on an  $\mathbb{R}$ -tree  $T$ , a set

$$L^2(T) \subset \partial^2 F_N := \partial F_N \times \partial F_N \setminus \Delta,$$

where  $\partial F_N$  denotes the Gromov boundary of the free group  $F_N$ , and  $\Delta$  is the diagonal. The set  $L^2(T)$  is empty if the  $F_N$ -action on  $T$  is free and discrete (simplicial), and it is an algebraic lamination otherwise. There are several competing natural approaches to define  $L^2(T)$ , which we present briefly below. Working out the precise relationship between them is the core content of this paper.

**1.** (see §4) The lamination  $L_\Omega^2(T)$  is defined for all isometric actions of  $F_N$  on an  $\mathbb{R}$ -tree  $T$ : For every  $\varepsilon > 0$  we consider the set  $\Omega_\varepsilon(T)$  of all elements  $g \in F_N$  with translation length on  $T$  that satisfies:

$$\|g\|_T < \varepsilon$$

The set  $\Omega_\varepsilon(T)$  generates an algebraic lamination  $L_\varepsilon^2(T)$  which is the smallest lamination that contains every  $(g^{-1}g^{-1}g^{-1}\dots, ggg\dots) \in \partial^2 F_N$  with  $g \in \Omega_\varepsilon(T)$ . We define  $L_\Omega^2(T)$  to be the intersection of all  $L_\varepsilon^2(T)$ .

We are most interested in  $\mathbb{R}$ -trees  $T$  where every  $F_N$ -orbit of points is dense in  $T$ . To any such  $T$  we associate in this paper two more laminations, which are of rather different nature:

**2.** (see §5) In order to define the lamination  $L_\infty^2(T)$  one first fixes a basis  $\mathcal{A}$  of  $F_N$ . Then one picks an arbitrary point  $P \in T$  and considers the set  $L_{\mathcal{A}}^1(T)$  of one-sided infinite words  $x_1x_2\dots$  in some basis  $\mathcal{A}$  of  $F_N$  such that the set of all  $x_1x_2\dots x_kP$  has bounded diameter in  $T$ . One immediately observes that the set  $L_{\mathcal{A}}^1(T)$  is independent of the choice of  $P$ . As next step, one considers the language  $\mathcal{L}_{\mathcal{A}}(L_{\mathcal{A}}^1(T))$  derived from  $L_{\mathcal{A}}^1(T)$ , i.e. the set of all finite subwords of any  $x_1x_2\dots \in L_{\mathcal{A}}^1(T)$ , and its *recurrent* sublanguage  $\mathcal{L}_{\mathcal{A}}^\infty(L_{\mathcal{A}}^1(T))$ : The latter consists precisely of those words which occur infinitely often as subword in some  $x_1x_2\dots \in L_{\mathcal{A}}^1(T)$ . The advantage of this language is that it is *laminary*, and thus it defines canonically an algebraic lamination  $L_\infty^2(T)$  (see [CHL-I]). We prove in detail in §5 that this algebraic laminaton does not depend on the basis  $\mathcal{A}$  used in the construction sketched above.

*Aside:* The passage from  $L^1(T)$  to the recurrent language  $\mathcal{L}_{\mathcal{A}}^\infty(L_{\mathcal{A}}^1(T))$  is subtle but rather important: It turns out that  $L^1(T)$  depends not just on the topology, but actually on the metric of  $T$ . This is discussed in detail in [CHL05] and [CHLL].

**3.** (see §8) If  $T$  is an  $\mathbb{R}$ -tree dual to a measured lamination  $\mathfrak{L}$  on surface  $S$ , then a leaf  $l$  of the lift  $\tilde{\mathfrak{L}}$  of  $\mathfrak{L}$  to the universal covering  $\tilde{S}$  determines on one hand a point  $x_l$  in the dual tree  $T$ , and on the other hand two limit points  $P_+, P_- \in \partial\tilde{S}$  on the boundary at infinity  $\partial\tilde{S}$  of  $\tilde{S}$ :

$$\{P_+, P_-\} = \partial l$$

Note that, in the case where  $S$  has non-empty boundary,  $\partial\tilde{S}$  is canonically identified with the Gromov boundary  $\partial F_N$  of  $F_N = \pi_1 S$ .

In [LL03] this correspondence has been generalized to an  $F_N$ -equivariant map  $\mathcal{Q} : \partial F_N \rightarrow \overline{T} \cup \partial T$ , for any  $\mathbb{R}$ -tree  $T$  with dense  $F_N$ -orbits, where  $\overline{T}$  denotes the metric completion of  $T$ , and  $\partial T$  the Gromov boundary. The definition of the map  $\mathcal{Q}$  is reviewed in §6, and the geometric meaning of  $\mathcal{Q}$  is explained in more detail in §7. In the above special case one gets:

$$\mathcal{Q}(P_+) = \mathcal{Q}(P_-) = x_l \in T$$

This motivates the definition of the lamination  $L_{\mathcal{Q}}^2(T)$ , which consists of all pairs  $(X, X') \in \partial^2 F_N$  that determine the same limit point  $\mathcal{Q}(X) = \mathcal{Q}(X') \in \overline{T}$ .

The main result of this paper, proved in two steps (Propositions 5.8 and 8.5), is:

**Theorem 1.1.** *For every very small  $\mathbb{R}$ -tree with  $F_N$ -action that has dense orbits the above described three algebraic laminations coincide:*

$$L^2_{\Omega}(T) = L^2_{\infty}(T) = L^2_{\mathcal{Q}}(T)$$

*This defines a dual algebraic lamination  $L^2(T)$  canonically associated to  $T$ .*

In a subsequent third paper we go one step further and consider invariant measures  $\mu$  (called *currents*) on  $L^2(T)$ . We show in [CHL-III] that such a current defines a *dual metric*  $d_{\mu}$  on  $T$ : If  $T$  is dual to a surface lamination  $\mathfrak{L}$  as in the beginning of this introduction, and if  $\mu$  comes from a transverse measure on  $\mathfrak{L}$ , then  $d_{\mu}$  is indeed the  $\mathbb{R}$ -tree metric on the dual tree  $T$ . In general, however, it is shown in [CHL-III] that this dual metric can have very exotic properties.

Note also that the lamination  $L^2(T)$  introduced in this paper has been used successfully in [CHL05] to characterize the underlying topological structure of  $\mathbb{R}$ -trees which stays invariant when the metric is  $F_N$ -equivariantly changed (so called “non-uniquely ergometric  $\mathbb{R}$ -trees”).

The dual lamination  $L^2(T)$  plays also a crucial role in establishing a continuous and  $\text{Out}(F_N)$ -equivariant map from a large part of the boundary of Outer space into the space of projectivized currents, see [CHL3] and [KL]. It is also the basis for work in progress of the third author with I. Kapovich on perpendicular ( $\mathbb{R}$ -tree, current)-pairs.

*Acknowledgements: This paper originates from a workshop organized at the CIRM in April 05, and it has greatly benefited from the discussions started there and continued around the weekly Marseille seminar “Teichmüller” (partially supported by the FRUMAM).*

## 2 Free group actions on $\mathbb{R}$ -trees

In this section we recall some of the known properties of actions of a free group  $F_N$  on an  $\mathbb{R}$ -tree  $T$ . For details and background see [Vog02, Sha87] and the references given there.

An  $\mathbb{R}$ -tree is a metric space which is 0-hyperbolic and geodesic. Alternatively, an  $\mathbb{R}$ -tree is a metric space  $(T, d)$  where any two points  $P, Q \in T$  are joined by a unique arc and this arc is isometric to the interval  $[0, d(P, Q)] \subset \mathbb{R}$ .

In this paper an  $\mathbb{R}$ -tree  $T$  always comes with a left action of  $F_N$  on  $T$  by isometries. Any isometry  $w$  of  $T$  is either *elliptic*, in which case it fixes at least one point of  $T$ , or else it is *hyperbolic*, in which case there is an *axis*  $\text{Ax}(w)$  in  $T$ , isometric to  $\mathbb{R}$ , which is  $w$ -invariant, and along which  $w$  acts as translation. The *translation length*

$$\|w\|_T = \inf\{d(P, wP) \mid P \in T\}$$

agrees in the hyperbolic case with  $d(Q, wQ)$  for any point  $Q$  in  $\text{Ax}(w)$ , while in the elliptic case it is 0. We always assume that  $T$  is a *minimal*  $\mathbb{R}$ -tree, i.e. there is no non-empty  $F_N$ -invariant proper subtree of  $T$ . Another  $\mathbb{R}$ -tree  $T'$  with isometric  $F_N$ -action is  $F_N$ -equivariantly isometric to  $T$  if and only if one has

$$\|w\|_T = \|w\|_{T'}$$

for every element  $w \in F_N$ . The set of such trees (or rather, of such tree actions) inherits a topology from its image in  $\mathbb{R}^{F_N}$  under the map

$$T \mapsto (\|w\|_T)_{w \in F_N} \in \mathbb{R}^{F_N}.$$

A tree (or a tree action) is called *small* if any two group elements that fix pointwise a non-trivial arc in  $T$  do commute. It is called *very small* if moreover (i) the fixed set  $\text{Fix}(g) \subset T$  of any elliptic element  $1 \neq g \in F_N$  is a segment or a single point (i.e. “no branching”), and (ii)  $\text{Fix}(g) = \text{Fix}(g^m)$  for all  $g \in F_N$  and  $m \geq 1$ .

The particular case of *simplicial*  $\mathbb{R}$ -trees  $T$  with isometric  $F_N$ -actions arises from graphs  $\Gamma$  with a *marking isomorphism*  $F_N \cong \pi_1\Gamma$ , where the edges of  $\Gamma$  are given a non-negative length, which is for at least one of them strictly positive: The simplicial tree  $T$  is then given by the universal covering  $\tilde{\Gamma}$ , equipped with the action of  $F_N$  by deck transformations. If every edge length of  $\Gamma$  is non-zero, then the action of  $F_N$  on  $\tilde{\Gamma}$  is free. The subspace  $\text{cv}_N$  of free simplicial actions has been introduced by M. Culler and K. Vogtmann. Its closure  $\overline{\text{cv}}_N$  in the space of  $F_N$ -actions on  $\mathbb{R}$ -trees is precisely the set of all of the above mentioned very small  $\mathbb{R}$ -trees. The *boundary*  $\overline{\text{cv}}_N \setminus \text{cv}_N$  is denoted by  $\partial\text{cv}_N$ . One often normalizes  $\Gamma$  to have volume 1, thus obtaining the subspace  $\text{CV}_N$  of  $\text{cv}_N$ , which has been named *Outer space* by P. Shalen. Alternatively, one can *projectivize* the space of tree actions: two trees  $T$  and  $T'$  are in the same equivalence class  $[T]$  if they are  $F_N$ -equivariantly homothetic. This projectivization maps  $\overline{\text{cv}}_N$  onto a compact space  $\overline{\text{CV}}_N$ ,

which contains a homeomorphic copy of  $CV_N$ , called the *interior*, and the projectivized image  $\partial CV_N$  of  $\partial cv_N$ , called the *boundary*. Both  $CV_N$  and  $\overline{CV_N}$  are contractible and finite dimensional. For more information see [Vog02, CV86].

The group  $\text{Out}(F_N)$  acts by homeomorphisms on  $cv_N$  and  $\partial cv_N$ , as well as on  $CV_N$  and  $\partial CV_N$ , and the action on  $CV_N$  is properly discontinuous (though not free). These actions, specified in §9, provide valuable information about the group  $\text{Out}(F_N)$ . Note also that there is a strong similarity between Outer space  $CV_N$  with the  $\text{Out}(F_N)$ -action on one side, and Teichmüller space with the action of the mapping class group on the other. The only substantial difference is that  $CV_N$  is not a manifold.

In the second half of this paper we will concentrate on the particularly interesting case where some (and hence every)  $F_N$ -orbit of points is dense in  $T$ . That this “dense orbits” hypothesis is not very restrictive follows from the following consideration:

The trees in  $cv_N \cup \partial cv_N$  have only finitely many  $F_N$ -orbits of *branch points*, i.e. points with more than 2 complementary components in  $T$  (see [GL95]). Every such minimal tree decomposes canonically into two disjoint  $F_N$ -invariant (possibly empty) subsets  $T_d$  and  $T_c$ , where the former is given as the union of all points  $P \in T$  such that the orbit  $F_N P$  is a discrete subset of  $T$ , and the latter is the complement  $T \setminus T_d$ . The subset  $T_c \subset T$  is easily seen to be closed, and each connected component of it is a subtree  $T'$  of  $T$  with the property that the subgroup  $U$  of  $F_N$  that stabilizes  $T'$  acts on  $T'$  with *dense orbits*:  $T' = \overline{UP}$  for any  $P \in T'$ . Unless  $T_c$  is empty, in which case the set of branch points is a discrete subset of  $T$  and thus  $T$  is simplicial, we can contract the closure of every connected component of  $T_d$  to a single point, to get the canonical maximal non-trivial quotient tree  $T/T_d$  on which now all of  $F_N$  acts minimally and with dense orbits. Compare [Gui98].

### 3 Bounded Back Tracking

Every small action on an  $\mathbb{R}$ -tree is known to have the *bounded backtracking property (BBT)* (see [GJLL98]), which is of great use in this paper:

Let  $\Gamma$  be any (non-metric) graph with a marking isomorphism  $\pi_1 \Gamma \cong F_N$ , and let  $\tilde{\Gamma}$  be its universal covering. Let  $i : \tilde{\Gamma} \rightarrow T$  be any  $F_N$ -equivariant map. Then the map  $i$  *satisfies BBT* if and only if for every pair of points

$P, Q \in \tilde{\Gamma}$  the  $i$ -image of the geodesic segment  $[P, Q] \subset \tilde{\Gamma}$  is contained in the  $C$ -neighborhood of  $[i(P), i(Q)] \subset T$ , where  $C \geq 0$  is an a priori constant independent of the choice of  $P$  and  $Q$ . We denote by  $\text{BBT}(i) \geq 0$  the smallest such constant.

It is easy to see that every  $\mathbb{R}$ -tree  $T$  with isometric  $F_N$ -action admits a map  $i$  as above, and that  $i$  satisfies BBT if and only if any other such map  $i' : \tilde{\Gamma}' \rightarrow T$  also satisfies BBT. Hence the property BBT is indeed a well defined property of the tree  $T$ .

We can assume that the above map  $i : \tilde{\Gamma} \rightarrow T$  is *edge-geodesic*:  $i$  maps every edge  $e \subset \tilde{\Gamma}$  to the geodesic segment that connects the images of the endpoints of  $e$ . One can make  $\Gamma$  into a metric graph by giving each edge of  $\Gamma$  and each of its lifts  $e$  to  $\tilde{\Gamma}$  the length of  $i(e)$ . Without loss of generality one can assume that the metric on every edge  $e$  is properly distributed so that  $i$  is actually *edge isometric*, i.e.  $i$  maps every edge of  $\tilde{\Gamma}$  isometrically onto its image. In this case the inequality

$$\text{BBT}(i) \leq \text{vol}(\Gamma)$$

has been proved in [GJLL98], where the *volume*  $\text{vol}(\Gamma)$  of  $\Gamma$  is the sum of the lengths of its edges.

A particular choice of  $\Gamma$ , for any base  $\mathcal{A}$  of  $F_N$ , is the rose  $\mathcal{R}_{\mathcal{A}}$  with  $n$  leaves that are in one-to-one correspondence with the elements of  $\mathcal{A}$ . In this case the universal covering  $\tilde{\mathcal{R}}_{\mathcal{A}}$  is canonically identified with the Cayley graph of  $F_N$  with respect to the generating system  $\mathcal{A}$ , and the edge-geodesic map  $i = i_{Q, \mathcal{A}}$  is uniquely determined by the choice of a *base point*  $P = i(\mathbf{1})$ , where  $\mathbf{1}$  is the vertex of the Cayley graph that corresponds to the neutral element  $1 \in F_N$ . In this case we denote the BBT-constant  $\text{BBT}(i_{P, \mathcal{A}})$  by  $\text{BBT}(\mathcal{A}, P)$ , and the volume of  $\mathcal{R}_{\mathcal{A}}$  by  $\text{vol}(\mathcal{A}, P)$ .

**Lemma 3.1.** *Let  $T$  be an  $\mathbb{R}$ -tree with isometric  $F_N$ -action that satisfies BBT. Let  $\mathcal{A}$  be any basis of  $F_N = F(\mathcal{A})$ , and let  $P$  be any point of  $T$ . Then the constants  $\text{BBT}(\mathcal{A}, P) \geq 0$  and  $\text{vol}(\mathcal{A}, P) > 0$  satisfy:*

(a) *For any reduced word  $w = x_1 x_2 \dots x_n$  in  $F(\mathcal{A})$  and any prefix  $v = x_1 x_2 \dots x_m$  of  $w$  the point  $vP$  is contained in the  $\text{BBT}(\mathcal{A}, P)$ -neighborhood of the geodesic segment  $[P, wP] \subset T$ .*

(b) *For any cyclically reduced word  $w$  in  $F(\mathcal{A})$  one has:*

$$d(wP, P) \leq 2\text{BBT}(\mathcal{A}, P) + \|w\|_T$$



(c) Any subword  $u$  of a cyclically reduced word  $w \in F(\mathcal{A})$  satisfies:

$$d(uP, P) \leq 2BBT(\mathcal{A}, P) + \|w\|_T$$

(d) Every  $x \in \mathcal{A} \cup \mathcal{A}^{-1}$  satisfies:

$$d(P, xP) \leq \text{vol}(\mathcal{A}, P)$$

*Proof of Lemma 3.1.* (a) We only need to observe that a reduced word in  $\mathcal{A}$  defines a geodesic segment in  $\tilde{\mathcal{R}}_{\mathcal{A}}$ .

(b) As  $w$  is cyclically reduced,  $wP$  and  $w^2P$  are contained in the  $BBT(\mathcal{A}, P)$ -neighborhood of the segment  $[P, w^3P]$ , and thus  $P$  and  $wP$  are contained in the  $BBT(\mathcal{A}, P)$ -neighborhood of the segment  $[w^{-1}P, w^2P]$ . As the axis  $Ax(w)$  contains  $[w^{-1}P, wP] \cap [P, w^2P]$ , and the latter is the fundamental domain with respect to the action of  $w$  on  $Ax(w)$  and hence has length  $\|w\|_T$ , the desired inequality follows.

(c) As in (b) we see that  $P$  and  $uP$  are contained in the  $BBT(\mathcal{A}, P)$ -neighborhood of  $[w^{-1}P, wP] \cap [P, w^2P]$  and thus of  $Ax(w)$ .

(d) This is a direct consequence of the above definition of the volume.  $\square$

It has been shown in [LL03], Remark 2.6, that

**Lemma 3.2.** *Let  $T$  be an  $\mathbb{R}$ -tree with a very small action with dense orbits of  $F_N$ . For any point  $P$  in  $T$ , there exists a sequence of bases  $\mathcal{A}_k$  such that the two sequences of constants  $BBT(P, \mathcal{A}_k)$  and  $\text{vol}(\mathcal{R}_{\mathcal{A}_k})$  both tend to 0, for  $k \rightarrow \infty$ .  $\square$*

## 4 The dual lamination associated to an $\mathbb{R}$ -tree

In this section and also some of the following we will assume familiarity of the reader with the concepts and also with the notation of [CHL-I], which we freely use without further explanation.

Let  $T$  be an  $\mathbb{R}$ -tree with isometric  $F_N$ -action. For every  $\varepsilon > 0$  we consider the set

$$\Omega_\varepsilon(T) = \{w \in F_N \mid \|w\|_T < \varepsilon\} \subset F_N$$

(which is invariant under conjugation and inversion in  $F_N$ ), and the  $F_N$ -action- and flip-invariant set

$$\Omega_\varepsilon^2(T) = \bigcup_{w \in \Omega_\varepsilon(T)} L^2(w)$$

in  $\partial^2 F_N$ . We note that either the  $F_N$ -action on  $T$  is free simplicial, i.e.  $T$  belongs to  $cv_N$ , or else  $\Omega_\varepsilon(T)$  and hence  $\Omega_\varepsilon^2(T)$  are non-empty for any  $\varepsilon > 0$ . In the latter case the closure  $L_\varepsilon^2(T)$  of  $\Omega_\varepsilon^2(T)$  in  $\partial^2 F_N$  is an algebraic lamination, and by Lemma 4.2 of [CHL-I] we can define:

**Definition 4.1.** Let  $T$  be an  $\mathbb{R}$ -tree on which  $F_N$  acts by isometries. If  $T$  belongs to  $cv_N$ , then we define  $L_\Omega^2(T)$  to be the empty set (which is not an algebraic lamination !). Otherwise we define the *dual algebraic lamination* associated to  $T$  as follows:

$$L_\Omega^2(T) = \bigcap_{\varepsilon > 0} L_\varepsilon^2(T)$$

We note that  $L_\Omega^2(T)$  depends only on the projective class  $[T]$ .

For any basis  $\mathcal{A}$  of  $F_N$  we define in a similar spirit, for any  $T$  which belongs to  $\partial cv_N$ , the laminary languages

$$\mathcal{L}_\mathcal{A}^\varepsilon(T) = \bigcup_{w \in \Omega_\varepsilon(T)} \mathcal{L}_\mathcal{A}(L^2(w))$$

and

$$\mathcal{L}_\mathcal{A}^\Omega(T) = \bigcap_{\varepsilon > 0} \mathcal{L}_\mathcal{A}^\varepsilon(T).$$

Hence  $u \in F(\mathcal{A})$  belongs to  $\mathcal{L}_\mathcal{A}^\Omega(T)$  if and only if for all  $\varepsilon > 0$  there exists a cyclically reduced word  $w \in F(\mathcal{A})$  with  $\|w\|_T < \varepsilon$  and an exponent  $m \geq 1$  such that  $u$  is a subword of  $w^m$ . However, it is obvious that one needs only to consider exponents which satisfy  $m \leq |u|_\mathcal{A}$  (= the word length of  $u$  in  $\mathcal{A}^{\pm 1}$ ). Thus we obtain as direct consequence:

**Remark 4.2.** A word  $u \in F(\mathcal{A})$  belongs to  $\mathcal{L}_\mathcal{A}^\Omega(T)$  if and only if for all  $\varepsilon > 0$  there exists a cyclically reduced word  $w \in F(\mathcal{A})$  with  $\|w\|_T < \varepsilon$  such that  $u$  is a subword of  $w$ .

It follows easily that these laminary languages correspond precisely to the algebraic laminations  $L_\varepsilon^2(T)$  and  $L_\Omega^2(T)$ , under the bijection  $\rho_{\mathcal{L}}^A \rho_{\mathcal{A}}^2$  established in Theorem 1.1 of [CHL-I]. Indeed, except for the passage from  $\Omega_\varepsilon^2(T)$  to  $L_\varepsilon^2(T)$ , i.e. closing up in  $\partial^2 F_N$ , the identity between the corresponding laminary languages is definitory. But as the language  $\mathcal{L}_{\mathcal{A}}^\varepsilon(T)$  is already laminary (see Remark 5.3 of [CHL-I]), it follows that it agrees with the laminary language  $\rho_{\mathcal{L}}^A \rho_{\mathcal{A}}^2(L_\varepsilon^2(T))$  associated canonically to the closure of  $\Omega_\varepsilon^2(T)$  in  $\partial^2 F_N$ .

## 5 One-sided infinite words

Let  $T$  be, as before, a real tree with a left action by isometries of the free group  $F_N$ . We fix a basis  $\mathcal{A}$  of  $F_N$  and a point  $P$  of  $T$ . The choice of the basis  $\mathcal{A}$  gives us an identification between the boundary  $\partial F_N$ , and the space  $\partial F(\mathcal{A})$  of (one-sided) infinite reduced words in  $\mathcal{A}^{\pm 1}$ .

Following [LL03] we denote by  $L_{\mathcal{A}}^1(T)$  the set of (one-sided) infinite reduced words  $X = x_1 x_2 \dots \in \partial F(\mathcal{A})$  such that for some  $P \in T$  the sequence  $(X_i P)_{i \in \mathbb{N}}$  is bounded, where  $X_i$  is the prefix of length  $i$  of  $X$ . We observe:

**Remark 5.1.** (1) If  $X$  belongs to  $L_{\mathcal{A}}^1(T)$ , then for *any*  $P \in T$  the sequence  $(X_i P)_{i \in \mathbb{N}}$  is bounded.

(2) If  $X$  does not belong to  $L_{\mathcal{A}}^1(T)$ , then for any  $C > 0$ , any  $P \in T$  and any integer  $K$  there exist  $l > k \geq K$  such that  $d(X_k P, X_l P) > C$ .

(3) If in addition  $T$  satisfies BBT, then for any  $X = x_1 x_2 \dots \in L_{\mathcal{A}}^1(T)$  there exists an integer  $K \geq 0$  such that for all  $k, l \geq K$  one has  $d(X_k P, X_l P) = d(x_{k+1} \dots x_l P, P) < 3BBT(\mathcal{A}, P)$ .

**Proposition 5.2 ([LL03]).** *The canonical identification  $\partial F(\mathcal{A}) = \partial F_N$  associates to the subset  $L_{\mathcal{A}}^1(T) \subset \partial F(\mathcal{A})$  a set  $L^1(T) \subset \partial F_N$  that does not depend on the choice of  $\mathcal{A}$  (which justifies the notation  $L^1(T)$ ).*

*Proof.* Let  $\mathcal{B}$  be another basis for  $F_N$  and  $X$  be in  $\partial F_N$ . Denote by  $X_{\mathcal{A}}$  and  $X_{\mathcal{B}}$  the corresponding (one-sided) infinite reduced words in  $\partial F(\mathcal{A})$  and  $\partial F(\mathcal{B})$ . The prefix sequences  $(X_{\mathcal{A},i})_{i \in \mathbb{N}}$  and  $(X_{\mathcal{B},i})_{i \in \mathbb{N}}$  (which are sequences of elements of  $F_N$ ) both converge to  $X$ . Geometrically they are two quasi-geodesics in  $F_N$ . If we fix the word metric  $d_{\mathcal{A}}$  on  $F_N$ , then the first sequence lies on a geodesic, and the second on a quasi-geodesic. In particular, as follows from Cooper's cancellation bound (see section 7 of [CHL-I]), the two

sequences have the property that for any positive integer  $j$  there exists a positive integer  $i$  such that  $d_{\mathcal{A}}(X_{\mathcal{B},j}, X_{\mathcal{A},i}) < C$ , where  $C = \text{BBT}(\mathcal{B}, \mathcal{A})$  is Cooper's cancellation bound between basis  $\mathcal{A}$  and  $\mathcal{B}$ , see Lemma 7.1 of [CHL-I] and the subsequent discussion.

This shows that the sequence of points  $X_{\mathcal{A},i}P$  is of bounded diameter if and only if the sequence of the  $X_{\mathcal{B},j}P$  is.  $\square$

Let us now state some properties of  $L^1(T)$ : Note first that the last proposition and thus the definition of  $L^1(T)$  does not require that  $T$  satisfies BBT. Furthermore, the subset  $L^1(T)$  of  $\partial F_N$  is  $F_N$ -invariant, and, unless it is empty, it is dense in  $\partial F_N$  (as is any  $F_N$ -invariant non-empty subset of  $\partial F_N$ ). The set  $L^1(T)$  is empty if and only if the action of  $F_N$  on  $T$  is free and discrete, i.e.  $T$  is a simplicial tree with quotient graph  $T/F_N$  that satisfies  $\pi_1(T/F_N) \cong F_N$ . Thus such a tree  $T$  defines a point of the (unprojectivized) Outer space  $\text{cv}_N$  and hence of the projectivized Outer space  $\text{CV}_N$  (see §2). In any case, we note that the set  $L^1(T)$  only depends on the projective class  $[T]$  of  $T$ .

From the algebraic lamination  $L_{\Omega}^2(T)$  defined in §4 we can alternatively deduce a second  $F_N$ -invariant subset of  $\partial F_N$  associated to  $T$ , namely

$$\pi(L_{\Omega}^2(T)) = \{X \in \partial F_N \mid (X, X') \in L_{\Omega}^2(T) \text{ for some } X' \in \partial F_N\}.$$

**Proposition 5.3.** *For any  $\mathbb{R}$ -tree  $T$  with isometric  $F_N$ -action that satisfies BBT, one has:*

$$\pi(L_{\Omega}^2(T)) \subset L^1(T)$$

*Proof.* Let  $\mathcal{A}$  be a basis of  $F_N$  and  $X = x_1x_2\dots$  a reduced infinite word in  $\partial F(\mathcal{A}) = \partial F_N$ . If  $X$  does not belong to  $L^1(T)$ , then it follows from Remark 5.1 (2) that for any point  $P \in T$ , for any  $C > 0$  and for any  $K > 0$ , there exist  $k, l$  with  $K \leq k \leq l$  such that  $d_{\mathcal{A}}(X_kP, X_lP) > C$  where  $X_i$  is the prefix of  $X$  of length  $i$ . It follows directly from Lemma 3.1 (c) that, choosing  $C$  large enough (depending on the given  $P$  and  $\mathcal{A}$ ), the word  $X_k^{-1}X_l$  cannot be a subword of any cyclically reduced word that represents an element  $w \in \Omega_{\varepsilon}(T)$ , for small  $\varepsilon > 0$ . Hence  $X_k^{-1}X_l$  and its inverse  $X_l^{-1}X_k$  do not belong to  $\mathcal{L}_{\mathcal{A}}^{\Omega}(T)$ .

But if there is some  $X' = x'_1x'_2\dots \in \partial F_N$  such that  $(X, X')$  belongs to  $L_{\Omega}^2(T)$ , then the reduced biinfinite word  $\rho_{\mathcal{A}}(X, X') = X^{-1}X' = \dots x_{j+2}^{-1}x_{j+1}^{-1} \cdot x'_{j+1}x'_{j+2}\dots$  in the symbolic lamination  $\rho_{\mathcal{A}}^2(L_{\Omega}^2(T)) = L_{\mathcal{A}}^{\Omega}(T) \subset \Sigma_{\mathcal{A}}$  associated to  $L_{\Omega}^2(T)$  (compare Proposition 4.4 of [CHL-I]) will contain any of these

$X_l^{-1}X_k = x_l^{-1} \dots x_{k+1}^{-1}$  with sufficiently large  $k < l$  as subword, in contradiction to the statement at the end of the last paragraph. This shows that  $X$  cannot belong to  $\pi(L_\Omega^2(T))$ .  $\square$

The converse inclusion with respect to Proposition 5.3 does not hold in general, as will be seen in section 7. In fact, one has to regard  $L^1(T)$  as a finer invariant of  $T$  than the algebraic lamination  $L_\Omega^2(T)$ , which only depends on a weakened version of the topology of  $T$ , compare [CHL05], while  $L^1(T)$  may change when different  $\mathbb{R}$ -tree structures are varying on a given topological tree  $T$ . The fact that one can derive  $L_\Omega^2(T)$  from  $L^1(T)$  will be shown below: it is a direct consequence of Proposition 5.6.

For any basis  $\mathcal{A}$  of  $F_N$  and any set  $S \subset \partial F_N$  we denote by  $\mathcal{L}_\mathcal{A}^\infty(S)$  the set of words  $u \in F(\mathcal{A})$  and their inverses such that  $u$  occurs infinitely often as subword in some of the reduced infinite words  $X_\mathcal{A}$  that represent an element  $X \in S$  (we say  $u$  is *recurrent* in  $X_\mathcal{A}$ ). It is obvious that for any non-empty  $S$  the language  $\mathcal{L}_\mathcal{A}^\infty(S)$  is laminary (because we artificially added to  $\mathcal{L}_\mathcal{A}^\infty(S)$  the inverses of any recurrent  $u$ ): we call it the *recurrent laminary language in  $\mathcal{A}^{\pm 1}$  associated to  $S$* .

**Definition 5.4.** For any basis  $\mathcal{A}$  of  $F_N$  and any non-empty set  $S \subset \partial F_N$  let  $L_\infty^2(S)$  the algebraic lamination defined by the recurrent language in  $\mathcal{A}^{\pm 1}$  associated to  $S$ :

$$L_\infty^2(S) = L^2(\mathcal{L}_\mathcal{A}^\infty(S))$$

Here (and below) we denote by  $L^2(\mathcal{L})$  the algebraic lamination  $(\rho_\mathcal{L}^A \rho_\mathcal{A}^2)^{-1}(\mathcal{L})$  defined in [CHL-I] for any laminary language  $\mathcal{L}$ .

Proposition 5.6 below justifies the absence of mentioning explicitly the basis  $\mathcal{A}$  in the notation  $L_\infty^2(S)$ . But first we observe:

For any basis  $\mathcal{A}$  of  $F_N$ , any  $k \in \mathbb{N}$  and any  $w \in F(\mathcal{A})$ , we denote by  $w \dagger_k \in F(\mathcal{A})$  the subword of  $w$  obtained from chopping off the initial and finite subword of length  $k$ . For any second basis  $\mathcal{B}$  of  $F_N$  consider Cooper's cancellation bounds  $C = \text{BBT}(\mathcal{B}, \mathcal{A})$  and  $C' = \text{BBT}(\mathcal{A}, \mathcal{B})$ , as well as  $C'' = C' + \lambda C$ , where  $\lambda$  is the maximal length of the elements of  $\mathcal{A}$  written as words in  $\mathcal{B}^{\pm 1}$ .

**Lemma 5.5.** *Consider a word  $u_\mathcal{B} \in F(\mathcal{B})$  in  $\mathcal{B}^{\pm 1}$ , and let  $u_\mathcal{A} \in F(\mathcal{A})$  be the word in  $\mathcal{A}^{\pm 1}$  representing the same element of  $F_N$ . Let  $v_\mathcal{A} = u_\mathcal{A} \dagger_C$ , and let  $w_\mathcal{A} \in F(\mathcal{A})$  be any word that contains  $v_\mathcal{A}$  as a subword. Let  $w_\mathcal{B} \in F(\mathcal{B})$  be*

the word in  $\mathcal{B}^{\pm 1}$  representing the same element of  $F_N$  as  $w_{\mathcal{A}}$ . Then  $u_{\mathcal{B}\dagger_{C''}}$  is a subword of  $w_{\mathcal{B}\dagger_{C'}}$ .

*Proof.* This follows from a straight forward calculation.  $\square$

**Proposition 5.6.** *Let  $S$  be a non-empty subset of  $\partial F_N$ , and  $\mathcal{A}$  and  $\mathcal{B}$  be two basis of  $F_N$ . Then one has:*

$$L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(S)) = L^2(\mathcal{L}_{\mathcal{B}}^{\infty}(S))$$

*Proof.* It suffices to prove for any  $Y \in S$  the equality

$$L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(Y)) = L^2(\mathcal{L}_{\mathcal{B}}^{\infty}(Y)).$$

Thus, for any  $(X, X')$  in  $L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(Y))$ , we have to prove that the biinfinite word  $Z_{\mathcal{B}} = \rho_{\mathcal{B}}(X, X')$  in  $\mathcal{B}^{\pm 1}$  is contained in the symbolic lamination  $L(\mathcal{L}_{\mathcal{B}}^{\infty}(Y)) = (\rho_{\mathcal{L}}^{\mathcal{B}})^{-1}(\mathcal{L}_{\mathcal{B}}^{\infty}(Y))$ . Let  $u_{\mathcal{B}} \in F(\mathcal{B})$  be a finite subword of  $Z_{\mathcal{B}}$ , and let  $u_{\mathcal{A}} \in F(\mathcal{A})$  be the word in  $\mathcal{A}^{\pm 1}$  representing the same element of  $F_N$  as  $u_{\mathcal{B}}$ . Using Cooper's cancellation bound  $C = \text{BBT}(\mathcal{B}, \mathcal{A})$ , we see that  $v_{\mathcal{A}} = u_{\mathcal{A}\dagger_C}$  is a subword of the biinfinite word  $Z_{\mathcal{A}} = \rho_{\mathcal{A}}(X, X')$ . By definition of  $L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(Y))$ ,  $v_{\mathcal{A}}$  is recurrent in the infinite word  $Y_{\mathcal{A}} \in \partial F(\mathcal{A})$  representing  $Y$ . Let  $w_{\mathcal{A}}$  be a subword of  $Y_{\mathcal{A}}$  such that  $v_{\mathcal{A}}$  is a subword of  $w_{\mathcal{A}}$ , sufficiently far from the beginning and the end of  $w_{\mathcal{A}}$ .

Let  $w_{\mathcal{B}} \in F(\mathcal{B})$  be the word in  $\mathcal{B}^{\pm 1}$  representing the same element of  $F_N$  as  $w_{\mathcal{A}}$ . Using Cooper's cancellation bound  $C' = \text{BBT}(\mathcal{A}, \mathcal{B})$ , we get that  $w_{\mathcal{B}\dagger_{C'}}$  is a subword of the infinite word  $Y_{\mathcal{B}} \in \partial F(\mathcal{B})$  representing  $Y$ , and that  $u_{\mathcal{B}\dagger_{C''}}$  is a subword of  $w_{\mathcal{B}\dagger_{C'}}$ , for  $C'' > 0$  depending only on  $\mathcal{A}$  and  $\mathcal{B}$  as specified in Lemma 5.5. This proves that  $u_{\mathcal{B}\dagger_{C''}}$  is recurrent in  $Y_{\mathcal{B}}$ . As  $u_{\mathcal{B}} \in \mathcal{L}(Z_{\mathcal{B}})$  was chosen arbitrarily and  $\mathcal{L}(Z_{\mathcal{B}})$  is a laminary language, this proves that the biinfinite word  $Z_{\mathcal{B}} = \rho_{\mathcal{B}}(X, X')$ , for any  $(X, X') \in L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(Y))$ , is contained in the symbolic lamination  $L(\mathcal{L}_{\mathcal{B}}^{\infty}(Y))$ . Thus we have

$$L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(Y)) \subset L^2(\mathcal{L}_{\mathcal{B}}^{\infty}(Y)),$$

which proves the proposition.  $\square$

**Definition 5.7.** For an  $\mathbb{R}$ -tree  $T$  and a basis  $\mathcal{A}$  of  $F_N$ , we define  $L_{\infty}^2(T)$  to be the empty set if  $T$  belongs to  $cv_N$ . Otherwise  $L_{\infty}^2(T)$  is the algebraic lamination defined by the recurrent language associated to  $L^1(T)$ :

$$L_{\infty}^2(T) = L^2(\mathcal{L}_{\mathcal{A}}^{\infty}(L^1(T)))$$

It follows from Proposition 5.2 and Proposition 5.6 that  $L_{\infty}^2(T)$  does not depend on the choice of the basis  $\mathcal{A}$ .

**Proposition 5.8.** *Let  $T$  be an  $\mathbb{R}$ -tree with very small action of  $F_N$  with dense orbits. The lamination  $L_\Omega^2(T)$  of Definition 4.1 and the lamination  $L_\infty^2(T)$  of Definition 5.7 are equal:*

$$L_\Omega^2(T) = L_\infty^2(T)$$

*Proof.* Let  $\mathcal{A}$  be a basis of  $F_N$ . We will prove that the laminary languages associated to these laminations (via the canonical map  $\rho_{\mathcal{L}}^{\mathcal{A}}\rho_{\mathcal{A}}^2 : \Lambda^2(F_N) \rightarrow \Lambda_{\mathcal{L}}(\mathcal{A})$  from [CHL-I]) are equal.

We first prove that  $\mathcal{L}_{\mathcal{A}}^\infty(L^1(T)) \subset \mathcal{L}_{\mathcal{A}}^\Omega(T)$ : Let  $X = x_1x_2\dots$  be a reduced infinite word in  $\mathcal{A}^{\pm 1}$  that belongs to  $L_{\mathcal{A}}^1(T)$ . We know that, according to the above definition of the associated recurrent laminary language, if  $X_{k,l} = x_k\dots x_l \in \mathcal{L}_{\mathcal{A}}^\infty(X)$ , then there are arbitrary large  $k', l'$  with  $X_{k,l} = X_{k',l'}$ . Hence Cooper's cancellation bound, see §7 of [CHL-I], assures us that, when writing  $X$  as a reduced word in the basis  $\mathcal{B}$ , say  $X_{\mathcal{B}} = y_1y_2\dots$ , there is a recurrent subword  $Y_{r,s} = y_r\dots y_s$  of  $X_{\mathcal{B}}$  which has the property that, when written as word  $Y_{\mathcal{A}}$  in  $\mathcal{A}^{\pm 1}$ , the latter contains  $X_{k,l}$  as subword. On the other hand, since  $\mathcal{B}$  contains at least two elements (by the general assumption that the rank  $N$  of  $F_N$  is at least 2), there is a  $y \in \mathcal{B}^{\pm 1}$  such that  $Y_{r,s}y$  is cyclically reduced. Hence we obtain from Remark 5.1 (3), from Lemma 3.1 (d) and from the assumption  $\text{BBT}(\mathcal{B}, P) < \varepsilon$  and  $\text{vol}(\mathcal{B}, P) < \varepsilon$  that  $\|Y_{r,s}y\|_T \leq d(P, Y_{r,s}P) + d(P, yP) \leq 4\text{BBT}(\mathcal{B}, P) < 4\varepsilon$ . But for  $s - r$  sufficiently large, the subword  $X_{k,l}$  of  $Y_{\mathcal{A}}$  will not be cancelled when  $Y_{r,s}y$  is written in  $\mathcal{A}^{\pm 1}$  and subsequently cyclically reduced. This implies that  $X_{k,l}$  belongs to  $\mathcal{L}_{\mathcal{A}}^{4\varepsilon}(T)$ , which proves the assertion.

We now turn to the proof of the converse inclusion, namely  $\mathcal{L}_{\mathcal{A}}^\Omega(T) \subset \mathcal{L}_{\mathcal{A}}^\infty(L^1(T))$ : For any word  $w$  in  $F(\mathcal{A})$  we distinguish between its conjugating part  $v \in F(\mathcal{A})$  and its cyclically reduced part  $w' \in F(\mathcal{A})$ , where  $w = vw'v^{-1}$  in reduced form, and with  $w'$  cyclically reduced.

Let  $u \in \mathcal{L}_{\mathcal{A}}^\Omega(T)$  be a word in  $\mathcal{A}^{\pm 1}$ ,  $P$  a point in  $T$  and  $\varepsilon > 0$ . We first want to show that there exists a word  $w \in F(\mathcal{A})$  that contains  $u$  as a subword of its cyclically reduced part and satisfies  $d(P, wP) < \varepsilon$ .

Indeed, by definition of  $\mathcal{L}_{\mathcal{A}}^\Omega(T)$  there exists a cyclically reduced word  $w$  in  $F(\mathcal{A})$  of which  $u$  is a subword and such that  $\|w\|_T < \varepsilon/6$ . Then  $u$  is a subword of any cyclic conjugate of  $w^2$ . As the action of  $F_N$  on  $T$  has dense orbits there exists a word  $v$  of  $F(\mathcal{A})$  such that

$$d(v^{-1}P, \text{Ax}(w)) < \frac{\varepsilon}{3}.$$

An easy calculation then shows that  $vw^2v^{-1}$  satisfies  $d(P, vw^2v^{-1}P) < \varepsilon$ , which is what we claimed.

Thus there exists a sequence of words  $u_k \in F(\mathcal{A})$  where each of them contains  $u$  or  $u^{-1}$  as a subword of its cyclically reduced part, such that  $d(P, u_k P) < \frac{1}{2^k}$ . We apply Lemma 5.9 stated below to obtain a sequence of  $w_n = u_{k_n}^{d_n}$ , with  $d_n = \pm 1$ , and with the further property that in each of the products  $w_n w_{n+1}$  the cancellation in  $w_n w_{n+1}$  does not go further than the conjugating parts of  $w_n$  and  $w_{n+1}$ . Then  $X = w_1 w_2 w_3 \cdots$  is a word in  $L_{\mathcal{A}}^1(T)$  in which  $u$  or  $u^{-1}$  is recurrent.

This concludes the proof of Proposition 5.8, modulo the proof of the subsequent lemma.  $\square$

**Lemma 5.9.** *Let  $(u_k)_{k \in \mathbb{N}}$  be a sequence of words from  $F(\mathcal{A})$ . Then there is an infinite subsequence  $(u_{k_n})_{n \in \mathbb{N}}$  and exponents  $d_n = \pm 1$ , such that the sequence of words  $w_n = u_{k_n}^{d_n}$  possesses the additional property that in each of the products  $w_n w_{n+1}$  the cancellation in  $w_n w_{n+1}$  does not exceed the conjugating parts of either  $w_n$  or  $w_{n+1}$ .*

*Proof.* If the given words  $u_k$  are almost all cyclically reduced, then we can build the sequence  $w_n$  by choosing inductively  $w_{n+1} = u_{k_{n+1}}$  or  $w_{n+1} = u_{k_{n+1}}^{-1}$  according to the previous choice to avoid any cancellation in  $w_n w_{n+1}$ .

If the given sequence  $(u_k)_{k \in \mathbb{N}}$  contains a subsequence of words  $u_{k_n}$  with bounded length of their conjugating part, then there exists a subsequence with constant conjugating part and we can use the previous construction.

In the remaining case there exists a subsequence of words  $u_{k_n}$  with strictly increasing length of their conjugating part. Assume that  $w_1, \dots, w_n$  were already chosen,  $w_i = u_{k_i}^{d_i}$  with  $d_i = \pm 1$ , and with the property that the cancellation in  $w_i w_{i+1}$  is not more than the conjugating part of  $w_i$  and strictly less than the conjugating part of  $w_{i+1}$ . Then replacing the last word  $w_n$  by its inverse  $w_n^{-1}$  does not change this property. If the cancellation in  $w_n u_{k_{n+1}}$  is bigger than the conjugating part of  $w_n$  we replace  $w_n$  by its inverse  $w_n^{-1}$ . It follows that the cancellation in both  $w_n u_{k_{n+1}}$  and  $w_n u_{k_{n+1}}^{-1}$  is then not more than the conjugating part of  $w_n$  and strictly less than the conjugating part of  $u_{k_{n+1}}$ , as the length of the latter is strictly bigger than that of the conjugating part of  $w_n$  (by our original “strictly increasing” condition for this case).  $\square$

**Remark 5.10.** The precise relationship between the various  $F_N$ -invariant sets of one-sided infinite words associated to a lamination or to an  $\mathbb{R}$ -tree is rather intricate, and it seems difficult to express the algebraic lamination



associated to an  $\mathbb{R}$ -tree properly in terms of such a set. An attempt, however confusing or misleading it may be, is made in the subsequent paragraph:

For any algebraic lamination  $L^2$  we denote by  $L_\infty^1(L^2) \subset \partial F_N$  the set of all infinite words  $X = x_1 x_2 \dots$ , in some basis  $\mathcal{A}$  of  $F_N$ , which have their associated recurrent laminary language contained in  $\mathcal{L}_{\mathcal{A}}(L^2)$ . Very possibly one can use Cooper's cancellation bound (or perhaps a variation of Proposition 5.6) to show that  $L_\infty^1(L^2)$  does not depend on the choice of  $\mathcal{A}$ . We believe that Proposition 5.3 can be extended to show that  $\pi(L^2)$  is always a subset of  $L_\infty^1(L^2)$ . The converse, however, seems in general to be wrong: For example, for  $\mathbb{R}$ -tree laminations  $L_\Omega^2(T)$ , where  $T$  satisfies BBT and has dense orbits, we know that both of the inclusions  $\pi(L_\Omega^2(T)) \subset L^1(T) \subset L_\infty^1(L_\Omega^2(T))$  hold, but we strongly suspect that, for any such  $T$ , they both are proper inclusions. On the other hand, we have seen above that the three recurrent laminary languages associated to these three  $F_N$ -invariant sets are equal.

## 6 Bounded Back Tracking property and the map $\mathcal{Q}$

Throughout this section we assume that  $T$  satisfies the property BBT (see §2), which follows for example if  $T$  is small. It is an easy exercise (compare [GJLL98]) to show that BBT ensures that every element  $X \in \partial F_N \setminus L^1(T)$  determines, through picking any point  $P \in T$  and any sequence of elements  $X_i \in F_n$  that converges towards  $X$ , a well defined point

$$\mathcal{Q}(X) = \lim_{i \rightarrow \infty} X_i P$$

of the Gromov boundary  $\partial T$  of  $T$ .

We suppose from now on that  $T$  is an  $\mathbb{R}$ -tree with very small  $F_N$ -action with dense orbits, and that for some (arbitrary) point  $P \in T$  one has given a sequence of bases  $\mathcal{A}_k$  of  $F_N$  that satisfies the properties assured by Lemma 3.2: Both  $vol(\mathcal{R}_{\mathcal{A}_k})$  and  $BBT(\mathcal{A}_k)$  tend to 0, for  $k \rightarrow \infty$ .

For any infinite reduced word  $X = x_1 x_2 \dots \in \partial F(\mathcal{A}_k)$  that represents an element of  $L^1(T)$ , the sequence of points  $(x_1 \dots x_i P)_{i \in \mathbb{N}}$  eventually stays in a bounded region  $R(X, k)$  of diameter  $3BBT(\mathcal{A}_k)$  (compare Remark 5.1 (c)), so that we can associate to  $X$  a well defined point  $\mathcal{Q}(X) = \lim_{k \rightarrow \infty} R(X, k)$ . It has been shown in [LL03] that  $\mathcal{Q}(X)$  depends only on  $X \in \partial F_N$  and not

on the above choice of  $P$  and of the  $\mathcal{A}_k$ . It is important to note that  $\mathcal{Q}(X)$  may well be contained in the metric completion  $\overline{T}$  of  $T$ , but not in  $T$  itself.

Alternative definitions of the point  $\mathcal{Q}(X)$ , for any  $X \in \partial F_N$ , which do not need to consider an infinite change of bases of  $F_N$ , have been given in [LL] and in [LL03], Lemma 3.4:

**Lemma 6.1.** *For all  $X$  in  $\partial F_N$ , for any sequence of points  $X_i \in F_N$  which converge towards  $X$ , and for any point  $P$  of  $T$ , one has in  $\overline{T} \cup \partial T$ :*

$$[P, \mathcal{Q}(X)] = \overline{\bigcup_{n \in \mathbb{N}} \bigcap_{i \geq n} [P, X_i P]}$$

**Lemma 6.2.** *For all  $X$  in  $\partial F_N$  and for all  $P$  in  $T$ , the point  $\mathcal{Q}(X)$  is the only point of  $\overline{T} \cup \partial T$  such that there exists a sequence of elements  $X_i \in F_n$  which converge towards  $X$  and a point  $P$  in  $T$  such that the points  $X_i P$  converge to  $\mathcal{Q}(X)$ .*

From Lemma 6.1 and Remark 5.1 (3) it follows directly:

**Lemma 6.3.** *Let  $P$  be a point in an  $\mathbb{R}$ -tree  $T \in \partial cv_N$  with dense  $F_N$ -orbits, and let  $\mathcal{A}$  be a basis of  $F_N$ . Then for every  $X = x_1 x_2 \dots \in L^1_{\mathcal{A}}(T)$  there exists a bound  $K(X) \geq 0$  such that for every  $X_k = x_1 \dots x_k$  with  $k \geq K$  one has*

$$d(X_k P, \mathcal{Q}(X)) < 3 \text{BBT}(\mathcal{A}, P).$$

□

Summarizing the above discussion, we observe that for every very small  $\mathbb{R}$ -tree  $T$  with dense orbits every boundary point  $X \in \partial F_N \setminus L^1(T)$  determines a point  $\mathcal{Q}(X) \in \partial T$ , while  $X \in L^1(T)$  determines a point  $\mathcal{Q}(X)$  in  $T$  or in its metric completion  $\overline{T}$ . This defines a map

$$\mathcal{Q} : \partial F_N \rightarrow \overline{T} \cup \partial T, X \mapsto \mathcal{Q}(X)$$

which is  $F_N$ -equivariant and surjective [LL03], but it is injective and continuous (with respect to the canonical boundary topologies) only on  $\partial F_N \setminus L^1(T)$ . In particular, on  $L^1(T)$  the map  $\mathcal{Q}$  is not continuous with respect to the metric on  $T$  (it does though possess on  $L^1(T)$  a kind of lower semi-continuous property, see Proposition 3.8 of [LL03]). In [CHL05] it is proved that  $\mathcal{Q}$  is continuous if we replace the metric topology on  $T$  by the weaker *observer's*

*topology* (this is the topology for which a basis of open subsets is given by connected components of  $T \setminus \{P\}$  for all points  $P$  of  $T$ ).

The basic phenomenon for the lack of continuity of the map  $\mathcal{Q}$  is illustrated as follows: If  $X_k$  is a converging sequence of elements from  $\partial F_N$  with the property that for some point  $Q \in T$  the segments  $[Q, \mathcal{Q}(X_k)]$  have pairwise intersections of lengths converging to 0, then  $X = \lim X_k$  satisfies  $Q = \mathcal{Q}(X)$ , while the lengths of the segments  $[Q, \mathcal{Q}(X_k)]$  may well not converge to 0 or even converge to  $\infty$ .

However, one can prove that the map  $\mathcal{Q}$  has the “closed graph property” (which will not be used in the sequel):

**Remark 6.4.** Let  $T$  be an  $\mathbb{R}$ -tree in  $\partial cv_N$  which has dense  $F_N$ -orbits, and, consider a sequence of boundary points  $X_k \in \partial F_N$  that converge to some  $X \in \partial F_N$ . Assume that the image points  $\mathcal{Q}(X_k) \in \overline{T} \cup \partial T$  converge to a point  $R \in \overline{T} \cup \partial T$ . Then one has:

$$R = \mathcal{Q}(X).$$

## 7 Geodesic lamination on a surface

To gain some geometric intuition, let us consider in this section the special case of an  $\mathbb{R}$ -tree that is dual to a measured lamination in a surface: As in §3 of [CHL-I] we denote by  $S$  a surface with non-empty boundary and with negative Euler characteristic, provided with a hyperbolic structure. The latter is given by an identification of the universal covering  $\tilde{S}$  with a convex part of the hyperbolic plane  $\mathbb{H}^2$ , which realizes the deck transformation action of  $F_N = \pi_1 S$  on  $\tilde{S}$  by hyperbolic isometries. Then any geodesic lamination  $\mathcal{L}$  on  $S$  defines, by taking the full preimage, a geodesic lamination  $\tilde{\mathcal{L}}$  in  $\tilde{S} \subset \mathbb{H}^2$ , on which  $F_N = \pi_1 S$  acts canonically.

There is a canonical *dual tree*  $T_{\mathcal{L}}$  with  $F_N$ -action by homeomorphisms associated to  $\tilde{\mathcal{L}}$ , associating to every non-boundary leaf of  $\tilde{\mathcal{L}}$  a point of  $T_{\mathcal{L}}$  which is not a branch point, and to the closure of any complementary component of  $\tilde{\mathcal{L}}$  in  $\tilde{S}$  a branch point of  $T_{\mathcal{L}}$ . This association is a bijective and can be made continuous: If the lamination  $\mathcal{L}$  is finite, then  $T_{\mathcal{L}}$  is simplicial, so that there is no ambiguity. If  $\mathcal{L}$  is infinite, then defining properly the topology of  $T_{\mathcal{L}}$  is much more delicate; we refer the interested reader to [CHL05] where this problem has been dealt with properly.

We now assume that the lamination  $\mathfrak{L}$  is provided with a transverse measure  $\mu$  (see [FLP91]). Then the lift  $\tilde{\mu}$  of  $\mu$  to  $\tilde{\mathfrak{L}}$  gives rise to a metric on  $T_{\mathfrak{L}}$  by defining for any points  $x, y \in T_{\mathfrak{L}}$ , corresponding to leaves  $l_x, l_y \in \tilde{\mathfrak{L}}$ , the distance  $d(x, y) = \tilde{\mu}(\alpha)$ , where  $\alpha$  is any arc in  $\tilde{S}$  with one endpoint on  $l_x$  and the other on  $l_y$ , and  $\alpha$  is assumed to be geodesic and hence transverse to  $\tilde{\mathfrak{L}}$ . This makes  $T_{\mathfrak{L}}$  into an  $\mathbb{R}$ -tree  $T_{\mu}$  with isometric  $F_N$ -action. It is noteworthy that, in the exceptional but fascinating case where  $\mathfrak{L}$  is not uniquely ergodic, projectively different transverse measures  $\mu$  will produce projectively distinct  $\mathbb{R}$ -trees, and that the simplex of projective measures on  $\mathfrak{L}$  gives rise to an analogous simplex of  $\mathbb{R}$ -trees in  $\partial CV_n$ .

We now consider an arbitrary point  $Q$  on the boundary  $\partial\tilde{S} \subset S_{\infty}^1 = \partial\mathbb{H}^2$ , where  $\partial\tilde{S}$  also coincides via our above identifications with the Gromov boundary  $\partial F_N$ . Let  $\beta$  be the geodesic arc which connects some arbitrary chosen point  $P$  in  $\tilde{S}$  to  $Q$ . We distinguish 3 cases:

1.  $Q$  is the endpoint of a leaf  $l$  of  $\tilde{\mathfrak{L}}$ . Then  $\tilde{\mu}(\beta)$  is finite. In fact,  $\beta$  projects to a segment in  $T_{\mu}$  of length  $\tilde{\mu}(\beta)$ . Denote by  $\hat{Q} \in T_{\mu}$  the image of  $Q$  under this projection.
2. The measure  $\tilde{\mu}(\beta)$  is infinite. Then  $\beta$  projects to an infinite arc in  $T_{\mu}$ , and  $Q$  defines a point  $\hat{Q}$  in the Gromov boundary  $\partial T_{\mu}$  of  $T_{\mu}$  (which is independent of the choice of  $\beta$ ).
3. In the remaining case the point  $Q$  defines a point  $\hat{Q}$  in the metric completion  $\bar{T}_{\mu}$  of  $T_{\mu}$ , and the arc  $\beta$  projects to a finite open arc in  $T_{\mu}$  which becomes closed when adding the point  $\hat{Q}$ .

In each of the three cases the geometrically described point  $\hat{Q}$  is precisely the image  $\mathcal{Q}(Q)$  of the point  $Q$ , if  $Q$  is viewed as element of  $\partial F_N$  via the above identification  $\partial\tilde{S} = \partial F_N$ .

We would like to note that this third class is non-empty for many (probably every non-rational) laminations  $\mathfrak{L}$ : For example, if  $\mathfrak{L}$  is the contracting (or expanding) lamination fixed by a pseudo-Anosov automorphism  $\varphi$  of  $S$ , then it suffices to consider a lift  $\tilde{\varphi}$  of  $\varphi$  to  $\tilde{S}$  that does not fix any leaf (or permute finitely many leaves) of  $\tilde{\mathfrak{L}}$ . The repulsive fixed point of  $\tilde{\varphi}$  on  $\partial\tilde{S}$  will then define such a point  $Q$ . Note that the existence of such lifts  $\tilde{\varphi}$  of  $\varphi$  has been proved in [LL00].

The distinction of these three cases is illuminating in that it shows that the set  $L^1(T_{\mu})$ , given here by the cases 1 and 3, may well be strictly larger

than the  $F_N$ -invariant set  $\pi(L^2(T_\mu)) \subset \partial F_N$  canonically associated to  $L^2(T_\mu)$ , given here by case 1. Indeed, while  $L^2(T_\mu)$  (and accordingly, the occurrence of the case 1 above) depends only on  $\mathfrak{L}$  and not on  $\mu$ , we do not know whether (but suspect that) the partition of the complement into cases 2 and 3 may actually depend on  $\mu$ .

The distinction of  $L^1(T_\mu)$  into cases 1 and 3 was the original motivation behind the definition given in this paper of the algebraic lamination  $L^2(T)$ , as is explained more directly by the content of the next section.

## 8 The lamination $L_{\mathcal{Q}}(T)$ and the map $\mathcal{Q}^2$

By restricting the domain and the range of the map  $\mathcal{Q}$  introduced in section 6, one obtains the map:

$$\mathcal{Q}^1 : L^1(T) \rightarrow \overline{T}, X \mapsto \mathcal{Q}(X)$$

Recall from §6 that the map  $\mathcal{Q}^1$  is surjective (see [LL03]), but in general not injective. Unless  $\mathcal{Q}(X)$  has a non-trivial stabilizer in  $F_N$ , the map  $\mathcal{Q}^1$  is conjectured to be finite to one (see Remark 3.6 of [LL03]).

**Definition 8.1.** To every very small  $\mathbb{R}$ -tree  $T$  with dense orbits we associate the following  $F_N$ - and flip-invariant subset of  $\partial^2 F_N$ :

$$L_{\mathcal{Q}}^2(T) = \{(X, X') \in \partial^2 F_N \mid \mathcal{Q}(X) = \mathcal{Q}(X')\}$$

Note that, as  $\mathcal{Q}$  is injective on  $\partial F_N \setminus L^1(T)$ , for  $(X, X')$  in  $L_{\mathcal{Q}}^2(T)$  one has that  $X$  and  $X'$  belong to  $L^1(T)$ . Equivalently, we know that  $\mathcal{Q}(X) = \mathcal{Q}(X')$  lies in  $\overline{T}$  and not in  $\partial T$ .

**Definition 8.2.** We define a map

$$\mathcal{Q}^2 : L_{\mathcal{Q}}^2(T) \rightarrow \overline{T}, (X, X') \mapsto \mathcal{Q}(X) = \mathcal{Q}(X')$$

which is  $F_N$ -equivariant and flip-invariant.

Just as remarked in §6 for the map  $\mathcal{Q} : \partial F_N \rightarrow \overline{T} \cup \partial T$ , the above map  $\mathcal{Q}^1 : L^1(T) \rightarrow \overline{T}$  is in general not continuous. The relevance of the set  $L_{\mathcal{Q}}^2(T)$  is underlined by the following:

**Proposition 8.3.** *Let  $T$  be an  $\mathbb{R}$ -tree with a very small action of  $F_N$  with dense orbits. Then the subset  $L_{\mathcal{Q}}^2(T)$  of  $\partial^2 F_N$  is closed, and the map  $\mathcal{Q}^2 : L_{\mathcal{Q}}^2(T) \rightarrow \overline{T}$  is continuous (for the metric topology on  $\overline{T}$ ).*

*Proof.* Let  $(X_n, Y_n)_{n \in \mathbb{N}}$  be a sequence of points from  $L_{\mathcal{Q}}^2(T)$  that converge to  $(X, Y) \in \partial^2 F_N$ . Let  $\mathcal{A}$  be a basis of  $F_N$ ,  $P$  be any point in  $T$ , and define  $C = 3\text{BBT}(\mathcal{A}, P)$ .

Let  $x_n$  the largest common prefix of the infinite reduced words in  $\mathcal{A}^{\pm 1}$  representing  $X_n$  and  $X$ ,  $y_n$  that of  $Y_n$  and  $Y$ , and  $h$  that of  $X$  and  $Y$ . From the assumption  $(X, Y) \in \partial^2 F_N$  we know that  $X$  and  $Y$  are different, so that  $h$  is a finite word:  $h \in F(\mathcal{A})$ . The assumption  $(X_n, Y_n) \xrightarrow{n \rightarrow \infty} (X, Y)$  implies that the  $X_n$  and the  $Y_n$  converge to  $X$  and  $Y$  respectively. Hence, for  $n$  big enough one obtains that  $h$  is a prefix of both,  $x_n$  and  $y_n$ . Indeed, since  $h$  is the longest common prefix of  $X$  and  $Y$ , it must also be that of  $X_n$  and  $Y_n$ . By hypothesis one has  $\mathcal{Q}(X_n) = \mathcal{Q}(Y_n)$ , so that the BBT property together with Lemma 6.1 ensures that  $hP$  lies in a  $C$ -neighborhood of  $\mathcal{Q}(X_n) = \mathcal{Q}(Y_n)$ . But then, by the definition of  $L^1(T)$ , the hypothesis  $X_n \xrightarrow{n \rightarrow \infty} X$  and  $Y_n \xrightarrow{n \rightarrow \infty} Y$  implies that  $X$  and  $Y$  belong to  $L^1(T)$ . Hence Lemma 6.3 shows that  $\mathcal{Q}(X)$  as well as  $\mathcal{Q}(Y)$  are contained in a  $2C$ -neighborhood of  $hP$ . Hence passing over to  $P$  and  $\mathcal{A}$  with arbitrary small  $\text{BBT}(\mathcal{A}, P)$  proves that  $\mathcal{Q}(X_n) = \mathcal{Q}(Y_n)$  converges to  $\mathcal{Q}(X) = \mathcal{Q}(Y)$ .  $\square$

As a direct consequence of Proposition 8.3 we obtain:

**Corollary 8.4.** *The set  $L_{\mathcal{Q}}^2(T) \subset \partial^2 F_N$  is an algebraic lamination.*

We remark that, contrary to the surface lamination case, the lamination  $L_{\mathcal{Q}}^2(T)$  contains all *diagonal leaves*: From the definition it follows directly that, if  $(X, X')$  and  $(X', X'')$  are in  $L_{\mathcal{Q}}^2(T)$  and  $X \neq X''$ , then  $(X, X'')$  is also in  $L_{\mathcal{Q}}^2(T)$ .

**Proposition 8.5.** *Let  $T$  be an  $\mathbb{R}$ -tree with a very small  $F_N$ -action with dense orbits. Then the lamination  $L_{\Omega}^2(T)$  of Definition 4.1 and the lamination  $L_{\mathcal{Q}}^2(T)$  of Definition 8.1 satisfy:*

$$L_{\Omega}^2(T) = L_{\mathcal{Q}}^2(T)$$

*Proof.* In order to show the inclusion  $L_{\mathcal{Q}}^2(T) \subset L_{\Omega}^2(T)$  it suffices, in view of Theorem 1.1 of [CHL-I], to show that for some basis  $\mathcal{A}$  of  $F_N$  any word  $z = x_1 \dots x_s$  in the laminary language  $\mathcal{L}_{\mathcal{A}}^{\mathcal{Q}}(T)$  associated to the algebraic

lamination  $L_{\mathcal{Q}}^2(T)$  (via the canonical map  $\rho_{\mathcal{L}}^A \rho_{\mathcal{A}}^2 : \Lambda^2(F_N) \rightarrow \Lambda_{\mathcal{L}}(\mathcal{A})$ , see [CHL-I]) is also contained in the laminary language  $\mathcal{L}_{\mathcal{A}}^{\mathcal{Q}}(T)$  from §4: By extendability of any word in a laminary language we find, for any  $k \geq 0$ , a “superword”  $u_{\mathcal{A}} = x_{-k} \dots x_1 \dots x_s \dots x_{s+k}$  in  $\mathcal{L}_{\mathcal{A}}^{\mathcal{Q}}(T)$ . Thus, for  $k$  sufficiently large, we can use Cooper’s cancellation bound and pass to another basis  $\mathcal{B}$  with  $\text{BBT}(\mathcal{B}, P) < \varepsilon$  and  $\text{vol}(\mathcal{B}, P) < \varepsilon$ , for small  $\varepsilon > 0$ , such that the word  $u_{\mathcal{B}}$  in the basis  $\mathcal{B}$ , which represents the same element of  $F_N$  as  $u_{\mathcal{A}}$ , contains a subword  $v_{\mathcal{B}}$  with the following properties (compare the similar situation considered in Lemma 5.5):

On the one hand  $v_{\mathcal{B}}$  belongs to the laminary language  $\mathcal{L}_{\mathcal{B}}^{\mathcal{Q}}(T)$ , which implies by Lemma 3.1 (a) and Lemma 6.1 that  $d(P, v_{\mathcal{B}}P) \leq 3\text{BBT}(\mathcal{B}, P) < 3\varepsilon$ . As the rank of our free group  $F_N$  satisfies  $N \geq 2$ , we find an element  $y \in \mathcal{B}^{\pm 1}$  such that  $w_{\mathcal{B}} = v_{\mathcal{B}}y$  is cyclically reduced and satisfies furthermore, by Lemma 3.1 (d), that  $\|w_{\mathcal{B}}\|_T \leq d(P, v_{\mathcal{B}}P) + d(P, yP) \leq 3\text{BBT}(\mathcal{B}, P) + \text{vol}(\mathcal{B}, P) < 4\varepsilon$ .

On the other hand, by a double application of Cooper’s cancellation bound we obtain, since  $w_{\mathcal{B}} = v_{\mathcal{B}}y$  is cyclically reduced, for sufficiently large  $k$ , that the word  $w_{\mathcal{A}}$  in the basis  $\mathcal{A}$ , which represents the same element of  $F_N$  as  $w_{\mathcal{B}}$ , contains the originally chosen word  $z = x_1 \dots x_s \in \mathcal{L}_{\mathcal{A}}^{\mathcal{Q}}(T)$  as subword in its cyclically reduced part. This shows that  $z = x_1 \dots x_s$  belongs to  $\mathcal{L}_{\mathcal{A}}^{4\varepsilon}(T)$ , and hence, as  $\varepsilon > 0$  was picked arbitrarily small, to  $\mathcal{L}_{\mathcal{A}}^{\Omega}(T)$ .

For the converse inclusion  $L_{\Omega}^2(T) \subset L_{\mathcal{Q}}^2(T)$  we consider any  $(X, X') \in L_{\Omega}^2(T)$  and apply Proposition 5.3 to deduce that both,  $X$  and  $X'$  belong to  $L^1(T)$ . Hence there are well defined points  $R = \mathcal{Q}(X)$  and  $Q = \mathcal{Q}(X')$  in  $\bar{T}$ . If  $R \neq Q$ , then for any small  $\varepsilon > 0$  we can pass to a basis  $\mathcal{B}$  of  $F_N$  with  $12\text{BBT}(\mathcal{B}, P) < 4\varepsilon < d(R, Q)$ , and we consider the biinfinite word  $Z_{\mathcal{B}} = \dots y_{i-1}y_i y_{i+1} \dots = \rho_{\mathcal{B}}(X, X')$  in this basis. Any subword  $w = y_{-k} \dots y_l$  of  $Z_{\mathcal{B}}$  with  $k$  and  $l$  sufficiently large satisfies  $d(y_1 \dots y_l P, R) < 3\text{BBT}(\mathcal{B}, P) < \varepsilon$ , as well as  $d((y_{-k} \dots y_0)^{-1} P, Q) < 3\text{BBT}(\mathcal{B}, P) < \varepsilon$ , as follows from Lemma 6.3.

As  $d(P, wP) = d(y_1 \dots y_l P, (y_{-k} \dots y_0)^{-1} P)$ , this gives  $4\varepsilon < d(R, Q) \leq d(y_1 \dots y_l P, R) + d(P, wP) + d((y_{-k} \dots y_0)^{-1} P, Q) < d(P, wP) + 2\varepsilon$ , and thus  $d(P, wP) > 2\varepsilon$ . Hence, if  $w$  is subword of any cyclically reduced word  $W$  in  $\mathcal{B}$ , we obtain from Lemma 3.1 (c) the inequalities  $\|W\|_T \geq d(P, wP) - 2\text{BBT}(\mathcal{B}, P) \geq \varepsilon$ . This shows that  $w$  does not belong to  $\mathcal{L}_{\mathcal{B}}^{\varepsilon}(T)$ , for  $\varepsilon > 0$  small, and hence not to  $\mathcal{L}_{\mathcal{B}}(T)$ , contradicting the assumption  $Z_{\mathcal{B}} \in \rho_{\mathcal{B}}^2(L_{\Omega}^2(T))$ . Thus we have proved that  $R = Q$ , which shows that  $(X, X')$  belongs to  $L_{\mathcal{Q}}(T)$ .  $\square$

## 9 From the boundary of Outer space to the space of laminations

The group  $\text{Out}(F_N)$  acts canonically (from the left) on the space  $\Lambda^2(F_N)$  of algebraic laminations (see [CHL-I]), but it also acts (from the right !) on the space  $cv_N$  and on its ‘‘Thurston boundary’’  $\partial cv_N$ , as well as on  $CV_N \cup \partial CV_N$  (see §2). This right action is defined as follows: For any  $\alpha \in \text{Aut}(F_N)$  and any tree  $T \in \overline{cv}_N$ , the length function of the image tree  $T\alpha_*$  is given by

$$\|w\|_{T\alpha_*} = \|\alpha(w)\|_T,$$

for every  $w \in F_N$ . This gives:

**Proposition 9.1.** *The map*

$$\lambda^2 : \partial CV_N \rightarrow \Lambda^2(F_N), \quad T \mapsto L^2(T)$$

*is  $\text{Out}(F_N)$ -anti-equivariant: For any automorphism  $\alpha$  of  $F_N$  and any  $T \in CV_N$  one has  $\alpha^{-1}(L^2(T)) = L^2(T\alpha_*)$ .*

*Proof.* This follows directly from the definition of  $L^2(T) = L_\Omega^2(T) = \bigcap_{\varepsilon > 0} L_\varepsilon^2(T)$  in §4, since we observe, for every  $\varepsilon > 0$ , that

$$\alpha^{-1}(\Omega_\varepsilon(T)) = \Omega_\varepsilon(T\alpha_*)$$

and hence

$$\alpha^{-1}(L_\varepsilon^2(T)) = L_\varepsilon^2(T\alpha_*).$$

□

However, it is important to point out that the map  $\lambda^2$  is not continuous. For example, consider the case  $F_3 = F(\{a, b, c\})$ , and let  $D$  be the *Dehn twist* automorphism given by  $a \mapsto bab^{-1}, b \mapsto b, c \mapsto c$ . Then in the last section of [CL95] it is shown that every tree  $T \in cv_3 \cup \partial cv_3$  with  $\|b\|_T > 0$  converges projectively under iteration of  $D$  to the simplicial tree  $\tilde{\Gamma}_b$  which is the Bass-Serre tree of the graph of groups decomposition  $F(a, b, c) = \langle a, b \rangle *_{\langle b=b' \rangle} \langle b', c \rangle$ . Thus, if  $T$  is, for example, the simplicial tree obtained from the rose  $\mathcal{R}_{\{a, b, c\}}$  by contracting in the universal covering the edge labelled  $c$  equivariantly (while leaving all other edges of same length), we see that  $L^2(T) = L^2(c)$ . Replacing  $a$  by  $b^m a b^{-m}$  defines a family of new



trees  $T_m \in \partial cv_3$ , which projectivize to points  $[T_m] \in \partial CV_3$  that give precisely the  $D$ -orbit of  $[T] \in \partial CV_3$ , which has  $[\tilde{\Gamma}_b]$  as forward and backward limit point. On the other hand, we see easily that  $L^2(T_m) = L^2(c)$  for all  $m \in \mathbb{Z}$ . Since  $L^2(\tilde{\Gamma}_b)$  contains  $L^2(c)$ , but is much larger (for example  $L^2(a)$  is equally contained in  $L^2(\tilde{\Gamma}_b)$ ), this example illustrates that there exist sequences of trees  $T_k$  converging to  $T$  such that  $L^2(T_k)$  converges into  $L^2(T)$ , but not to  $L^2(T)$ .

## References

- [BF92] M. Bestvina and M. Feighn. A combination theorem for negatively curved groups. *J. Differential Geom.* **35**, 85–101, 1992.
- [BF95] M. Bestvina and M. Feighn. Stable actions of groups on real trees. *Invent. Math.* **121**, 287–321, 1995.
- [BF96] M. Bestvina and M. Feighn. Addendum and correction to: “A combination theorem for negatively curved groups” [*J. Differential Geom.* **35** (1992), no. 1, 85–101]. *J. Differential Geom.* **43**, 783–788, 1996.
- [BF00] M. Bestvina and M. Feighn. The topology at infinity of  $\text{Out}(F_n)$ . *Invent. Math.* **140**, 651–692, 2000.
- [BFH97] M. Bestvina, M. Feighn, and M. Handel. Laminations, trees, and irreducible automorphisms of free groups. *Geom. Funct. Anal.* **7**, 215–244, 1997.
- [CHLL] T. Coulbois, A. Hilion, G. Levitt, and M. Lustig. In preparation.
- [CHL05] T. Coulbois, A. Hilion, and M. Lustig. Non-uniquely ergodic  $\mathbb{R}$ -trees are topologically determined by their algebraic lamination. Preprint, 2005. (To appear in *Illinois J. Math.*)
- [CHL-I] T. Coulbois, A. Hilion, and M. Lustig.  $\mathbb{R}$ -trees and laminations for free groups I: Algebraic laminations. Preprint, 2006.
- [CHL-III] T. Coulbois, A. Hilion, and M. Lustig.  $\mathbb{R}$ -trees and laminations for free groups III: Currents and dual  $\mathbb{R}$ -tree metrics. Preprint, 2006.

- [CHL3] T. Coulbois, A. Hilion and M. Lustig. Which  $\mathbb{R}$ -trees can be mapped continuously to a current ? Preliminary preprint, 2006.
- [CL95] M. M. Cohen and M. Lustig. Very small group actions on  $\mathbf{R}$ -trees and Dehn twist automorphisms. *Topology* **34**, 575–617, 1995.
- [CV86] M. Culler and K. Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.* **84**, 91–119, 1986.
- [FLP91] A. Fathi, F. Laudenbach and V. Poénaru, editors *Travaux de Thurston sur les surfaces*. Séminaire Orsay, Astérisque **66-67**, 1991. Reprint of *Travaux de Thurston sur les surfaces*, Soc. Math. France, Paris, 1979.
- [GJLL98] D. Gaboriau, A. Jaeger, G. Levitt, and M. Lustig. An index for counting fixed points of automorphisms of free groups. *Duke Math. J.* **93**, 425–452, 1998.
- [GL95] D. Gaboriau and G. Levitt. The rank of actions on  $\mathbf{R}$ -trees. *Ann. Sci. École Norm. Sup.* **28**, 549–570, 1995.
- [GLP94] D. Gaboriau, G. Levitt, and F. Paulin. Pseudogroups of isometries of  $\mathbf{R}$  and Rips’ theorem on free actions on  $\mathbf{R}$ -trees. *Israel J. Math.* **87**, 403–428, 1994.
- [Gui98] V. Guirardel. *Action de groupes sur des arbres réels et dynamique sur la frontière de l’outre-espace*. PhD thesis, Toulouse 1998.
- [Gui00] V. Guirardel. Dynamics of  $\text{Out}(F_n)$  on the boundary of outer space. *Ann. Sci. École Norm. Sup.* **33**, 433–465, 2000.
- [KL] I. Kapovich and M. Lustig. The actions of  $\text{Out}(F_n)$  on the boundary of Outer space and on the space of currents: minimal sets and equivariant incompatibility. ArXiv GR/0605548. (To appear in *Ergodic Theory Dynam. Systems*)
- [LL] G. Levitt and M. Lustig. Automorphisms of free groups have asymptotically periodic dynamics. ArXiv math.GR/0407437.
- [LL00] G. Levitt and M. Lustig. Periodic ends, growth rates, Hölder dynamics for automorphisms of free groups. *Comment. Math. Helv.* **75**, 415–429, 2000.

- [LL03] G. Levitt and M. Lustig. Irreducible automorphisms of  $F_n$  have north-south dynamics on compactified outer space. *J. Inst. Math. Jussieu* **2**, 59–72, 2003.
- [LP97] G. Levitt and F. Paulin, Geometric group actions on trees. *Am. J. Math.* **119**, 83–102, 1997.
- [Sha87] P. Shalen. Dendrology of groups: an introduction. In *Essays in group theory*, Math. Sci. Res. Inst. Publ. **8**, 265–319. Springer, New York, 1987.
- [Sko96] R. Skora. Splittings of surfaces. *J. Amer. Math. Soc.* **9**, 605–616, 1996.
- [Vog02] K. Vogtmann. Automorphisms of free groups and outer space. *Geom. Dedicata* **94**, 1–31, 2002.

Thierry.Coulbois@univ-cezanne.fr  
 Arnaud.Hilion@univ-cezanne.fr  
 Martin.Lustig@univ-cezanne.fr  
 Mathématiques (LATP)  
 Université Paul Cézanne – Aix-Marseille III  
 av. escadrille Normandie-Niémen  
 13397 Marseille 20  
 France