

# THE DIRICHLET PROBLEM FOR SOME NONLOCAL DIFFUSION EQUATIONS

EMMANUEL CHASSEIGNE<sup>1</sup>

**ABSTRACT.** We study the Dirichlet problem for the non-local diffusion equation  $u_t = \int \{u(x+z, t) - u(x, t)\} d\mu(z)$ , where  $\mu$  is a  $L^1$  function and “ $u = \varphi$  on  $\partial\Omega \times (0, \infty)$ ” has to be understood in a non-classical sense. We prove existence and uniqueness results of solutions in this setting. Moreover, we prove that our solutions coincide with those obtained through the standard “vanishing viscosity method”, but show that a boundary layer occurs: the solution does not take the boundary data in the classical sense on  $\partial\Omega$ , a phenomenon related to the non-local character of the equation. Finally, we show that in a bounded domain, some regularization may occur, contrary to what happens in the whole space.

**Keywords:** integro-differential equations ; Dirichlet problem ; Lvy operators ; general nonlocal operators.

**Mathematics Subject Classification:** 47G20, 35D05, 35D10 35B05, 35B50, 35B65;

## 1. INTRODUCTION

In this paper we investigate a suitable notion of Dirichlet Problems in a bounded set  $\Omega \subset \mathbb{R}^n$  for a class of non-local equations, starting from remarks and ideas developed in [5] to cover a more general framework.

To begin with, let us consider our model equation posed in the whole space  $\mathbb{R}^n$  :

$$(1.1) \quad \frac{\partial u}{\partial t}(x, t) = \int_{\mathbb{R}^n} \{u(x+z, t) - u(x, t)\} d\mu(z),$$

where  $\mu \in \mathcal{M}(\mathbb{R}^n)$  is a finite measure. Such non-local diffusion operators arise in various problems like diffusion of population (see [5] and the references therein for more on this topic), Lévy processes [4, 6, 8], fractional Laplace diffusion [7], and more generally, integro-differential pde’s (see [1] and the references therein). Note that in the fractional Laplacian case,  $\mu(z) = |z|^{-N-\lambda}$  with  $0 < \lambda < 2$ , a case in which  $\mu$  is not a finite measure that we shall not consider here (we refer to [3] for a general study of the Dirichlet problem in such situations).

Now, if the equation has a rather clear sense when posed in the entire space  $\mathbb{R}^N$ , the question of what is a Dirichlet problem in this context becomes less obvious. Indeed, due to the nonlocal character of the equation, we need to assign the values of  $u$  not only on

<sup>1</sup>Laboratoire de Mathématiques et Physique Théorique, Université F. Rabelais, Parc de Grandmont, 37200 Tours (FRANCE) - e-mail : emmanuel.chasseigne@lmpt.univ-tours.fr

the topological boundary  $\partial\Omega$ , but on the complement of  $\Omega$  (see more precise statements below).

Let us mention that a viscosity approach of such Dirichlet problems can be found in [2] and [6], but it concerns mainly the case of singular (*i.e.*, non-integrable) measures. We shall prefer here a similar approach to that which can be found in [5] where we considered an homogenous Dirichlet problem with  $\mu$ , a continuous bounded symmetric function. But we want to go further in two directions: first to consider non-homogeneous problems (to be suitably defined), and second to include functions  $\mu$  that may be singular, but yet integrable.

## AIMS AND ORGANIZATION OF THE PAPER

The aim of this paper is thus to give a rather complete view of the case when  $\mu \geq 0$  is an integrable function in  $\mathbb{R}^n$ .

We shall first define a suitable notion of what is a Dirichlet problem for our equation in Section 2, and then we investigate solvability of the problem in Section 3. More precisely, we prove that for any continuous and bounded initial data  $u_0$  and “boundary” data  $\varphi$ , there exists a unique solution  $u$ , continuous and bounded in  $\bar{\Omega} \times [0, \infty)$ . We also prove that comparison holds in this class of solutions.

Section 4 is devoted to study the vanishing viscosity method which is another way to try to solve the problem : we add a viscous term  $-\epsilon\Delta u$  to the equation in order to force the boundary data to be taken, and then try to see what happens in the limit as  $\epsilon \rightarrow 0$ . Well, it turns out that we get indeed in the limit the unique solution obtained in Section 3, but this solution is strictly positive on  $\partial\Omega$  whereas  $u_\epsilon$  is zero on the boundary for any  $\epsilon > 0$ . Hence, we are facing a boundary layer formation which proves that in general the boundary data will not be taken in the usual sense.

In Section 5 we investigate some fine regularity properties of the solutions. It is known that in the case  $\Omega = \mathbb{R}^n$ , no regularization occurs for such non-local equations, in the sense that the solution at time  $t > 0$  cannot have a better regularity than that of  $u_0$  (this was proven in [5]). But in the case of bounded domains, it appears that however some kind of smoothing does occur, provided  $\mu$  is not too much concentrated: the modulus of continuity of the solution decreases exponentially as time increases (see Section 5 for precise statements).

We will focus on some other qualitative aspects in Section 6. In particular, we prove that the equation enjoys an positivity property similar to the heat equation (if  $u_0 \geq 0$  and not identically zero, then  $u(t)$  is automatically positive in all  $\Omega$ ). But in the present situation, this effect is not purely diffusive as for the Laplace operator, it is due to the nonlocal character of the equation. Moreover we show that solutions are immediately positive also on the boundary  $\partial\Omega$ , which is consistent with the layer formation in the vanishing viscosity method.

Finally we collect some other qualitative remarks, examples and possible extensions in the last Section.

*This note benefited a lot from discussions with Guy Barles and Ch. Georgelin, whom I warmly thank.*

## 2. DEFINITION OF THE DIRICHLET PROBLEM

We shall assume throughout this paper that the measure  $\mu$  has a density (still denoted by  $\mu$ ) which is a nonnegative, integrable function of total mass  $\mu(\mathbb{R}^n) = 1$ . In fact, if the density is integrable, we can always assume that the total mass is 1 by invoking scaling arguments: simply put  $v(x, t) = u(x, \lambda t)$  where  $\lambda$  is the total mass of  $\mu$ . The set  $\Omega \subset \mathbb{R}^n$  will be open, bounded, connected and regular (to simplify).

Now, as we noticed above, to give sense to a Dirichlet problem one needs to prescribe the “boundary” value not only on the topological frontier  $\partial\Omega$ , but in fact in the complement of  $\Omega$  or to be more precise, on the set where the measure  $\mu$  will read some information. So let  $\tau_x^{-1}$  be the translated measure defined by  $\tau_x^{-1}\mu(z) = \mu(x + z)$ , let  $\text{supp}(\mu)$  denote the support of the measure  $\mu$  and let us define the following subsets of  $\mathbb{R}^n$  :

$$\bar{\Omega}^\mu = \overline{\bigcup_{x \in \Omega} \text{supp}(\tau_x^{-1}\mu)}, \text{ and } \partial^\mu\Omega = \bar{\Omega}^\mu \setminus \Omega.$$

In other words,  $\bar{\Omega}^\mu$  is what we call the *domain of influence* of  $\Omega$ . In short, if you want to know how things evolve in  $\Omega$ , you need to read information in the bigger set  $\bar{\Omega}^\mu \supset \Omega$ , because the operator will look at it. In this direction, the natural notion of frontier  $\partial\Omega$  has to be replaced by the *extended frontier*  $\partial^\mu\Omega$ , which is of course not a negligible set since it is regular (*i.e.*, the closure of an open set in  $\mathbb{R}^N$ ). Then in this setting, we may construct a general theory for the following Dirichlet problem :

$$(\mathcal{P}) \quad \begin{cases} \frac{\partial u}{\partial t}(x, t) = \int_{\mathbb{R}^n} \{u(x + z, t) - u(x, t)\} d\mu(z) & \text{in } \Omega \times (0, \infty), \\ u(x, t) = \varphi(x, t) & \text{in } \partial^\mu\Omega \times [0, \infty), \\ u(x, 0) = u_0(x) & \text{in } \Omega. \end{cases}$$

The preceeding formulation requires the function  $u$  to be defined in  $\Omega^\mu \times [0, \infty)$ , extended by  $\varphi$ , in order to give sense to the integral term. But as we shall see below,  $u$  will not be continuous when crossing the boundary  $\partial\Omega$  so that practically, we shall often prefer a different formulation, keeping the function  $u$  to be defined only within  $\Omega \times [0, \infty)$ . Let us denote by  $K_\varphi(u)$  the following non-local operator : for all  $(x, t) \in \Omega \times (0, \infty)$ ,

$$K_\varphi(u)(x, t) = \int_{\{(x+z) \in \Omega\}} u(x + z, t) d\mu(z) + \int_{\{(x+z) \in \partial^\mu\Omega\}} \varphi(x + z, t) d\mu(z) - u(x, t).$$

Thus we arrive at the following definition :













Hence using Gronwall's lemma we get that  $\omega_v(\eta, t) \leq \omega_0(\eta)e^t$ , and going back to  $u(x, t) = e^{-t}v(x, t)$  gives the result.  $\square$

In this generality, the Theorem is optimal : take  $N = 1$  and for instance  $\mu(z) = 1/2$  on  $[-1; 1]$ ,  $\mu(z) = 0$  if  $|z| > 1$ . Then if  $u_0(x) = ax$ , the solution is stationary and so the modulus of continuity is constant.

The situation is a little bit more complicated in a bounded domain  $\Omega$  since the boundary terms may induce some sort of discontinuity. So let us first restrict ourselves to the case  $\varphi = 0$  and let us introduce some notation :

For any  $x \in \mathbb{R}^n$ , let  $\tau_x^{-1}\Omega = \{z : (x + z) \in \Omega\}$  and for any two sets  $A, B$ , let  $A \Delta B = (A \setminus B) \cup (B \setminus A)$ . Now, for any  $\eta > 0$ , let us define the following quantities :

$$\begin{aligned}\lambda(\eta) &= \sup_{\substack{(x,y) \in \Omega^2 \\ |x-y| < \eta}} \mu(\tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega), \\ \gamma(\eta) &= \sup_{\substack{(x,y) \in \Omega^2 \\ |x-y| < \eta}} \mu(\tau_x^{-1}\Omega \Delta \tau_y^{-1}\Omega).\end{aligned}$$

Notice that obviously,  $\lambda(\eta), \gamma(\eta) \in [0, 1]$  and that  $\gamma(\eta)$  goes to zero as  $\eta \rightarrow 0$ . Roughly speaking,  $\lambda(\eta)$  is related to the total mass of  $\mu$  in  $\tilde{\Omega} = \bigcup_{x \in \Omega} \tau_x^{-1}\Omega$ , while  $\gamma(\eta)$  is related to the mass of  $\mu$  involved at the boundary.

**Theorem 5.2.** *Let us assume that  $\Omega$  is bounded and regular,  $\mu \geq 0$  is a  $L^1$  measure,  $\varphi = 0$  and that  $u$  is a solution with initial data  $u_0 \in C_0(\bar{\Omega})$ . Then the following estimates hold :*

- If  $\lambda(\eta) < 1$ , then for any  $t > 0$

$$\omega(\eta, t) \leq \left( \omega_0(\eta) + \gamma(\eta) \cdot \|u_0\|_\infty \frac{e^{(1-\lambda(\eta))t} - 1}{1 - \lambda(\eta)} \right) e^{(\lambda(\eta)-1)t}.$$

- If  $\lambda(\eta) = 1$ , then for any  $t > 0$ ,

$$\omega(\eta, t) \leq \left( \omega_0(\eta) + \gamma(\eta) \cdot \|u_0\|_\infty \cdot t \right).$$

*Proof.* The proof proceeds with the same ideas as for the case  $\Omega = \mathbb{R}^n$  : integrating in time the equation for  $v(x, t) = e^t u(x, t)$  and again denoting by  $\omega_v$  the modulus of continuity for  $v$  one gets :

$$v(x, t) - v(y, t) = \int_0^t \int_{\tau_x^{-1}\Omega} v(x + z, s) d\mu(z) ds - \int_0^t \int_{\tau_y^{-1}\Omega} v(y + z, s) d\mu(z) ds,$$

so that if  $|x - y| < \eta$ , we can estimate the modulus of continuity as follows :

$$|v(x, t) - v(y, t)| \leq \int_0^t \int_{\tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega} |v(x + z, s) - v(y + z, s)| d\mu(z) ds + R(x, y, t)$$

where  $R(x, y, t)$  contains the integral over  $(\tau_x^{-1}\Omega \Delta \tau_y^{-1}\Omega)$ , that we estimate uniformly using the fact that  $\|v(s)\|_\infty \leq \|u_0\|_\infty \cdot e^s$  :

$$R(x, y, t) \leq \|u_0\|_\infty \cdot \gamma(\eta) \cdot \int_0^t e^s ds.$$

Thus, going back to the estimate, we have for any  $|x - y| < \eta$ ,

$$|v(x, t) - v(y, t)| \leq \omega_v(\eta, 0) + \int_0^t \{ \lambda(\eta) \cdot \omega_v(\eta, s) + \gamma(\eta) \cdot \|u_0\|_\infty \cdot e^s \} ds.$$

Passing to the supremum over all  $(x, y)$  such that  $|x - y| < \eta$ , and using Gronwall's lemma we obtain:

$$\omega_v(\eta, t) \leq (\omega_v(\eta, 0) + \gamma(\eta) \cdot \|u_0\|_\infty \frac{e^{(1-\lambda(\eta))t} - 1}{1 - \lambda(\eta)}) e^{\lambda(\eta)t},$$

and finally going back to  $u(x, t) = e^{-t}v(x, t)$  yields the desired estimate. When  $\lambda(\eta) = 1$ , then the estimate is easier to obtain, using also Gronwall's lemma.  $\square$

Before stating a corollary, let us give a heuristic idea of the meaning of those estimates: if  $\lambda = 1$ , we see that the estimate on the modulus of continuity deteriorates as time increases, because of the discontinuity at the boundary (the jump between  $u$  and  $\varphi = 0$ ) which propagates in the interior. On the contrary, if  $\lambda(\eta) < 1$ , this effect is compensated by the fact that the measure  $\mu$  is not fully integrated, so that we gain an exponential decay of the modulus of continuity : some kind of regularization occurs ! (See end Section for some examples).

Let us give two sufficient conditions that explain when both cases may appear (recall that  $\tilde{\Omega} = \bigcup_{x \in \Omega} \tau_x^{-1}\Omega$ ) :

**Proposition 5.1.** *If  $\mu(\mathbb{R}^n \setminus \tilde{\Omega}) > 0$ , then for any  $\eta > 0$ ,*

$$\lambda(\eta) \leq 1 - \mu(\mathbb{R}^n \setminus \tilde{\Omega}) < 1.$$

*On the other hand, if there exists  $x \in \Omega$  such that  $\text{supp}(\mu) \subset \tau_x^{-1}\Omega$  (with strict inclusion), then  $\lambda(\eta) = 1$  for  $\eta > 0$  sufficiently small.*

*Proof.* Let us first notice that obviously, for any  $x, y \in \Omega$ ,  $\tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega \subset \tilde{\Omega}$  so that  $\mu(\tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega) \leq \mu(\tilde{\Omega})$ , which is independent of  $\eta > 0$  and  $x, y \in \Omega$ . Thus passing to the supremum and using the fact that the total mass of  $\mu$  is 1, one gets the first result.

Now for the second statment: if such an  $x$  exists, then for  $\eta$  small enough, any  $y \in \Omega$  such that  $|x - y| < \eta$ , will satisfy :  $\text{supp}(\mu) \subset \tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega$ . Hence  $\mu(\tau_x^{-1}\Omega \cap \tau_y^{-1}\Omega) = 1$  and passing to the supremum for such  $x, y$  yields the result.  $\square$

We refer to the final Section where explicit examples are given, by let us mention that more or less,  $\lambda > 1$  if  $\mu$  has a huge support with respect to the shape of  $\Omega$ , whereas  $\lambda = 1$  if  $\mu$  is rather concentrated (again with respect to the shape of  $\Omega$ ). The limit being when  $\mu = \delta_0$  is so concentrated that nothing moves.

**Remark 5.1.** *If  $\varphi \neq 0$  is bounded and continuous, the estimate involves another term depending on the integral of  $\varphi$  on  $A_\mu(\eta) = \partial^\mu \Omega_\eta \setminus \partial^\mu \Omega$  : if*

$$\theta(\eta, t) = \max_{\Omega \times (0, t)} \int_{\{(x+z) \in A_\mu(\eta)\}} \varphi(x+z) d\mu(z),$$

*then we can estimate the modulus of continuity using again Gronwall's Lemma, taking into account  $\theta(\eta, t)$ . The details are left to the reader.*

## 6. QUALITATIVE PROPERTIES OF SOLUTIONS

**Theorem 6.1.** *Let us assume that  $\mu \geq 0$  is a measure such that for some  $\eta > 0$ ,*

$$\text{supp}(\mu) \supset \{|z| \leq \eta\}.$$

*Then if  $u(x, t) \in X$  is a solution of problem  $(\mathcal{P})$  with  $\varphi = 0$  and  $u_0 \in C(\Omega)$  non-negative and compactly supported in  $\Omega$ , we have*

$$\forall (x, t) \in \bar{\Omega} \times (0, \infty), \quad u(x, t) > 0.$$

*Proof.* Let us first observe that if we set  $v(x, t) = e^t \tilde{u}(x, t)$  where  $\tilde{u}$  is the extension of  $u$  defined by (2.2), then  $v$  satisfies

$$\partial_t v = \int_{\mathbb{R}^n} v(x+z) d\mu(z) \geq 0,$$

so that once  $v$  (hence  $u$  also) is positive, it stays as such for later times. Let  $A = \text{supp}(u_0) \subset \Omega$  which is compact and notice first that obviously  $u \geq 0$  in  $\Omega \times [0, \infty)$ , using the comparison principle. Now, let  $x_0 \in \bar{\Omega} \setminus A$ , and assume that  $\text{dist}(x_0; A) < \eta$ . Using the equation at  $(x_0, 0)$  (remember that the equation holds even on the boundary), one gets

$$v_t(x_0, 0) \geq \int_{|z| < \eta} v_0(x+z, 0) d\mu(z) > 0.$$

Thus we get that necessarily, for any  $t > 0$ ,  $u(x, t)$  will be positive in  $A_\eta = \{x \in \bar{\Omega} : \text{dist}(x, A) < \eta\}$ . Iterating the same argument at  $t > 0$  arbitrary small, we get that  $u(x, t) > 0$  in  $A_{k\eta}$  for any  $k \in \mathbb{N}$ . Thus it is clear that  $u(x, t) > 0$  for any  $x \in \bar{\Omega}$ , for any time  $t > 0$ .  $\square$

As a corollary of this result, we have obtained that the boundary data is never taken, more precisely :

**Corollary 6.1.** *In the vanishing viscosity method, a boundary layer occurs and the limit solution is strictly positive on  $\partial\Omega$  for any time  $t > 0$ . Thus the boundary data is never taken in general.*

## 7. CONCLUSIONS, REMARKS AND EXAMPLES

We have proposed a general framework to deal with Dirichlet problems for the non-local equation (1.1). If  $\mu \in L^1(\mathbb{R}^n)$ , we have proved existence and uniqueness of a continuous solution for any continuous initial and boundary data. Furthermore, under some condition on the support of  $\mu$  near the origin, we have also proved a positivity property, and that the boundary values will not be taken in the classical sense. Let us end this note with some remarks and extensions :

1. If  $\mu \in L^\infty$ , a similar existence and uniqueness result as in Theorem 3.1 may be obtained for data  $u_0, \varphi$  in  $L^1$ , using similar fixed-point arguments. But if  $\mu$  is a finite measure with singular part (like Dirac deltas), then the above construction does not work since the integral term may not be continuous. However, we think that similar things could maybe be done for measures less concentrated than the delta.

2. An interesting feature of the equation is that it holds true everywhere on  $\partial\Omega$  (see Proposition 3.1). This explains why, although the boundary data is not taken in the usual sense, there would not be any problem with the viscosity definition of boundary values. Indeed, passing to the limit in the viscous equation can be done also in the framework of viscosity solutions since the convergence holds in  $L^\infty$ -weak\*.

3. Theorem 6.1 shows that propagation with infinite speed occurs like for the Heat Equation, although the phenomenon is of different nature (due to the non-local term and not instantaneous diffusion as it is the case for the heat equation). But even more,  $u$  also becomes positive at the boundary  $x \in \partial\Omega$ . So, even if both  $\varphi$  and  $u_0$  are continuous, in general  $u(x, t)$  will not take the boundary data on  $\partial\Omega$  in the usual sense (this was proved for stationary solutions in [5]). And again, we recover the necessity to define a suitable notion of boundary for non-local operators,  $\partial^\mu\Omega$  here.

4. Note that some property on the support of  $\mu$  near the origin is necessary to have this propagation property, although our criteria is maybe not optimal. Indeed, if  $\mu$  is too much off-centered, then the operator will not see anything in  $\Omega$  itself: consider for instance  $\Omega = ]-10; +10[$ ,  $\varphi = 0$  and let  $\mu$  be the characteristic function of the set  $] -1, 0[$ . If  $u(0) \in C_0(\Omega)$ , it will remain compactly supported on the left for any  $t > 0$ .

5. We showed a kind of regularizing effect in the case of bounded domains, provided the measure  $\mu$  satisfies  $\lambda(\eta) < 1$ . Let us consider the case  $\Omega = ]-1; 1[$  and let us give an example of a measure  $\mu$  satisfying this property :  $\mu(z) = ce^{-|z|^2}$ , where  $c > 0$  is such that  $\mu(\mathbb{R}) = 1$ . Then clearly, since  $\tilde{\Omega} = ]-2; 2[$ ,  $\lambda(\eta) \leq 1 - \int_{-2}^{+2} d\mu(z) < 1$ . One can also have a good estimate of the rate in the exponential decay. Now if  $\mu(z)$  is the characteristic function of the set  $] -1/2; 1/2[$ , we have clearly that for any  $\eta < 1/2$ ,  $\lambda(\eta) = 1$ .

6. A different story happens when  $\mu$  is not integrable near the origin, which is the case for the fractional Laplace operator. In this case, the boundary data are taken since the measure is so singular that discontinuities at the boundary are not allowed (see [3]).

## APPENDIX : THE PROBABILISTIC INTERPRETATION

In this Appendix, we briefly explain the probabilistic context in which the Cauchy problem may be understood. Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a measured space and  $Y_0 : \Omega \rightarrow \mathbb{R}$  be a random variable with law  $u_0 \in L^1(\mathbb{R}^N)$  (we assume that  $\int u_0 = 1$ ). Now let  $(Y_i)_{i \in \mathbb{N}}$  be a sequence of independent, identically distributed random variables of law  $\mu$ , and  $t \mapsto N_t$  be a random variable on  $\mathbb{N}$  with Poisson law of intensity  $t > 0$  :

$$\mathbb{P}(N_t = k) = e^{-t} \frac{t^k}{k!}.$$

Then we consider the stochastic process  $X_t = Y_0 + \sum_{i=1}^{N_t} Y_i$ . The term  $Y_0$  represents the initial state of the stochastic process  $X_t$ , while the sum represents a random number of jumps  $N_t$ , each jump being independent of the others, with probability distribution  $\mu$ . Such a process is named a *composed Poisson process*, which is a special case of the more general Lévy processes. If we consider the density  $u(t)$  of the probability distribution of  $X_t$ , it can be easily shown that precisely,

$$\frac{\partial u}{\partial t} = \int_{\mathbb{R}^N} \{u(x+z, t) - u(x, t)\} d\mu(z).$$

Now, by definition the characteristic function of  $X_t$  is nothing but the Fourier transform of  $u(t)$ , and it is known that for any Lévy process, there exists a uniquely determined triplet  $(A, \gamma, \nu)$  such that :

$$\mathbb{E}(e^{i\xi X_t}) = e^{t\varphi(z)} \text{ with } \varphi(z) = -\frac{1}{2}z \cdot Az + \gamma \cdot z + \int_{\mathbb{R}^N} (e^{iz\xi} - 1 - z\xi \mathbf{1}_{|z| \leq 1}(z)) d\nu(z).$$

In this decomposition,  $A$  is a positive-definite matrix,  $\gamma \in \mathbb{R}^n$  and  $\nu$  is the Lévy measure, satisfying the following properties :  $\int_{|z| \leq 1} z^2 d\nu$ ,  $\int_{|z| \geq 1} d\nu < \infty$ . The infinitesimal generator of the Feller semi-group associated to  $(X_t)$  is the following :

$$Lf := -\frac{1}{2} \sum_{i,j=1}^N a_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j} + \sum_{i=1}^N \gamma_i \frac{\partial f}{\partial x_i} + \int_{\mathbb{R}^N} (f(x+z) - f(x) - \nabla f(x) \cdot z \mathbf{1}_{|z| \leq 1}(z)) d\nu(z).$$

In the special case of composed Poisson processes, the matrix  $A$  is nul,  $\gamma = 0$  and the corrector term involving the gradient of  $f$  vanishes for symmetric measures.

## REFERENCES

- [1] N. ALIBAUD, "Existence, Uniqueness and Regularity for nonlinear degenerate parabolic equations with nonlocal terms", *Preprint* (2006).
- [2] M. ARISAWA, "A new definition of viscosity solutions for a class of second-order degenerate elliptic integro-differential equations", *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **23** (2006), no. 5, 695–711.
- [3] G. BARLES, E. CHASSEIGNE & C. IMBERT, "On the Dirichlet Problem for Second-Order Elliptic Integro-Differential Equations", *preprint*.

- [4] J. BERTOIN, *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [5] E. CHASSEIGNE, M. CHAVES & J. ROSSI, “Asymptotic behavior for nonlocal diffusion equations”, *J. Math. Pures Appl.* **86** (2006) 271-291.
- [6] R. CONT, E. VOLTCHKOVA, “Integro-differential equations for option prices in exponential Levy models”, *Finance Stoch.* **9** (2005), no. 3, 299–325.
- [7] L. SILVESTRE, “Hölder estimates for solutions of integro-differential equations like the fractional Laplace”, *Indiana Univ. Math. J.*, **55**(3), 1155-1174 (2006).
- [8] WOJBOR A. WOYCZYŃSKI, “Lévy processes in the physical sciences”, in *Lévy processes*, pages 241-266. Birkhäuser Boston, Boston, MA, 2001.