



**HAL**  
open science

## Dynamics random walks on Heisenberg groups

Nadine Guillotin-Plantard, René Schott

► **To cite this version:**

Nadine Guillotin-Plantard, René Schott. Dynamics random walks on Heisenberg groups. Journal of Theoretical Probability, 2006, 19, pp.377-395. hal-00135062

**HAL Id: hal-00135062**

**<https://hal.science/hal-00135062>**

Submitted on 6 Mar 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Dynamic Random Walks on Heisenberg groups

Nadine Guillotin-Plantard \*      René Schott†

Key words: Dynamic random walks, group theory, dynamical systems, Lévy stochastic area.

AMS Subject classification: 60G50, 60J15.

## Abstract

We prove a Guivarc'h law of large numbers and a central limit theorem for dynamic random walks on Heisenberg groups. The limiting distribution is explicitly given. To our knowledge this is the first study of dynamic random walks on non-commutative Lie groups.

## 1 Introduction

Random walks on Lie groups have been extensively studied over the last decades ([11, 12, 15]). Among these groups Heisenberg groups play a special role. These groups have their origin in quantum mechanics where they can be interpreted as the Lie algebras generated by the location operator, the momentum operator, and the identity operator. They are simply connected nilpotent Lie groups of rank 2 and one-dimensional center. Heisenberg groups are often considered as the simplest non-commutative Lie groups. The geometry of these groups has been investigated by A. Korányi ([13, 14]). Limit theorems for random walks on Heisenberg groups have been proved by P. Crepel, B. Roynette ([4]) and D. Neuenschwander ([15]) in connection with the resolution of Kesten's conjecture on the classification of recurrent and transient groups. A central limit theorem for nilpotent Lie groups has been proved by P. Crepel and A. Raugi ([3]). The novelty of this paper is in the dynamic model of random walks which we define on Heisenberg groups. The theory of dynamic random walks has been done by the first author in a commutative setting ([6, 7, 8, 9]). This paper is a first step/attempt in extending this theory to non-commutative algebraic structures.

---

\*Université Claude Bernard- Lyon 1 - 50, av. Tony-Garnier, 69366 Lyon Cedex 07, France, e-mail: nadine.guillotin@univ-lyon1.fr

†IECN and LORIA, Université Henri Poincaré-Nancy 1, 54506 Vandoeuvre-lès-Nancy, France, e-mail: schott@loria.fr

The organization is as follows: Section 2 provides some introductory material on Heisenberg groups and dynamic random walks. Section 3 is devoted to the proof of a strong law of large numbers and a central limit theorem. The limiting distribution is explicitly calculated. Our strong law of large numbers extends Guivarc'h's result [10] to dynamic random walks on Heisenberg groups.

## 2 Generalities on Heisenberg groups and dynamic random walks

The continuous Heisenberg group  $H_d$  is the group with underlying manifold  $\mathbb{R}^{2d+1}$  and group operation

$$X \cdot Y = \left( x_1 + y_1, x_2 + y_2, \dots, x_{2d} + y_{2d}, z_1 + z_2 + \frac{1}{2} \left( \sum_{i=1}^{2d-1} (x_i y_{i+1} - x_{i+1} y_i) \right) \right)$$

where  $X = (x_1, \dots, x_{2d}, z_1) \in \mathbb{R}^{2d+1}$  and  $Y = (y_1, \dots, y_{2d}, z_2) \in \mathbb{R}^{2d+1}$ .

$H_d$  is a nilpotent Lie group of rank  $2d$  and with one-dimensional center. In this paper we identify the Heisenberg group  $H_d$  with its Lie algebra  $\mathcal{H}_d$ .

For simplicity we focus on  $H_1$  but all results of this paper remain true for  $H_d$ . Let us remember that another representation  $H_1$  is as a group of upper triangular  $(3 \times 3)$ -matrices with 1's on the diagonal:

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}, x, y, z \in \mathbb{R}$$

If  $x, y, z$  are integers (in the ring  $\mathbb{Z}$ ) then we get the discrete Heisenberg group which is known to be the simplest non-abelian nilpotent group.

Let  $g = (x, y, z) \in H_1$  and

$$|g| = [(x^2 + y^2)^2 + z^2]^{1/4}.$$

Let  $\delta_r(g) = (rx, ry, r^2z)$  where  $r > 0$ . The mapping  $\delta_r$  is called dilation of ratio  $r$  on  $H_1$ . The mapping  $g \rightarrow |g|$  from  $H_1$  into  $\mathbb{R}^+$  is an homogeneous norm. This means that:

- i)  $|g| = 0$  if and only if  $g = 0$ ,
- ii)  $|g| = |-g|$ ,
- iii)  $|\delta_r(g)| = r |g|$ .

### Remark.

Let  $V$  be a compact neighborhood of the identity in  $H_1$ . The mapping defined by  $|g| = \inf\{n \in \mathbb{N}, g \in V^n\}$  is also an homogeneous norm.

It is known [10] that all homogeneous norms on  $H_1$  are equivalent.

$B(0, r) = \{g \in H_1; |g| \leq r\}$  ( $r > 0$ ) is called a Korányi ball. It replaces the traditional euclidean ball in the geometry developed by A. Korányi ([13, 14]).

We introduce now our model of dynamic random walks on  $H_1$ .

Let  $(X_n, Y_n, Z_n)_{n \geq 1}$  be a sequence of independent random variables with values in  $H_1$  defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . Let  $(E, \mathcal{A}, \mu, T)$  be a dynamical system where  $(E, \mathcal{A}, \mu)$  is a probability space and  $T$  is a transformation defined on  $E$  preserving the measure  $\mu$ . Let  $f_i, i = 1, 2, 3$  be functions defined on  $E$  with values in  $[0, \frac{1}{3}]$ . Let  $x \in E$  and  $(e_j)_{1 \leq j \leq 3}$  be the unit coordinate vectors of  $\mathbb{R}^3$ . For each  $i \geq 1$ , the distribution of the random vector  $M_i := (X_i, Y_i, Z_i)$  is given by

$$\mathbb{P}(M_i = z) = \begin{cases} f_j(T^i x) & \text{if } z = e_j \\ \frac{1}{3} - f_j(T^i x) & \text{if } z = -e_j \\ 0 & \text{otherwise} \end{cases}$$

We are interested in the right dynamic random walk

$$S_n := (X_1, Y_1, Z_1) \cdot (X_2, Y_2, Z_2) \cdots (X_n, Y_n, Z_n), n \geq 1.$$

We prove in the next sections a strong law of large numbers and a central limit theorem for  $(S_n)_{n \geq 1}$ .

### 3 Limit theorems

#### 3.1 Law of large numbers

Y. Guivarc'h [10] proved a strong law of large numbers for random walks on Lie groups. We prove a similar result but for dynamic random walks on Heisenberg groups.

In this section, we assume that the dynamical system  $(E, \mathcal{A}, \mu, T)$  is *ergodic* i.e. there exists a unique invariant measure  $\mu$ . The functions  $f_i, i = 1, 2, 3$  are all with integral  $1/6$ . Let  $d(0, \frac{S_n}{n}) = \frac{1}{n} |S_n|$ ,  $d$  will be called Korányi distance.

**Theorem 3.1** *For  $\mu$ -almost every  $x \in E$ , as  $n \rightarrow +\infty$ ,*

$$d\left(0, \frac{S_n}{n}\right) \rightarrow 0 \quad \mathbb{P} - a.s..$$

**Proof:**

From the definition of the Korányi distance,

$$d\left(0, \frac{S_n}{n}\right) = \left( \left( \left( \frac{S_n^{(1)}}{n} \right)^2 + \left( \frac{S_n^{(2)}}{n} \right)^2 \right)^2 + \left( \frac{S_n^{(3)}}{n^2} \right)^2 \right)^{1/4}.$$

The two first coordinates of  $S_n$  are given by

$$S_n^{(1)} = X_1 + X_2 + \dots + X_n$$

and

$$S_n^{(2)} = Y_1 + Y_2 + \dots + Y_n.$$

From Birkhoff's theorem, the sequence

$$\frac{\mathbb{E}(S_n^{(1)})}{n} = \frac{1}{n} \sum_{k=1}^n (2f_1(T^k x) - \frac{1}{3})$$

converges for  $\mu$ -almost every  $x \in E$  to  $\int_E (2f_1 - \frac{1}{3}) d\mu = 0$ . With the same arguments, the sequence

$$\frac{\mathbb{E}(S_n^{(2)})}{n} = \frac{1}{n} \sum_{k=1}^n (2f_2(T^k x) - \frac{1}{3})$$

is proved to converge for  $\mu$ -almost every  $x \in E$  to  $\int_E (2f_2 - \frac{1}{3}) d\mu = 0$ . Thus, from Kolmogorov's theorem, the sequences  $(\frac{S_n^{(1)}}{n})_n$  and  $(\frac{S_n^{(2)}}{n})_n$  converge  $\mathbb{P}$ -almost surely to 0 for  $\mu$ -almost every  $x \in E$ .

The third component of the dynamic random walk on  $H_1$  is given by

$$S_n^{(3)} = Z_1 + Z_2 + \dots + Z_n + \frac{1}{2} \{X_1 Y_2 + (X_1 + X_2) Y_3 + \dots + (X_1 + \dots + X_{n-1}) Y_n \\ - Y_1 X_2 - (Y_1 + Y_2) X_3 - \dots - (Y_1 + \dots + Y_{n-1}) X_n\}.$$

For  $\mu$ -almost every  $x \in E$ , the sequence

$$\frac{1}{n^2} (Z_1 + Z_2 + \dots + Z_n)$$

converges  $\mathbb{P}$ -almost surely to 0 so we are only interested in the asymptotic behavior of the sequence

$$\frac{T_n}{n^2} = \frac{1}{2n^2} (T_n^{(1)} - T_n^{(2)})$$

where

$$T_n^{(1)} = X_1 Y_2 + (X_1 + X_2) Y_3 + \dots + (X_1 + \dots + X_{n-1}) Y_n \\ T_n^{(2)} = Y_1 X_2 + (Y_1 + Y_2) X_3 + \dots + (Y_1 + \dots + Y_{n-1}) X_n.$$

Let us prove that  $T_n^{(1)}/n^2$  converges almost surely to 0. Since the  $Y$ 's can only take values 0 and  $\pm 1$ , we can bound  $|\frac{T_n^{(1)}}{n^2}|$  by

$$\frac{1}{\left(\sum_{i=2}^n (i-1) |Y_i|\right)} \sum_{i=2}^n (i-1) |Y_i| \frac{1}{i-1} \left| \sum_{j=1}^{i-1} X_j \right|.$$

We conclude by using Toeplitz's Theorem (see for instance Theorem 2.34 in [5]) since the sequence  $\sum_{j=1}^n X_j/n$  converges  $\mathbb{P}$ -almost surely to 0 for  $\mu$ -almost every  $x \in E$  as  $n$  goes to infinity. By inverting the random variables  $X_j$ 's and  $Y_j$ 's, we evidently get that  $T_n^{(2)}/n^2$  converges  $\mathbb{P}$ -almost surely to 0 for  $\mu$ -almost every  $x \in E$  as  $n$  goes to infinity. Consequently,  $T_n/n^2$  converges  $\mathbb{P}$ -almost surely to 0 for  $\mu$ -almost every  $x \in E$  as  $n$  goes to infinity as well as the third component of the dynamic random walk. So the theorem is proved.

### 3.2 Central limit theorem

In order to prove a central limit theorem for the dynamic random walk on the Heisenberg group  $H_1$  we need to add assumptions on the functions  $f_i, i = 1, 2, 3$ . Let  $\mathcal{C}$  denote the class of functions  $f \in L^1(E, \mu)$  satisfying the following condition

$$\sup_{x \in E} \left| \sum_{i=1}^n \left( f(T^i x) - \int_E f(x) d\mu(x) \right) \right| = o(\sqrt{n}).$$

We denote by  $A = (a_{ij})_{1 \leq i, j \leq 3}$  the matrix with coefficients

$$\begin{aligned} a_{jj} &= \frac{4}{9} \int_E (1 - 9f_j^2(x)) d\mu(x) \\ a_{ij} &= a_{ji} = \frac{1}{9} \int_E (1 - 36f_i(x)f_j(x)) d\mu(x). \end{aligned}$$

Let us assume that  $f_i, f_i f_j \in \mathcal{C}$  for every  $i, j = 1, 2, 3$  and  $\int_E f_i d\mu = \frac{1}{6}$ . For every  $x \in E$ , the two first coordinates of  $S_n$  converge  $\mathbb{P}$ -almost surely to 0 (see the proof of Theorem 3.1). In the third component given by

$$\begin{aligned} Z_1 + Z_2 + \dots + Z_n + \frac{1}{2} \{ X_1 Y_2 + (X_1 + X_2) Y_3 + \dots + (X_1 + \dots + X_{n-1}) Y_n \\ - Y_1 X_2 - (Y_1 + Y_2) X_3 - \dots - (Y_1 + \dots + Y_{n-1}) X_n \}, \end{aligned}$$

for every  $x \in E$ , the sequence

$$\frac{1}{n} (Z_1 + Z_2 + \dots + Z_n)$$

also converges  $\mathbb{P}$ -almost surely to 0 so we are only interested in the asymptotic distribution of the sequence

$$T_n = \frac{1}{2} \{ X_1 Y_2 + (X_1 + X_2) Y_3 + \dots + (X_1 + \dots + X_{n-1}) Y_n - Y_1 X_2 - (Y_1 + Y_2) X_3 - \dots - (Y_1 + \dots + Y_{n-1}) X_n \}.$$

We assume that the matrix  $C$  defined by  $(a_{ij})_{1 \leq i, j \leq 2}$  is diagonal, namely

$$C = \begin{pmatrix} a_{11} & 0 \\ 0 & a_{22} \end{pmatrix}$$

with  $a_{11} = 4[\frac{1}{9} - \int_E f_1^2 d\mu]$  and  $a_{22} = 4[\frac{1}{9} - \int_E f_2^2 d\mu]$ .

Since our random variables  $(X_i)_{i \geq 1}$  and  $(Y_i)_{i \geq 1}$  are not centered, we need to make the following assumption **(A)**:

$$\sup_{x \in E} \left| \sum_{i=2}^k \sum_{j=1}^{i-1} \left( f_2(T^i x) - \frac{1}{6} \right) \left( f_1(T^j x) - \frac{1}{6} \right) \right| = o(k).$$

**Theorem 3.2** For every  $x \in E$ , for every  $t > 0$ , the sequence

$$\left( \frac{T_{[nt]}}{n} \right)_{n \geq 1}$$

converges in distribution to the random variable

$$A(t) = \frac{1}{2} \left\{ \int_0^t B_s^{(1)} dB_s^{(2)} - \int_0^t B_s^{(2)} dB_s^{(1)} \right\}$$

where  $B_t = (B_t^{(1)}, B_t^{(2)})$  is a two-dimensional Brownian motion with zero mean and Covariance matrix  $Ct$ . The density of  $A(t)$  is given by

$$d_t(x) = \frac{1}{t\sqrt{a_{11}a_{22}} \cosh\left(\frac{\pi x}{t\sqrt{a_{11}a_{22}}}\right)}.$$

The random variable  $A(t)$  is usually called the Lévy stochastic area. In our context  $A(t)$  is driven by dynamic random walks, therefore we will call  $A(t)$ , by analogy, the dynamic Lévy stochastic area.

**Remarks:**

1- The hypotheses on the functions  $f_1, f_2$  and  $f_1 f_2$  imply that the assumption **(A)** is satisfied as soon as for every  $i, j \geq 1$ , for every  $x \in E$ ,

$$(f_2(T^i x) - \frac{1}{6})(f_1(T^j x) - \frac{1}{6}) = (f_2(T^j x) - \frac{1}{6})(f_1(T^i x) - \frac{1}{6})$$

i.e., for instance, in the following cases:

- a)-  $f_1 = f_2$  (but not necessarily equal to  $1/6$ ).
- b)-  $f_1$  or  $f_2$  equal to  $1/6$ .
- c)-  $f_2 = \frac{1}{3} - f_1$ .

Moreover, this assumption is necessary. Given a dynamical system  $(E, \mathcal{A}, \mu, T)$ , the fact that the functions  $f_1, f_2, f_1 f_2$  belong to the class  $\mathcal{C}$ ,  $\int_E f_1 d\mu = \int_E f_2 d\mu = 1/6$  and  $\int_E f_1 f_2 d\mu = 1/36$  does not imply that the assumption **(A)** is necessarily verified. For instance, take the rotation  $T$  on the one-dimensional torus  $\mathbb{T}^1$  with rational angle equal to  $1/4$  and the particular functions  $f_1 = \frac{1}{3}\mathbf{1}_{[0, \frac{1}{4}]} + \frac{1}{6}\mathbf{1}_{[\frac{1}{2}, 1]}$  and  $f_2 = \frac{1}{3}\mathbf{1}_{[0, \frac{1}{2}]}$ . The functions  $f_1, f_2$  and  $f_1 f_2$  are clearly integrable,  $\int f_1 dx = \int f_2 dx = 1/6$  and  $\int f_1 f_2 dx = 1/36$  and they belong to the class  $\mathcal{C}$  associated to the dynamical system. A calculation also gives that for every  $x \in \mathbb{T}^1$ ,

$$\sum_{i=2}^k \sum_{j=1}^{i-1} (f_2(T^i x) - \frac{1}{6})(f_1(T^j x) - \frac{1}{6}) \sim \frac{k}{4} \frac{1}{36}, \quad k \rightarrow +\infty.$$

So assumption **(A)** is not satisfied and for these exotic cases, Theorem (3.2) holds by adding a drift to the limit process.

2- Given a dynamical system  $(E, \mathcal{A}, \mu, T)$ , the class  $\mathcal{C}$  is quite difficult to determine. However, for

particular cases, it is possible to find a large sub-class of  $\mathcal{C}$ . We refer to papers [6, 7, 8] where this question has been studied in details. When  $T$  is an irrational rotation on the one-dimensional torus  $\mathbb{T}^1$ , it was proved that every function with bounded variation belongs to the class  $\mathcal{C}$ . For example, we can choose  $f_1(x) = 1/6$  and  $f_2(x) = 1/3 \cos^2(2\pi x)$  which takes its values in  $[0, 1/3]$ . The hypothesis **(A)** is clearly satisfied thanks to remark 1. The integral of  $f_2$  is equal to  $1/6$  so  $a_{12} = a_{21} = 0$ . After simple computations,  $a_{11} = 1/3$  and  $a_{22} = 5/18$ . In this particular example, the density of the limit distribution  $A(t)$  in Theorem 3.2 is then given by

$$d_t(x) = \frac{1}{t\sqrt{5/54}} \cosh\left(\frac{\pi x}{t\sqrt{5/54}}\right), \quad x \in \mathbb{R}.$$

### 3.2.1 A preliminary result

In order to prove Theorem 3.2, we shall need a central limit theorem for the dynamic  $\mathbb{Z}^2$ -random walk  $(S_n^{(1)}, S_n^{(2)})_{n \geq 1}$ .

**Proposition 3.1** *For every  $x \in E$ , the sequence of random vectors  $\frac{1}{\sqrt{n}}(S_n^{(1)}, S_n^{(2)})$ ,  $n \geq 1$  converges in distribution, as  $n \rightarrow +\infty$ , to the centered Gaussian random variable  $G_0$  with Covariance matrix  $C$ .*

**Proof:**

Let us introduce the characteristic function  $\phi_n$  of

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n (X_k, Y_k), \quad n \geq 1.$$

By independence of the random vectors  $(X_k, Y_k)$ ,  $k \geq 1$ ,

$$\begin{aligned} \phi_n(u) &= \prod_{k=1}^n \mathbb{E} \left( \exp \left( i \frac{u_1 X_k + u_2 Y_k}{\sqrt{n}} \right) \right) \\ &= \prod_{k=1}^n Q_n^{(k)}(u_1, u_2) \end{aligned}$$

where

$$Q_n^{(k)}(u_1, u_2) = \frac{1}{3} \left( 1 + \sum_{j=1}^2 \cos\left(\frac{u_j}{\sqrt{n}}\right) \right) + 2i \sum_{j=1}^2 \left( f_j(T^k x) - \frac{1}{6} \right) \sin\left(\frac{u_j}{\sqrt{n}}\right)$$

A direct calculation gives

$$|Q_n^{(k)}(u)|^2 = \frac{1}{9} \left( 1 + \sum_{j=1}^2 \left( 1 - \frac{u_j^2}{2n} \right) + \mathcal{O}(n^{-2}) \right)^2 + \frac{1}{9} \sum_{j=1}^2 \left( 6f_j(T^k x) - 1 \right)^2 \frac{u_j^2}{n}$$



$$\begin{aligned}
& + \frac{2}{9}(6f_1(T^k x) - 1)(6f_2(T^k x) - 1)\frac{u_1 u_2}{n} + \mathcal{O}(n^{-2}) \\
& = 1 - \frac{1}{3n} \sum_{j=1}^2 u_j^2 + \frac{1}{9n} \sum_{j=1}^2 u_j^2 (6f_j(T^k x) - 1)^2 \\
& + \frac{2}{9n}(6f_1(T^k x) - 1)(6f_2(T^k x) - 1)u_1 u_2 + \mathcal{O}(n^{-2})
\end{aligned}$$

and then

$$\begin{aligned}
|\phi_n(u)| & = \prod_{k=1}^n |Q_n^{(k)}(u)| \\
& = \exp\left(-\frac{1}{2} \langle u, Cu \rangle + o(1)\right)
\end{aligned}$$

The imaginary part of the characteristic function can be rewritten as

$$\begin{aligned}
& \prod_{k=1}^n \exp\left(i \arctan\left(\frac{\sum_{j=1}^2 (6f_j(T^k x) - 1) \sin(\frac{u_j}{\sqrt{n}})}{(1 + \sum_{j=1}^2 \cos(\frac{u_j}{\sqrt{n}}))}\right)\right) \\
& = \exp\left(\frac{i}{3} \sum_{k=1}^n \sum_{j=1}^2 (6f_j(T^k x) - 1) \frac{u_j}{\sqrt{n}} + o(1)\right) = 1 + o(1)
\end{aligned}$$

using the fact that  $f_1, f_2$  belong to the class  $\mathcal{C}$  and for every  $j = 1, 2$ , the integral of  $f_j$  is equal to  $1/6$ .

### 3.2.2 Proof of the central limit theorem 3.2

We only prove Theorem 3.2 for the particular value  $t$  equal to 1. The proof can easily be adapted to get the result for every  $t > 0$ . The proof of the theorem is decomposed into four parts.

a) Straightforward calculations give us

$$\begin{aligned}
2T_{nk} & = [N_0^{(k)} + \dots + N_{n-1}^{(k)}] - [\tilde{N}_0^{(k)} + \dots + \tilde{N}_{n-1}^{(k)}] \\
& + [M_0^{(k)}(Q_1^{(k)} - Q_0^{(k)}) + \dots + M_{n-2}^{(k)}(Q_{n-1}^{(k)} - Q_{n-2}^{(k)})] \\
& - [Q_0^{(k)}(M_1^{(k)} - M_0^{(k)}) + \dots + Q_{n-2}^{(k)}(M_{n-1}^{(k)} - M_{n-2}^{(k)})]
\end{aligned}$$

where

$$\begin{aligned}
N_p^{(k)} & = X_{pk+1}Y_{pk+2} + (X_{pk+1} + X_{pk+2})Y_{pk+3} + \dots + (X_{pk+1} + \dots + X_{(p+1)k-1})Y_{(p+1)k} \\
\tilde{N}_p^{(k)} & = Y_{pk+1}X_{pk+2} + (Y_{pk+1} + Y_{pk+2})X_{pk+3} + \dots + (Y_{pk+1} + \dots + Y_{(p+1)k-1})X_{(p+1)k} \\
M_p^{(k)} & = X_1 + \dots + X_{(p+1)k}
\end{aligned}$$

$$Q_p^{(k)} = Y_1 + \dots + Y_{(p+1)k}.$$

The choice of this decomposition can appear artificial but it will be justified in the sequel.

b) Let  $x \in E$  fixed. From Proposition 3.1, for every  $p \geq 0$ , the sequence of random vectors

$$W_p^{(k)} := \frac{1}{\sqrt{k}} \sum_{j=pk+1}^{(p+1)k} (X_j, Y_j), \quad k \geq 1$$

converges in distribution, as  $k \rightarrow +\infty$ , to the centered Gaussian random vector  $G_0$  with covariance matrix  $C$ . Since the random vectors  $(W_p^{(k)})_{p \geq 0}$  are independent, we deduce that  $(W_0^{(k)}, W_1^{(k)}, \dots, W_{n-1}^{(k)})$  converges in distribution, as  $k \rightarrow +\infty$ , to the centered Gaussian random vector  $(G_0, G_1, \dots, G_{n-1})$  where the  $G_i$ 's are independent copies of  $G_0$ . Since  $C$  is diagonal, each random vector  $G_i$  can be decomposed in two independent centered Gaussian random variables  $G_i^{(1)}$  and  $G_i^{(2)}$  with variance  $a_{11}$  and  $a_{22}$  respectively. So it implies that  $(W_0^{(k)}, W_1^{(k)}, \dots, W_{n-1}^{(k)})$  converges in distribution, as  $k \rightarrow +\infty$ , to the centered Gaussian random vector  $(G_0^{(1)}, G_0^{(2)}, G_1^{(1)}, G_1^{(2)}, \dots, G_{n-1}^{(1)}, G_{n-1}^{(2)})$ . Now,  $W_p^{(k)} = (M_p^{(k)} - M_{p-1}^{(k)}, Q_p^{(k)} - Q_{p-1}^{(k)})/\sqrt{k}$  (with the convention  $M_{-1}^{(k)} = Q_{-1}^{(k)} = 0$ ). The convergence in distribution being preserved by linear transformation we get that

$$\frac{1}{\sqrt{k}}(M_0^{(k)}, M_1^{(k)}, \dots, M_{n-1}^{(k)}, Q_0^{(k)}, Q_1^{(k)}, \dots, Q_{n-1}^{(k)})$$

converges in distribution, as  $k \rightarrow +\infty$ , to the centered Gaussian random vector  $(G_0^{(1)}, G_0^{(1)} + G_1^{(1)}, \dots, G_0^{(1)} + \dots + G_{n-1}^{(1)}, G_0^{(2)}, G_0^{(2)} + G_1^{(2)}, \dots, G_0^{(2)} + \dots + G_{n-1}^{(2)})$ . Then,

$$\begin{aligned} & \frac{1}{k}[M_0^{(k)}(Q_1^{(k)} - Q_0^{(k)}) + \dots + M_{n-2}^{(k)}(Q_{n-1}^{(k)} - Q_{n-2}^{(k)})] \\ & - \frac{1}{k}[Q_0^{(k)}(M_1^{(k)} - M_0^{(k)}) + \dots + Q_{n-2}^{(k)}(M_{n-1}^{(k)} - M_{n-2}^{(k)})] \end{aligned}$$

converges in distribution, as  $k \rightarrow +\infty$ , to the random variable

$$\begin{aligned} & [G_0^{(1)}G_1^{(2)} + (G_0^{(1)} + G_1^{(1)})G_2^{(2)} + \dots + (G_0^{(1)} + \dots + G_{n-2}^{(1)})G_{n-1}^{(2)}] \\ & - [G_0^{(2)}G_1^{(1)} + (G_0^{(2)} + G_1^{(2)})G_2^{(1)} \dots + (G_0^{(2)} + \dots + G_{n-2}^{(2)})G_{n-1}^{(1)}]. \end{aligned}$$

c) Let us now prove that

$$\text{Var} \left( N_0^{(k)} + \dots + N_{n-1}^{(k)} \right) = \mathcal{O}(nk^2) \quad (1)$$

and

$$\text{Var} \left( \tilde{N}_0^{(k)} + \dots + \tilde{N}_{n-1}^{(k)} \right) = \mathcal{O}(nk^2). \quad (2)$$

The above result is quite evident in the classical case studied in [11] where the random variables  $X_i$  and  $Y_i$  are centered and uncorrelated but due to the temporal inhomogeneity of our model it is **not at all the case** and we need the following lemma.

**Lemma 3.1** *Uniformly in  $p \geq 0$ ,*

$$\text{Var} (N_p^{(k)}) = \mathcal{O}(k^2) \quad (3)$$

and

$$\text{Var} (\tilde{N}_p^{(k)}) = \mathcal{O}(k^2). \quad (4)$$

**Proof:**

Let us recall that

$$N_p^{(k)} = \sum_{l=1}^{k-1} Y_{pk+l+1} \left( \sum_{i=1}^l X_{pk+i} \right).$$

For every random variable  $X$  such that  $\mathbb{E}(|X|) < +\infty$ , we define  $\bar{X} = X - \mathbb{E}(X)$ . The random variable  $N_p^{(k)}$  can be rewritten

$$N_p^{(k)} = \Sigma_1(k) + \Sigma_2(k) + \Sigma_3(k) - \Sigma_4(k)$$

where

$$\begin{aligned} \Sigma_1(k) &= \sum_{l=1}^{k-1} \bar{Y}_{pk+l+1} \left( \sum_{i=1}^l \bar{X}_{pk+i} \right) \\ \Sigma_2(k) &= \sum_{l=1}^{k-1} \mathbb{E}(Y_{pk+l+1}) \left( \sum_{i=1}^l X_{pk+i} \right) \\ \Sigma_3(k) &= \sum_{l=1}^{k-1} Y_{pk+l+1} \left( \sum_{i=1}^l \mathbb{E}(X_{pk+i}) \right) \\ \Sigma_4(k) &= \sum_{l=1}^{k-1} \mathbb{E}(Y_{pk+l+1}) \left( \sum_{i=1}^l \mathbb{E}(X_{pk+i}) \right). \end{aligned}$$

Using that the random variables  $\bar{X}_i$  and  $\bar{Y}_j$  are independent when  $i \neq j$  and centered, a direct computation gives

$$\text{Var} (\Sigma_1(k)) \leq Ck^2$$

where  $C > 0$  is a constant independent of  $p$ . Furthermore, the random variable  $\Sigma_2(k)$  can be rewritten as

$$\Sigma_2(k) = \sum_{l=1}^{k-1} X_{pk+l} \left( \sum_{i=l}^{k-1} \mathbb{E}(Y_{pk+i+1}) \right)$$

thus

$$\begin{aligned} \text{Var} (\Sigma_2(k)) &= \sum_{l=1}^{k-1} \left( \sum_{i=l}^{k-1} \mathbb{E}(Y_{pk+i+1}) \right)^2 \text{Var} (X_{pk+l}) \\ &= 4 \sum_{l=1}^{k-1} \left( \sum_{i=l}^{k-1} (f_2(T^{pk+i+1}x) - \frac{1}{6}) \right)^2 \text{Var} (X_{pk+l}) \end{aligned}$$

Since the function  $f_2$  belongs to the class  $\mathcal{C}$  and  $\int_E f_2 d\mu = 1/6$ , there exists a constant  $C$  which does not depend on  $p$  and  $k$  such that for every  $l \geq 1$ ,

$$\left| \sum_{i=1}^{k-l} \left( f_2(T^{pk+l+i}x) - \frac{1}{6} \right) \right| \leq C\sqrt{k}.$$

Then,

$$\text{Var}(\Sigma_2(k)) \leq C \sum_{l=1}^{k-1} \left| \sum_{j=1}^{k-l} \left( f_2(T^{pk+l+i}x) - \frac{1}{6} \right) \right|^2 = \mathcal{O}(k^2).$$

The variance of the third sum  $\Sigma_3(k)$  is estimated in the same manner using the fact that the function  $f_1$  belongs to the class  $\mathcal{C}$  and  $\int_E f_1 d\mu = 1/6$ . So, we get

$$\text{Var}(\Sigma_3(k)) = \mathcal{O}(k^2).$$

The variance of the last sum is evidently zero.

The following inequality holds for square integrable random variables  $X$  and  $Y$

$$\text{Var}(X + Y) \leq 2(\text{Var}(X) + \text{Var}(Y)).$$

So, by applying twice this inequality, we get

$$\text{Var}(N_p^{(k)}) \leq 4 \sum_{i=1}^4 \text{Var}(\Sigma_i(k)) = \mathcal{O}(k^2)$$

uniformly in  $p$ . The estimation (4) is obtained in the same manner by inverting the  $X$ 's and the  $Y$ 's.

From Lemma 3.1 we can now establish (1) as follows: the random variables  $N_p^{(k)}$  and  $N_{p'}^{(k)}$  are independent when  $p \neq p'$ , so we have

$$\text{Var}(N_0^{(k)} + \dots + N_{n-1}^{(k)}) = \sum_{p=0}^{n-1} \text{Var}(N_p^{(k)}) \leq Cnk^2$$

using Lemma 3.1, so (1) is proved and (2) can be obtained by inverting the  $X$ 's and the  $Y$ 's.

Let  $\varepsilon > 0$ ,

$$\begin{aligned} & \mathbb{P}\left(\frac{1}{2nk} \left| \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] - \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right| \geq \varepsilon\right) \\ & \leq \mathbb{P}\left(\frac{1}{nk} \left| \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] \right| \geq \varepsilon\right) + \mathbb{P}\left(\frac{1}{nk} \left| \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right| \geq \varepsilon\right) \\ & \leq \frac{Cnk^2}{\varepsilon^2 n^2 k^2} = \frac{C}{\varepsilon^2 n} \end{aligned}$$

Consequently,

$$\mathbb{P}\left(\frac{1}{2nk} \left| \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] - \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right| \geq \varepsilon\right) \rightarrow 0$$

as  $n$  goes to infinity, uniformly in  $k$ . Using that  $|e^{i\theta} - 1| \leq |\theta| \wedge 2$ , we obtain

$$\begin{aligned} & \mathbb{E}\left(\left| \exp\left\{i\frac{\theta}{2nk} \left( \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] - \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right)\right\} - 1 \right| \right) \\ & \leq \varepsilon + 2\mathbb{P}\left(\frac{1}{2nk} \left| \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] - \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right| \geq \varepsilon\right) \rightarrow 0 \end{aligned} \quad (5)$$

as  $n$  goes to infinity, uniformly in  $k$ .

Let us remark that for every  $\theta \in \mathbb{R}$ ,

$$\begin{aligned} & \left| \mathbb{E}\left(e^{i\theta \frac{T_{nk}}{nk}}\right) - \mathbb{E}\left(\exp\left[i\frac{\theta}{2nk} \left\{ M_0^{(k)}(Q_1^{(k)} - Q_0^{(k)}) + \dots + M_{n-2}^{(k)}(Q_{n-1}^{(k)} - Q_{n-2}^{(k)}) \right. \right. \right. \right. \\ & \left. \left. \left. - Q_0^{(k)}(M_1^{(k)} - M_0^{(k)}) - \dots - Q_{n-2}^{(k)}(M_{n-1}^{(k)} - M_{n-2}^{(k)}) + \sum_{p=0}^{n-1} [\mathbb{E}(N_p^{(k)}) - \mathbb{E}(\tilde{N}_p^{(k)})] \right\} \right]\right) \Big| \\ & \leq \mathbb{E}\left(\left| \exp\left\{i\frac{\theta}{2nk} \left( \sum_{p=0}^{n-1} [N_p^{(k)} - \mathbb{E}(N_p^{(k)})] - \sum_{p=0}^{n-1} [\tilde{N}_p^{(k)} - \mathbb{E}(\tilde{N}_p^{(k)})] \right)\right\} - 1 \right| \right) \end{aligned}$$

converging to 0 as  $n$  goes to infinity, uniformly in  $k$ .

Let  $\varepsilon > 0$ , we can choose  $n_0$  such that for every  $k$ ,

$$\begin{aligned} & \left| \mathbb{E}\left(e^{i\theta \frac{T_{n_0 k}}{n_0 k}}\right) - \mathbb{E}\left(\exp\left[i\frac{\theta}{2n_0 k} \left\{ M_0^{(k)}(Q_1^{(k)} - Q_0^{(k)}) + \dots + M_{n_0-2}^{(k)}(Q_{n_0-1}^{(k)} - Q_{n_0-2}^{(k)}) \right. \right. \right. \right. \\ & \left. \left. \left. - Q_0^{(k)}(M_1^{(k)} - M_0^{(k)}) - \dots - Q_{n_0-2}^{(k)}(M_{n_0-1}^{(k)} - M_{n_0-2}^{(k)}) + \sum_{p=0}^{n_0-1} [\mathbb{E}(N_p^{(k)}) - \mathbb{E}(\tilde{N}_p^{(k)})] \right\} \right]\right) \Big| \leq \varepsilon. \end{aligned}$$

The characteristic function of the random variable

$$\begin{aligned} & [G_0^{(1)}G_1^{(2)} + (G_0^{(1)} + G_1^{(1)})G_2^{(2)} + \dots + (G_0^{(1)} + \dots + G_{n-2}^{(1)})G_{n-1}^{(2)}] \\ & - [G_0^{(2)}G_1^{(1)} + (G_0^{(2)} + G_1^{(2)})G_2^{(1)} + \dots + (G_0^{(2)} + \dots + G_{n-2}^{(2)})G_{n-1}^{(1)}] \end{aligned}$$

can be written as

$$I_n(\theta) := \frac{1}{(2\pi\sqrt{a_{11}a_{22}})^n} \int_{\mathbb{R}^{2n}} e^{g_n(x,y)} dx_1 \dots dx_n dy_1 \dots dy_n$$

where  $g_n$  is the function defined for every  $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$  by

$$\begin{aligned} g_n(x, y) &= \frac{i\theta}{2n} [x_1 y_2 + (x_1 + x_2) y_3 + \dots + (x_1 + \dots + x_{n-1}) y_n] \\ &- y_1 x_2 - (y_1 + y_2) x_3 - \dots - (y_1 + \dots + y_{n-1}) x_n \\ &- \frac{1}{2a_{11}} (x_1^2 + \dots + x_n^2) - \frac{1}{2a_{22}} (y_1^2 + \dots + y_n^2). \end{aligned}$$

Now from Lemma 37 in [11], we know that

$$\lim_{n \rightarrow +\infty} \frac{1}{(2\pi)^n} \int_{\mathbb{R}^{2n}} e^{g_n(x\sqrt{a_{11}}, y\sqrt{a_{22}})} dx dy = \frac{1}{\cosh(\frac{\theta\sqrt{a_{11}a_{22}}}{2})}$$

uniformly in  $\theta$  on a compact set of  $\mathbb{R}$ . So we also impose  $n_0$  to be large enough so as to have

$$\left| I_{n_0}(\theta) - \frac{1}{\cosh(\frac{\theta\sqrt{a_{11}a_{22}}}{2})} \right| \leq \varepsilon$$

uniformly in  $\theta$  on a compact set.

Under the assumption **(A)**, it is possible to control the drift coming from the non-centered random variables  $N_p^{(k)}$  and  $\tilde{N}_p^{(k)}$ .

**Lemma 3.2** *Under hypothesis **(A)**,*

$$\frac{1}{n_0 k} \sum_{p=0}^{n_0-1} [\mathbb{E}(N_p^{(k)}) - \mathbb{E}(\tilde{N}_p^{(k)})] = o(k). \quad (6)$$

**Proof:**

From the definition of the law of the random variables  $X_i$  and  $Y_i$ , the above sum can be explicitly calculated and bounded as follows:

$$\begin{aligned} & \left| \frac{1}{n_0 k} \sum_{p=0}^{n_0-1} [\mathbb{E}(N_p^{(k)}) - \mathbb{E}(\tilde{N}_p^{(k)})] \right| \\ & \leq \frac{1}{k} \sup_{x \in E} \left| \sum_{l=1}^k \sum_{i=1}^l [(2f_2(T^l x) - \frac{1}{3})(2f_1(T^i x) - \frac{1}{3}) - (2f_1(T^l x) - \frac{1}{3})(2f_2(T^i x) - \frac{1}{3})] \right| \end{aligned}$$

Using Abel's summation by parts,

$$\begin{aligned} \sum_{l=1}^k \sum_{i=1}^l (2f_1(T^l x) - \frac{1}{3})(2f_2(T^i x) - \frac{1}{3}) &= \sum_{l=1}^k (2f_1(T^l x) - \frac{1}{3}) \sum_{l=1}^k (2f_2(T^l x) - \frac{1}{3}) \\ &\quad - \sum_{l=2}^k (2f_2(T^l x) - \frac{1}{3}) \sum_{i=1}^{l-1} (2f_1(T^i x) - \frac{1}{3}) \end{aligned}$$

Then,

$$\begin{aligned} & \sum_{l=1}^k \sum_{i=1}^l [(2f_2(T^l x) - \frac{1}{3})(2f_1(T^i x) - \frac{1}{3}) - (2f_1(T^l x) - \frac{1}{3})(2f_2(T^i x) - \frac{1}{3})] \\ &= 2 \sum_{l=2}^k (2f_2(T^l x) - \frac{1}{3}) \sum_{i=1}^{l-1} (2f_1(T^i x) - \frac{1}{3}) - \sum_{l=1}^k (2f_1(T^l x) - \frac{1}{3}) \sum_{l=1}^k (2f_2(T^l x) - \frac{1}{3}) \\ &\quad + \sum_{l=1}^k (2f_1(T^l x) - \frac{1}{3})(2f_2(T^l x) - \frac{1}{3}) \end{aligned}$$

In the right hand side the second sum divided by  $k$  goes to 0 uniformly in  $x$  since  $f_1$  and  $f_2$  belong to the class  $\mathcal{C}$  and  $\int_E f_i d\mu = 1/6, i = 1, 2$ . The third one divided by  $k$  also goes to 0 uniformly in  $x \in E$  since  $\int_E f_1 f_2 d\mu = 1/36$ . So, as soon as **(A)** is satisfied, (6) holds.

From item b) and the previous lemma, the characteristic function of the random variable

$$\begin{aligned} & \frac{1}{2n_0k} \{M_0^{(k)}(Q_1^{(k)} - Q_0^{(k)}) + \dots + M_{n_0-2}^{(k)}(Q_{n_0-1}^{(k)} - Q_{n_0-2}^{(k)}) \\ & - Q_0^{(k)}(M_1^{(k)} - M_0^{(k)}) - \dots - Q_{n_0-2}^{(k)}(M_{n_0-1}^{(k)} - M_{n_0-2}^{(k)}) + \sum_{p=0}^{n_0-1} [\mathbb{E}(N_p^{(k)}) - \mathbb{E}(\tilde{N}_p^{(k)})]\} \end{aligned}$$

converges, as  $k$  goes to infinity, to  $I_{n_0}(\theta)$ . Hence there exists  $k_0$  such that for every  $k \geq k_0$ ,

$$\left| \mathbb{E} \left( e^{i\theta \frac{T_{n_0k}}{n_0k}} \right) - \frac{1}{\cosh\left(\frac{\theta \sqrt{a_{11} a_{22}}}{2}\right)} \right| \leq 3\varepsilon. \quad (7)$$

d) Let  $p = n_0k + q, 0 \leq q < n_0, k \geq k_0$  (remark that  $p \geq n_0k_0$ ) and define

$$\begin{aligned} V_{n_0}^{(k)} &= T_{n_0k+q} - T_{n_0k} \\ &= \frac{1}{2} \{ (X_1 + \dots + X_{n_0k}) Y_{n_0k+1} + \dots + (X_1 + \dots + X_{n_0k+q-1}) Y_{n_0k+q} \\ &\quad - (Y_1 + \dots + Y_{n_0k}) X_{n_0k+1} - \dots - (Y_1 + \dots + Y_{n_0k+q-1}) X_{n_0k+q} \} \end{aligned}$$

From Tchebychev's inequality, for every  $\varepsilon > 0$ ,

$$\mathbb{P} \left( \left| \frac{V_{n_0}^{(k)} - \mathbb{E}(V_{n_0}^{(k)})}{n_0k} \right| \geq \varepsilon \right) \leq \frac{\text{Var}(V_{n_0}^{(k)})}{\varepsilon^2 n_0^2 k^2}.$$

Now, using the same techniques than the ones developed in the proof of Lemma 3.1, it can be proved that

$$\text{Var}(V_{n_0}^{(k)}) = \mathcal{O}(n_0^2 k).$$

Assumption **(A)** implies (see the proof of Lemma 3.2) that

$$\lim_{k \rightarrow +\infty} \frac{\mathbb{E}(V_{n_0}^{(k)})}{n_0k} = 0.$$

Then, we deduce that

$$\lim_{k \rightarrow +\infty} \mathbb{P} \left( \left| \frac{V_{n_0}^{(k)}}{n_0k} \right| \geq \varepsilon \right) = 0.$$

Straightforward calculations lead to

$$\begin{aligned} & \left| \mathbb{E} \left( e^{i\theta \frac{T_p}{p}} \right) - \mathbb{E} \left( e^{i\theta \frac{T_{n_0k}}{n_0k}} \right) \right| \\ & \leq \left| \mathbb{E} \left( e^{i\theta \frac{n_0k}{n_0k+q} \frac{T_{n_0k}}{n_0k} + \frac{V_{n_0}^{(k)}}{n_0k+q}} \right) - \mathbb{E} \left( e^{i\theta \frac{T_{n_0k}}{n_0k}} \right) \right| \end{aligned}$$

$$\begin{aligned}
&\leq \left| \mathbb{E} \left( e^{i\theta \frac{n_0 k}{n_0^{k+q}} \frac{T_{n_0 k}}{n_0^k} + \frac{V_{n_0}^{(k)}}{n_0^{k+q}}} \right) - \mathbb{E} \left( e^{i\theta \frac{T_{n_0 k}}{n_0^k} + \frac{V_{n_0}^{(k)}}{n_0^{k+q}}} \right) \right| \\
&+ \left| \mathbb{E} \left( e^{i\theta \frac{T_{n_0 k}}{n_0^k} + \frac{V_{n_0}^{(k)}}{n_0^{k+q}}} \right) - \mathbb{E} \left( e^{i\theta \frac{T_{n_0 k}}{n_0^k}} \right) \right| \\
&\leq \mathbb{E} \left( \left| e^{i\theta \frac{q}{n_0^{k+q}} \frac{T_{n_0 k}}{n_0^k} - 1 \right| \right) + \mathbb{E} \left( \left| e^{i\theta \frac{V_{n_0}^{(k)}}{n_0^{k+q}}} - 1 \right| \right).
\end{aligned}$$

Now, we have (since  $|e^{i\theta} - 1| \leq 2 \wedge |\theta|$ ) for  $k$  large enough

$$\mathbb{E} \left( \left| e^{i\theta \frac{V_{n_0}^{(k)}}{n_0^{k+q}}} - 1 \right| \right) \leq \varepsilon + 2\mathbb{P} \left( \left| \frac{\theta V_{n_0}^{(k)}}{n_0^{k+q}} \right| \geq \varepsilon \right) \leq 3\varepsilon.$$

From (7), for  $k$  large enough,

$$\left| \mathbb{E} \left( e^{i\theta \frac{q}{n_0^{k+q}} \frac{T_{n_0 k}}{n_0^k}} \right) - 1 \right| \leq 3\varepsilon. \quad (8)$$

On a compact set of  $\mathbb{R}$  (for  $\theta$ ), for  $p$  large enough,

$$\begin{aligned}
\left| \mathbb{E} \left( e^{i\theta \frac{T_p}{p}} \right) - \frac{1}{\cosh\left(\frac{\theta \sqrt{a_{11} a_{22}}}{2}\right)} \right| &\leq \left| \mathbb{E} \left( e^{i\theta \frac{T_p}{p}} \right) - \mathbb{E} \left( e^{i\theta \frac{T_{n_0 k}}{n_0^k}} \right) \right| \\
&+ \left| \mathbb{E} \left( e^{i\theta \frac{T_{n_0 k}}{n_0^k}} \right) - \frac{1}{\cosh\left(\frac{\theta \sqrt{a_{11} a_{22}}}{2}\right)} \right| \\
&\leq 9\varepsilon,
\end{aligned}$$

so Theorem 3.2 is proved.

### 3.3 Concluding remarks

As stated in the introduction, this paper is a first attempt in extending the theory of dynamic random walks developed by the first author to non-commutative algebraic structures. More results on limit theorems on Heisenberg groups are expected. The law of the iterated logarithm is presently under investigation.

## References

- [1] BERTHUET, R. *Loi du logarithme itéré pour certaines intégrales stochastiques*. C. R. Acad. Sci. Paris Sr. (1979), 289, No. 16, 813 -815
- [2] BERTHUET, R. *Etude de processus généralisant l'aire de Lévy*. (French) [Study of a process generalizing the Levy stochastic area] Probab. Theory Related Fields (1986), 73, No. 3, 463 - 480.



- [3] CREPEL, P. AND RAUGI, A. *Théorème central limite sur les groupes nilpotents*. Ann. Inst. H. Poincaré. Statist. (1978) 14, 145-162.
- [4] CREPEL, P. AND ROYNETTE, B. *Une loi du logarithme itéré pour le groupe d'Heisenberg* Z. Wahrscheinlichkeitstheorie und verw. Geb. (1977), 14, 217-229.
- [5] DRMOTA, M. AND TICHY, R. *Sequences, discrepancies and applications*. Lecture Notes in Mathematics, Vol. 1651, Springer-Verlag, 1997.
- [6] GUILLOTIN, N. *Asymptotics of a dynamic random walk in a random scenery I. A law of large numbers*. Annales de l'Institut Henri Poincaré - Probabilités et Statistiques (2000), Vol. **36**, No 2, 127–151.
- [7] GUILLOTIN, N. *Asymptotics of a dynamic random walk in a random scenery II. A functional limit theorem*. Markov Processes and Related Fields (1999), Vol. **5**, No 2, 201–218.
- [8] GUILLOTIN-PLANTARD, N. *Dynamic  $\mathbb{Z}^d$ -random walks in a random scenery: a strong law of large numbers*. J. Theoret. Probab. (2001), Vol. **14**, No 1, 241–260.
- [9] GUILLOTIN-PLANTARD, N. AND SCHOTT, R. *Dynamic random walks: Theory and applications* . 200 pages, (book in press).
- [10] GUIVARC'H, Y. *Sur la loi des grands nombres et le rayon spectral d'une marche aléatoire*. Astérisque (1980), 74, 47-98.
- [11] GUIVARC'H, Y.; KEANE, M.; ROYNETTE, B. *Marches aléatoires sur les groupes de Lie*. Lecture Notes in Mathematics, Vol. 624, Springer-Verlag, 1977.
- [12] HEYER, H. *Probability measures on locally compact groups*. Springer Verlag, 1977.
- [13] KORÁNYI, A. *Geometric properties of Heisenberg-type groups*. Adv. Math. (1985), 56, 1, 28-38.
- [14] KORÁNYI, A. *Geometric aspects of analysis on the Heisenberg group*. In: *De Michele, L. Ricci, F. (ed.)*. Topics in modern harmonic analysis vol.1, Istituto Nazionale di Alta Matematica Francesco Severi, Roma (1983), 209-258.
- [15] NEUENSCHWANDER, D. *Probabilities on the Heisenberg group*. Lecture Notes in Mathematics, Vol. 1630, Springer-Verlag, Berlin-New York, 1996.
- [16] NEUENSCHWANDER, D. AND SCHOTT, R. *On the local and asymptotic behavior of Brownian motion on simply connected Lie groups* J. Theoret. Probab. (1995), 8, 4, 795-806.
- [17] SCHOTT, R. *Une loi du logarithme itéré pour certaines intégrales stochastiques*. (French) C. R. Acad. Sci. Paris Sr. I Math. (1981), 292, No. 4, 295 -298.