



**HAL**  
open science

# Schur-Weyl duality and the heat kernel measure on the unitary group.

Thierry Lévy

► **To cite this version:**

Thierry Lévy. Schur-Weyl duality and the heat kernel measure on the unitary group.. 2007. hal-00137982v1

**HAL Id: hal-00137982**

**<https://hal.science/hal-00137982v1>**

Preprint submitted on 23 Mar 2007 (v1), last revised 15 Jan 2008 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SCHUR-WEYL DUALITY AND THE HEAT KERNEL MEASURE ON THE UNITARY GROUP

THIERRY LÉVY

**ABSTRACT.** We establish a convergent power series expansion for the expectation of a product of traces of powers of a random unitary matrix under the heat kernel measure. These expectations turn out to be the generating series of certain paths in the Cayley graph of the symmetric group. We then compute the asymptotic distribution of a random unitary matrix under the heat kernel measure on the unitary group  $U(N)$  as  $N$  tends to infinity, and prove a result of asymptotic freeness for independent large unitary matrices, thus recovering results obtained previously by Xu [17] and Biane [1]. We give an interpretation of our main expansion in terms of random ramified coverings of a disk. Our approach is based on the Schur-Weyl duality and we extend some of our results to the orthogonal and symplectic cases.

## 1. INTRODUCTION

In this paper, we are concerned with the asymptotics of large random unitary matrices distributed according to the heat kernel measure. This problem has been studied about ten years ago by P. Biane [1] and F. Xu [17]. It shares some similarities with the case of unitary matrices distributed under the Haar measure, studied by B. Collins and P. Śniady [4, 5]. The origin of our interest in this problem is the hypothetical existence of a large  $N$  limit to the two-dimensional  $U(N)$  Yang-Mills theory. This limit has been investigated in different directions by physicists, in particular V. Kazakov and V. Kostov [8], D. Gross and W. Taylor [7] and D. Gross and A. Matytsin [6]. In [15], I. Singer has given the name of "Master field" to this limit, which still has to be constructed. A. Sengupta has given in [14] a very clear account of the relationship between Yang-Mills theory and large unitary matrices. We refer the interested reader to this paper and will not develop this motivation further. Sengupta's work also contains some results whose study was at the origin of this paper (see Proposition 2.2 and the discussion thereafter).

Our approach relies on the fact that Schur-Weyl duality determines a (non-bijective) correspondence between conjugation-invariant objects on the unitary group on one hand and on the symmetric group on the other hand. To be specific, let  $n, N \geq 1$  be integers. Let  $\rho_{n,N} : \mathfrak{S}_n \times U(N) \longrightarrow GL((\mathbb{C}^N)^{\otimes n})$  be the usual representation. The set  $\mathcal{P}_{n,N}$  of partitions of  $n$  with at most  $N$  parts indexes irreducible representations of both  $\mathfrak{S}_n$  and  $U(N)$ . If  $\lambda$  is such a partition, let  $\chi^\lambda$  (resp.  $\chi_\lambda$ ) denote the corresponding character on  $\mathfrak{S}_n$  (resp.  $U(N)$ ). Let  $Z$  be an element of the centre of  $\mathbb{C}[\mathfrak{S}_n]$ . Let  $D$  be a conjugation-invariant distribution on  $U(N)$ . Then the equalities

$$(1) \quad \forall \lambda \in \mathcal{P}_{n,N}, \quad \frac{\chi^\lambda(Z)}{\chi^\lambda(\text{id})} = \frac{\chi_\lambda(D)}{\chi_\lambda(I_N)},$$

where  $\chi_\lambda(D) = D\chi_\lambda$ , imply  $\rho_{n,N}(Z \otimes 1) = \rho_{n,N}(1 \otimes D)$ . The main observation, implicit in [7], is the following: the element  $Z = Nn + n(n-1) - 2 \sum_{1 \leq k < l \leq n} (kl) \in \mathbb{C}[\mathfrak{S}_n]$  and the distribution  $D$  on  $U(N)$  defined by  $D\varphi = \frac{1}{2} \Delta_{U(N)} \varphi(I_N)$ , where  $\Delta_{U(N)}$  is the Laplace operator, satisfy (1).

---

*Date:* March 2007.

Now  $Z$  is almost the generator of the most natural random walk on  $\mathfrak{S}_n$  and it follows from this discussion that this random walk is closely related to the Brownian motion on the unitary group.

By elaborating on this observation, we prove a convergent power series expansion for the expected value of products of traces of powers of unitary matrices under the heat kernel measure. We use this expansion to give the asymptotic distribution of these random matrices, thus recovering a result of P. Biane [1]. We also recover a result of F. Xu [17] on the asymptotic factorization of the expected values of products of traces. As another application, we use Speicher's criterion of freeness to prove the asymptotic freeness of independent unitary matrices under the heat kernel measure. Finally, we give an interpretation of our formula in terms of random branched coverings and prove rigorously a formula which has been described by Gross and Taylor in [7].

It is a pleasure to thank Philippe Biane for several enlightening conversations.

## 2. PROBABILISTIC ASPECTS OF SCHUR-WEYL DUALITY

In this first section, we establish formulae which relate the heat kernel measures on  $U(N)$ ,  $SU(N)$ ,  $SO(N)$  and  $Sp(N)$ , to natural random walks in the symmetric group and the Brauer monoid. At a general level, the fact that such formulae exist is a consequence of the Schur-Weyl duality, as we explain below. Nevertheless, our approach is elementary and does not rely logically on Schur-Weyl duality.

**2.1. Unitary group.** Let  $n$  and  $N$  be two positive integers. There is a natural action of each of the groups  $U(N)$  and  $\mathfrak{S}_n$  on the vector space  $(\mathbb{C}^N)^{\otimes n}$ , defined as follows: for all  $U \in U(N)$ ,  $\sigma \in \mathfrak{S}_n$  and  $x_1, \dots, x_n \in \mathbb{C}^N$ ,

$$(2) \quad U \cdot (x_1 \otimes \dots \otimes x_n) = Ux_1 \otimes \dots \otimes Ux_n \text{ and } \sigma \cdot (x_1 \otimes \dots \otimes x_n) = x_{\sigma^{-1}(1)} \otimes \dots \otimes x_{\sigma^{-1}(n)}.$$

It is a basic observation that these actions commute to each other. In particular, they determine an action  $\rho_{n,N}$  of  $\mathfrak{S}_n \times U(N)$  on  $(\mathbb{C}^N)^{\otimes n}$  by

$$\rho_{n,N}(\sigma, U)(x_1 \otimes \dots \otimes x_n) = Ux_{\sigma^{-1}(1)} \otimes \dots \otimes Ux_{\sigma^{-1}(n)}.$$

**Definition 2.1.** Let  $M_1, \dots, M_n$  be  $N \times N$  complex matrices. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . We denote by  $p_\sigma^{st}(M_1, \dots, M_n)$  the complex number

$$p_\sigma^{st}(M_1, \dots, M_n) = \text{Tr}_{(\mathbb{C}^N)^{\otimes n}}((M_1 \otimes \dots \otimes M_n) \circ \rho_{n,N}(\sigma, U)) = \prod_{\substack{c=(i_1 \dots i_r) \\ \text{cycle of } \sigma}} \text{Tr}(M_{i_1} \dots M_{i_r}).$$

We set  $p^{st}(M) = p_\sigma^{st}(M, \dots, M)$ .

The upper index  $st$  indicates that we use the standard trace rather than the normalized one in the definition. The letter  $p$  stand for "power sums", since  $p_\sigma^{st}(M)$ , as a symmetric function of the eigenvalues of  $M$ , is the product of power sums corresponding to the partition determined by  $\sigma$ . Observe that, by definition, the character of the representation  $\rho_{n,N}$  is the function  $\chi_{\rho_{n,N}}(\sigma, U) = p_\sigma^{st}(U)$ .

The core result of Schur-Weyl duality is that the two subalgebras of  $\text{End}((\mathbb{C}^N)^{\otimes n})$  generated respectively by the actions of  $U(N)$  and  $\mathfrak{S}_n$  are each other's commutant. Let us explain briefly why this makes a relation between the Brownian motion on  $U(N)$  and some element of the centre of the group algebra of  $\mathfrak{S}_n$  unavoidable.

Let  $\mathfrak{u}(N)$  denote the Lie algebra of  $U(N)$ . It consists exactly of the  $N \times N$  anti-Hermitian complex matrices. Let  $\mathcal{U}(\mathfrak{u}(N))$  denote the enveloping algebra of  $\mathfrak{u}(N)$ , which is canonically isomorphic to the algebra of left-invariant differential operators on  $U(N)$ . Let also  $\mathbb{C}[\mathfrak{S}_n]$  denote

the group algebra of  $\mathfrak{S}_n$ . The representation  $\rho_{n,N}$  determines a homomorphism of associative algebras  $\mathbb{C}[\mathfrak{S}_n] \otimes \mathcal{U}(\mathfrak{u}(N)) \longrightarrow \text{End}((\mathbb{C}^N)^{\otimes n})$ . The centre  $\mathcal{Z}(\mathfrak{u}(N))$  of  $\mathcal{U}(\mathfrak{u}(N))$  consists in the space of bi-invariant differential operators on  $U(N)$ . Since  $\rho_{n,N}(1 \otimes \mathcal{Z}(\mathfrak{u}(N)))$  commutes to  $\rho_{n,N}(1, U)$  for every  $U \in U(N)$ , the Schur-Weyl duality asserts in particular that

$$\rho_{n,N}(1 \otimes \mathcal{Z}(\mathfrak{u}(N))) \subset \rho_{n,N}(\mathbb{C}[S_n] \otimes 1).$$

We are primarily interested in the Laplace operator, which is defined as follows. The  $\mathbb{R}$ -bilinear form  $\langle X, Y \rangle = \text{Tr}(X^*Y) = -\text{Tr}(XY)$  is a scalar product on  $\mathfrak{u}(N)$ . Let  $(X_1, \dots, X_{N^2})$  be an orthonormal basis of  $\mathfrak{u}(N)$ . Identifying the elements of  $\mathfrak{u}(N)$  with left-invariant vector fields on  $U(N)$ , thus with first-order differential operators on  $U(N)$ , the Laplace operator  $\Delta_{U(N)}$  is the differential operator  $\sum_{i=1}^{N^2} X_i^2$ . It corresponds to the Casimir element  $\sum_{i=1}^{N^2} X_i \otimes X_i$  of the enveloping algebra of  $\mathfrak{u}(N)$ . This element is central and does not depend on the choice of the orthonormal basis. Hence,  $\Delta_{U(N)}$  is well defined and bi-invariant. The discussion above shows that, in the representation  $\rho_{n,N}$ , the Laplace operator of  $U(N)$  can be expressed as an element of  $\mathbb{C}[S_n]$ . This is exactly what the main formula of this section does, in an explicit way.

The element of  $\mathbb{C}[\mathfrak{S}_n]$  which plays the main role is the following. Denote by  $T_n$  the subset of  $\mathfrak{S}_n$  consisting of all transpositions and set  $\Delta_{\mathfrak{S}_n} = -\frac{n(n-1)}{2} + \sum_{\tau \in T_n} \tau$ . The formula for the unitary group is the following.

**Proposition 2.2.** *For all integers,  $n, N \geq 1$ , one has*

$$(3) \quad \rho_{n,N} \left( \Delta_{\mathfrak{S}_n} \otimes 1 + 1 \otimes \frac{1}{2} \Delta_{U(N)} \right) = -\frac{Nn + n(n-1)}{2}.$$

Before we prove this formula, let us derive some of its consequences.

**Proposition 2.3.** *For each  $\sigma \in \mathfrak{S}_n$ , the function  $p_\sigma^{st} : U(N) \longrightarrow \mathbb{C}$  satisfies the following relation:*

$$(4) \quad \frac{1}{2} \Delta_{U(N)} p_\sigma^{st} = -\frac{Nn}{2} p_\sigma^{st} - \sum_{\tau \in T_n} p_{\sigma\tau}^{st}.$$

More generally, let  $M_1, \dots, M_n$  be arbitrary  $N \times N$  matrices. Then, regarding  $p_\sigma^{st}(M_1 U, \dots, M_n U)$  as a function of  $U \in U(N)$ ,

$$(5) \quad \frac{1}{2} \Delta_{U(N)} p_\sigma^{st}(M_1 U, \dots, M_n U) = -\frac{Nn}{2} p_\sigma^{st}(M_1 U, \dots, M_n U) - \sum_{\tau \in T_n} p_{\sigma\tau}^{st}(M_1 U, \dots, M_n U).$$

**Proof** – Recall that  $p_\sigma^{st}(M_1 U, \dots, M_n U) = \text{Tr}((M_1 \otimes \dots \otimes M_n) \circ \rho_{n,N}(\sigma, U))$ . Let us use the shorthand notation  $M = M_1 \otimes \dots \otimes M_n$ . We have

$$\begin{aligned} \frac{1}{2} \Delta_{U(N)} p_\sigma^{st}(M_1 U, \dots, M_n U) &= \text{Tr}(M \circ \rho_{n,N}(\sigma, U) \circ \rho_{n,N}(1 \otimes \frac{1}{2} \Delta_{U(N)})) \\ &= -\frac{Nn + n(n-1)}{2} p_\sigma^{st}(M_1 U, \dots, M_n U) - \text{Tr}(M \circ \rho_{n,N}(\sigma, U) \circ \rho_{n,N}(\Delta_{S_n} \otimes 1)). \end{aligned}$$

The result follows immediately from the definition of  $\Delta_{\mathfrak{S}_n}$ .  $\square$

The function  $p_\sigma^{st}$  depends only on the cycle structure of  $\sigma$ . In concrete terms, if the lengths of the cycles of  $\sigma$  are  $m_1, \dots, m_r$ , then  $p_\sigma^{st}(U) = \text{Tr}(U^{m_1}) \dots \text{Tr}(U^{m_r})$ . This redundant labelling is however nicely adapted to our problem, as equation (4) shows. Let us spell out the right hand side of this equality. The permutation  $\sigma$  being fixed, the cycle structure of  $\sigma\tau$  depends on the two points exchanged by the transposition  $\tau$ . If they belong to the same cycle of  $\sigma$ , then this cycle is split into two cycles. A cycle of length  $m$  can be split into a cycle of length  $s$  and a

cycle of length  $m - s$  by  $m$  distinct transpositions, unless  $m = 2s$ , in which case only  $\frac{m}{2}$  of these transpositions are distinct. If on the contrary the points exchanged by  $\tau$  belong to two distinct cycles of  $\sigma$ , these two cycles are merged into a single cycle. Two cycles of lengths  $m$  and  $m'$  can be merged by  $mm'$  distinct permutations. Altogether, we find the following equation, which was already present in papers of Xu [17] and Sengupta [14].

$$\begin{aligned} \Delta_{U(N)}(\mathrm{Tr}(U^{m_1}) \dots \mathrm{Tr}(U^{m_r})) &= -Nn \mathrm{Tr}(U^{m_1}) \dots \mathrm{Tr}(U^{m_r}) \\ &+ \sum_{i=1}^r m_i \mathrm{Tr}(U^{m_1}) \dots \widehat{\mathrm{Tr}(U^{m_i})} \dots \mathrm{Tr}(U^{m_r}) \sum_{s=1}^{m_i-1} \mathrm{Tr}(U^s) \mathrm{Tr}(U^{m_i-s}) \\ &+ \sum_{\substack{i,j=1 \\ i \neq j}}^r m_i m_j \mathrm{Tr}(U^{m_1}) \dots \widehat{\mathrm{Tr}(U^{m_i})} \dots \widehat{\mathrm{Tr}(U^{m_j})} \dots \mathrm{Tr}(U^{m_r}) \mathrm{Tr}(U^{m_i+m_j}). \end{aligned}$$

The fact that  $\rho_{n,N}(1 \otimes \Delta_{U(N)})$  can be expressed in terms of permutations is, as explained before, a consequence of Schur-Weyl duality. What is not is the remarkable fact that the element of  $\mathbb{C}[\mathfrak{S}_n]$  which appears in (3) has coefficients of the same sign on the elements which are not the identity. Hence, up to an additive constant, it can be interpreted as the generator of a Markov chain on  $\mathfrak{S}_n$ . This leads us to the following simple probabilistic interpretation of (3).

Let us introduce the standard random walk on the Cayley graph of the symmetric group generated by the set of transpositions. It is the continuous-time Markov chain on  $\mathfrak{S}_n$  with generator  $\Delta_{\mathfrak{S}_n}$ , that is, the chain which jumps at rate  $\binom{n}{2}$  from its current position  $\sigma$  to  $\sigma\tau$ , where  $\tau$  is chosen uniformly at random among the  $\binom{n}{2}$  transpositions.

If  $\sigma$  is a permutation, we denote by  $\ell(\sigma)$  the number of cycles of  $\sigma$ . For example,  $\tau$  is a transposition if and only if  $\ell(\tau) = n - 1$ .

**Proposition 2.4.** *Let  $N, n \geq 1$  be integers. Let  $(B_t)_{t \geq 0}$  be a Brownian motion on  $U(N)$  starting at the identity and corresponding to the scalar product  $(X, Y) \mapsto -\mathrm{Tr}(XY)$  on  $\mathfrak{u}(N)$ . Let  $(\pi_t)_{t \geq 0}$  be a standard random walk on the Cayley graph of the symmetric group  $\mathfrak{S}_n$ , independent of  $(B_t)_{t \geq 0}$ . Then*

$$(6) \quad \mathbb{E} [p_{\pi_t}^{st}(B_t)] = e^{-\frac{Nn+n(n-1)}{2}t} \mathbb{E} [N^{\ell(\pi_0)}].$$

More generally, let  $M_1, \dots, M_n$  be arbitrary  $N \times N$  complex matrices. Then

$$(7) \quad \mathbb{E} [p_{\pi_t}^{st}(M_1 B_t, \dots, M_n B_t)] = e^{-\frac{Nn+n(n-1)}{2}t} \mathbb{E} [p_{\pi_0}^{st}(M_1, \dots, M_n)].$$

**Proof** – The process  $(\pi, B)$  is a Markov process on  $\mathfrak{S}_n \times U(N)$  with generator  $\Delta_{\mathfrak{S}_n} \otimes 1 + 1 \otimes \frac{1}{2}\Delta_{U(N)}$ . By Proposition 2.3,

$$\begin{aligned} \frac{d}{dt} \mathbb{E} [p_{\pi_t}^{st}(M_1 B_t, \dots, M_n B_t)] &= \mathbb{E} \left[ \left( \Delta_{\mathfrak{S}_n} \otimes 1 + 1 \otimes \frac{1}{2}\Delta_{U(N)} \right) p_{\pi_t}^{st}(M_1 B_t, \dots, M_n B_t) \right] \\ &= -\frac{Nn + n(n-1)}{2} \mathbb{E} [p_{\pi_t}^{st}(M_1 B_t, \dots, M_n B_t)]. \end{aligned}$$

The result follows immediately from the fact that  $B_0 = I_N$  a.s.  $\square$

Let us turn to the proof of Proposition 2.2.

**Proof of Proposition 2.2** – The action of  $\mathfrak{u}(N)$  on  $(\mathbb{C}^N)^{\otimes n}$  extends by complexification to  $\mathfrak{gl}(N, \mathbb{C}) = \mathfrak{u}(N) \oplus i\mathfrak{u}(N)$ . Let  $(X_1, \dots, X_{N^2})$  be a real basis of  $\mathfrak{u}(N)$ . It is also a complex basis

of  $\mathfrak{gl}(N, \mathbb{C})$ . Define a  $N \times N$  matrix  $g$  by  $g_{ij} = -\text{Tr}(X_i X_j)$ . Since  $-\text{Tr}(\cdot)$  is non-degenerate on  $\mathfrak{gl}_N(\mathbb{C})$ , the matrix  $g$  has an inverse  $g^{-1}$ , the entries of which we denote by  $g^{ij}$ . Then it is easy to check that the element  $\sum_{i,j=1}^{N^2} g^{ij} X_i \otimes X_j$  of the enveloping algebra is independent of the choice of the basis. Of course, by choosing our original basis of  $\mathfrak{u}(N)$  orthonormal, we find that this element is simply  $\Delta_{U(N)}$ .

In order to compute  $\rho_{n,N}(1 \otimes \Delta_{U(N)})$ , we prefer to use another complex basis of  $\mathfrak{gl}(N, \mathbb{C}) = \mathbb{M}_N(\mathbb{C})$ , namely the canonical basis  $(E_{ij})_{i,j \in \{1, \dots, N\}}$ . For this basis,  $g_{ij,kl} = -\delta_{jk} \delta_{il}$  and  $g = g^{-1}$ .

First, notice that  $\rho_{n,N}(1 \otimes E_{ij})(x_1 \otimes \dots \otimes x_n) = \sum_{k=1}^n x_1 \otimes \dots \otimes E_{ij}(x_k) \otimes \dots \otimes x_n$ . Hence,

$$\begin{aligned} \rho_{n,N} \left( 1 \otimes \sum_{i,j=1}^N E_{ij} \otimes E_{ji} \right) &= 2 \sum_{i,j=1}^N \sum_{1 \leq k < l \leq n} \text{Id}^{\otimes k-1} \otimes E_{ij} \otimes \text{Id}^{\otimes l-k-1} \otimes E_{ji} \otimes \text{Id}^{\otimes n-l-1} + \\ &\quad + \sum_{i,j=1}^N \sum_{k=1}^n \text{Id}^{\otimes k-1} \otimes E_{ii} \otimes \text{Id}^{\otimes n-k-1}. \end{aligned}$$

The last term is simply  $Nn$  times the identity. For the first part of the right hand side, observe that  $\sum_{i,j=1}^N E_{ij} \otimes E_{ji} \in \text{End}((\mathbb{C}^N)^{\otimes 2})$  is the transposition operator  $x \otimes y \mapsto y \otimes x$ , that is, the operator  $\rho_{2,N}((12), I_N)$ . Finally, we have found that

$$-\rho_{n,N}(1 \otimes \Delta_{U(N)}) = Nn \text{Id} + \sum_{1 \leq k \neq l \leq n} \rho_{n,N}((kl), I_N).$$

The result follows immediately.  $\square$

The results of this section still hold, after a minor modification, when  $U(N)$  is replaced by  $SU(N)$ . Indeed, the orthogonal complement of  $\mathfrak{su}(N)$  in  $\mathfrak{u}(N)$  is the line generated by  $\frac{i}{\sqrt{N}} I_N$ . Since  $\rho_{n,N}(1 \otimes (I_N \otimes I_N)) = n^2 \text{Id}$ , the Casimir operator of  $\mathfrak{su}(N)$  satisfies the relation

$$\rho_{n,N}(1 \otimes \Delta_{SU(N)}) = \rho_{n,N}(1 \otimes \Delta_{U(N)}) + \frac{n^2}{N} \text{Id}.$$

This modifies only the exponential factors in (6) and (7).

We will explore further consequences of Proposition 2.4 in the rest of the paper. For the moment, we derive similar results for the orthogonal and symplectic group.

**2.2. Orthogonal group.** Let us consider the action of  $SO(N)$  on  $(\mathbb{C}^N)^{\otimes n}$  defined by analogy with (2). The action of  $\mathfrak{S}_n$  still commutes to that of  $SO(N)$ , but, unless  $n = 1$ , the subalgebra of  $\text{End}((\mathbb{C}^N)^{\otimes n})$  generated by the image of  $\mathbb{C}[\mathfrak{S}_n]$  is strictly smaller than the commutant of the image of  $SO(N)$ . Let us review briefly the operators which are classically used to describe this commutant. We denote by  $\{e_1, \dots, e_N\}$  the canonical basis of  $\mathbb{C}^N$ .

**Definition 2.5.** Let  $\beta$  be a partition of  $\{1, \dots, 2n\}$  into pairs. Define  $\rho_{n,N}(\beta) \in \text{End}((\mathbb{C}^N)^{\otimes n})$  by setting, for all  $i_1, \dots, i_n \in \{1, \dots, N\}$ ,

$$\rho_{n,N}(\beta)(e_{i_1} \otimes \dots \otimes e_{i_n}) = \sum_{i_{n+1}, \dots, i_{2n} \in \{1, \dots, n\}} \prod_{\{k,l\} \in \beta} \delta_{i_k i_l} e_{i_{n+1}} \otimes \dots \otimes e_{i_{2n}}.$$

Observe that the partition  $\{\{1, n+1\}, \dots, \{n, 2n\}\}$  is sent to the identity operator by  $\rho_{n,N}$ . Let  $B_n$  denote the set of partitions of  $\{1, \dots, 2n\}$  into pairs. The composition of the operators  $\rho_{n,N}(\beta)$  corresponds to a monoid structure on  $B_n$  which is easiest to understand on a picture. An element of  $B_n$  is represented in a box with  $n$  dots on its top edge and  $n$  dots on its bottom edge. The dots on the top are labelled from 1 to  $n$ , from the left to the right. The dots on the

bottom are labelled from  $n + 1$  to  $2n$ , from the left to the right too. A pairing is then simply represented by  $n$  chords which join the appropriate dots. Multiplication of pairings is done in the intuitive topological way by superposing boxes.




FIGURE 1. Multiplication of two diagrams in the Brauer monoid.

The monoid  $B_n$  is sometimes called the Brauer monoid. Its elements are then called Brauer diagrams. The group  $\mathfrak{S}_n$  is naturally a submonoid of  $B_n$ , by the identification of a permutation  $\sigma$  with the pairing  $\{\{1, \sigma(1) + n\}, \dots, \{n, \sigma(n) + n\}\}$ . It is not difficult to check that  $\mathfrak{S}_n \subset B_n$  is exactly the subset of invertible elements. Indeed, for  $\beta \in B_n$ , let  $T(\beta)$  be the set of pairs  $\{k, l\} \in \beta$  such that  $1 \leq k, l \leq n$ . In words,  $T(\beta)$  is the set of chords in the diagram of  $\beta$  which join two dots on the top edge of the box. It is clear that  $T(\beta_1\beta_2) \subset T(\beta_1)$  for all  $\beta_1, \beta_2 \in B_n$ . Hence,  $T(\beta)$  must be empty for  $\beta$  to be invertible. More generally, it is not difficult to check that, given  $\beta$  and  $\beta'$  in  $B_n$ , there exists  $\beta'' \in B_n$  such that  $\beta\beta'' = \beta'$  if and only if  $T(\beta) \subset T(\beta')$ .

The identification of  $\mathfrak{S}_n$  with a subset of  $B_n$  is compatible with our previous definition of  $\rho_{n,N}$  in the sense that  $\rho_{n,N}(\sigma)$  is the same if we consider  $\sigma$  as a permutation or as a Brauer diagram.

The correct statement of Schur-Weyl duality is that the subalgebras of  $\text{End}((\mathbb{C}^N)^{\otimes n})$  generated by  $SO(N)$  and  $B_n$  are each other's commutant. Let  $\rho_{n,N}$  denote the morphism of monoids

$$\rho_{n,N} : B_n \times SO(N) \longrightarrow GL((\mathbb{C}^N)^{\otimes n}).$$

Just as in the unitary case, this action determines a morphism of associative algebras  $\rho_{n,N} : \mathbb{C}[B_n] \otimes \mathcal{U}(\mathfrak{so}(N)) \longrightarrow \text{End}((\mathbb{C}^N)^{\otimes n})$ .

By analogy to the unitary case, let us define "power sums" functions associated to Brauer diagrams. Given  $\beta \in B_n$  and  $M_1, \dots, M_n \in \mathbb{M}_N(\mathbb{C})$ , set


$$p_\beta^{st}(M_1, \dots, M_n) = \text{Tr}((M_1 \otimes \dots \otimes M_n) \circ \rho_{n,N}(\beta)).$$

In particular, the character of  $\rho_{n,N}$  is given by  $\chi_{\rho_{n,N}}(\beta, R) = p_\beta^{st}(R)$ .

The number  $p_\beta^{st}(M_1, \dots, M_n)$  is a product of traces of words in the matrices  $M_1, {}^tM_1, \dots, M_n, {}^tM_n$ . Let us describe in more detail how to compute  $p_\beta^{st}(I_N)$ . Let  $\beta$  be a Brauer diagram. Consider the graph with vertices  $\{1, \dots, n\}$  and unoriented edges  $\{k, l\}$ , where  $k$  and  $l$  are such that there exist  $k' \in \{k, k + n\}$  and  $l' \in \{l, l + n\}$  with  $\{k', l'\} \in \beta$ . This is the graph obtained by identifying the top edge with the bottom edge in the graphical representation of  $\beta$ . Then each vertex has degree 2 in this graph. Hence, it is a union of disjoint unoriented cycles. If  $\beta$  belongs to  $\mathfrak{S}_n \subset B_n$ , this cycle structure is of course that of  $\beta$  as a permutation, apart from the orientation which is lost. In general, let  $\ell(\beta)$  denote the number of cycles in this graph. Then  $p_\beta^{st}(I_N) = N^{\ell(\beta)}$ .

Let us define an element of  $\mathbb{C}[B_n]$  as follows. Given  $k$  and  $l$  two integers such that  $1 \leq k < l \leq n$ , we define the element  $\langle kl \rangle$  of  $B_n$  as the following pairing:

$$\langle kl \rangle = \{\{k, l\}, \{n + k, n + l\}\} \cup \bigcup_{i \in \{1, \dots, n\} - \{k, l\}} \{\{i, n + i\}\}.$$

FIGURE 2. The elements  $\langle 24 \rangle$  and  $(24)$  of  $B_6$ .

Let  $C_n$  be the subset of  $B_n$  consisting of all the element of the form  $\langle kl \rangle$ . We now define  $\Delta_{B_n} = -\frac{n(n-1)}{2} + \sum_{\alpha \in C_n} \alpha$ . Thanks to the inclusion  $\mathfrak{S}_n \subset B_n$ , we still see  $\Delta_{\mathfrak{S}_n}$  as an element of  $\mathbb{C}[B_n]$ . The formula for the orthogonal group is the following.

**Proposition 2.6.** *For all integers,  $n, N \geq 1$ , one has*

$$(8) \quad \rho_{n,N}(\Delta_{S_n} \otimes 1 + 1 \otimes \Delta_{SO(N)}) = -\frac{(N-1)n}{2} + \rho_{n,N}(\Delta_{B_n} \otimes 1).$$

**Proof** – The computation is very similar to that of the unitary case. Endow  $\mathfrak{so}(N)$  with the scalar product  $\langle X, Y \rangle = -\text{Tr}(XY)$ . The basis  $(A_{ij})_{1 \leq i < j \leq N}$ , with  $A_{ij} = E_{ij} - E_{ji}$ , is orthogonal and  $\langle A_{ij}, A_{ij} \rangle = 2$  for all  $i < j$ . Hence,  $\Delta_{SO(N)} = \frac{1}{2} \sum_{1 \leq i < j \leq N} A_{ij} \otimes A_{ij}$ . We have

$$\begin{aligned} \rho_{n,N}(1 \otimes \Delta_{SO(N)}) &= \sum_{1 \leq k < l \leq n} \sum_{1 \leq i < j \leq N} \text{Id}^{\otimes k-1} \otimes A_{ij} \otimes \text{Id}^{\otimes l-k-1} \otimes A_{ij} \otimes \text{Id}^{\otimes n-l} + \\ &\quad + \frac{1}{2} \sum_{k=1}^n \sum_{1 \leq i < j \leq N} \text{Id}^{\otimes k-1} \otimes A_{ij}^2 \otimes \text{Id}^{\otimes n-k-1} \\ &= \sum_{1 \leq k < l \leq n} \sum_{i,j=1}^N \text{Id}^{\otimes k-1} \otimes E_{ij} \otimes \text{Id}^{\otimes l-k-1} \otimes E_{ij} \otimes \text{Id}^{\otimes n-l} \\ &\quad - \sum_{1 \leq k < l \leq n} \sum_{i,j=1}^N \text{Id}^{\otimes k-1} \otimes E_{ij} \otimes \text{Id}^{\otimes l-k-1} \otimes E_{ji} \otimes \text{Id}^{\otimes n-l} - \frac{(N-1)n}{2} \text{Id} \\ &= \sum_{1 \leq k < l \leq n} \rho_{n,N}(\langle kl \rangle - (kl) \otimes 1) - \frac{(N-1)n}{2} \text{Id}. \end{aligned}$$

The result follows immediately.  $\square$

**Proposition 2.7.** *For all  $\beta \in B_n$ , the following relation holds:*

$$(9) \quad \Delta_{SO(N)} p_{\beta}^{st} = -\frac{(N-1)n}{2} p_{\beta}^{st} - \sum_{\tau \in T_n} p_{\beta\tau}^{st} + \sum_{\alpha \in C_n} p_{\beta\alpha}^{st}.$$

More generally, let  $M_1, \dots, M_n$  be arbitrary  $N \times N$  matrices. Then, regarding  $p_{\beta}^{st}(M_1 R, \dots, M_n R)$  as a function of  $R \in SO(N)$ ,

$$(10) \quad \begin{aligned} \Delta_{SO(N)} p_{\beta}^{st}(M_1 R, \dots, M_n R) &= -\frac{(N-1)n}{2} p_{\beta}^{st}(M_1 R, \dots, M_n R) - \sum_{\tau \in T_n} p_{\beta\tau}^{st}(M_1 R, \dots, M_n R) + \\ &\quad + \sum_{\alpha \in C_n} p_{\beta\alpha}^{st}(M_1 R, \dots, M_n R). \end{aligned}$$



It seems more difficult to find a probabilistic interpretation of (8) than in the unitary case, because the element of  $\mathbb{C}[B_n]$  which appears does not have coefficients of the same sign on all elements not equal to 1.

**2.3. Symplectic group.** Nothing really new is needed to treat the case of the symplectic group. Let us describe briefly the results.

Let  $J \in \mathbb{M}_{2N}(\mathbb{C})$  denote the matrix  $\begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}$ . The symplectic group is defined by  $Sp(N) = \{S \in U(2N) : {}^tSJS = J\}$ . It acts naturally on  $((\mathbb{C}^{2N})^{\otimes n})$ . The action of the Brauer monoid needs to be slightly modified to fit the symplectic case. If  $\beta$  belongs to  $B_n$ , we define the operator  $\rho_{n,2N}(\beta)$  by setting, for all  $i_1, \dots, i_n \in \{1, \dots, 2N\}$ ,

$$\rho_{n,2N}(\beta)(e_{i_1} \otimes \dots \otimes e_{i_n}) = \sum_{i_{n+1}, \dots, i_{2n} \in \{1, \dots, n\}} \prod_{\{k,l\} \in \beta} J_{i_k i_l} e_{i_{n+1}} \otimes \dots \otimes e_{i_{2n}}.$$

Then we have an action  $\rho_{n,2N} : B_n \times Sp(N) \longrightarrow \text{End}((\mathbb{C}^{2N})^{\otimes n})$  and the images of  $B_n$  and  $Sp(N)$  generate two algebras which are each other's commutant.

The Lie algebra  $\mathfrak{sp}(N)$  is endowed with the scalar product  $\langle X, Y \rangle = -\text{Tr}(XY)$  and we denote by  $\Delta_{Sp(N)}$  the corresponding Laplace operator. The main formula is the following.

**Proposition 2.8.** *For all integers,  $n, N \geq 1$ , one has*

$$(11) \quad \rho_{n,2N}(\Delta_{S_n} \otimes 1 + 1 \otimes 2\Delta_{Sp(N)}) = -(2N+1)n + \rho_{n,2N}(\Delta_{B_n} \otimes 1).$$

**Proof** – Like in the unitary case, it is more convenient to use complexification. The Lie algebra  $\mathfrak{sp}(N, \mathbb{C}) = \mathfrak{sp}(N) \oplus i\mathfrak{sp}(N)$  is the Lie subalgebra of  $\mathfrak{gl}(2N, \mathbb{C})$  defined by the relation  ${}^tXJ = -JX$ . It consists of the matrices  $\begin{pmatrix} A & B \\ C & -{}^tA \end{pmatrix}$ , where  $A$  is an arbitrary  $N \times N$  matrix and  $B, C$  are two symmetric  $N \times N$  matrices. We use the following basis of  $\mathfrak{sp}(N, \mathbb{C})$ :

$$\begin{aligned} A_{ij} &= E_{ij} - E_{j+N, i+N} & 1 \leq i, j \leq N \\ B_{ij} &= E_{i, j+N} + E_{j, i+N} & 1 \leq i < j \leq N \\ C_{ij} &= E_{i+N, j} + E_{j+N, i} & 1 \leq i < j \leq N \\ D_i &= E_{i, i+N} & 1 \leq i \leq N \\ D_{i+N} &= E_{i+N, i} & 1 \leq i \leq N. \end{aligned}$$

The bilinear form  $\langle \cdot, \cdot \rangle$  takes the following values on this basis:

$$\begin{aligned} \langle A_{ij}, A_{ji} \rangle &= -2 & 1 \leq i, j \leq N \\ \langle B_{ij}, C_{ij} \rangle &= -2 & 1 \leq i < j \leq N \\ \langle D_i, D_{i+N} \rangle &= -1 & 1 \leq i \leq N. \end{aligned}$$

The other values are zero. It follows that the Casimir element of  $\mathfrak{sp}(N, \mathbb{C})$  is equal to

$$\Delta_{\mathfrak{sp}(N, \mathbb{C})} = -\frac{1}{2} \sum_{1 \leq i, j \leq N} A_{ij} \otimes A_{ji} - \frac{1}{2} \sum_{1 \leq i < j \leq N} (B_{ij} \otimes C_{ij} + C_{ij} \otimes B_{ij}) - \sum_{1 \leq i \leq N} (D_i \otimes D_{i+N} + D_{i+N} \otimes D_i).$$

The formula follows now by a direct computation. In order to recognize operators of the form  $\langle kl \rangle$  and  $(kl)$ , observe that, when  $n = 2$  for example,

$$\rho_{2,2N}((12), I_N) = \sum_{i,j=1}^{2N} E_{ij} \otimes E_{ji},$$

$$\rho_{2,2N}(\langle 12 \rangle, I_N) = \sum_{i,j=1}^N E_{ij} \otimes E_{i+N,j+N} + E_{i+N,j+N} \otimes E_{ij} - E_{i,j+N} \otimes E_{i+N,j} - E_{i+N,j} \otimes E_{i,j+N}.$$

□

**Proposition 2.9.** *For all  $\beta \in B_n$ , the following relation holds:*

$$(12) \quad 2\Delta_{Sp(N)} p_\beta^{st} = -(2N+1)n p_\beta^{st} - \sum_{\tau \in T_n} p_{\beta\tau}^{st} + \sum_{\alpha \in C_n} p_{\beta\alpha}^{st}.$$

More generally, let  $M_1, \dots, M_n$  be arbitrary  $2N \times 2N$  matrices. Then, regarding  $p_\beta^{st}(M_1 S, \dots, M_n S)$  as a function of  $R \in Sp(N)$ ,

$$(13) \quad \begin{aligned} 2\Delta_{Sp(N)} p_\beta^{st}(M_1 S, \dots, M_n S) &= -(2N+1)n p_\beta^{st}(M_1 S, \dots, M_n S) - \sum_{\tau \in T_n} p_{\beta\tau}^{st}(M_1 S, \dots, M_n S) + \\ &+ \sum_{\alpha \in C_n} p_{\beta\alpha}^{st}(M_1 S, \dots, M_n S). \end{aligned}$$

### 3. POWER SERIES EXPANSION

Let us denote by  $\text{Tr}_N = \frac{1}{N} \text{Tr}$  the normalized trace on  $\mathbb{M}_N(\mathbb{C})$ . Let  $M_1, \dots, M_n$  be  $N \times N$  matrices. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . We denote by  $p_\sigma(M_1, \dots, M_n)$  the number

$$p_\sigma(M_1, \dots, M_n) = \prod_{\substack{c=(i_1 \dots i_r) \\ \text{cycle of } \sigma}} \text{Tr}_N(M_{i_1} \dots M_{i_r}).$$

If  $\ell(\sigma)$  denotes the number of cycles of  $\sigma$ , then  $p_\sigma = N^{-\ell(\sigma)} p_\sigma^{st}$ .

In this section, we exploit the result of Proposition 2.4 and derive a convergent power series expansion of  $\mathbb{E} \left[ p_\sigma(B_{\frac{t}{N}}) \right]$  when  $B$  is a Brownian motion on  $U(N)$ . This expansion involves combinatorial coefficients, which count paths in the Cayley graph of  $\mathfrak{S}_n$ . We start by discussing these paths and introducing some notation.

**3.1. The Cayley graph of the symmetric group.** Fix  $n \geq 1$ . The Cayley graph of  $\mathfrak{S}_n$  generated by  $T_n$  can be described as follows. The vertices of this graph are the elements of  $\mathfrak{S}_n$  and two permutations  $\sigma_1$  and  $\sigma_2$  are joined by an edge if and only if  $\sigma_1 \sigma_2^{-1}$  is a transposition. A finite sequence  $(\sigma_0, \dots, \sigma_k)$  of permutations such that  $\sigma_i$  is joined to  $\sigma_{i+1}$  by an edge for each  $i \in \{0, \dots, k-1\}$  is called a path of length  $k$ . The distance between two permutations is the smallest length of a path which joins them. This distance can be computed explicitly as follows.

Let us introduce the notation  $|\sigma| = n - \ell(\sigma)$ . We have  $|\sigma| \in \{0, \dots, n-1\}$  and  $|\sigma| = 0$  (resp. 1, resp.  $n-1$ ) if and only if  $\sigma$  is the identity (resp. a transposition, resp. a  $n$ -cycle). Other values of  $|\sigma|$  do not characterize uniquely the conjugacy class of  $\sigma$ . It is well-known and easy to check that  $|\sigma|$  is the minimal number of transpositions required to write  $\sigma$ . In other words, the graph distance between two permutations  $\sigma_1$  and  $\sigma_2$  in the Cayley graph is given by  $|\sigma_1^{-1} \sigma_2|$ .

It turns out that the paths which play the most important role in our problem are those which tend to get closer to the identity. Let  $\gamma = (\sigma_0, \dots, \sigma_k)$  be a path. We call defect of  $\gamma$  and denote by  $d(\gamma)$  the number of steps which increase the distance to the identity. In symbols,

$$d(\gamma) = \#\{i \in \{0, \dots, k-1\} : |\sigma_{i+1}| = |\sigma_i| + 1\} = \#\{i \in \{0, \dots, k-1\} : \ell(\sigma_{i+1}) = \ell(\sigma_i) - 1\}.$$

We will count explicitly the number of paths with defect zero, given length and given origin in the next section. For the moment, we need only the following simple lemma, whose proof by induction on  $k$  is straightforward.

**Lemma 3.1.** *Let  $\gamma = (\sigma_0, \dots, \sigma_k)$  be a path. Then  $2d(\gamma) = k - (\ell(\sigma_k) - \ell(\sigma_0))$ .*

For  $\sigma, \sigma'$  in  $\mathfrak{S}_n$  and  $k \geq 0$ , let us denote by  $\Pi_k(\sigma \rightarrow \sigma')$  the set of paths of length  $k$  which start at  $\sigma$  and finish at  $\sigma'$ . Let us also denote by  $\Pi_k(\sigma)$  the set of all paths of length  $k$  starting at  $\sigma$  and by  $\Pi(\sigma \rightarrow \sigma')$  the set of all paths from  $\sigma$  to  $\sigma'$ . Notice that the cardinality of  $\Pi_k(\sigma)$  is equal to  $\binom{n}{2}^k \leq n^{2k}$ . Let us finally define the coefficients which appear in the expansion.

**Definition 3.2.** *Consider  $\sigma \in \mathfrak{S}_n$  and two integers  $k, d \geq 0$ . We set*

$$S(\sigma, k, d) = \#\{\gamma \in \Pi_k(\sigma) : d(\gamma) = d\}.$$

*In words,  $S(\sigma, k, d)$  is the number of paths in the Cayley graph of  $\mathfrak{S}_n$  starting at  $\sigma$ , of length  $k$  and with defect  $d$ .*

Observe that the adjoint action of  $\mathfrak{S}_n$  on itself determines an action of  $\mathfrak{S}_n$  on its Cayley graph by automorphisms. Thus,  $S(\sigma, k, d)$  depends only on the conjugacy class of  $\sigma$ .

### 3.2. The main expansion.

**Theorem 3.3.** *Let  $N, n \geq 1$  be integers. Let  $(B_t)_{t \geq 0}$  be a Brownian motion on  $U(N)$  starting at the identity and corresponding to the scalar product  $(X, Y) \mapsto -\text{Tr}(XY)$  on  $\mathfrak{u}(N)$ . Let  $M_1, \dots, M_n$  be arbitrary  $N \times N$  complex matrices. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . Then, for all  $t \geq 0$ , we have the following series expansions:*

(14)

$$\mathbb{E} \left[ p_\sigma(M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \right] = e^{-\frac{nt}{2}} \sum_{k, d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} \sum_{|\sigma'| = |\sigma| - k + 2d} \#\Pi_k(\sigma \rightarrow \sigma') p_{\sigma'}(M_1, \dots, M_n).$$

*In particular, if  $m_1, \dots, m_r$  denote the lengths of the cycles of  $\sigma$ , then*

$$(15) \quad \mathbb{E}[\text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \dots \text{Tr}_N(B_{\frac{t}{N}}^{m_r})] = e^{-\frac{nt}{2}} \sum_{k, d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} S(\sigma, k, d).$$

*For all  $T \geq 0$ , both expansions converge uniformly on  $(N, t) \in \mathbb{N}^* \times [0, T]$ .*

In order to understand the role of the defect of a path in our problem, let us write down the result corresponding to Proposition 2.3 for the functions  $p_\sigma$ . As explained earlier, the number of cycles of  $\sigma\tau$  can be either  $\ell(\sigma) + 1$  or  $\ell(\sigma) - 1$ , respectively when the two points exchanged by  $\tau$  belong to the same cycle of  $\sigma$  or to two distinct cycles. For each permutation  $\sigma \in \mathfrak{S}_n$ , we are led to partition  $T_n$  into two classes  $F(\sigma)$  and  $C(\sigma)$ , those which fragment a cycle of  $\sigma$  and those which coagulate two cycles. More precisely,

$$F(\sigma) = \{\tau \in T(n) : \ell(\sigma\tau) = \ell(\sigma) + 1\} \text{ and } C(\sigma) = \{\tau \in T(n) : \ell(\sigma\tau) = \ell(\sigma) - 1\}.$$

Observe that  $F(\text{id}) = C((1 \dots n)) = \emptyset$ . More generally, if the lengths of the cycles of  $\sigma$  are  $m_1, \dots, m_{\ell(\sigma)}$ , then

$$\#F(\sigma) = \frac{1}{2} \left( \sum m_i^2 - n \right) \text{ and } \#C(\sigma) = \frac{1}{2} \left( n^2 - \sum m_i^2 \right).$$

The following result is now a straightforward consequence of Proposition 2.3.

**Proposition 3.4.** *Let  $\sigma$  be a permutation in  $\mathfrak{S}_n$ . Let  $M_1, \dots, M_n$  be  $N \times N$  matrices. Then the following relation holds:*

$$\begin{aligned} \frac{1}{2N} \Delta_{U(N)} p_\sigma(M_1 U, \dots, M_n U) &= -\frac{n}{2} p_\sigma(M_1 U, \dots, M_n U) + \sum_{\tau \in F(\sigma)} p_{\sigma\tau}(M_1 U, \dots, M_n U) + \\ &+ \frac{1}{N^2} \sum_{\tau \in C(\sigma)} p_{\sigma\tau}(M_1 U, \dots, M_n U). \end{aligned}$$

This explains the role played by the defect of a path. Each step which increases the distance to the identity is penalized by a weight  $N^{-2}$ . In the proof of the power series expansion, we use the following lemma.

**Lemma 3.5.** *Let  $t \geq 0$  and  $N > 0$  be real numbers. For all  $\sigma, \sigma' \in \mathfrak{S}_n$  and  $\varepsilon \in \{-1, 1\}$ , define*

$$M_{\sigma, \sigma'}^\varepsilon = \sum_{k=0}^{+\infty} \frac{\varepsilon^k t^k}{k!} \frac{\#\Pi_k(\sigma \rightarrow \sigma')}{N^{k - (\ell(\sigma') - \ell(\sigma))}}.$$

*Then the matrices  $(M_{\sigma, \sigma'}^1)_{\sigma, \sigma' \in \mathfrak{S}_n}$  and  $(M_{\sigma, \sigma'}^{-1})_{\sigma, \sigma' \in \mathfrak{S}_n}$  are each other's inverse.*

**Proof** – If  $\gamma$  is a path in the Cayley graph of  $\mathfrak{S}_n$ , let  $L(\gamma)$  denote its length. First observe that

$$M_{\sigma, \sigma'}^\varepsilon = \sum_{\gamma \in \Pi(\sigma \rightarrow \sigma')} \frac{(\varepsilon t)^{L(\gamma)}}{L(\gamma)! N^{2d(\gamma)}}.$$

Now choose  $\sigma, \sigma'$  in  $S_n$ . Then

$$\begin{aligned} \sum_{\sigma'' \in \mathfrak{S}_n} M_{\sigma, \sigma''}^1 M_{\sigma'', \sigma'}^{-1} &= \sum_{\gamma \in \Pi(\sigma \rightarrow \sigma''), \gamma' \in \Pi(\sigma'', \sigma)} \frac{(-1)^{L(\gamma')} t^{L(\gamma) + L(\gamma')}}{L(\gamma)! L(\gamma')! N^{2d(\gamma) + 2d(\gamma')}} \\ &= \sum_{\gamma \in \Pi(\sigma \rightarrow \sigma')} \sum_{l=0}^{L(\gamma)} \binom{L(\gamma)}{l} (-1)^l \frac{t^{L(\gamma)}}{L(\gamma)! N^{2d(\gamma)}} \\ &= \sum_{\gamma \in \Pi(\sigma \rightarrow \sigma')} \delta_{L(\gamma), 0} \frac{t^{L(\gamma)}}{L(\gamma)! N^{2d(\gamma)}} = \delta_{\sigma, \sigma'}. \end{aligned}$$

The result is proved.  $\square$

In fact,  $M_{\sigma, \sigma'}^\varepsilon$  is the kernel of the operator  $e^{\varepsilon t L}$  on  $\mathbb{C}[\mathfrak{S}_n]$ , where  $(Lf)(\sigma) = \sum_{\tau \in F(\sigma)} f(\sigma\tau) + \frac{1}{N^2} \sum_{\tau \in C(\sigma)} f(\sigma\tau)$ . From this point of view, the lemma above becomes obvious.

**Proof of Theorem 3.3** – We claim that the right-hand side of (14) is an absolutely convergent series. Indeed, since  $N \geq 1$ , the sum of the absolute values is smaller than

$$e^{-\frac{nt}{2}} \max_{\sigma' \in \mathfrak{S}_n} |p_{\sigma'}(M_1, \dots, M_n)| \sum_{k=0}^{+\infty} \frac{t^k}{k!} \sum_{d=0}^{+\infty} S(\sigma, k, d) = e^{-\frac{nt}{2} + n^2 t} \max_{\sigma' \in \mathfrak{S}_n} |p_{\sigma'}(M_1, \dots, M_n)|.$$

The assertion on the uniform convergence of the expansions follows.

In order to prove (14), we start from the expression given by Proposition 2.4, at time  $\frac{t}{N}$  and with an arbitrary deterministic initial condition  $\pi_0 = \sigma$ . It reads

$$(16) \quad \forall \sigma \in \mathfrak{S}_n, \quad \mathbb{E} \left[ p_{\pi_{\frac{t}{N}}}^{st} (M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \middle| \pi_0 = \sigma \right] = e^{-\frac{nt}{2} - \frac{n(n-1)t}{2N}} p_\sigma^{st} (M_1, \dots, M_n).$$

We expand the left hand side by using the properties of  $(\pi_t)_{t \geq 0}$ . This chain jumps at rate  $\binom{n}{2}$  and its jump chain is a standard discrete-time random walk on the Cayley graph of  $\mathfrak{S}_n$ , independent of the jump times. Thus, the left-hand side of (16) is equal to

$$\sum_{k=0}^{\infty} e^{-\binom{n}{2} \frac{t}{N}} \binom{n}{2}^k \frac{t^k}{k! N^k} \frac{1}{\binom{n}{2}^k} \sum_{\sigma' \in S_n} \sum_{\gamma \in \Pi_k(\sigma \rightarrow \sigma')} \mathbb{E} \left[ p_{\sigma'}^{st} (M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \right],$$

where the expectation is now only with respect to the Brownian motion. After simplification, (16) becomes

$$\forall \sigma \in S_n, \quad \sum_{\sigma' \in S_n} \mathbb{E} \left[ p_{\sigma'} (M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \right] \sum_{k=0}^{\infty} \frac{t^k}{k!} \frac{\#\Pi_k(\sigma \rightarrow \sigma')}{N^{k - (\ell(\sigma') - \ell(\sigma))}} = e^{-\frac{nt}{2}} p_{\sigma} (M_1, \dots, M_n).$$

We recognize the expression of  $M_{\sigma, \sigma'}^1$  and, by Lemma 3.5, we conclude that for all  $\sigma \in S_n$ ,

$$\mathbb{E} \left[ p_{\sigma} (M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \right] = e^{-\frac{nt}{2}} \sum_{k=0}^{\infty} \frac{(-1)^k t^k}{k!} \sum_{\sigma' \in S_n} \frac{\#\Pi_k(\sigma \rightarrow \sigma')}{N^{k - (\ell(\sigma') - \ell(\sigma))}} p_{\sigma'} (M_1, \dots, M_n).$$

The first formula follows from the fact that  $|\sigma| = n - \ell(\sigma)$ . Setting  $M_1 = \dots = M_n = I_N$  yields the second formula.  $\square$

Most of the coefficients which appear in the expansion (15) are zero. More precisely, the situation is the following.

**Lemma 3.6.** *Let  $\gamma$  be a path of length  $k$  and defect  $d$  starting at  $\sigma$ . Then the following inequalities hold:*

$$0 \leq d \leq k \quad \text{and} \quad 2d - (\ell(\sigma) - 1) \leq k \leq 2d + (n - \ell(\sigma)).$$

*In particular,  $|k - 2d| \leq n - 1$ .*

*Moreover, let  $k$  and  $d$  be given such that  $0 \leq d \leq k$ . Then  $S(\sigma, 2d + (n - \ell(\sigma)), d) > 0$  and, if  $d \geq \ell(\sigma) - 1$ , then  $S(\sigma, 2d - (\ell(\sigma) - 1), d) > 0$ . Finally, if  $d \leq \ell(\sigma) - 1$ , then  $S(\sigma, d, d) > 0$ .*

**Proof** – Assume that the path finishes at  $\sigma_k$ . Then the first two inequalities reflect simply the fact that  $1 \leq \ell(\sigma_k) \leq n$ .

To prove the second part of the statement, consider  $d \geq 0$ . Recall that  $\sigma$  is fixed. Let us construct a longest possible path starting at  $\sigma$  with defect  $d$ . For this, we minimize the defect at each step. First, we build a path by going from  $\sigma$  down to the identity through a geodesic. This takes  $n - \ell(\sigma)$  steps and the defect of the path is still zero. Then the path must make one step up. Immediately after this, it can go down to the identity again. It can repeat this at most  $d$  times without its defect becoming larger than  $d$ . By then it has length  $2d + (n - \ell(\sigma))$ . Thus we have constructed a path of length  $2d + (n - \ell(\sigma))$  with defect  $d$ . A similar argument works for a shortest path of given defect.  $\square$

In particular, for all  $d \geq 0$ , the contribution of order  $N^{-2d}$  to  $\mathbb{E}[p_{\sigma}(B_{\frac{t}{N}})]$  is a polynomial function of  $t$  of degree  $2d + (n - \ell(\sigma))$  and valuation  $\max(d, 2d - (\ell(\sigma) - 1))$ .

**3.3. Examples.** Let us work out explicitly a few examples.

For  $n = 1$ : there is a single path in the Cayley graph of  $\mathfrak{S}_1$ . It has length and defect 0. Thus, we recover the well-known formula

$$\mathbb{E} \left[ \text{Tr}_N (B_{\frac{t}{N}}) \right] = e^{-\frac{t}{2}}.$$

For  $n = 2$ : for each  $k \geq 0$  there is a unique path of length  $k$  starting at the identity. It has defect  $\lfloor \frac{k+1}{2} \rfloor$ . Thus,

$$\mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}})^2 \right] = e^{-t} \sum_{k=0}^{\infty} \frac{(-1)^k t^k}{k! N^{2 \lfloor \frac{k+1}{2} \rfloor}} = e^{-t} \left( \cosh \frac{t}{N} - \frac{1}{N} \sinh \frac{t}{N} \right).$$

Similarly, for each  $k \geq 0$ , there is a unique path of length  $k$  starting at (12). It has defect  $\lfloor \frac{k}{2} \rfloor$ . Thus,

$$\mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^2) \right] = e^{-t} \sum_{k=0}^{\infty} \frac{(-1)^k t^k}{k! N^{2 \lfloor \frac{k}{2} \rfloor}} = e^{-t} \left( \cosh \frac{t}{N} - N \sinh \frac{t}{N} \right).$$

For  $n = 3$ : the situation is a bit more complicated but it is still possible to compute everything by hand. For a path starting at the identity of length  $k$  and defect  $d$ , we must have  $2d - 2 \leq k \leq 2d$ . Hence, if  $k$  is odd, it must be equal to  $2d - 1$ . So, for all  $l \geq 1$ ,  $S(\text{id}, 2l - 1, l) = 3^{2l-1}$ . If  $k$  is even, then two situations are possible. We leave it as an exercise to check that, for all  $l \geq 1$ ,  $S(\text{id}, 2l, l) = 3^{2l-1}$  and  $S(\text{id}, 2l, l + 1) = 2 \cdot 3^{2l-1}$ . Finally,  $S(\text{id}, 0, 0) = 1$ . We find

$$\mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}})^3 \right] = e^{-\frac{3t}{2}} \left( 1 + \frac{N^2 + 2}{3N^2} \left( \cosh \frac{3t}{N} - 1 \right) - \frac{1}{N} \sinh \frac{3t}{N} \right).$$

Similarly, we find

$$\mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^2) \text{Tr}_N(B_{\frac{t}{N}}) \right] = e^{-\frac{3t}{2}} \left( \cosh \frac{3t}{N} - \frac{N^2 + 2}{3N} \sinh \frac{3t}{N} \right),$$

$$\mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^3) \right] = e^{-\frac{3t}{2}} \left( 1 + \frac{N^2 + 2}{3} \left( \cosh \frac{3t}{N} - 1 \right) - N \sinh \frac{3t}{N} \right).$$

For  $n \geq 4$ , it seems difficult to determine all the coefficients at once and by hand. Nevertheless, the following diagram, which indicates how many edges join the various conjugacy classes of  $\mathfrak{S}_4$  in the Cayley graph allows one to compute specific values of  $S(\sigma, k, d)$ .




FIGURE 3. The Cayley graph of  $\mathfrak{S}_4$  modulo conjugation.

For instance, one can use it to prove the following formulae :

$$e^{2t} \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}})^4 \right] = 1 + \frac{1}{N^2} (-6t + 3t^2) + \frac{1}{N^4} (15t^2 - 20t^3 + 5t^4) + O\left(\frac{1}{N^6}\right),$$

$$e^{2t} \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^4) \right] = (1 - 6t + 8t^2 - \frac{8}{3}t^3) + \frac{1}{N^2} (10t^2 - \frac{58}{3}t^3 + \frac{71}{4}t^4 - \frac{16}{3}t^5) + O\left(\frac{1}{N^4}\right).$$

**3.4. The case of  $SU(N)$ .** Let us conclude this section by stating without proof the following analogue of Theorem 3.3 in the case of the the special unitary group. This theorem is proved exactly like its unitary version, by using the observation made at the end of Section 2.1.

**Theorem 3.7.** *Let  $N, n \geq 1$  be integers. Let  $(B_t)_{t \geq 0}$  be a Brownian motion on  $SU(N)$  starting at the identity and corresponding to the scalar product  $(X, Y) \mapsto -\text{Tr}(XY)$  on  $\mathfrak{su}(N)$ . Let  $M_1, \dots, M_n$  be arbitrary  $N \times N$  complex matrices. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . Then, for all  $t \geq 0$ , we have the following series expansions:*

$$(17) \quad \mathbb{E} \left[ p_\sigma(M_1 B_{\frac{t}{N}}, \dots, M_n B_{\frac{t}{N}}) \right] = e^{-\frac{nt}{2} + \frac{n^2 t}{2N^2}} \sum_{k, d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} \sum_{|\sigma'| = |\sigma| - k + 2d} \#\Pi_k(\sigma \rightarrow \sigma') p_{\sigma'}(M_1, \dots, M_n).$$

In particular, if  $m_1, \dots, m_r$  denote the lengths of the cycles of  $\sigma$ , then

$$(18) \quad \mathbb{E}[\text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \dots \text{Tr}_N(B_{\frac{t}{N}}^{m_r})] = e^{-\frac{nt}{2} + \frac{n^2 t}{2N^2}} \sum_{k, d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} S(\sigma, k, d).$$

For all  $T \geq 0$ , both expansions converge uniformly on  $(N, t) \in \mathbb{N}^* \times [0, T]$ .

#### 4. A REPRESENTATION-THEORETIC DERIVATION OF THE POWER SERIES EXPANSIONS

In this section, we give an alternative derivation of the expansions (15) and (18), based on the representation theory of the unitary and symmetric groups and the relations between symmetric functions.

**4.1. Expansion for the unitary group.** The integer  $N, n \geq 1$  are fixed throughout this section. We write  $\lambda \vdash n$  if  $\lambda = (\lambda_1 \geq \dots \geq \lambda_r > 0)$  is a partition of  $n$ . The integer  $r$  is called the length of  $\lambda$  and we denote it by  $\ell(\lambda)$ . We denote the set of all partitions by  $\mathcal{P}$ .

Let  $\lambda$  partition of  $n$ . We denote by  $s_\lambda$  the Schur function associated to the partition  $\lambda$ , whose definition is given in [12, I.3]. For all  $U \in U(N)$ , the number  $s_\lambda(U)$  is defined as the value of  $s_\lambda$  on the eigenvalues of  $U$ . We will use the fact that, if  $\ell(\sigma) > N$ , then the symmetric polynomial in  $N$  variables determined by  $s_\lambda$  is the zero polynomial. This follows for example from the expression of  $s_\lambda$  as a determinant in the elementary symmetric functions [12, I.3, (3.5)].

Recall the definition of the power sums, that is, the functions  $p_\sigma^{st} : U(N) \rightarrow \mathbb{C}$  for  $\sigma \in \mathfrak{S}_n$  (see Definition 2.1).

The Schur functions and the power sums are related as follows. Let  $\chi^\lambda : \mathfrak{S}_n \rightarrow \mathbb{C}$  denote the character of the irreducible representation of  $\mathfrak{S}_n$  associated with  $\lambda$ . Then one has the following pair of relations [12, I.7, (7.7)]:

$$(19) \quad \forall \lambda \vdash n, s_\lambda = \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \chi^\lambda(\sigma) p_\sigma^{st},$$

$$(20) \quad \forall \sigma \in \mathfrak{S}_n, p_\sigma^{st} = \sum_{\lambda \vdash n} \chi^\lambda(\sigma) s_\lambda.$$

The set  $\widehat{U(N)}$  of isomorphism classes of irreducible representations (irreps) of  $U(N)$  is in one-to-one correspondence with the set  $\mathbb{Z}_+^N$  of non-increasing sequences  $\alpha = (\alpha_1 \geq \dots \geq \alpha_N)$  of elements of  $\mathbb{Z}$ . Even when  $\alpha_N < 0$ , the Schur function  $s_\alpha$  is well-defined and the character of the irrep  $\alpha$  is  $\chi_\alpha(U) = s_\alpha(U)$ .

Let  $(B_t)_{t \geq 0}$  be the Brownian motion on  $U(N)$  of Theorem 3.3. Let  $dU$  denote the Haar measure on  $U(N)$ . For each  $t > 0$ , let  $Q_t$  denote the heat kernel at time  $t$  on  $U(N)$ , that is,

the density of the distribution of  $B_t$  with respect to the Haar measure. Our main result is the following reformulation of (15).

**Theorem 4.1.** *Let  $N, n \geq 1$  be integers. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . Then, for all  $t \geq 0$ ,*

$$(21) \quad N^{-\ell(\sigma)} \int_{U(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = e^{-\frac{nt}{2}} \sum_{k,d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} S(\sigma, k, d).$$

In the course of the proof, we admit two lemmas which we prove afterwards. We have preferred this order to the strict logical order to make the proof easier to follow.

**Proof** – If  $t = 0$ , the result is clearly true. When  $t > 0$ , the proof consists in expanding  $Q_t$  into the sum of its Fourier series and turning all quantities related to  $U(N)$  into quantities related to  $\mathfrak{S}_n$ . For all  $t > 0$ , the function  $Q_t$  is smooth on  $U(N)$  and invariant by conjugation. It admits the following uniformly convergent Fourier expansion [11, Thm 4.4]:

$$(22) \quad Q_t(U) = \sum_{\alpha \in \mathbb{Z}_+^N} e^{-\frac{c_2(\alpha)t}{2}} s_\alpha(I_N) \overline{s_\alpha(U)},$$

where the number  $c_2(\alpha)$  is characterized by the equality  $\Delta_{U(N)} \chi_\alpha = -c_2(\alpha) \chi_\alpha$ . Using the relation (20) to expand  $p_\sigma^{st}(U)$ , we find the following expression for the left-hand side of (21):

$$\int_{U(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = \sum_{\alpha \in \mathbb{Z}_+^N, \mu \vdash n} e^{-\frac{c_2(\alpha)t}{2N}} s_\alpha(I_N) \chi^\mu(\sigma) \int_{U(N)} \overline{s_\alpha(U)} s_\mu(U) dU.$$

By the orthogonality properties of the characters of irreps, the integral in the right-hand side is zero unless  $\alpha = \mu$ . Hence, we can replace the sum over  $\alpha$  and  $\mu$  by a sum over the partitions  $\mu$  such that  $\mu \vdash n$  and  $\ell(\mu) \leq N$ :

$$(23) \quad \int_{U(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = \sum_{\mu \vdash n, \ell(\mu) \leq N} e^{-\frac{c_2(\mu)t}{2N}} s_\mu(I_N) \chi^\mu(\sigma).$$

We still need to express  $s_\mu(I_N)$  and  $c_2(\mu)$  in terms of quantities related to the symmetric group.

In order to compute  $s_\mu(I_N)$ , we use the relation (19). Let us define  $\Omega = \sum_{\sigma \in \mathfrak{S}_n} N^{\ell(\sigma)} \sigma$ . This notation is borrowed from [11]. Then (19) implies the equality

$$(24) \quad s_\mu(I_N) = \frac{1}{n!} \chi^\mu(\Omega).$$

In Lemma 4.3, we will prove directly that  $\chi^\mu(\Omega) = 0$  if  $\ell(\mu) > N$ . This allows us to drop the restriction  $\ell(\mu) \leq N$  in the summation.

Let us compute  $c_2(\mu)$ , the eigenvalue of  $\Delta_{U(N)}$  associated to  $s_\mu$ . Thanks to (4), we know the value of  $\Delta_{U(N)} p_\sigma^{st}$  for all  $\sigma$  and (19) expresses  $s_\mu$  as a linear combination of power sums. Combining these two equations, we find

$$(25) \quad \Delta_{U(N)} s_\mu = -N n s_\mu - \frac{2}{n!} \sum_{\sigma \in \mathfrak{S}_n} \sum_{\tau \in T_n} \chi^\mu(\sigma \tau) p_\sigma^{st},$$

where  $T_n$  is the set of the transpositions of  $\mathfrak{S}_n$ . We now use the following instance of the orthogonality relations between irreducible characters: whenever  $x$  belongs to the group algebra  $\mathbb{C}[\mathfrak{S}_n]$  and  $y$  to the center of the group algebra,

$$(26) \quad \forall \mu \vdash n, \quad \chi^\mu(xy) = \frac{\chi^\mu(x) \chi^\mu(y)}{\chi^\mu(1)}.$$



This relation implies that the last term of (25) is equal to  $n(n-1)\frac{\chi^\mu((12))}{\chi^\mu(1)}s_\mu$ . Hence,

$$(27) \quad c_2(\mu) = nN + n(n-1)\frac{\chi^\mu((12))}{\chi^\mu(1)}.$$

Combining (23), (24) and (27), we find

$$\int_{U(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = e^{-\frac{nt}{2}} \sum_{k \geq 0} \frac{(-t)^k}{k!} \sum_{\mu \vdash n} \frac{\chi^\mu(\Omega)\chi^\mu(\sigma)}{n!} \left( \frac{n(n-1)}{2N} \frac{\chi^\mu((12))}{\chi^\mu(1)} \right)^k.$$

By Lemma 4.2 below, the sum over  $\mu$  is equal to  $N^{\ell(\sigma)} \sum_{d \geq 0} N^{-2d} S(\sigma, k, d)$ . The result follows immediately.  $\square$

**Lemma 4.2.** *Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . Let  $k, d \geq 0$  be integers. Then*

$$(28) \quad \sum_{d \geq 0} \frac{S(\sigma, k, d)}{N^{2d}} = N^{-\ell(\sigma)-k} \sum_{\mu \vdash n} \frac{\chi^\mu(\Omega)\chi^\mu(\sigma)}{n!} \left( \frac{n(n-1)}{2} \frac{\chi^\mu((12))}{\chi^\mu(1)} \right)^k.$$

**Proof** – There is no issue of convergence, since the sum on the left-hand side is finite. Let  $\gamma = (\sigma_0, \dots, \sigma_k)$  be a path of defect  $d$ . By Lemma 3.1,  $\ell(\sigma_k) = \ell(\sigma_0) + k - 2d$ . Hence,

$$\sum_{d \geq 0} \frac{S(\sigma, k, d)}{N^{2d}} = N^{-\ell(\sigma)-k} \sum_{\sigma' \in \mathfrak{S}_n} \#\Pi_k(\sigma \rightarrow \sigma') N^{\ell(\sigma')}.$$

Now,  $\#\Pi_k(\sigma \rightarrow \sigma')$  is the number of  $k$ -tuples  $(\tau_1, \dots, \tau_k) \in T_n^k$  such that  $\sigma\tau_1 \dots \tau_k = \sigma'$ . A standard computation based on the fact that  $\sum_{\mu \vdash n} \chi^\mu(1)\chi^\mu(\sigma) = (n!)^2 \delta_{\sigma, \text{id}}$  and on (26) leads to

$$\#\Pi_k(\sigma \rightarrow \sigma') = \sum_{\mu \vdash n} \frac{\chi^\mu(1)\chi^\mu(\sigma^{-1}\sigma')}{n!} \left( \frac{n(n-1)}{2} \frac{\chi^\mu((12))}{\chi^\mu(1)} \right)^k.$$

The result follows by summing over  $\sigma'$  and applying (26) again.  $\square$

**Lemma 4.3.** *Define  $\Omega \in \mathbb{C}[\mathfrak{S}_n]$  by  $\Omega = \sum_{\sigma \in \mathfrak{S}_n} N^{\ell(\sigma)} \sigma$ . Then the following relations hold.*

1. *For all  $\mu \vdash n$  such that  $\ell(\mu) \leq N$ ,  $\chi^\mu(\Omega) = n! s_\mu(I_N)$ .*
2. *For all  $\mu \vdash n$  such that  $\ell(\mu) > N$ ,  $\chi^\mu(\Omega) = 0$ .*

**Proof** – The first assertion follows immediately from (19).

In order to prove the second assertion, let us introduce the Jucys-Murphy elements  $X_1, \dots, X_n$  of  $\mathbb{C}[\mathfrak{S}_n]$ , defined by  $X_1 = 0$  and  $X_i = (1 \ i) + (2 \ i) + \dots + (i-1 \ i)$  for  $i \in \{2, \dots, n\}$ . They generate a maximal Abelian subalgebra of  $\mathbb{C}[\mathfrak{S}_n]$ . In particular, they can be simultaneously diagonalized in every irreducible representation of  $\mathfrak{S}_n$ . We borrow the following statements from [13].

Let  $\mu = (\mu_1, \dots, \mu_r)$  be a partition of  $n$ . The subset  $D_\mu = \{(i, j) \in (\mathbb{N}^*)^2 : j \leq \mu_i\}$  of  $\mathbb{Z}^2$  is called the diagram of  $\mu$ . An element  $(i, j)$  of  $D_\mu$  is called a box and its content is defined as the integer  $c(i, j) = j - i$ .

The space of the irreducible representation of  $\mathfrak{S}_n$  associated to  $\mu$  admits a basis which diagonalizes the Jucys-Murphy elements and is indexed by the bijections  $t : D_\mu \rightarrow \{1, \dots, n\}$  which are increasing in each variable. These bijections are usually called tableaux. The eigenvalue of the Jucys-Murphy element  $X_k$  on the vector associated to the tableau  $t$  is the content of the box  $t^{-1}(k)$ .

We need also the following well-known fact: for every  $k \in \{0, \dots, n-1\}$ , the  $k$ -th elementary symmetric function of the Jucys-Murphy elements is equal to  $\sum_{\sigma \in \mathfrak{S}_n} \mathbf{1}_{|\sigma|=k} \sigma$ , where  $|\sigma| = n - \ell(\sigma)$ . This can be proved as follows. For each  $m \in \{1, \dots, n\}$ , let us imbed  $\mathfrak{S}_m$  into  $\mathfrak{S}_n$  as the subgroup which leaves  $\{m+1, \dots, n\}$  invariant. For all  $m \in \{1, \dots, n\}$  and  $k \in \{0, \dots, m-1\}$ , set  $\Sigma_{m,k} = \sum_{\sigma \in \mathfrak{S}_m} \mathbf{1}_{|\sigma|=k} \sigma$ . Let us also define  $e_{m,k} = \sum_{i_1 < \dots < i_k \leq m} X_{i_1} \dots X_{i_k}$ . We need to prove that  $e_{m,k} = \Sigma_{m,k}$ . This is clearly true if  $k \in \{0, 1\}$ . The general case follows by induction on  $m$ , each inductive step being proved by induction on  $k$ , thanks to the relations

$$\Sigma_{m,k} = \Sigma_{m-1,k} + \Sigma_{m-1,k-1} X_k \quad \text{and} \quad e_{m,k} = e_{m-1,k} + e_{m-1,k-1} X_k.$$

From the equality proved in the last paragraph and the relation  $|\sigma| = n - \ell(\sigma)$ , we deduce the following equality in the polynomial ring  $\mathbb{C}[\mathfrak{S}_n][z]$ :

$$\prod_{i=1}^n (z + X_i) = \sum_{\sigma \in \mathfrak{S}_n} z^{\ell(\sigma)} \sigma.$$

Evaluating at  $z = N$  and applying  $\chi^\mu$ , we find

$$\chi^\mu(\Omega) = \chi^\mu \left( \prod_{i=1}^n (N + X_i) \right) = \sum_{t \text{ tableau}} \prod_{i=1}^n (N + c(t^{-1}(i))) = \chi^\mu(1) \prod_{i=1}^{\ell(\mu)} \prod_{j=1}^{\mu_i} (N + j - i).$$

If  $\ell(\mu) \geq N+1$ , then  $(N+1, 1)$  is a box of  $D_\mu$ , whose content is  $-N$ . It follows that  $\chi^\mu(\Omega) = 0$  in this case.  $\square$

**4.2. Expansion for the special unitary group.** Let us apply a similar analysis to the special unitary group in order to derive (18) in another way.

By restriction, any irrep of  $U(N)$  determines an irrep of  $SU(N)$  and the restrictions of  $\alpha, \alpha' \in \mathbb{Z}_\downarrow^N$  are isomorphic if and only if there exists  $k \in \mathbb{Z}$  such that  $\alpha' = (\alpha_1 + k, \dots, \alpha_N + k)$ . Hence, the set of irreps of  $SU(N)$  is in one-to-one correspondence with the set of partitions of length at most  $N-1$  and the character of the irreducible representation corresponding to a partition  $\lambda$  is given by the Schur function  $s_\lambda$ .

Let  $(B_t)_{t \geq 0}$  be the Brownian motion on  $SU(N)$  of Theorem 18. Let  $dU$  denote the Haar measure on  $SU(N)$ . For each  $t > 0$ , let  $Q_t$  denote the heat kernel at time  $t$  on  $SU(N)$ , that is, the density of the distribution of  $B_t$  with respect to the Haar measure. We reformulate (18) as follows.

**Theorem 4.4.** *Let  $N, n \geq 1$  be integers. Let  $\sigma$  be an element of  $\mathfrak{S}_n$ . Then, for all  $t \geq 0$ ,*

$$(29) \quad N^{-\ell(\sigma)} \int_{SU(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = e^{-\frac{nt}{2} + \frac{tn^2}{2N^2}} \sum_{k,d=0}^{+\infty} \frac{(-1)^k t^k}{k! N^{2d}} S(\sigma, k, d).$$

**Proof** – If  $t = 0$ , both sides are equal to 1. Assume that  $t > 0$ . The Fourier expansion of  $Q_t$  is then the following:

$$(30) \quad Q_t(U) = \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda) \leq N-1}} e^{-\frac{c_2(\lambda)t}{2}} s_\lambda(I_N) \overline{s_\lambda(U)},$$

where now  $c'_2(\lambda)$  is defined by the equality  $\Delta_{SU(N)}\chi_\lambda = -c'_2(\lambda)\chi_\lambda$ . Combined with the relation (20), it implies

$$\int_{SU(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = \sum_{\substack{\lambda \in \mathcal{P}, \mu \vdash n \\ \ell(\lambda) \leq N-1}} e^{-\frac{c'_2(\lambda)t}{2N}} s_\lambda(I_N) \chi^\mu(\sigma) \int_{SU(N)} \overline{s_\lambda(U)} s_\mu(U) dU.$$

When  $\ell(\mu) > N$ ,  $s_\mu$  is identically zero on  $SU(N)$ . When  $\ell(\mu) \leq N-1$ , then the integral in the right-hand side of the last equation is equal to  $\delta_{\lambda, \mu}$ . Let us consider the terms of the sum for which  $\ell(\mu) = N$ . In this case, let us write  $\mu = \mu' + (\mu_N, \dots, \mu_N)$ , so that  $\ell(\mu') \leq N-1$ . Then the integral is equal to  $\delta_{\lambda, \mu'}$  and we may assume that  $\lambda = \mu'$ . In this case,  $s_\lambda(I_N) = s_{\mu'}(I_N) = s_\mu(I_N)$  and  $c'_2(\lambda) = c'_2(\mu')$ . Hence,

$$\int_{SU(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = \sum_{\mu \vdash n, \ell(\mu) \leq N-1} e^{-\frac{c'_2(\mu)t}{2N}} s_\mu(I_N) \chi^\mu(\sigma) + \sum_{\mu \vdash n, \ell(\mu) = N} e^{-\frac{c'_2(\mu')t}{2N}} s_\mu(I_N) \chi^\mu(\sigma).$$

Let us first compute  $c'_2(\mu)$  when  $\ell(\mu) \leq N-1$ . For this, we use the fact that  $s_\mu$  is an eigenvector of  $\Delta_{U(N)}$  whose restriction to  $SU(N)$  is  $s_\mu$ . Since  $\Delta_{SU(N)} p_\sigma^{st} = \left( \Delta_{U(N)} + \frac{n^2}{N} \right) p_\sigma^{st}$  whenever  $\sigma \in \mathfrak{S}_n$ , we find

$$\Delta_{SU(N)} s_\mu = \left( -Nn - n(n-1) \frac{\chi^\mu((12))}{\chi^\mu(1)} + \frac{n^2}{N} \right) s_\mu.$$

When  $\ell(\mu) = N$ , we are interested in  $s_{\mu'}$  but  $s_\mu$  is still an eigenvector of  $\Delta_{U(N)}$  whose restriction to  $SU(N)$  is  $s_{\mu'}$ . Hence,

$$\Delta_{SU(N)} s_{\mu'} = \Delta_{SU(N)} s_\mu = \left( -Nn - n(n-1) \frac{\chi^\mu((12))}{\chi^\mu(1)} + \frac{n^2}{N} \right) s_\mu = -c'_2(\mu') s_{\mu'}.$$

Thus, in both sums, the argument of the exponential is  $-\frac{nt}{2} + \frac{n^2 t}{2N^2} - t \frac{n(n-1)}{2N} \frac{\chi^\mu((12))}{\chi^\mu(1)}$ . Using this fact and the first assertion of Lemma 4.3, we find

$$\int_{SU(N)} p_\sigma^{st}(U) Q_{\frac{t}{N}}(U) dU = e^{-\frac{nt}{2} + \frac{n^2 t}{2N^2}} \sum_{\mu \vdash n, \ell(\mu) \leq N} e^{-t \frac{n(n-1)}{2N} \frac{\chi^\mu((12))}{\chi^\mu(1)}} \frac{\chi^\mu(\Omega) \chi^\mu(\sigma)}{n!}.$$

The second assertion of Lemma 4.3 tells us that we can remove the restriction  $\ell(\mu) \leq N$ , since the other terms are zero. After expanding the exponential, Lemma 4.2 allows us to finish the proof just as in the unitary case.  $\square$

## 5. ASYMPTOTIC DISTRIBUTION

One of the consequences of Theorem 3.3 is that the limit as  $N$  tends to infinity of  $\mathbb{E}[p_\sigma(B_{\frac{t}{N}})]$  exists. Using Lemma 3.6, we get the following result.

**Proposition 5.1.** *Consider  $\sigma \in \mathfrak{S}_n$ . The limit of  $\mathbb{E}[p_\sigma(B_{\frac{t}{N}})]$  as  $N$  tends to infinity exists and it is equal to*

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[ p_\sigma(B_{\frac{t}{N}}) \right] = e^{-\frac{nt}{2}} \sum_{k=0}^{|\sigma|} (-1)^k \frac{S(\sigma, k, 0)}{k!} t^k.$$

In this section, we determine  $S(\sigma, k, 0)$  for all  $\sigma$  and  $k \geq 0$ . For this, we prove the factorization property and use the relation between the metric geometry of the Cayley graph of  $S_n$  and the lattice of non-crossing partitions of the cycle  $(1 \dots n)$ .

**5.1. The factorization property.** The factorization property is the following result. It reduces the problem of the determination of  $S(\sigma, k, 0)$  to the case where  $\sigma$  is a cycle.

**Proposition 5.2.** *Let  $m_1, \dots, m_r$  be positive integers. Then*

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \dots \text{Tr}_N(B_{\frac{t}{N}}^{m_r}) \right] = \lim_{N \rightarrow \infty} \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \right] \dots \lim_{N \rightarrow \infty} \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_r}) \right].$$

More precisely,

$$(31) \quad \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \dots \text{Tr}_N(B_{\frac{t}{N}}^{m_r}) \right] - \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_1}) \right] \dots \mathbb{E} \left[ \text{Tr}_N(B_{\frac{t}{N}}^{m_r}) \right] = O(N^{-2}),$$

uniformly in  $t$  on bounded intervals.

We start by proving the following property of the numbers  $S(\sigma, k, 0)$ . It is in fact equivalent to the proposition.

**Proposition 5.3.** *Consider  $\sigma \in S_n$ . Assume that  $\sigma = c_1 \dots c_{\ell(\sigma)}$  is the decomposition of  $\sigma$  as a product of cycles with disjoint support. Then*

$$(32) \quad \forall k \geq 0, \quad S(\sigma, k, 0) = \sum_{l_1 + \dots + l_{\ell(\sigma)} = k} \frac{k!}{l_1! \dots l_{\ell(\sigma)}!} S(c_1, l_1, 0) \dots S(c_{\ell(\sigma)}, l_{\ell(\sigma)}, 0).$$

**Proof** – The number  $S(\sigma, k, 0)$  is the number of paths of length  $k$  starting at  $\sigma$  and which at each step move towards a permutation with one more cycle than their current position. As we already observed several times, each step of such a path corresponds to the multiplication by a transposition which exchanges two points which belong to the same cycle of  $\sigma$ . There is thus a natural partition of the set of all steps of such a path, according to the cycle of  $\sigma$  in which their support is contained. Let us introduce some notation. Let  $(\sigma_0 = \sigma, \sigma_1, \dots, \sigma_k)$  be a path with defect zero. For each  $i \in \{1, \dots, k\}$ , set  $\tau_i = \sigma_{i-1}^{-1} \sigma_i$ . Let  $(C_1, \dots, C_{\ell(\sigma)})$  be the partition of  $\{1, \dots, k\}$  determined by the fact that  $i \in C_j$  if and only if the support of  $\tau_i$  is contained in the support of  $c_j$ . Then it is clear that, for all  $j \in \{1, \dots, \ell(\sigma)\}$ , the transpositions  $(\tau_i, i \in C_j)$  are the steps of a path with defect zero starting from  $c_j$ .

Hence, constructing a path of length  $k$  starting at  $\sigma$  and with defect zero is equivalent to constructing a collection of  $\ell(\sigma)$  paths with defect zero starting at  $c_1, \dots, c_{\ell(\sigma)}$  respectively, whose lengths  $l_1, \dots, l_{\ell(\sigma)}$  add up to  $k$ , and a shuffling of the steps of these paths, that is, a sequence  $(C_1, \dots, C_{\ell(\sigma)})$  of subsets of  $\{1, \dots, k\}$  which partition  $\{1, \dots, k\}$  and whose cardinals are  $l_1, \dots, l_{\ell(\sigma)}$  respectively. The equation (32) is just the translation in symbols of the last sentence.  $\square$

**Proof of Proposition 5.2** – By Theorem 3.3 and Proposition 5.3, the terms of degree  $N^0$  of the difference on the left hand side of (31) vanish. Hence, this difference is of the form  $N^{-2}F(t, N^{-2})$  for some entire function  $F$ . The result follows.  $\square$

We have observed after Definition 3.2 that  $S(\sigma, k, d)$  depends only on the conjugacy class of  $\sigma$ . Hence, we need to compute  $S((1 \dots m), k, 0)$ . The arguments of the proof of Proposition 5.3 show that the paths of defect 0 starting at  $(1 \dots m)$  stay in  $\mathfrak{S}_m$  if we identify  $\mathfrak{S}_m$  with the subgroup of  $\mathfrak{S}_n$  which leaves  $\{m+1, \dots, n\}$  invariant. Hence, we are reduced to the computation of  $S((1 \dots n), k, 0)$  for all  $n \geq 1$  and  $k \in \{0, \dots, n-1\}$ . This computations involves non-crossing partitions. For the sake of being self-contained, we give a brief review of the properties of non-crossing partitions that we use.

**5.2. Non-crossing partitions.** Let  $P = \{P_1, \dots, P_\ell\}$  be a partition of  $\{1, \dots, n\}$ . The partition  $P$  is said to be non-crossing if there does not exist  $i, j, k, l \in \{1, \dots, n\}$  such that  $i < j < k < l$  and  $r, s \in \{1, \dots, \ell\}$  with  $r \neq s$  such that  $i, k \in P_r$  and  $j, l \in P_s$ . Another way to formulate the fact that  $P$  is non-crossing is the following. For each class  $P_j$  of the partition, let  $H_j$  denote the convex hull in  $\mathbb{C}$  of  $\{e^{\frac{2ik\pi}{n}} : k \in P_j\}$ . Then  $P$  is non-crossing if and only if for all  $i, j \in \{1, \dots, \ell\}$ ,  $i \neq j \Rightarrow H_i \cap H_j = \emptyset$ . This notion is relative to the cyclic order on  $\{1, \dots, n\}$  determined by  $(1 \dots n)$ . We denote by  $NC(n)$  the set of non-crossing partitions of the cycle  $(1 \dots n)$ . This set has been first considered by Kreweras in [10].

The fineness relation between partitions restricted to  $NC(n)$  makes  $NC(n)$  a poset. More precisely, we say that  $P_1 \preceq P_2$  if every class of  $P_1$  is contained in a class of  $P_2$ . The poset  $(NC(n), \preceq)$  is in fact a lattice, which means that suprema and infima exist. There is in particular a maximum,  $\{\{1, \dots, n\}\}$ , which we denote by  $1_n$ , and a minimum,  $\{\{1\}, \dots, \{n\}\}$ , which we denote by  $0_n$ . The poset  $NC(n)$  can be made into a graph by joining two partitions  $P$  and  $Q$  if they are distinct and comparable, say  $P \prec Q$ , and the interval  $[P, Q] = \{R \in NC(n) : P \preceq R \preceq Q\}$  is reduced to  $\{P, Q\}$ .

A non-crossing partition  $P = (P_1, \dots, P_\ell)$  of the cycle  $(1 \dots n)$  determines an element  $\sigma_P$  of  $\mathfrak{S}_n$  as follows: take the cycles of  $\sigma_P$  to be the classes of  $P$  with the cyclic order induced by  $(1 \dots n)$ . In symbols, if  $i \in P_j$ , then

$$\sigma_P(i) = (1 \dots n)^k(i), \text{ where } k = \min\{l \geq 1 : (1 \dots n)^l(i) \in P_j\}.$$

In particular,  $\sigma_{0_n} = \text{id}$  and  $\sigma_{1_n} = (1 \dots n)$ . The partial order on  $NC(n)$  corresponds via the mapping  $P \mapsto \sigma_P$  to the following partial order on  $\mathfrak{S}_n$ .

Consider  $\sigma_1, \sigma_2 \in \mathfrak{S}_n$ . Recall that  $|\sigma_1| = n - \ell(\sigma_1)$ , the minimal number of terms of a decomposition of  $\sigma_1$  in a product of transpositions, is the distance from  $\text{id}$  to  $\sigma_1$  in the Cayley graph of  $\mathfrak{S}_n$  generated by  $T_n$ . By definition, we say that  $\sigma_1 \preceq \sigma_2$  if  $|\sigma_2| = |\sigma_1| + |\sigma_1^{-1}\sigma_2|$ . In words,  $\sigma_1 \preceq \sigma_2$  if and only if there exists a geodesic path from  $\text{id}$  to  $\sigma_2$  through  $\sigma_1$ . The identity is the minimum of  $\mathfrak{S}_n$  for this partial order, and the  $n$ -cycles the (pairwise incomparable) maximal elements. The next lemma is well known and its proof is left to the reader.

**Lemma 5.4.** *The mapping from  $NC(n)$  to  $\mathfrak{S}_n$  which sends a partition  $P$  to the permutation  $\sigma_P$  is an isomorphism of posets from  $NC(n)$  onto  $[\text{id}, (1 \dots n)] = \{\sigma \in \mathfrak{S}_n : \sigma \preceq (1 \dots n)\}$ .*

As a consequence of this Lemma,  $S((1 \dots n), k, 0)$  is the number of decreasing paths of length  $k$  starting at  $1_n$  in  $NC(n)$ . It turns out to be easier to count increasing paths in  $NC(n)$  starting at  $0_n$ . They are in one-to-one correspondence by the following duality property of  $NC(n)$  discovered by Kreweras.

For  $\sigma \in [\text{id}, (1 \dots n)]$ , let us introduce  $K(\sigma) = \sigma^{-1}(1 \dots n)$ . It is readily checked that  $K$  is a decreasing bijection of  $[\text{id}, (1 \dots n)]$ . The corresponding decreasing bijection of  $NC(n)$  is called the Kreweras complementation. It can be described combinatorially at the level of non-crossing partitions as follows.

Given a partition  $R$  of  $\{1, \dots, n\}$  and a partition  $S$  of  $\{1, \dots, n\} \simeq \{\bar{1}, \dots, \bar{n}\}$ , let  $R \cup S$  denote the partition of  $\{1, \bar{1}, 2, \bar{2}, \dots, n, \bar{n}\}$  obtained by merging  $R$  and  $S$ . Even if  $R$  and  $S$  are non-crossing,  $R \cup S$  may be crossing with respect to the cyclic order  $(1, \bar{1}, 2, \bar{2}, \dots, n, \bar{n})$ . Now let  $P$  be a non-crossing partition of  $\{1, \dots, n\}$ . The partition  $K(P)$  is by definition the largest element of  $NC(n)$  such that  $P \cup K(P)$  is non-crossing.

The following result summarizes this discussion of non-crossing partitions in relation to our problem.

**Proposition 5.5.** *For all  $n \geq 1$  and  $k \geq 0$ ,  $S((1 \dots n), k, 0)$  is the number of increasing paths of length  $k$  starting at  $\{\{1\}, \dots, \{n\}\}$  in the lattice of non-crossing partitions of  $(1 \dots n)$ .*




FIGURE 4. The Kreweras complement of  $\{\{1, 3, 12\}, \{2\}, \{4, 8, 9\}, \{5, 6, 7\}, \{10, 11\}\}$  is  $\{\{1, 2\}, \{3, 9, 11\}, \{4, 7\}, \{5\}, \{6\}, \{8\}, \{10\}, \{12\}\}$ .

It remains to count these paths. To do this, we use the fact that  $S((1 \dots n), n-1, 0) = n^{n-2}$ . This is a classical result of combinatorics, since  $S((1 \dots n), n-1, 0)$  is the number of ways to write a  $n$ -cycle as a product of  $n-1$  transpositions.

**Proposition 5.6.** *For all  $n \geq 1$  and  $k \geq 0$ ,*

$$S((1 \dots n), k, 0) = \binom{n}{k+1} n^{k-1}.$$

*It is understood that this number is zero if  $k \geq n$ .*

**Proof** – We count the increasing paths of length  $k$  in  $NC(n)$  starting at  $\{\{1\}, \dots, \{n\}\}$  by first regrouping them according to their terminal point. The possible terminal points of these paths are exactly the non-crossing partitions of  $(1 \dots n)$  into  $n-k$  classes. Such partitions may be classified according to the number of singletons they contain, the number of pairs, and so on.

Let  $s_1, \dots, s_n$  be non-negative integers such that  $s_1 + \dots + s_n = n-k$  and  $s_1 + 2s_2 + \dots + ns_n = n$ . We say that a partition is of type  $(s_1, \dots, s_n)$  if it contains exactly  $s_i$  classes of cardinal  $i$  for each  $i \in \{1, \dots, n\}$ . The number of non-crossing partitions of type  $(s_1, \dots, s_n)$  has been computed by Kreweras [10]. It is equal to  $\frac{n!}{(k+1)!s_1! \dots s_n!}$ .

Let  $P$  be a non-crossing partition of type  $(s_1, \dots, s_n)$ . An increasing path from  $\{\{1\}, \dots, \{n\}\}$  to  $P$  looked at in the reverse direction is a decreasing path from  $P$  to  $\{\{1\}, \dots, \{n\}\}$ . There are as many such paths as there are geodesic paths from the permutation  $\sigma_P$  induced by  $P$  to the identity, that is,  $S(\sigma_P, k, 0)$ . Let us apply Lemma 5.3 to compute this number. The only non-zero term in the sum corresponds to the situation where  $l_i = |c_i|$  for each  $i \in \{1, \dots, \ell(\sigma_P)\}$ . Since  $S(c, |c|, 0) = m^{m-2}$  for every cycle  $c$  of size  $m$ , we find the formula

$$S((1 \dots n), k, 0) = \sum_{(s_1, \dots, s_n)} \frac{n!}{(k+1)!s_1! \dots s_n!} \frac{k!}{1!s_2! \dots (n-1)!s_n} 2^{0s_2} 3^{1s_3} \dots n^{(n-2)s_n},$$

where the sum is extended to all possible types of partitions of  $\{1, \dots, n\}$  into  $n-k$  classes. Let us enumerate the possible types by enumerating the partitions themselves. If we do this, each type will appear as many times as the number of partitions of this specific type. The number of partitions of type  $(s_1, \dots, s_n)$  is

$$\frac{n!}{1!^{s_1} s_1! \dots n!^{s_n} s_n!}.$$

Hence, we find the following expression:

$$S((1 \dots n), k, 0) = \frac{1}{k+1} \sum_P 1^{(1-1)s_1} 2^{(2-1)s_2} \dots n^{(n-1)s_n},$$

where the sum runs over all partitions of  $\{1, \dots, n\}$  into  $n - k$  classes. Now the right hand side has been computed by Kreweras in [9] and it is equal to

$$S((1 \dots n), k, 0) = \frac{1}{k+1} \binom{n-1}{n-k-1} n^k = \binom{n}{k+1} n^{k-1}.$$

This is the expected result.  $\square$

Let us state separately the following result which has been used in the course of this proof.

**Lemma 5.7.** *Let  $P \in NC(n)$  be a partition of type  $(s_1, \dots, s_n)$ . There are exactly*

$$\frac{(s_2 + \dots + (n-1)s_n)!}{1!^{s_2} \dots (n-1)!^{s_n}} 2^{0s_2} \dots n^{(n-2)s_n}$$

increasing paths from  $0_n$  to  $P$ .

The explicit expression of the large  $N$  limit obtained in this section has been stated by Singer in [15] and proved by Biane in [1].

**Remark 5.8.** *The asymptotic distribution of  $B_{\frac{t}{N}}$  as  $N$  tends to infinity is the unique probability measure  $\mu_t$  on the group  $\mathbb{U}$  of complex numbers of modulus 1 such that, for all  $n \in \mathbb{N}$ ,  $\int_{\mathbb{U}} z^n \mu(dz) = \int_{\mathbb{U}} z^{-n} \mu_t(dz) = e^{-\frac{nt}{2}} \sum_{k=0}^{n-1} \frac{(-nt)^k}{nk!} \binom{n}{k+1}$ . This expression of the moments of  $\mu_t$  is not very easy to handle, if only numerically, because it is an alternate sum of large numbers. Let us point out two analytical ways of studying  $\mu_t$ .*

The  $S$ -transform of  $\mu_t$  is its moment generating function, defined by  $S_t(z) = \int_{\mathbb{U}} \frac{\xi z}{1-\xi z} d\mu_t(\xi) = \sum_{n=1}^{\infty} m_{n,t} z^n$ , where  $m_{n,t}$  is the  $n$ -th moment of  $\mu_t$ . Since  $|m_{n,t}| \leq 1$  for all  $n$  and  $t$ , the function  $S_t$  is holomorphic on the unit disk  $\mathbb{D} = \{z : |z| < 1\}$ . Moreover, since  $S_t(z) = z + O(z^2)$ , there exist a reciprocal function to  $S_t$  in a neighbourhood of 0, which we denote by  $\chi_t$ . It turns out that  $\chi_t$  is much simpler than  $S_t$ :  $\chi_t(z) = \frac{z}{z+1} e^{t(z+\frac{1}{2})}$ , as one can easily check by using Lagrange's inversion formula. By studying  $\chi_t$ , Biane proved in [3] the following facts. For each  $t > 0$ , the measure  $\mu_t$  has a continuous density with respect to the uniform measure on  $\mathbb{U}$ . For  $t \in (0, 4]$ , this density is zero exactly on the set  $\left\{ e^{i\theta} : |\theta - \pi| \leq 2 \arctan \sqrt{\frac{4-t}{t}} - \frac{1}{2} \sqrt{t(4-t)} \right\}$ . For  $t > 4$ , the density of  $\mu_t$  is positive. Finally, for all  $t > 0$ , the density of  $\mu_t$  at  $e^{i\theta}$  is a real analytic function of  $\theta$  on the relative interior of its support.

Another way of studying  $\mu_t$  is to observe that  $m_{n,t} = \frac{e^{-\frac{nt}{2}}}{2in\pi} \oint e^{-ntz} \left(1 + \frac{1}{z}\right)^n dz$ , the integral being along any contour of index 1 with respect to 0. In [6], Gross and Matytsin use this expression and the saddle point method to exhibit the following phase transition with respect to  $t$ : if  $t \in (0, 4)$ ,  $m_{n,t}$  decays with  $n$  like  $n^{-\frac{3}{2}}$ , whereas for  $t \in (4, +\infty)$ , it decays exponentially. Along the same lines, one can check that for  $t = 4$ ,  $m_{n,4}$  decays like  $n^{-\frac{4}{3}}$ . This indicates that the density of  $\mu_4$  is less regular than the density of  $\mu_t$  for  $t \in (0, 4)$ . This behaviour is consistent with a general result of Biane about additive convolution with the semi-circle law [2].

**5.3. Almost sure convergence.** The material gathered so far allows us to prove very easily the following result.

**Proposition 5.9.** *Consider  $\sigma \in \mathfrak{S}_n$ . Then, uniformly in  $t$  on bounded intervals,*

$$\text{Var} \left[ p_{\sigma} \left( B_{\frac{t}{N}} \right) \right] = O(N^{-2}).$$

In particular, on any probability space on which a Brownian motion on  $U(N)$  is defined for  $N$  large enough, the following convergence holds almost surely and in  $L^2$  :

$$\lim_{N \rightarrow \infty} p_\sigma(B_{\frac{t}{N}}) = e^{-\frac{nt}{2}} \sum_{k=0}^{n-\ell(\sigma)} (-1)^k S(\sigma, k, 0) \frac{t^k}{k!}.$$

**Proof** – Let us denote by  $\sigma \times \sigma$  the element of  $\mathfrak{S}_{2n}$  which sends  $i$  on  $\sigma(i)$  and  $n+i$  on  $n+\sigma(i)$  for each  $i \in \{1, \dots, n\}$ . With this notation,  $p_\sigma^2 = p_{\sigma \times \sigma}$ , so that

$$\begin{aligned} \text{Var} \left[ p_\sigma(B_{\frac{t}{N}}) \right] &= \mathbb{E} \left[ p_{\sigma \times \sigma}(B_{\frac{t}{N}}) \right] - \mathbb{E} \left[ p_\sigma(B_{\frac{t}{N}}) \right]^2 \\ &= \sum_{k,d=0}^{\infty} \frac{(-1)^k t^k}{k! N^{2d}} \left[ S(\sigma \times \sigma, k, d) - \sum_{\substack{l_1+l_2=k \\ d_1+d_2=d}} \frac{k!}{l_1! l_2!} S(\sigma, l_1, d_1) S(\sigma, l_2, d_2) \right], \end{aligned}$$

where the last expression follows after simplification from Theorem 3.3. In the term corresponding to  $d = 0$ , both  $d_1$  and  $d_2$  must be equal to 0. By the same argument of support as in the proof of Proposition 5.3, we find

$$S(\sigma \times \sigma, k, 0) = \sum_{l_1+l_2=k} \frac{k!}{l_1! l_2!} S(\sigma, l_1, 0) S(\sigma, l_2, 0).$$

The result follows immediately.  $\square$

## 6. ASYMPTOTIC FREENESS

In this section, we prove that independent Brownian motions on  $U(N)$  converge in distribution, as  $N$  tends to infinity, towards free non-commutative random variables. We do not consider  $*$ -freeness, that is, we do not consider products involving  $B_{\frac{t}{N}}^{-1}$ . In fact, the asymptotic  $*$ -freeness of independent Brownian motions is a consequence of the conjugation-invariance of their distribution, as proved in [1, Lemma 6]. Here we use Speicher's characterization of freeness by the vanishing of mixed free cumulants. This combinatorial characterization is very well suited to the approach we have adopted in this paper.

**6.1. The factorization property.** Let us start by slightly improving Proposition 5.2, by including the extra deterministic matrices  $M_1, \dots, M_n$  of Theorem 3.3. We consider these matrices as elements of the non-commutative probability space  $(\mathbb{M}_N(\mathbb{C}), \text{Tr}_N)$  and speak of their distribution accordingly.

**Proposition 6.1.** *Let  $(M_1^{(N)}, \dots, M_n^{(N)})_{N \geq 1}$  be a sequence of families of  $N \times N$  matrices. Assume that this sequence converges in distribution. Consider  $\sigma \in \mathfrak{S}_n$ . Write  $\sigma$  as a product of cycles :  $\sigma = (i_{1,1} \dots i_{1,m_1}) \dots (i_{\ell(\sigma),1} \dots i_{\ell(\sigma),m_{\ell(\sigma)}})$ . Then*

$$(33) \quad \lim_{N \rightarrow \infty} \mathbb{E} \left[ p_\sigma(M_1^{(N)} B_{\frac{t}{N}}, \dots, M_n^{(N)} B_{\frac{t}{N}}) \right] = \prod_{r=1}^{\ell(\sigma)} \lim_{N \rightarrow \infty} \mathbb{E} \text{Tr}_N \left( M_{i_{r,1}}^{(N)} B_{\frac{t}{N}} \dots M_{i_{r,m_r}}^{(N)} B_{\frac{t}{N}} \right).$$

More precisely,

$$(34) \quad \mathbb{E} \left[ p_\sigma(M_1^{(N)} B_{\frac{t}{N}}, \dots, M_n^{(N)} B_{\frac{t}{N}}) \right] - \prod_{r=1}^{\ell(\sigma)} \mathbb{E} \text{Tr}_N \left( M_{i_{r,1}}^{(N)} B_{\frac{t}{N}} \dots M_{i_{r,m_r}}^{(N)} B_{\frac{t}{N}} \right) = O(N^{-2}),$$



uniformly in  $t$  on bounded intervals. It is understood that all limits exist.

**Proof** – According to Theorem 3.3, the left hand side of (33) is equal to

$$(35) \quad e^{-\frac{nt}{2}} \sum_{k=0}^{\infty} \frac{(-t)^k}{k!} \sum_{|\sigma'|=|\sigma|-k} \#\Pi_k(\sigma \rightarrow \sigma') \lim_{N \rightarrow \infty} p_{\sigma'}(M_1^{(N)}, \dots, M_n^{(N)}).$$

This last limit exists for all  $\sigma'$  by the assumption that the family  $(M_1^{(N)}, \dots, M_n^{(N)})$  converges in distribution. All permutations  $\sigma'$  which contribute to the sum satisfy on one hand  $|\sigma'| = |\sigma| - k$ , hence  $|\sigma'\sigma^{-1}| \geq k$ , and  $\#\Pi_k(\sigma \rightarrow \sigma') > 0$ , hence  $|\sigma'\sigma^{-1}| \leq k$ . Hence, only permutations  $\sigma'$  such that  $\sigma' \preceq \sigma$  contribute and (35) can be rewritten as

$$(36) \quad e^{-\frac{nt}{2}} \sum_{\sigma' \preceq \sigma} \frac{(-t)^{|\sigma'\sigma^{-1}|}}{|\sigma'\sigma^{-1}|!} \#\Pi_{|\sigma'\sigma^{-1}|}(\sigma \rightarrow \sigma') \lim_{N \rightarrow \infty} p_{\sigma}(M_1^{(N)}, \dots, M_n^{(N)}).$$

Let  $c_1, \dots, c_{\ell(\sigma)}$  denote the cycles of  $\sigma$ . It is not difficult to check that the interval  $[\text{id}, \sigma]$  in the poset  $\mathfrak{S}_n$  is isomorphic to the product of intervals  $\prod_{r=1}^{\ell(\sigma)} [\text{id}, c_r]$  by the mapping  $(\alpha_1, \dots, \alpha_{\ell(\sigma)}) \mapsto \alpha_1 \dots \alpha_{\ell(\sigma)}$ . Consider  $\sigma' \preceq \sigma$  and write  $\sigma' = \alpha_1 \dots \alpha_{\ell(\sigma)}$  accordingly. Then  $|\sigma'\sigma^{-1}| = |\alpha_1 c_1^{-1}| + \dots + |\alpha_{\ell(\sigma)} c_{\ell(\sigma)}^{-1}|$ . Moreover, by the same argument of shuffling used in the computation of  $S((1 \dots n), k, 0)$ ,

$$\#\Pi_{|\sigma'\sigma^{-1}|}(\sigma \rightarrow \sigma') = \frac{|\sigma'\sigma^{-1}|!}{\prod_{r=1}^{\ell(\sigma)} |\alpha_r c_r^{-1}|!} \prod_{r=1}^{\ell(\sigma)} \#\Pi_{|\alpha_r c_r^{-1}|}(c_r \rightarrow \alpha_r).$$

Hence, (36) can be written as

$$(37) \quad \prod_{r=1}^{\ell(\sigma)} e^{-\frac{(|c_r|+1)t}{2}} \sum_{\alpha_r \preceq c_r} \frac{(-t)^{|\alpha_r c_r^{-1}|}}{|\alpha_r c_r^{-1}|!} \#\Pi_{|\alpha_r c_r^{-1}|}(c_r \rightarrow \alpha_r) \lim_{N \rightarrow \infty} \text{Tr}_N(M_{i_r,1}^{(N)} \dots M_{i_r,m_r}^{(N)}),$$

and this is just the right-hand side of (33).

The equation (34) follows from (33) just as in Proposition 5.2.  $\square$

**6.2. Free cumulants.** As a preliminary to the proof of the asymptotic freeness, we compute the free cumulants of the limiting distribution of  $B_{\frac{t}{N}}$ . Let  $(\mathcal{A}, \varphi)$  be a non-commutative probability space and  $u_t$  an element of  $\mathcal{A}$  such that  $B_{\frac{t}{N}}$  converges in distribution to  $u_t$  as  $N$  tends to infinity. We have spent a substantial part of this paper proving that the moments of  $u_t$  are given by

$$(38) \quad \varphi(u_t^n) = e^{-\frac{nt}{2}} \sum_{k=0}^{n-1} \binom{n}{k+1} \frac{(-nt)^k}{nk!}.$$

Given  $a \in \mathcal{A}$  and  $\sigma \in \mathfrak{S}_n$  with cycle lengths  $(m_1, \dots, m_{\ell(\sigma)})$ , let us use the notation  $\varphi_{\sigma}(a) = \varphi(a^{m_1}) \dots \varphi(a^{m_{\ell(\sigma)}})$ . The free cumulants of  $u_t$  form a family of complex numbers  $(k_{\pi}(u_t))_{\pi \in \cup_{n \geq 1} \mathfrak{S}_n}$  and they are characterized by the identity

$$(39) \quad \forall n \geq 1, \forall \sigma \in \mathfrak{S}_n, \quad \varphi_{\sigma}(u_t) = \sum_{\sigma' \preceq \sigma} k_{\sigma'}(u_t).$$

Let us use the notation  $k_n = k_{(1 \dots n)}$ . It is an elementary property of the free cumulants that they are multiplicative, in that  $k_{\sigma} = k_{m_1} \dots k_{m_{\ell(\sigma)}}$  when  $(m_1, \dots, m_{\ell(\sigma)})$  are the cycle lengths of  $\sigma$ .

**Proposition 6.2.** *The free cumulants of  $u_t$  are given by*

$$(40) \quad k_n(u_t) = e^{-\frac{nt}{2}} \frac{(-nt)^{n-1}}{n(n-1)!}.$$

More generally, if  $\sigma \in \mathfrak{S}_n$ , then

$$(41) \quad k_\sigma(u_t) = e^{-\frac{nt}{2}} \frac{(-t)^{|\sigma|}}{|\sigma|!} \#\Pi_{|\sigma|}(\text{id} \rightarrow \sigma).$$

**Proof** – Let us put  $\varphi(u_t^n)$  under the form of the right hand side of (39). Applying Theorem 3.3, using Kreweras complementation and using Lemma 5.7, we find

$$\begin{aligned} \varphi(u_t^n) &= e^{-\frac{nt}{2}} \sum_{\sigma \preccurlyeq (1\dots n)} \frac{(-t)^{|\sigma(1\dots n)^{-1}|}}{|\sigma(1\dots n)^{-1}|!} \#\Pi_{|\sigma(1\dots n)^{-1}|}((1\dots n) \rightarrow \sigma) \\ &= e^{-\frac{nt}{2}} \sum_{\sigma \preccurlyeq (1\dots n)} \frac{(-t)^{|\sigma|}}{|\sigma|!} \#\Pi_{|\sigma|}(\text{id} \rightarrow \sigma) \\ &= \sum_{\sigma \preccurlyeq (1\dots n)} \frac{(e^{-\frac{1t}{2}}(-t)^{0}1^{-1})^{s_1} (e^{-\frac{2t}{2}}(-t)^{1}2^0)^{s_2} \dots (e^{-\frac{nt}{2}}(-t)^{n-1}n^{n-2})^{s_n}}{0!^{s_1} 1!^{s_2} \dots (n-1)!^{s_n}}, \end{aligned}$$

where  $s_1, \dots, s_n$  are respectively the number of fixed points of  $\sigma$ , and the numbers of transpositions, 3-cycles,  $\dots$ ,  $n$ -cycles in the decomposition of  $\sigma$ . By comparing this expression with (39), we find the desired expression for the cumulants of  $u_t$ .  $\square$

Let us recall briefly Speicher's characterization of freeness by the vanishing of mixed free cumulants [16]. Let  $a_1, \dots, a_n$  be non-commutative random variables on a space  $(\mathcal{A}, \varphi)$ , where  $\varphi$  is a tracial state. For  $\sigma \in \mathfrak{S}_n$ , the number  $m_\sigma(a_1, \dots, a_n)$  is defined by

$$m_\sigma(a_1, \dots, a_n) = \prod_{\substack{c \text{ cycle of } \sigma \\ c=(i_1 \dots i_r)}} \varphi(a_{i_1} \dots a_{i_r}).$$

It is well defined thanks to the fact that  $\varphi$  is tracial. The numbers  $m_\sigma(a_1, \dots, a_n)$  are the mixed moments of  $a_1, \dots, a_n$ . The relation

$$m_\sigma(a_1, \dots, a_n) = \sum_{\sigma' \preccurlyeq \sigma} k_{\sigma'}(a_1, \dots, a_n)$$

characterizes the family of numbers  $k_\sigma(a_1, \dots, a_n)$ , the mixed free cumulants of  $a_1, \dots, a_n$ .

Speicher's characterization of freeness is the following. Let  $(a_k)_{k \geq 1}$  be a family of elements of  $\mathcal{A}$ . Then this family is free if and only if, for all  $n \geq 2$  and all  $i_1, \dots, i_n \geq 1$  such that  $i_r \neq i_s$  for some  $r, s \in \{1, \dots, n\}$ ,  $k_{(1\dots n)}(a_{i_1}, \dots, a_{i_n}) = 0$ .

**Theorem 6.3.** *Let  $(B^{(N,k)})_{N,k \geq 1}$  be a family of Brownian motions, such that, for all  $N, k \geq 1$ ,  $B^{(N,k)}$  is a Brownian motion on  $U(N)$  and, for all  $N \geq 1$ , the Brownian motions  $(B^{(N,k)})_{k \geq 1}$  are independent. Let  $(t_k)_{k \geq 1}$  be a sequence of non-negative real numbers.*

*Then, as  $N$  tends to infinity, the family of non-commutative random variables  $(B_{\frac{t_k}{N}}^{(N,k)})_{k \geq 1}$  converges in distribution towards a free family  $(b^{(k)})_{k \geq 1}$  of non-commutative random variables such that, for all  $k \geq 1$ ,  $b^{(k)}$  has the distribution of  $u_{t_k}$ .*

**Proof** – We prove the result for finite families of Brownian motion  $(B^{(N,k)})_{N \geq 1, k \in \{1, \dots, K\}}$  for some finite  $K$ , by induction on  $K$ . The case  $K = 1$  is settled by our computation of the asymptotic distribution of  $B_{\frac{t}{N}}^{(N)}$ , that is, Propositions 5.1 and 5.6.

Let  $K \geq 2$  be an integer and let us assume that the property is proved for  $K - 1$  independent Brownian motions. We need to prove that the mixed free cumulants of  $B_{\frac{t}{N}}^{(N,1)}, \dots, B_{\frac{t}{N}}^{(N,K)}$  tend to zero as  $N$  tends to infinity. We regard  $B_{\frac{t}{N}}^{(N,k)}$  as elements of the non-commutative probability space  $(L^\infty(\Omega, \mathbb{P}) \otimes \mathbb{M}_N(\mathbb{C}), \mathbb{E} \otimes \text{Tr}_N)$  and we use the notation  $m_\sigma$  and  $k_\sigma$  accordingly. In particular, with our previous notation,  $m_\sigma = \mathbb{E} p_\sigma$ .

What we need to prove is that, for all  $n \geq 2$ , all  $\sigma \in \mathfrak{S}_n$ , all  $i_1, \dots, i_n \in \{1, \dots, K\}$  not all equal,

$$\lim_{N \rightarrow \infty} m_\sigma(B_{\frac{t}{N}}^{(N,i_1)}, \dots, B_{\frac{t}{N}}^{(N,i_n)}) = \sum_{\substack{\sigma' \preceq \sigma \\ \forall r \in \{1, \dots, n\}, i_{\sigma'(r)} = i_r}} \lim_{N \rightarrow \infty} k_{\sigma'}(B_{\frac{t}{N}}^{(N,i_1)}, \dots, B_{\frac{t}{N}}^{(N,i_n)}).$$

By the factorization property (33) and Fubini's theorem, the left hand side is multiplicative with respect to the cycle decomposition of  $\sigma$ . The right hand side is also clearly multiplicative, hence, it suffices to consider the case where  $\sigma = (1 \dots n)$ . In this case, we are looking at the expected trace of a product  $B_{\frac{t}{N}}^{(N,i_1)} \dots B_{\frac{t}{N}}^{(N,i_n)}$  where at least two factors are distinct. Of course, the case where one of the  $K$  possible factors does not appear is treated by the induction hypothesis. Let us assume that the  $K$  factors appear, in particular  $B^{(N,1)}$ . Up to cyclic permutation, which does not affect its trace, the product above can be put under the form  $W_1^{(N)} B_{\frac{t}{N}}^{(N,1)} \dots W_r^{(N)} B_{\frac{t}{N}}^{(N,1)}$ , for some  $r \geq 1$  and some products  $W_1^{(N)}, \dots, W_r^{(N)}$  of factors among  $B^{(N,2)}, \dots, B^{(N,K)}$ . Our previous results show that

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathbb{E} \text{Tr}_N \left( B_{\frac{t}{N}}^{(N,i_1)} \dots B_{\frac{t}{N}}^{(N,i_n)} \right) &= e^{-\frac{nt}{2}} \sum_{\sigma \preceq (1 \dots r)} \frac{(-t)^{|\sigma(1 \dots r)^{-1}|}}{|\sigma(1 \dots r)^{-1}|!} \# \Pi_{|\sigma(1 \dots r)^{-1}|}((1 \dots r) \rightarrow \sigma) \cdot \\ &\quad \cdot \lim_{N \rightarrow \infty} m_\sigma(W_1^{(N)}, \dots, W_r^{(N)}), \\ &= e^{-\frac{nt}{2}} \sum_{\sigma \preceq (1 \dots r)} \frac{(-t)^{|\sigma|}}{|\sigma|!} \# \Pi_{|\sigma|}(\text{id} \rightarrow \sigma) \lim_{N \rightarrow \infty} m_{K(\sigma)}(W_1^{(N)}, \dots, W_r^{(N)}) \\ &= \sum_{\sigma \preceq (1 \dots r)} k_\sigma(u_{t_1}) \lim_{N \rightarrow \infty} m_{K(\sigma)}(W_1^{(N)}, \dots, W_r^{(N)}), \end{aligned}$$

where we have changed  $\sigma$  in  $K(\sigma) = \sigma^{-1}(1 \dots r)$  between the first and the second line.

The term  $\lim_{N \rightarrow \infty} m_{K(\sigma)}(W_1^{(N)}, \dots, W_r^{(N)})$  is equal to a sum of limits of free cumulants of the factors appearing in  $W_1^{(N)}, \dots, W_r^{(N)}$  in this order. By induction, only pure free cumulants appear, those which do not involve more than one Brownian motion in each cycle of the permutation. Moreover, by the combinatorial description of  $K(\sigma)$  as a non-crossing partition, only such cumulants appear that remain non-crossing when they are merged with  $\sigma$ . In symbols,

$$\lim_{N \rightarrow \infty} \mathbb{E} \text{Tr}_N \left( B_{\frac{t}{N}}^{(N,i_1)} \dots B_{\frac{t}{N}}^{(N,i_n)} \right) = \sum_{\substack{\sigma \preceq (1 \dots n) \\ \forall k \in \{1, \dots, n\}, i_k = i_{\sigma(k)}}} k_\sigma(B_{\frac{t}{N}}^{(N,i_1)}, \dots, B_{\frac{t}{N}}^{(N,i_n)}).$$

This is exactly what we expected.  $\square$

## 7. LARGE N YANG-MILLS THEORY ON A DISK AND BRANCHING COVERS

In this section, we explain how our main expansion relates the Brownian motion on the unitary group to a natural model of random branching covers on this disk. In doing this, we give a rigorous proof of results which are stated in [7].

Let  $D$  be the closed disk of radius 1 centred at the origin  $O$  of  $\mathbb{R}^2$ . Let  $n \geq 1$  be an integer. Let  $\lambda$  be a partition of  $n$ . We define the set  $\mathcal{R}_{n,\lambda}$  as the set of isomorphism classes of ramified coverings  $\pi : R \rightarrow D$  which satisfy the following properties.

1.  $R$  is a ramified covering of degree  $n$ .
2. For each ramification point  $x \in D$  of  $R$  which is not the origin  $O$ ,  $R$  has a generic ramification type at  $x$ , in that  $\#\pi^{-1}(x) = n - 1$ .
3. The monodromy of  $R$  along the boundary of  $D$  belongs to the conjugacy class of  $\mathfrak{S}_n$  corresponding to  $\lambda$ .

An element of  $\mathcal{R}_{n,\lambda}$  is allowed to be ramified over  $O$ , with any kind of ramification. The set ramification its points distinct from  $O$  is called its locus of generic ramification. It is contained in the interior of  $D - \{0\}$ , which we denote by  $D^*$ .

Let  $X$  be a finite subset of  $D^*$ . We define  $\mathcal{R}_{n,\lambda,X}$  as the subset of  $\mathcal{R}_{n,\lambda}$  formed by the coverings whose locus of generic ramification is  $X$ .

The set  $\mathcal{R}_{n,\lambda,X}$  is in natural one-to-one correspondence with a set of equivalence classes of paths in the Cayley graph of  $\mathfrak{S}_n$  as follows. Assume that  $X = \{x_1, \dots, x_k\}$ . Choose a point  $b$  on the boundary of  $D$ . By the interior of a simple closed continuous curve based at  $b$ , we mean the bounded connected component of the complement of its range. Let  $C, C_1, \dots, C_k$  be simple closed curves in  $D$  based at  $b$  with pairwise disjoint interiors such that the interior of  $C$  contains  $O$  and, for each  $i \in \{1, \dots, k\}$ , the interior of  $C_i$  contains  $x_i$ . We assume that this is done in such a way that the curve  $CC_k \dots C_1$  is homotopic to the boundary of  $D$  in  $D - (X \cup \{O\})$ . Then the monodromies  $\sigma_O, \tau_1, \dots, \tau_k$  of an element  $R \in \mathcal{R}_{n,\lambda,X}$  along the curves  $C, C_1, \dots, C_k$  are defined in  $\mathfrak{S}_n$  up to simultaneous conjugation and their orbit characterizes  $R$ . The assumptions made on  $R$  imply that  $\tau_1, \dots, \tau_k$  are transpositions and  $\sigma_O \tau_1 \dots \tau_k$  belongs to  $\lambda$ .

Let  $\mathcal{P}_{n,\lambda,k}$  be the set of paths in the Cayley graph of  $\mathfrak{S}_n$  which start at an element of the conjugacy class determined by  $\lambda$ . The symmetric group acts on  $\mathcal{P}_{n,\lambda,k}$  by conjugation. The mapping from  $\mathcal{R}_{n,\lambda,X}$  which associates to  $R$  the orbit of the path  $(\sigma_O \tau_k \dots \tau_1, \sigma_O \tau_k \dots \tau_2, \dots, \sigma_O \tau_k, \sigma_O)$  is a bijection. Moreover, the cardinal of the stabilizer of this orbit is equal to the cardinal of the automorphism group  $\text{Aut}(R)$  of  $R$ . Hence, the image on  $\mathcal{R}_{n,\lambda,X}$  of the counting measure on  $\mathcal{P}_{n,\lambda,k}$  by the mapping  $\mathcal{P}_{n,\lambda,k} \rightarrow \mathcal{P}_{n,\lambda,k}/\mathfrak{S}_n \simeq \mathcal{R}_{n,\lambda,X}$  is the measure

$$\rho_{n,\lambda,X} = \sum_{R \in \mathcal{R}_{n,\lambda}(X)} \frac{n!}{\#\text{Aut}(R)} \delta_R.$$

This measure is finite and satisfies  $\rho_{n,\lambda,X}(1) = \binom{n}{2}^k$ .

There is a natural topology on  $\mathcal{R}_{n,\lambda}$ , which is generated by the sets

$$\mathcal{O}(R, U) = \{R' \in \mathcal{R}_{n,\lambda} : R|_{D-U} \simeq R'|_{D-U}\},$$

where  $R$  spans  $\mathcal{R}_{n,\lambda}$  and  $U$  the set of neighbourhoods of the locus of generic ramification of  $R$ . This is a fairly coarse topology: for example, the cardinal of the locus of generic ramification is not continuous, but only lower semi-continuous in this topology. However, the ramification type at  $O$  is continuous. On the set  $\mathcal{X}$  of finite subsets of  $D^*$ , we put the topology which makes

the bijection  $\mathcal{X} \simeq \bigsqcup_{n \geq 0} ((D^*)^n - \Delta_n) / \mathfrak{S}_n$  a homeomorphism, where  $\Delta_n$  is the subset of  $(D^*)^n$  where at least two components coincide. These topology do not make the ramification locus a continuous function of the ramified covering. Nevertheless, let  $\mathcal{M}(\mathcal{R}_{n,\lambda})$  denote the space of finite Borel measures on  $\mathcal{R}_{n,\lambda}$  endowed with the topology of weak convergence.

**Lemma 7.1.** *The mapping  $\mathcal{X} \longrightarrow \mathcal{M}(\mathcal{R}_{n,\lambda})$  which sends  $X$  to  $\rho_{n,\lambda,X}$  is continuous.*

**Proof** – By definition of the topology on  $\mathcal{X}$ , it suffices to prove that the mapping is continuous on  $(D^*)^k - \Delta_k$  for all  $k \geq 0$ . Consider  $k \geq 0$ ,  $X = \{x_1, \dots, x_k\}$  and a bounded continuous function  $f : \mathcal{R}_{n,\lambda,X} \longrightarrow \mathbb{R}$ . Choose  $\varepsilon > 0$ .

Since  $\mathcal{R}_{n,\lambda,X}$  is finite, the continuity of  $f$  implies the existence of  $r > 0$  such that the balls  $B(x_i, r)$  are contained in  $D^*$ , pairwise disjoint for  $i \in \{1, \dots, k\}$  and the neighbourhood  $U = B(x_1, r) \times \dots \times B(x_k, r)$  of  $X$  in  $(D^*)^k - \Delta_k$  satisfies

$$\forall R \in \mathcal{R}_{n,\lambda,X}, \forall R' \in \mathcal{O}(R, U), |f(R') - f(R)| < \varepsilon \binom{n}{2}^{-k}.$$

Let  $X' = \{x'_1, \dots, x'_k\}$  be an element of  $U$ . Let  $\phi$  be a homeomorphism of  $D$  such that  $\phi|_{D-U} = \text{id}_{D-U}$  and  $\phi(x_i) = x'_i$  for all  $i \in \{1, \dots, k\}$ . For each ramified covering  $\pi : R \longrightarrow D$  belonging to  $\mathcal{R}_{n,\lambda,X}$ , the covering  $\Phi(R) = (\phi \circ \pi : R \longrightarrow D)$  belongs to  $\mathcal{R}_{n,\lambda,X'}$ . Replacing  $\phi$  by its inverse in the definition of  $\Phi : \mathcal{R}_{n,\lambda,X} \longrightarrow \mathcal{R}_{n,\lambda,X'}$  yields the inverse mapping, hence  $\Phi$  is a bijection. Moreover, the conjugation by  $\phi$  determines an isomorphism between  $\text{Aut}(R)$  and  $\text{Aut}(\Phi(R))$ . Finally,  $R$  and  $\Phi(R)$  are isomorphic outside  $U$ . Altogether,

$$|\rho_{n,\lambda,X'}(f) - \rho_{n,\lambda,X}(f)| \leq \sum_{R \in \mathcal{R}_{n,\lambda,X}} \frac{n!}{\#\text{Aut}(R)} |f(\Phi(R)) - f(R)| < \varepsilon.$$

Since  $k$ ,  $X$ ,  $f$  and  $\varepsilon$  were arbitrary, the result follows.  $\square$

Let  $t \geq 0$  be a real number. Let  $\Xi_t$  be the distribution of a Poisson point process on  $D$  of intensity  $\frac{t}{\pi}$  times the Lebesgue measure on  $D$ . Under  $\Xi_t$ , a random subset of  $D$  is contained in  $D^*$  with probability 1 and the average number of points of such a random set is  $t$ . Thinking of  $\Xi_t$  as a Borel probability measure on  $\mathcal{X}$ , we define a measure on  $\mathcal{R}_{n,\lambda}$  by setting

$$\rho_{n,\lambda}^t = \int_{\mathcal{X}} \rho_{n,\lambda,X} \Xi_t(dX).$$

The measure  $\rho_{n,\lambda}^t$  is finite and satisfies  $\rho_{n,\lambda}^t(1) = e^{t \binom{n}{2}^{-1}}$ . We define a probability measure  $\mu_{n,\lambda}^t$  on  $\mathcal{R}_{n,\lambda}$  by normalizing  $\rho_{n,\lambda}^t$ .

Let us define two functions on  $\mathcal{R}_{n,\lambda}$ . Firstly, given  $R \in \mathcal{R}_{n,\lambda}$ , let us define  $k(R)$  as the number of ramification points of  $R$  distinct from  $O$ . We have observed that this is a lower semi-continuous, hence measurable function of  $R$ . Secondly, let  $\chi(R)$  be the Euler characteristic of  $R$ .

**Lemma 7.2.** *Let  $X$  be a subset of cardinal  $k$  of  $D^*$ . Let  $R$  be an element of  $\mathcal{R}_{n,\lambda,X}$  and  $\gamma = (\sigma_0, \dots, \sigma_k)$  a representative of the associated orbit of  $\mathcal{P}_{n,\lambda,k}$ . Then*

$$\chi(R) = \ell(\sigma_k) - k = \ell(\lambda) - 2d(\gamma).$$

*In particular,  $\chi : \mathcal{R}_{n,\lambda} \longrightarrow \mathbb{Z}$  is upper semi-continuous and measurable.*

**Proof** – The first equality follows from the Riemann-Hurwitz formula, the second from the definition of the defect of a path. The last assertion follows from the lower semi-continuity of  $k$  and the fact that  $\ell(\sigma_k)$ , which depends only on the ramification type at  $O$ , is a continuous

function of  $R$ . □

The main result is the following.

**Theorem 7.3.** *Let  $N, n \geq 1$  be two integers. Let  $(B_t)_{t \geq 0}$  be the Brownian motion on  $U(N)$  defined in Theorem 3.3. Let  $\lambda$  be a partition of  $n$  and  $\sigma$  an element of  $\mathfrak{S}_n$  which belongs to the conjugacy class determined by  $\lambda$ . Let  $t \geq 0$  be a real number. Then*

$$e^{nt - \frac{n^2 t}{2}} \mathbb{E} \left[ p_\sigma^{st} \left( B_{\frac{t}{N}} \right) \right] = \int_{\mathcal{R}_{n,\lambda}} (-1)^{k(R)} N^{\chi(R)} \mu_{n,\lambda}^t(dR).$$

We could have avoided the unpleasant exponential factor in the statement of this theorem if we had considered the signed measure  $\tilde{\rho}_{n,\lambda}^t = \int_{\mathcal{X}} (-1)^{\#X} \rho_{n,\lambda,X} \Xi_t(dX)$  instead of  $\rho_{n,\lambda}$ .

**Proof** – Let  $X$  be a finite subset of  $D^*$ . The set  $\mathcal{R}_{n,\lambda,X}$  is in bijection with the set of orbits of  $\Pi_k(\sigma)$  under the action of  $\mathfrak{S}_n$  by conjugation. Let  $R$  be an element of  $\mathcal{R}_{n,\lambda,X}$  and  $\gamma = (\sigma, \sigma_1, \dots, \sigma_k)$  a representative of the corresponding orbit. By Lemma 7.2,

$$\begin{aligned} \int_{\mathcal{R}_{n,\lambda}} (-1)^{k(R)} N^{\chi(R)} \rho_{n,\lambda,X}(dR) &= \sum_{\gamma \in \Pi_k(\sigma)} (-1)^{k(X)} N^{\ell(\lambda) - 2d(\gamma)} \\ &= N^{\ell(\lambda)} \sum_{d \geq 0} \frac{(-1)^{k(X)}}{N^{2d}} S(\sigma, k(X), d). \end{aligned}$$

Integrating with respect to  $\Xi_t(dX)$ , we find

$$\int_{\mathcal{R}_{n,\lambda}} (-1)^{k(R)} N^{\chi(R)} \rho_{n,\lambda}^t(dR) = e^{-t} N^{\ell(\sigma)} \sum_{k,d \geq 0} \frac{(-t)^k}{k! N^{2d}} S(\sigma, k, d).$$

By Theorem 3.3, the right-hand side of this equality is equal to  $e^{-t + \frac{nt}{2}} \mathbb{E} \left[ p_\sigma^{st} \left( B_{\frac{t}{N}} \right) \right]$ . The result follows after normalizing  $\rho_{n,\lambda}^t$ . □

## REFERENCES

- [1] Philippe Biane. Free Brownian motion, free stochastic calculus and random matrices. In *Free probability theory (Waterloo, ON, 1995)*, volume 12 of *Fields Inst. Commun.*, pages 1–19. Amer. Math. Soc., Providence, RI, 1997.
- [2] Philippe Biane. On the free convolution with a semi-circular distribution. *Indiana Univ. Math. J.*, 46(3):705–718, 1997.
- [3] Philippe Biane. Segal-Bargmann transform, functional calculus on matrix spaces and the theory of semi-circular and circular systems. *J. Funct. Anal.*, 144(1):232–286, 1997.
- [4] Benoît Collins. Moments and cumulants of polynomial random variables on unitary groups, the Itzykson-Zuber integral, and free probability. *Int. Math. Res. Not.*, (17):953–982, 2003.
- [5] Benoît Collins and Piotr Śniady. Integration with respect to the Haar measure on unitary, orthogonal and symplectic group. *Comm. Math. Phys.*, 264(3):773–795, 2006.
- [6] David J. Gross and Andrei Matytsin. Some properties of large- $N$  two-dimensional Yang-Mills theory. *Nuclear Phys. B*, 437(3):541–584, 1995.
- [7] David J. Gross and Washington Taylor. Twists and Wilson loops in the string theory of two dimensional QCD. *Nuclear Physics B*, 403:395, 1993.
- [8] V. A. Kazakov and I. K. Kostov. Nonlinear strings in two-dimensional  $U(\infty)$  gauge theory. *Nuclear Phys. B*, 176(1):199–215, 1980.
- [9] Germain Kreweras. Une famille d’identités mettant en jeu toutes les partitions d’un ensemble fini de variables en un nombre donné de classes. *C. R. Acad. Sci. Paris Sér. A-B*, 270:A1140–A1143, 1970.
- [10] Germain Kreweras. Sur les partitions non croisées d’un cycle. *Discrete Math.*, 1(4):333–350, 1972.

- [11] Ming Liao. *Lévy processes in Lie groups*, volume 162 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2004.
- [12] Ian G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [13] Andrei Okounkov and Anatoly Vershik. A new approach to representation theory of symmetric groups. *Selecta Math. (N.S.)*, 2(4):581–605, 1996.
- [14] Ambar N. Sengupta. Traces in two-dimensional QCD: the large- $N$  limit. *Preprint*.
- [15] Isadore M. Singer. On the master field in two dimensions. In *Functional analysis on the eve of the 21st century, Vol. 1 (New Brunswick, NJ, 1993)*, volume 131 of *Progr. Math.*, pages 263–281. Birkhäuser Boston, Boston, MA, 1995.
- [16] Roland Speicher. Free probability theory and non-crossing partitions. *Sém. Lothar. Combin.*, 39:Art. B39c, 38 pp. (electronic), 1997.
- [17] Feng Xu. A random matrix model from two-dimensional Yang-Mills theory. *Comm. Math. Phys.*, 190(2):287–307, 1997.